

A large and unusually colored new snake species of the genus *Tantilla* (Squamata; Colubridae) from the Peruvian Andes (#12701)

1

First submission

Please read the **Important notes** below, and the **Review guidance** on the next page.
When ready [submit online](#). The manuscript starts on page 3.

Important notes

Editor and deadline

Michael Wink / 11 Sep 2016

Files

Please visit the overview page to [download and review](#) the files not included in this review pdf.

Declarations

One or more DNA sequences were reported.
Involves vertebrate animals.
Involves a field study on animals or plants.
Describes a new species.



Please in full read before you begin

How to review

When ready [submit your review online](#). The review form is divided into 5 sections. Please consider these when composing your review:

1. BASIC REPORTING

2. EXPERIMENTAL DESIGN

3. VALIDITY OF THE FINDINGS

4. General comments

5. Confidential notes to the editor

 You can also annotate this **pdf** and upload it as part of your review

To finish, enter your editorial recommendation (accept, revise or reject) and submit.

BASIC REPORTING

-  Clear, unambiguous, professional English language used throughout.
-  Intro & background to show context. Literature well referenced & relevant.
-  Structure conforms to [PeerJ standard](#), discipline norm, or improved for clarity.
-  Figures are relevant, high quality, well labelled & described.
-  Raw data supplied (See [PeerJ policy](#)).

EXPERIMENTAL DESIGN

-  Original primary research within [Scope of the journal](#).
-  Research question well defined, relevant & meaningful. It is stated how research fills an identified knowledge gap.
-  Rigorous investigation performed to a high technical & ethical standard.
-  Methods described with sufficient detail & information to replicate.

VALIDITY OF THE FINDINGS

-  Impact and novelty not assessed. Negative/inconclusive results accepted. *Meaningful* replication encouraged where rationale & benefit to literature is clearly stated.
-  Data is robust, statistically sound, & controlled.
-  Conclusion well stated, linked to original research question & limited to supporting results.
-  Speculation is welcome, but should be identified as such.

The above is the editorial criteria summary. To view in full visit <https://peerj.com/about/editorial-criteria/>

A large and unusually colored new snake species of the genus *Tantilla* (Squamata; Colubridae) from the Peruvian Andes

Claudia Koch ^{Corresp., 1}, Pablo J Venegas ²

¹ Department of Herpetology, Zoologisches Forschungsmuseum Alexander Koenig (ZFMK), Bonn, Germany

² Department of Herpetology, Centro de Ornitología y Biodiversidad (CORBIDI), Lima, Peru

Corresponding Author: Claudia Koch

Email address: c.koch@zfmk.de

A new colubrid species of the genus *Tantilla* is described from the dry forest of the Northern Peruvian Andes on the basis of two specimens which exhibit a conspicuous sexual dimorphism. It represents the third species of the genus in Peru. The new species is easily distinguished from its congeners on the basis of mtDNA sequence divergence and by the combination of scalation characteristics and the unusual transversely-banded color pattern on the dorsum. A detailed description of the skull morphology of the new species is given based on micro-computed tomography images. The habitat of this new species is gravely threatened due to human interventions. Thorough investigation and conservation actions in the intern-andean valley Marañón River  is urgently needed.

1 **A large and unusually colored new snake species of the genus *Tantilla* (Squamata;**
 2 **Colubridae) from the Peruvian Andes**

3 Claudia Koch¹ & Pablo J. Venegas²

4

5 ¹Zoologisches Forschungsmuseum Alexander Koenig (ZFMK), Adenauerallee 160, 53113
 6 Bonn, Germany

7 ²Centro de Ornitología y Biodiversidad (CORBIDI), Santa Rita 117, Huertos de San Antonio,
 8 Surco, Lima, Perú

9 Corresponding author: Claudia Koch, E-mail: c.koch@zfmk.de

10

11 ABSTRACT

12 A new colubrid species of the genus *Tantilla* is described from the dry forest of the Northern
 13 Peruvian Andes on the basis of two specimens which exhibit a conspicuous sexual
 14 dimorphism. *Tantilla tjiasmantoi* sp. nov. represents the third species of the genus in Peru.
 15 The new species is easily distinguished from its congeners on the basis of mtDNA sequence
 16 divergence and by the combination of scalation characteristics and the unusual transversely-
 17 banded color pattern on the dorsum. A detailed description of the skull morphology of the
 18 new species is given based on micro-computed tomography images. The habitat of this new
 19 species is gravely threatened due to human interventions. Thorough investigation and
 20 conservation actions in the intern-andean valley Marañón River d urgently needed.

21
 22 **Key words:** Colubridae, dry forest, endemic, hemipods, intern-andean valley, La Libertad,
 23 Marañón, micro-computed tomography, osteology, skull morphology, new species, osteology,
 24 Peru, phylogenetic relationship, Reptilia, Serpentes, X-ray

25 26 INTRODUCTION

27 Seasonally dry tropical forests (SDTF) are characterised by a distinct seasonality with several
 28 months of arid-like conditions in which many plants lose their leaves (Murphy & Lugo,
 29 1986). In South America SDTFs are discontinuously distributed and can occupy large areas
 30 such as the Caatinga in north-eastern Brazil or small fragments as being found in inter-
 31 Andean valleys of Peru or Ecuador (Werneck et al., 2011). Nevertheless, the different areas of
 32 South American SDTFs are very divers and the species compositions differ substantially
 33 (Miles et al., 2006). However, SDTFs are considered as being one of the most threatened
 34 tropical ecosystems with a strong rate of annual deforestation (Janzen, 1988). The Equatorial
 35 dry forest is one representative of this forest type, which expands from south Ecuador to the
 36 northern part of Peru (Brack, 1986; Särkinen et al., 2011; Venegas, 2005). Although it has so
 37 far rarely been studied with respect to its flora and fauna, recent studies indicate that
 38 especially the inter-Andean part of this dry forest, along the valley of the Marañón River and
 39 its tributaries, is home to a high proportion of endemic species (Statterfield et al., 1998;
 40 Bridgewater et al., 2003; Koch, Venegas & Böhme, 2006, 2015; Koch et al., 2011, 2013;
 41 Venegas et al., 2008; Särkinen et al., 2011). Unfortunately, this part of the Equatorial dry

forest is facing multiple and complex threats due to logging, agriculture and narcotics plantations, mining activities, and above all due to several planned hydroelectric projects that will lead to flooding of vast portions of the inter-Andean dry forest (Q & V Ingenieros SAC, 2007; Finer & Jenkins, 2012). These threats, together with the enormous lack of knowledge that still exists about its biodiversity, and the fact that no protected area has so far been designated in the inter-Andean part of the Equatorial dry forest (IUCN & UNEP-WCMC, 2016), highlights the urgent priority for conservation and research activities in this area.

To contribute to the knowledge of the peculiar herpetofauna of the Andean dry forest, we conducted fieldwork in interandean valleys of the Northern Peruvian Departments of Amazonas, Cajamarca, and La Libertad and surveyed 28 localities along a stretch of more than 350 km of the Marañón River and its tributaries. In two of the southernmost investigation areas in the Department of La Libertad, we collected each a specimen of a new colubrid species, which we could hardly assign to any of the known genera based on outer morphological traits alone. However, a phylogenetic analysis based on 12S and 16S rRNA together with a comparison of the skull morphology via micro-computed tomography (micro-CT) scans revealed sufficient evidence to place it in the genus *Tantilla* Baird & Girard, 1853.

Currently, 61 species within the genus *Tantilla* are recognized (Wilson & Mata-Silva, 2015; Mata-Silva & Wilson, 2016). Twelve species occur in mainland South America, of which only two are found in Peru: *T. capistrata* Cope, 1875, and *T. melanocephala* (Linnaeus, 1758). Most species of the genus have either a plain-colored or a longitudinally striped dorsal color pattern. Only *T. shawi* Taylor, 1949 from Mexico, *T. semicincta* (Duméril, Bibron & Duméril, 1854) from Colombia and Venezuela, and *T. supracincta* (Peters, 1863) from Colombia, Costa Rica, Ecuador, Nicaragua, and Panama have a transversely-banded color pattern on the dorsal part of the body (Wilson, 1976; Wilson & Mata-Silva, 2016).

MATERIALS AND METHODS

Fieldwork, data sampling and Zoobank registration

The specimens were detected during visual encounter surveys (Crump & Scott, 1994) during our fieldwork conducted between March 2009 and November 2010. Altitudes above sea level and geographic coordinates were recorded with a GPS (Garmin GPSMap 60CSx) using the geodetic datum WGS84. Air temperatures and humidity were taken with a digital thermo-

hygrometer (Extech) using an external sensor. Specimens were collected and processed following the guidelines and protocols stated in the collecting permits. All the necessary research and collecting (0020-2009-AG-DGFFS-DGEFFS, 0424-2010-AG-DGFFS-DGEFFS) and export permits (003983-AG-DGFFS) for this study were issued by the Ministerio de Agricultura of the government of Peru (Ministerio de Agricultura). All efforts were made to minimize animal suffering. Both specimens were photographed in live and subsequently euthanized with the narcotic T61[®]. After taking tissue samples from the muscle of the lateral body side, the specimens were placed for about 12h in 10% formalin for

fixation, and subsequently stored in 70% ethanol. Finally the specimens were deposited in the collections of the Centro de Ornitología y Biodiversidad, Lima, Peru (CORBIDI), and the Zoologisches Forschungsmuseum Alexander Koenig, Bonn, Germany (ZFMK).

The electronic version of this article in Portable Document Format (PDF) will represent a published work according to the International Commission on Zoological Nomenclature (ICZN), and hence the new names contained in the electronic version are effectively published under that Code from the electronic edition alone. This published work and the nomenclatural acts it contains have been registered in ZooBank, the online registration system for the ICZN. The ZooBank LSIDs (Life Science Identifiers) can be resolved and the associated information viewed through any standard web browser by appending the LSID to the prefix <http://zoobank.org/>. The LSIDs for this publication is: urn:lsid:zoobank.org:pub:00EBF842-3AFA-4913-B381-95BDF86DAFAB. The online version of this work is archived and available from the following digital repositories: PeerJ, PubMed Central and CLOCKSS.

Comparative data for other South American colubrid species and genera were taken from Peracca (1904), Van Denburgh (1912), Werner (1924), Stuart & Bailey (1941), Slevin (1942), Smith & Taylor (1945), Stuart (1949, 1954), Peters (1960), Fritts & Smith (1969), Peters, Donoso-Barros & Orejas-Miranda (1970), Villa (1971, 1990), Myers (1973), Wilson (1976, 1987), Wilson & Mena (1980), Savages & Donnelly (1988), Perez-Santos & Moreno (1988), Savage & Crother (1989), McCranie & Wilson (1991a, 1991b, 1992), McCranie & Villa (1993), Cei (1993), Ferrarrezzi (1993), Puerto & Ferrarrezzi (1993), Myers & Cadle (1994), Smith & Campbell (1994), Campbell, Camarillo & Ustach (1995), Zaher (1996, 1999), Franco, Marques & Puerto (1997), Campbell (1998), Campbell & Smith (1998), Starace (1998), Kornacker (1999), Savage (2002), Sawaya & Sazima (2003), Köhler (2003, 2008), McCranie & Castaneda (2004), Solórzano (2004), Stafford (2004), Lema, D'Agostini & Cappelari (2005), Scott et al. (2006), Cacciali, Carreira & Scott (2007), Harvey et al. (2008),

Jansen & Köhler (2008), Zaher et al. (2009), Myers (2011), Passos, Ramos & Pereira (2012),

Moura, Caldeira Costa & Magalhães Pirani (2013), Myers & McDowell (2014), Wilson &

Mata-Silva (2015).

110

111 *Morphological analyses*

112 Measurements of head and scales were taken with a digital caliper and rounded to the nearest
113 0.1 mm, snout-vent-length and tail length were taken with a measuring tape and rounded to
114 the nearest 1 mm. Morphometric and meristic characters are abbreviated as follows: SVL
115 (snout–vent length, from tip of snout to cloaca); TL (length of tail); HW (width of head
116 across supraoculars); HH (height of head at highest part of head); HL (length of head); DSN
117 (distance from tip of snout to nostril); DNE (distance from nostril to anterior margin of eye);
118 ED (diameter of eye); MBD (body diameter at midbody); MTD (midtail diameter). The
119 number of ventral scales was counted in longitudinal row from mental to anal plate, and the
120 number of subcaudal scales was counted in longitudinal row from the cloaca to the tip of the
121 tail (Dowling, 1951). The number of dorsal scales rows around the body was counted at three
122 different points: (1) at a head's length behind the head; (2) at midbody; (3) at a head's length
123 before the cloaca. A light binocular was used to count and characterize small scales and to
124 identify the number of teeth in the male paratype.

125 For obtaining information on skeletal morphology, specimens were X-rayed in 2D (Faxitron
126 X-ray LX60) and in 3D by use of a micro-CT scanner (Bruker Skyscan 1272). Terminology
127 for the skull structures was adopted from Bulloch & Tanner (1966) and Cundall & Irish
128 (2008). The structures of the skull are abbreviated as follows: PMX (Premaxilla); NA (nasal);
129 SMX (septomaxilla); F (frontal); PFR (prefrontal); P (parietal); PO (postorbital); PRO
130 (prootic); ST (supratemporal); SO (supraoccipital); EXO (exoccipital); Q (quadrate); MX
131 (maxilla); ECP (ectopterygoid); PAL (palatine); MP (maxillary process of palatine); CHP
132 (chanal process of palatine); PT (pterygoid); BS (basisphenoid); BO (basioccipital).

133 The partially everted hemipenes of the male paratype were removed from the specimen and
134 prepared following Zaher & Prudente (2003). Finally the organs were scanned with the micro-
135 CT scanner. Left hemipenis was scanned dry, whereas right hemipenis was scanned in
136 alcohol. Terminology for hemipenes morphology follows Zaher (1999).

137

138 *Phylogenetic analysis*

139 Genomic DNA was extracted from the collected tissue samples at the Center for Molecular
 140 Biodiversity Research of the ZFMK, using the DNeasy Blood & Tissue Kit (Qiagen)
 141 following the standardized extraction protocol provided by the manufacturer. Two
 142 mitochondrial markers, 12S rRNA and 16S rRNA, and the nuclear gene RAG1
 143 (recombination-activating gene 1) were amplified using polymerase chain reactions (PCR)
 144 having a final volume of 20 µl and carried out either on a GeneAmp 2700 or on a Biometra
 145 thermal cycler. The QIAGEN® Multiplex PCR Kit (HotStarTaq® DNA Polymerase,
 146 Multiplex PCR Buffer with 6 mM MgCl₂, dNTP Mix, 2 µl Q-solution, 2.3 µl ultrapure H₂O)
 147 was used for the reaction for all three genes with 1.6 µl of each primer, and 2.5 µl of extracted
 148 DNA. To amplify the 12S fragment the roughly universal primer pair 12SA-L (light chain)
 149 and 12SB-H (heavy chain) of Kocher et al. (1989) was used. The 16S fragment was amplified
 150 using the likewise universal primer pair 16sar-L (light chain) and 16sbr-H (heavy chain) of
 151 Palumbi et al. (1991). For amplification of RAG1 the primer pair RAG1f2 (light chain) and
 152 RAG1r3 (heavy chain) of Schulte & Cartwright (2009) were used. Amplification with the 12S
 153 primer pair started with an initial denaturation step for 90 s at 94°C, and 38 cycles were run
 154 with denaturing for 45 s at 94°C, annealing for 60 s at 50°C, elongation for 120 s at 74°C, the
 155 final elongation for 300 s at 74°C, and cooling at 10°C. Amplification with the 16S primer
 156 pair started with an initial denaturation step for 900 s at 95°C, followed by 15 cycles of
 157 denaturation for 35 s at 94°C, annealing for 90 s at 60°C, elongation for 90 s at 72°C, plus 25
 158 cycles of denaturation for 35 s at 94°C, annealing for 90 s at 45°C, elongation for 90 s at
 159 72°C, the final elongation for 600 s at 72°C, and cooling at 10°C. Amplification with the
 160 RAG1 primer pair started with an initial denaturation step for 900 s at 95°C, and 40 cycles
 161 were run with denaturing for 20 s at 94°C, annealing for 50 s at 60°C, elongation for 90 s at
 162 72°C, the final elongation for 600 s at 72°C, and cooling at 10°C. After the PCR, each sample
 163 proving successful DNA amplification in an agarose gel electrophoresis, was purified for
 164 sequencing using the QIAquick PCR Purification Kit (Qiagen) and the standardized
 165 purification protocol provided by the manufacturer. Subsequently, the samples were
 166 sequenced by Macrogen Europe Laboratory (Amsterdam, Netherlands). Obtained sequences
 167 were checked with the original chromatograph data using BioEdit 7.5.2 (Hall, 1999). The 12S
 168 rRNA and 16S rRNA data was supplemented with sequences of 48 species representing 27
 169 genera of American colubrid snakes obtained from GenBank. Accession numbers are
 170 provided in Table 1. Sequence alignment was done with MAFFT (Katoh, Asimenos & Toh,
 171 2009) and manually corrected where necessary. Ambiguous sites (namely in the hypervariable

regions of the 16S rRNA) were identified with Gblocks (Castresana, 2000) and excluded from the alignment during subsequent analyses, resulting in 350 bp and 438 bp for 12S and 16S, respectively. For each gene, GTR+I+G was chosen as model of nucleotide substitution by the Akaike information criterion using Modeltest (Posada & Crandall, 1998) as implemented in the package ‘phangorn’ for Cran R. Phylogenetic trees were inferred using MrBayes 3.2.6 (Ronquist et al., 2012), estimating model parameters separately for each gene by partitioning the data set. We used a random starting tree and four independent runs with a maximum of 10 million generations each, sampled every 1000. Runs were stopped when the average standard deviation of split frequencies had reached 0.01. Convergence of the Markov chains and effective sample sizes were checked with Tracer v1.6 (Rambaut et al., 2014) and the initial 25% of each run were discarded prior to building a consensus tree. In addition to the Bayesian inference (BI), phylogenies were also calculated with Maximum Likelihood (ML) via the RAxML BlackBox (Stamatakis, Hoover & Rougemont, 2008) using the partitioned data, the Gamma model of rate heterogeneity, and 100 bootstraps.

RESULTS

The taxonomic conclusion of this study is based on the observation of morphological features and inferred phylogenetic relationships.

Morphological analyses

As typical for the genus *Tantilla*, the two specimens possess a number of 15 smooth dorsal scale rows throughout the body, one preocular, no loreal, no suboculars, 1+1 temporals, a divided cloacal shield, paired subcaudals (Table 2). Additionally, the skull of the new species is composed of similar bones and bone structures as other species of the genus *Tantilla*. A comparison with two other congeners (*T. capistrata* and *T. relictata*) reveals great similarity to our new species with only minor differences in the shape or size of some bones (Fig. 6).

Phylogenetic analysis

A pairwise analysis of a fragment of 526 bp of the mitochondrial gene 16S and a 1050 bp fragment of the nuclear gene RAG1 of both specimens showed no differences in the 16S fragment and only a single difference in one base position in the RAG1 fragment.

The Bayesian consensus tree (Fig. 1) obtained from 788 bp of mitochondrial DNA (12S and 16S rRNA) was based on 3600 sampled trees and effective sample sizes were > 1000, which indicated good mixing of the Markov chains. Although the topologies obtained from BI and  differ and the nodes were generally not very well resolved in terms of posterior probabilities and bootstrap values, the four species of *Tantilla* formed a well-supported clade including our two specimens (CORBIDI 7726 and ZFMK 95238). However, node support is lacking between the different species of *Tantilla*, obscuring the exact relationship between these taxa.

***Tantilla tjiasmantoi* sp. nov.**

urn:lsid:zoobank.org:act:15B70DA0-55D5-47D9-8383-453869F3B530

Holotype. CORBIDI 7726 from Pías, Province Pataz, Department of La Libertad, Peru (S 07°53'56.6'', W 077°34'43.8'', 1726 m a.s.l), collected by E. Hoyos Granda, A. Beraún and C. Koch on 15 January 2010.

Paratype. ZFMK 95238, from Santa Rosa/El Tingo (Marcamachay), Province Cajabamba, Department of La Libertad, Peru (07°21'59.3'' S, 77°53'53.0'' W, 1154 m a.s.l.), collected by M. Palacios Panta and C. Koch on 13 October 2010.

Diagnosis and definition. This comparatively large *Tantilla* is distinguished from its congeners by the following combination of characters: (1) maximum known SVL of 513 mm and total length of 638 mm; (2) 179-182 ventrals; (3) 57-65 paired subcaudals; (4) 7 supralabials; (5) eyes small, not visible from below, with round pupils; (6) dorsals smooth, without keels or apical pits, rhomboid, in 15 rows throughout the body; (7) loreals absent; (8) suboculars absent; (9) 2 postoculars; (10) 1 +1 temporals; (11) anal plate divided; (12) hypapophyses absent on posterior vertebrae; (13) hemipenes single; (14) body with dark bands that are not closed on ventral side; (15) conspicuous sexual dimorphism present.

With a maximum known total length of 638 mm *Tantilla tjiasmantoi* sp. nov. is among the largest species in the genus, only *T. shawi*, *T. impensa*, and *T. semicineta* reach similar or even larger total length. It can further be easily distinguished from all congeners except for *T. shawi*, *T. semicineta*, and *T. supracincta* by having a transversely-banded color pattern on the dorsal part of the body. The higher number of subcaudals (57-65) differentiates the new species from *T. shawi* (48-50) and the higher number of ventrals (179-182) further

distinguishes it from *T. semicineta* (161-176) and *T. supracincta* (138-151), as well as from the Peruvian species *T. capistrata* (130-156) and *T. melancocephala* (125-177). From the other species occurring in mainland South America, it can further be distinguished by a higher number of ventrals as compared to *T. alticola* (128-145), *T. andinista* (157), *T. boipiranga* (156-167), *T. miyatai* (165-172), *T. nigra* (137), *T. petersi* (172), *T. insulamontana* (144-157), *T. reticulata* (158-173).

Description of holotype. an adult female with a snout-vent-length (SVL) of 513 mm; tail length (TL) 125 mm; head length (HL) 16.3 mm; head width at broadest point (HW) 12.2 mm; head height (HH) 6.1 mm; TL/Total Length 0.2; SVL/HL 31.5; SVL/HW 42.1; SVL/HH 84.1; HW/HL 0.75; HH/HL 2.7; eye diameter (ED) 1.5 mm; HL/ED 10.9; HW/ED 8.1; midbody diameter (MBD) 11.5 mm; SVL/MBD 44.6; distance from tip of snout to nostril (DSN) 1.5 mm; distance from nostril to anterior margin of eye (DNE) 3 mm; midtail diameter (MTD) 2.9 mm.

Body robust, tail long, body and tail round in cross-section; dorsal scales in 15-15-15-rows, without reduction, rhomboid, smooth, lacking keels or apical pits; 182 ventrals; tail distinctly smaller in diameter than the body, long and tapering, tail spine pointed; 57 paired subcaudals; anal plate divided.

Head (Fig. 2) distinct from the body and horizontally compressed, laterally broadened behind the eyes. Rostral visible in dorsal view, slightly triangular with dorsal termination subacute, about 1.9 times wider than high. Snout rounded in dorsal view. Two internasals, subequal, about twice as broad as long, laterally in contact with anterior and posterior nasals. Prefrontals large, 1.2 times broader than long, laterally contacting postnasal, second supralabial and preocular. Frontal slightly hexagonal with dorsal termination acute, about 1.2 times longer than wide, laterally in contact with supraoculars, laterodorsally in contact with parietals. Supraoculars about 2.1 times wider than high. Parietals large, 1.6 times longer than wide, median suture about as long as frontal length. Nostrils located in posterior portion of anterior nasals. Anterior nasals about as high as broad, contacting rostral, internasal, postnasal, and first supralabial. Posterior nasals about 1.8 times longer than high, contacting anterior nasal, internasal, prefrontal, and first and second supralabial. No loreal. Eyes small with round pupils, partly visible in dorsal view, surrounded by one preocular, one supraocular, two postoculars, and third and fourth supralabial. No suboculars. Preoculars almost squarish. Superior postoculars, about 1.5 times longer than high; lower postoculars slightly pentagonal to almost circular. Supralabials 7/7, increasing in size posteriad, last one very large, about 3

times higher and 2.2 times broader than first supralabial, second contacting postnasal and preocular, third contacting prefrontal and eye, fourth contacting eye and lower postocular, fifth contacting lower postocular and anterior temporal, sixth contacting anterior temporal, seventh contacting anterior and posterior temporals, and first scale of dorsolateral and lateral body scale row, respectively. 1 +1 temporals, almost rectangular, separating supralabials 5-7 from parietals, anterior temporal about 2.5 times longer than high on both sides, 1.3 times longer than posterior one on left side and almost twice as long as posterior one on right side of head, posterior temporal 1.8 times longer than high on left side and about 1.1 times longer than high on right side of head. Mental subrhombical, 1.4 times broader than long, separated from chinshields by first pair of infralabials, which contact each other along the ventral midline. Infralabials 7/7, fourth largest, second smallest, first to fourth contacting anterior pair of chinshields; fourth infralabial contacting posterior chinshield and first gular scale. Two pairs of almost rectangular chinshields; anterior chinshields about 1.7 times longer than wide and 1.3 times longer than posterior chinshields; posterior chinshields about 2.2 times longer than wide, laterally contacting fourth infralabial, dorsally separated from the ventrals by four gular scales.

Trunk vertebrae 185; hypapophyses present on anterior 1/5 of trunk vertebrae, absent on posterior vertebrae; caudal vertebrae 56 (Fig. 4).

Coloration. In live, the dorsal ground color of head, body and tail is orange-yellowish, slightly paler laterally, most scales on body and tail with reddish-brown outlines; there are about 27 blackish dorsal crossbars on the body that are four to seven scales in length and are stretched across all dorsal scale rows except the most lateral row, slightly longer than ground color interspaces, fused in some parts of the body to form a zigzag band, slightly mottled in some parts with yellow. There are 12 dark tail blotches, reaching to subcaudals on both sides, fused in median part of the tail along the midline to form a zigzag band. Head with a large dark dorsal t-shirt-shaped blotch covering frontal, supraoculars, most of parietals except for the most posterior parts, and posterior part of prefrontals, the dark blotch is laterally extended at eye level, covering orbit, preocular, third supralabial, and adjacent parts of second and fourth supralabials, respectively. Infralabials, rostral and mental yellowish, except for blackish region surrounding the lingual groove. The ventral scales of head and body and subcaudal scales are cream-colored with dark dotted outlines in some parts. The coloration of the tongue is black to grayish-black (Fig. 3).

In preservative, the dorsal pattern on body and tail of a light ground color with dark crossbands remains unchanged, likewise the dark coloration of the head; the orange-yellowish dorsal ground color changed to cream and the darker outlines of most dorsal scales disappeared; the ventral coloration changed to grayish-white in some parts (Fig. 3). The coloration of the tongue changed to gray.

Variation. The single paratype is a small male with a conspicuous sexual dimorphism in body size and ventral coloration compared to the holotype. As the hemipenes ornamentation is as detailed as in other adult specimens of the genus *Tantilla* we assumed it to be already sexually mature. Intraspecific variation in scale counts and measurements is shown in Table 2. The paratype further differs from the holotype in the following characters: SVL/HL 24.4; HW/HL 0.58; ED 1.1 mm; HL/ED 7.4; HW/ED 4.3; MBD 3.5 mm; SVL/MBD 56.6; DSN 1.0 mm; DNE 1.3 mm; MTD 1.3 mm. On the right side this specimen has only 6 infralabials, of which the third one is the largest.

The dorsal ground coloration of the paratype (Fig. 3) in live is bright yellow and there are 35 blackish dorsal crossbars on the body and 16 dark tail blotches, some of the latter are fused to form a zigzag band, the black parts are usually not mottled with yellow. The t-shirt-shaped dark pattern on the dorsal surface of the head is also present in this specimen, but there is a dark vertical line on the left side of the head covering whole sixth and adjacent parts of seventh supralabial. On left side of the head posterior part of second infralabial and infralabials three to five are dark, on right side of head the posterior part of third infralabial and the infralabials four to six are dark. The ventral coloration differs from the holotype as the dark dorsal coloration of the paratype continues on the ventral scales, and thus sometimes forming complete rings around the body or creating a checkerboard pattern on the ventral surface of black and cream squares.

Cranial osteology. The snout is terminated by a small, single, toothless, median premaxilla. The triangular-shaped dorsal process of the premaxilla is not contacting the anterior portion of the two nasals. The paired nasals are slightly convex and oval-shaped in dorsal view, medially in contact and separate anteriorly to form an interspace. Their posterior edges are in contact with the anterior inner edges of the frontals. The paired frontals are dorsally flat, subrectangular, longer than broad, slightly larger than the nasals, and in close medial contact. The posterolateral edge of each frontal forms a supraorbital ridge which is perforated with two supraorbital foramina. The lateral surface of the frontal forms a major portion of the mesial wall of the orbit. The prefrontal is loosely attached to the anterolateral edge of the

frontal. In lateral view, the prefrontal is anteriorly contacting the posterior edge of the nasal and the septomaxilla, and the ventral border is contacting the dorsal surface of the maxilla. The frontals articulate posteriorly with the fused parietals. In dorsal view, the parietal is flat, oval-shaped except for the more or less straight anterior border along the suture with the frontals. It is the largest of the cranial elements and like the frontals, extends laterally far down, reaching ventrally to the basisphenoid. The parietal forms the posterior portion of each orbit. Its posterolateral borders suture with the prootics and its posterior border sutures with the supraoccipital. The postorbitals are narrow, elongate, flattened, slightly curved bones, larger than the prefrontals. They articulate with the anterolateral surface of the parietal and form the dorsoposterior boundary of each orbit. The single supraoccipital is flat, subpentagonal, and broader than long. Laterally it unites with the prootics and posteriorly with the exoccipitals. The paired prootics are largely separated by the broad supraoccipital. Each prootic is subtrapezoidal in dorsal view and is posteriorly bordered by the exoccipital. Laterally it extends far down, reaching ventrally the basisphenoid and the basioccipital. Each prootic has two large foramen of which the anterior one is anteriorly not enclosed by bone. The posterior foramen is larger than the anterior one. Other smaller foramina pierce each prootic. Posterodorsally each prootic is fused with the elongate, spine-like supratemporal, which connects the posterolateral dorsal part of the skull with the proximal end of the quadrate. Both supratemporals are almost parallel to each other, converging only slightly posteriorly. The exoccipitals are dorsally flat, form the posterolateral walls of the braincase and are joined together by a mid-dorsal suture. Ventrally they are resting upon the basioccipital. The basioccipital is pentagonal-shaped and joined anteroventrally with the basisphenoid along a straight transverse suture. The basisphenoid is applied anteriorly without a suture to the narrow, elongate parasphenoid, forming a single bone, which extends anteriorly into the rostrum area and becomes the floor of the orbit. Each maxilla is a curved bar with a small horizontal dorsoposteriorly-pointing process, about midway on the mesial border, that articulates with the ventral surface of the prefrontal. Anterodorsally articulates with the posteroventral surface of the septomaxilla. The posterior end of the maxilla is broadened and received by the flattened, pincers-shaped ectopterygoid, which connects it to the pterygoid. Each maxilla bears sockets for about 18 prediastemal, slightly recurved teeth, followed by two slightly enlarged, ungrooved fangs. The ectopterygoid do not bear teeth. The narrow palatine bears 13 or 14 teeth. The posterior end of each palatine articulates with the anterior end of the pterygoid. At about the height of the third and fourth tooth each palatine possesses a broad, horizontally flattened, lateral maxillary process. At about the height of the

sixth to eighth tooth each palatine possesses a similar but median choanal process with the apex directed anteriorly. The pterygoid is a slightly curved, flattened bar with about 17 teeth along the medial border. The subtriangular quadrate articulates with the lateral border of the supratemporal and its distal surface articulates with the mandibular condyle. The mandible is composed of two jaw bones. The anterolateral portion of the jaw is formed by the dentary, which contains a row of about 22, slightly recurved teeth. The longer proximal part of each jaw is without teeth. (Fig. 5). There are 179 trunk vertebrae and 70 caudal vertebrae (Fig. 4).

Hemipenial morphology. The hemipenes are unilobed, unicalyculate and noncapitate with a single sulcus spermaticus. The apical part is mostly uniformly spinulate. The hemipenial body is more or less uniformly ornamented with long and thin spines, which increase in size towards the base (Fig. 7).

Etymology. The species is dedicated to Wewin Tjiasmanto (Indonesia) in recognition of his support of nature conservation and taxonomic research through the BIOPAT initiative.

Distribution and natural history. This species is so far known from the southern portion of the seasonally dry forest along the Marañón River and its tributaries, from near Santa Rosa de Marcamachay at the Río Crisnejas, Province Cajabamba, and from near Laguna de Pías, Province Pataz, both Department of La Libertad, at elevations of 1154 m and 1726 m a.s.l, respectively (Fig. 8). The female CORBIDI 7726 was detected on 7th of January 2010 at 12.30 pm resting on a stone. The male ZFMK 95238 was detected on 12th of October 2010 at 8.15 pm on pebbly-clayey ground. Air temperature when animals were sighted was 33.3°C and 28.1°C, respectively.

DISCUSSION

Despite the comparatively close localities (> 70km air distance) and similarities in scale counts and arrangement of scales, the conspicuous differences of both specimens in body size and ventral coloration created some doubt if they represent the same species. A pairwise analysis of 1576 bp derived from the mitochondrial gene 16S and the nuclear gene RAG1 showed only a single difference in one base position in the RAG1 fragment, thus strongly supporting the assumption that both specimens belong to the same species. The collection and examination of further specimens is needed to determine whether the differences in size and color pattern are a product of sexual dimorphism or are referred to a different cause (e.g. age dependence, geographic variation).

In order to get a general idea of the phylogenetic position of the new species described herein, we performed phylogenetic analyses based on 12S rRNA and 16S rRNA. We did not attempt to conduct a taxonomically extensive analysis of South American Colubridae, instead we preferred to include only those species, for which the two gene regions sequenced in this study were available in GenBank. Our phylogenetic tree based on mitochondrial DNA (Fig. 1) corroborates the assignment of our new species into the genus *Tantilla*. Both the monophyly of the sampled *Tantilla* and the conspecificity of our two dimorphic specimens are well supported by the analyses. However, intrageneric relationships remain dubious due to the few species with genetic data available. Moreover, scutellation characteristics and the comparison of the skull morphology via micro-CT scans (Fig. 6) strongly support this hypothesis.

With currently 62 species assigned to this genus it represents the second largest genus of new world colubrid snakes, after the genus *Atractus*. However, taxonomic approaches are limited by the fact that pretty much nothing concrete is known about the phylogenetic relationships within the genus *Tantilla*, and its monophyly has not yet been tested adequately. Only few specimens representing less than ten species of *Tantilla* have been included in previous DNA-based studies (e.g. Vidal et al., 2000; Lawson et al., 2005; Burbrink & Myers, 2015; Schrey et al., 2015; Chambers & Hebert, 2016). Thus, a comprehensive study to include a large genomic sample, both numerical and geographical, of a large number of species of *Tantilla* is needed to clarify the phylogenetic relationship among the different species, to uncover their diversity and evolutionary history, and to further support our current generic assignation.

The habitat of this new species is gravely threatened due to human interventions, such as deforestation, mining activities and intended dam constructions for hydroelectric projects. To date no protected area has been established in the Marañón river valley. We hope that this beautiful and untypically colored new *Tantilla* could serve as a flagship species, together with several other endemic species of reptile and birds, for the establishment of conservation strategies in this region. If no such strategies will soon be developed and implemented the destruction of this unique habitat will proceed and will lead to a decline of its diversity, which may also have a negative impact on the endemic new species described herein.

ACKNOWLEDGEMENTS

Peter Rühr and Morris Flecks (both ZFMK) kindly helped with the micro-CT imaging and phylogenetic analysis, respectively. We are indebted to Hussam Zaher for sharing his

429 expertise on hemipenes morphology with us. CK thanks the Deutscher Akademischer
 430 Austauschdienst (DAAD), the Alexander Koenig Stiftung (AKS) and the Alexander Koenig
 431 Gesellschaft (AKG) for financial support. The Ministerio de Agricultura, Peru kindly
 432 provided collecting and export permits. We thank Alfredo Beraún, Erick Hoyos Granda, and
 433 Manuel Palacios Panta for assistance during the fieldwork. We are indebted to the residents of
 434 Santa Rosa and Laguna de Pías for their hospitality, support and assistance during the
 435 fieldwork.

436

437 **Table 1.** Taxa used for phylogenetic analysis and respective GenBank accession numbers.

Species	12S rRNA	16S rRNA
<i>Chironius bicarinatus</i>	HM565744	HM582206
<i>Chironius carinatus</i>	HM565745	HM582207
<i>Chironius exoletus</i>	HM565746	HM582208
<i>Chironius flavolineatus</i>	HM565747	HM582209
<i>Chironius foveatus</i>	HM565748	HM582210
<i>Chironius fuscus</i>	HM565749	HM582211
<i>Chironius laevicollis</i>	HM565751	HM582213
<i>Chironius monticola</i>	HM565753	HM582214
<i>Chironius quadricarinatus</i>	HM565755	HM582215
<i>Chironius scurrulus</i>	HM565756	HM582216
<i>Clelia bicolor</i>	GQ457787	GQ457729
<i>Coluber constrictor</i>	AY122667	
<i>Coluber flagellum</i>	AY122668	
<i>Coluber taeniatus</i>	AY122669	
<i>Dendrophidion percarinatus</i>	HM565757	HM582217
<i>Drymarchon corais</i>	HM565758	HM582218
<i>Drymobius rhombifer</i>	HM565761	HM582220
<i>Drymoluber brazili</i>	HM565760	HM582219
<i>Drymoluber dichrous</i>	HM565759	HM582221
<i>Erythrolamprus aesculapii</i>	GQ457795	GQ457736
<i>Gyalopion canum</i>	KR814624	KR814641
<i>Lampropeltis getula</i>	AY122821	
<i>Lampropeltis mexicana</i>	FJ623962	
<i>Lampropeltis triangulum</i>	FJ623963	
<i>Leptodrymus pulcherrimus</i>	KR814627	KR814649
<i>Leptophis ahaetulla</i>	HM565762	HM582222
<i>Liophis elegantissimus</i>	GQ457808	GQ457748
<i>Liophis jaegeri</i>	GQ457809	GQ457749
<i>Mastigodryas bifossatus</i>	HM565763	HM582223
<i>Mastigodryas boddaerti</i>	HM565764	HM582224
<i>Mastigodryas dorsalis</i>	KR814625	KR814650
<i>Oxybelis aeneus</i>	HM565765	HM582225
<i>Oxybelis wilsoni</i>	KR814626	KR814647
<i>Pituophis catenifer</i>	KU833245	KU833245
<i>Pituophis lineaticollis</i>	AF512746	AF512746
<i>Pseustes poecilonotus</i>		KR815895
<i>Rhinobothryum lentiginosum</i>	HM565767	HM582227
<i>Salvadora grahamiae</i>	AY122847	
<i>Scolecophis atrocinctus</i>	KR814619	KR814642
<i>Senticolis triaspis</i>	AY122848	
<i>Spilotes pullatus</i>	HM565768	HM582228
<i>Stenorrhina freminvillei</i>	HM565769	
<i>Symphimus leucostomus</i>	KR814618	KR814651
<i>Tantilla armillata</i>	KR814613	KR814644
<i>Tantilla impensa</i>	KR814614	KR814645
<i>Tantilla melanocephala</i>	AF158424	AF158491
<i>Tantilla vermiformis</i>	KR814615	KR814646
<i>Xenodon neuwiedi</i>	GQ457841	GQ457779
CORBIDI 7726	xxx	xxx
ZFMK 95238		xxx

438

Table 2. Scale counts and measurements of the type specimens of *Tantilla tjasmanoi* sp.

	♀ Holotype CORBIDI 7726	♂ Paratype ZFMK 95238
Dorsal scale rows	15-15-15	15-15-15
Ventrals	182	179
Subcaudals	57	65
Supralabials	7	7
Infralabials	7	6-7
Internasals	2	2
Prefrontals	2	2
Preoculars	1	1
Postoculars	2	2
Supraoculars	1	1
Suboculars	absent	absent
Loreal	absent	absent
Anterior temporals	1	1
Posterior temporals	1	1
Sublinguals (paired)	2	2
Snout-vent-length (SVL)	513 mm	198 mm
Tail length	125 mm	56 mm
Tail length/Total length	0.2	0.22
Head width (at widest point)	11.9 mm	4.7 mm
Head length (from tip of snout to end of quadrate)	16.3 mm	8.1 mm
Head width/ head length	0.73	0.58





Figure 1. Bayesian consensus tree based on 788 bp of mitochondrial DNA (12S and 16S rRNA) of our specimens (CORBIDI 7726 and ZFMK 95238) and 48 further species representing 27 genera of American colubrid snakes. Numbers at nodes are the Bayesian posterior probabilities (values < 0.95 not shown).




450

451 **Figure 2.** Dorsal (top), ventral (center) and lateral (bottom) views of head of female holotype
 452 of *Tantilla tjiasmantoi* sp. nov. from La Libertad, Peru (CORBIDI 7726).

453



Figure 3. *Tantilla tjiasmantoi* sp. nov. from La Libertad, Peru: female holotype CORBIDI 7726 (left) and male paratype ZFMK 95238 (right) in live in dorsal view (top), and in preservative in ventral view (bottom).



459

460 **Figure 4.** X-ray photographs of body of *Tantilla tjiasmantoi* sp. nov. from La Libertad,
 461 Peru (needle marks the cloaca): female holotype CORBIDI 7726 (left) and male paratype
 462 ZFMK 95238 (right).

463




Figure 5. Micro-CT images of the male paratype of *Tantilla tjiasmantoi* sp. nov. from La Libertad: dorsal (A), lateral (B), and ventral views of the skull (C, lower jaw removed), and dorsal view of lower jaw (D). See Materials and Methods section for explanation of the abbreviations of the skull structures.




Figure 6. Micro-CT images of the dorsal (above), lateral (center), and ventral (below) views of the skull of three species of *Tantilla*: *T. tjiasmantoi* sp. nov. from Peru (left), *T. capistrata* from Lambayeque, Peru (ZFMK 85028, middle), and *T. relictta* from Florida, USA (ZFMK 84387, right).





Figure 7. Micro-CT images of the hemipenes of the male paratype of *Tantilla tjiasmantoi* sp. nov. from La Libertad: sulcate (A), asulcate (B), and lateral views (C) of the left (left) and right (right) hemipenis. Left hemipenis was scanned dry, whereas right hemipenis was scanned in alcohol.



482

483 **Figure 8.** Type locality of *Tantilla tjiasmantoi* sp. nov. CORBIDI 7726 near Laguna de Pías,
 484 La Libertad, Peru (top), and locality of male paratype ZFMK 95238 near Santa Rosa de
 485 Marcamachay, La Libertad, Peru (bottom).

486

REFERENCES

- Brack A. 1986. Las Ecoregiones del Perú. *Boletín de Lima* 44: 57–70.
- Bridgewater S, Pennington RT, Reynel CA, Daza A, Pennington TD. 2003. A preliminary floristic and phytogeographic analysis of the woody flora of seasonally dry forests in northern Peru. *Candollea* 58: 129–141.
- Bullock RE, Tanner WW. 1966. A comparative osteological study of two species of Colubridae (*Pituophis* and *Thamnophis*). *Brigham Young University Science Bulletin, Biological Series* 8(3): 1–29.
- Burbrink FT, Myers EA. 2015. Both traits and phylogenetic history influence community structure in snakes over steep environmental gradients. *Ecography* 38: 1–13. DOI: 10.1111/ecog.01148.
- Cacciali P, Carreira S, Scott NJ. 2007. Redescription of *Phalotris nigrilatus* Ferrarezzi, 1993 (Serpentes: Colubridae: Xenodontinae). *Herpetologica* 63(4): 552–559. DOI: 10.1655/0018-0831(2007)63[552:ROPNFS]2.0.CO;2.
- Campbell JA. 1998. Comments on the identities of certain *Tantilla* (Squamata: Colubridae) from Guatemala, with descriptions of two new species. *Scientific Papers of the Natural History Museum, University of Kansas* 7:1–14. DOI: 10.5962/bhl.title.16234.
- Campbell JA, Camarillo JL, Ustach PC. 1995. Redescription and rediagnosis of *Tantilla shawi* (Serpentes: Colubridae) from the Sierra Madre Oriental of Mexico. *The Southwestern Naturalist* 40(1): 120–132.
- Campbell JA, Smith EN. 1998. A new genus and species of colubrid snake from the Sierra de las Minas of Guatemala. *Herpetologica* 54: 207–220.
- Castresana J. 2000. Selection of conserved blocks from multiple alignments for their use in phylogenetic analysis. *Molecular Biology and Evolution* 17: 540–552. DOI: 10.1093/oxfordjournals.molbev.a026334.

- Cei JM. 1993. *Reptiles del noroeste, nordeste y este de la Argentina*. Torino: Museo Regionale Scienze Naturali Torino, Monografie 14, 949 pp.
- Chambers EA, Hebert PDN. 2016. Assessing DNA barcodes for species identification in North American reptiles and amphibians in natural history collections. *PLoS ONE* 11(4): e0154363. DOI: 10.1371/journal.pone.0154363.
- Crump ML, Scott NJ. 1994. Visual encounter surveys. In: Heyer WR, Donnelly MA, McDiarmid RW, Hayek LC, Foster MS, eds. *Measuring and monitoring biological diversity, standard methods for amphibians*. Washington, DC: Smithsonian Institution Press, 84–92.
- Cundall D, Irish FJ. 2008. The snake skull. In: Gans C, Gaunt AS, Adler K, eds. *Biology of the Reptilia, volume 20, Morphology H. The skull of Lepidosauria*. Ithaca: Society for the Study of Amphibians and Reptiles, 349–692.
- Dowling HG. 1951. A proposed standard system of counting ventrals in snakes. *British Journal of Herpetology* 1: 97–99.
- Ferrarezzi H. 1993. Nota sobre o genero *Phalotris* com revisao do grupo nasutus e descricao de tres novas especies (Serpentes, Colubridae, Xenodontinae). *Memorias do Instituto Butantan, Sao Paulo* 55 (Suppl. 1): 21–38.
- Finer M, Jenkins CN. 2012. Proliferation of hydroelectric dams in the Andean Amazon and implications for Andes–Amazon connectivity. *PLoS ONE* 7(4): e35126. DOI: 10.1371/journal.pone.0035126.
- Franco FL, Marques OAV, Puerto G. 1997. Two new species of colubrid snakes of the genus *Clelia* from Brazil. *Journal of Herpetology* 31(4): 483–490. DOI: 10.2307/1565599.
- Fritts TH, Smith HM. 1969. A new genus and species of snake from western Ecuador. *Transactions of the Kansas Academy of Sciences* 72(1): 60–66. DOI: 10.2307/3627049.

- 554 Hall TA. 1999. BioEdit: a user-friendly biological sequence alignment editor and analysis
555 program for Windows 95/98/NT. *Nucleic Acids Symposium Series* 41: 95–98.
- 556
- 557 Harvey MB, Rivas Fuenmayor G, Rances Caicedo-Portilla J, Rueda-Alm JV. 2008.
558 Systematics of the enigmatic Dipsadine Snake *Tropidodipsas perijanensis* Alemán
559 (Serpentes: Colubridae) and review of morphological characters of Dipsadini. *Herpetological*
560 *Monographs* 22(1): 106–132. DOI: 10.1655/08-025.1.
- 561
- 562 IUCN, UNEP-WCMC. 2016. The World Database on Protected Areas (WDPA) [On-line].
563 Available at <http://www.protectedplanet.net> (accessed 27 June 2016).
- 564
- 565 Jansen M, Kohler G. 2008. A new species of *Phalotris* from the eastern lowlands of Bolivia
566 (Reptilia, Squamata, Colubridae). *Senckenbergiana Biologica* 88(1): 103–110.
- 567
- 568 Janzen DH. 1988. Tropical dry forests: the most endangered major tropical ecosystem. In:
569 Wilson EO, Peter FM, ed. *Biodiversity*. Washington, DC: National Academy Press: 130–137.
- 570
- 571 Katoh K, Asimenos G, Toh H. 2009. Multiple Alignment of DNA Sequences with MAFFT.
572 *Methods in Molecular Biology* 537: 39–64. DOI: 10.1007/978-1-59745-251-9_3.
- 573
- 574 Koch C, Venegas PJ, Böhme W. 2006. A remarkable discovery: Description of a big-growing
575 new gecko (Squamata: Gekkonidae: *Phyllopezus*) from northwestern Peru. *Salamandra*
576 42(2/3): 145–150.
- 577
- 578 Koch C, Venegas PJ, Böhme W. 2015. Three new endemic species of *Epictia* Gray, 1845
579 (Serpentes: Leptotyphlopidae) from the dry forest of northwestern Peru. *Zootaxa* 3964(2):
580 228–244. DOI: 10.11646/zootaxa.3964.2.4.
- 581
- 582 Koch C, Venegas PJ, Garcia-Bravo A, Böhme W. 2011. A new bush anole (Iguanidae:
583 Polychrotinae: *Polychrus*) from the upper Marañón basin, Peru, with a redescription of
584 *Polychrus peruvianus* (Noble, 1924) and additional information on *P. gutturosus* Berthold,
585 1845. *Zookeys* 141: 79–107. DOI: 10.3897/zookeys.141.1678.
- 586

- Koch C, Venegas PJ, Rödder D, Flecks M, Böhme W. 2013. Two new endemic species of *Ameiva* (Squamata: Teiidae) from the dry forest of northwestern Peru and additional information on *Ameiva concolor* Ruthven, 1924. *Zootaxa* 3745(2): 263–295. DOI: 10.11646/zootaxa.3745.2.6.
- Kocher TD, Thomas WK, Meyer A, Edwards SV, Pääbo S, Villablanca FX, Wilson AC. 1989. Dynamics of mitochondrial DNA evolution in animals: amplification and sequencing with conserved primers. *Proceedings of the National Academy of Sciences of United States of America* 86(16): 6196–6200. DOI: 10.1073/pnas.86.16.6196.
- Köhler G. 2003. *Reptiles of Central America. 1st Ed.* Offenbach: Herpeton-Verlag, 367 pp.
- Köhler G. 2008. *Reptiles of Central America. 2nd Ed.* Offenbach: Herpeton-Verlag, 400 pp.
- Kornacker PM. 1999. *Checklist and key to the snakes of Venezuela.* Rheinbach: PaKo-Verlag, Germany, 270 pp.
- Lawson R, Slowinski JB, Crother BI, Burbrink FT. 2005. Phylogeny of the Colubroidea (Serpentes): New evidence from mitochondrial and nuclear genes. *Molecular Phylogenetics and Evolution* 37: 581–601. DOI:10.1016/j.ympev.2005.07.016.
- Lema T, D’Agostini FM, Cappelari LH. 2005. Nova espécie de *Phalotris*, redescricao de *P. tricolor* e osteologia craniana (Serpentes, Elapomorphinae). *Iheringia, Série Zoologia* 95: 65–78. DOI: 10.1590/s0073-47212005000100010.
- Mata-Silva V, Wilson LD. 2016. The taxonomic status of *Tantilla marcovani* Lema 2004 (Squamata: Colubridae). *Zootaxa* 4092(3): 421–425. DOI: 10.11646/zootaxa.4092.3.7.
- McCranie JR, Castaneda FE. 2004. Notes on the second specimens of *Geophis damiani* Wilson, McCranie, and Williams and *Rhadinaea tolpanorum* Holm and Cruz D. (Colubridae). *Herpetological Review* 35(4): 341.
- McCranie JR, Villa J. 1993. A new genus for the snake *Enulius sclateri* (Colubridae: Xenodontinae). *Amphibia-Reptilia* 14(3): 261–267. DOI: 10.1163/156853893x00453.

621

622 McCranie JR, Wilson LD. 1991. *Rhadinaea montecristi* Mertens. *Catalogue of American*
623 *Amphibians and Reptiles* 524: 1–2.

624

625 McCranie JR, Wilson LD. 1991. *Rhadinaea kinkelini* Boettger. *Catalogue of American*
626 *Amphibians and Reptiles* 523: 1–2.

627

628 McCranie JR, Wilson LD. 1992. *Rhadinaea godmani*. *Catalogue of American Amphibians*
629 *and Reptiles* 546: 1–3.

630

631 Miles L, Newton AC, DeFries RS, Ravilious C, May I, Blyth S, Kapos V, Gordon JE. 2006.
632 A global overview of the conservation status of tropical dry forests. *Journal of Biogeography*
633 33: 491–505. DOI: 10.1111/j.1365-2699.2005.01424.x.

634 Moura Ribeiro M, Caldeira Costa H, Magalhães Pirani R. 2013. Rediscovery of *Phalotris*
635 *concolor* (Serpentes: Dipsadidae: Elapomorphini). *Zoologia* 30(4): 430–436. DOI:
636 10.1590/s1984-46702013000400009.

637

638 Murphy PG, Lugo AE. 1986. Ecology of tropical dry forest. *Annual Review of Ecology and*
639 *Systematics* 17: 67–88. DOI: 10.1146/annurev.es.17.110186.000435.

640 Myers CW, Cadle JE. 1994. A new genus for South American snakes related to *Rhadinaea*
641 *obtusa* Cope (Colubridae) and resurrection of *Taeniophallus* Cope for the "*Rhadinaea*"
642 *brevirostris* group. *American Museum Novitates* 3102: 1–33.

643

644 Myers CW, McDowell SB. 2014. New taxa and cryptic species of Neotropical snakes
645 (Xenodontinae), with commentary on hemipenes as generic and specific characters. *Bulletin*
646 *of the American Museum of Natural History* 385(1): 1–112. DOI: 10.1206/862.1.

647

648 Myers CW. 1973. A new genus for Andean snakes related to *Lygophis boursieri* and a new
649 species (Colubridae). *American Museum Novitates* 2522: 1–37.

650

651 Myers CW. 2011. A new genus and new tribe for *Enicognathus melanauchen* Jan, 1863, a
652 neglected South American snake (Colubridae: Xenodontinae), with taxonomic notes on some
653 Dipsadinae. *American Museum Novitates* 3715: 1–33. DOI: 10.1206/3715.2.

654

655 Palumbi SR, Martin A, Romano S, McMillan WO, Stice L, Grabowski G. 1991. *The Simple*
656 *Fool's Guide to PCR, Version 2*. Honolulu: University of Hawaii Zoology Department, 94 pp.

657

658 Passos P, Ramos L, Pereira DN. 2012. Distribution, natural history, and morphology of the
659 rare snake, *Caaeteboia amarali* (Serpentes: Dipsadidae). *Salamandra* 48(1): 51–58.

660

661 Peracca MG. 1904. Nouvelles espèces d'Ophidiens d'Asie et d'Amerique, faisant partie de la
662 collection du Museum d'histoire naturelle de Genève. *Revue Suisse de Zoologie* 12: 663–668.
663 DOI: 10.5962/bhl.part.10151.

664

665 Pérez-Santos C, Moreno AG. 1988. *Ofidios de Colombia*. Torino: Museo Regionale di
666 Scienze Naturali, Monographie VI, 517 pp.

667

668 Peters JA, Donoso-Barros R, Orejas-Miranda B. 1970. Catalogue of the Neotropical
669 Squamata: Part I Snakes. Part II Lizards and Amphisbaenians. *United States National*
670 *Museum Bulletin* 297: 1–347.

671

672 Peters JA. 1960. The snakes of the subfamily Dipsadinae. *Miscellaneous Papers of the*
673 *Museum of Zoology, University of Michigan* 114: 1–224.

674

675 Posada D, Crandall KA. 1998. MODELTEST: testing the model of DNA substitution.
676 *Bioinformatics* 14: 817–818. DOI: 10.1093/bioinformatics/14.9.817.

677

678 Puerto G, Ferrarezzi H. 1993. Uma nova especie de *Phalotris* Cope, 1862, com comentarios
679 sobre o grupo bilineatus (Serpentes: Colubridae: Xenodontinae). *Memorias do Instituto*
680 *Butantan, Sao Paulo* 55 (Supl. 1): 39–46.

681

682 Q&V Ingenieros SAC. 2007. *Elaboración de resúmenes ejecutivos y fichas de estudio de las*
683 *centrales hidroeléctricas con potencial para exportación a Brasil*. Ministerio de Energía y
684 Minas, Dirección General de Electricidad, 79 pp.

685

686 Rambaut A, Suchard MA, Xie D, Drummond AJ. 2014. Tracer v1.6, Available at
687 <http://beast.bio.ed.ac.uk/Tracer> (accessed on 08 August 2016).

688

689 Ronquist F, Teslenko M, van der Mark P, Ayres DL, Darling A, Höhna S, Larget B, Liu L,
690 Suchard MA, Huelsenbeck JP. 2012. MrBayes 3.2: efficient Bayesian phylogenetic inference
691 and model choice across a large model space. *Systematic Biology* 61: 539–542. DOI:
692 10.1093/sysbio/sys029.

693

694 Särkinen TE, Marcelo-Peña JL, Daza Yomona A, Simon MF, Pennington RT, Hughes CE.
695 2011. Underestimated endemic species diversity in the dry inter-Andean valley of the Río
696 Marañón, northern Peru: An example from *Mimosa* (Leguminosae, Mimosoideae). *Taxon*
697 60(1): 139–150.

698

699 Savage JM. 2002. *The amphibians and reptiles of Costa Rica: A herpetofauna between two*
700 *continents, between two seas*. Chicago: University of Chicago Press, 934 pp.

701

702 Savage JM, Crother BI. 1989. The status of *Pliocercus* and *Urotheca* (Serpentes: Colubridae),
703 with a review of included species of coral snake mimics. *Zoological Journal of the Linnean*
704 *Society* 95: 335–362. DOI: 10.1111/j.1096-3642.1989.tb02315.x.

705

706 Savage JM, Donnelly MA. 1988. Variation and systematics in the colubrid snakes of the
707 genus *Hydromorphus*. *Amphibia-Reptilia* 9: 289–300. DOI: 10.1163/156853888x00378.

708

709 Sawaya RJ, Sazima I. 2003. A new species of *Tantilla* (Serpentes: Colubridae) from
710 southeastern Brazil. *Herpetologica* 59(1): 119–126. DOI: 10.1655/0018-
711 0831(2003)059[0119:ansots]2.0.co;2.

712

713 Schrey AW, Evans RK, Netherland M, Ashton KG, Mushinsky HR, McCoy ED. 2015.
714 Phylogeography of the Peninsula Crowned Snake (*Tantilla relictata relictata*) on the Lake Wales
715 Ridge in Central Florida. *Journal of Herpetology* 49(3): 415–419. DOI: 10.1670/13-190.

716

717

718 Schulte II JA, Cartwright EM. 2009. Phylogenetic relationships among iguanian lizards using
719 alternative partitioning methods and TSHZ1: a new phylogenetic marker for reptiles.
720 *Molecular Phylogenetics and Evolution* 50: 391–396. DOI: 10.1016/j.ympev.2008.10.018.

721

722 Scott NJ, Giraudo AR, Scrocchi G, Aquino AL, Cacciali P, Motte M. 2006. The genera
723 *Boiruna* and *Clelia* (Serpentes: Pseudoboini) in Paraguay and Argentina. *Papéis Avulsos de*
724 *Zoologia (São Paulo)* 46(9): 77–105. DOI: 10.1590/S0031-10492006000900001.

725

726 Slevin JR. 1942. Notes on a collection of reptiles from Boquete, Panama, with the description
727 of a new species of *Hydromorphus*. *Proceedings of the California Academy Sciences* 23: 463–
728 480.

729

730 Smith EN, Campbell JA. 1994. A new species of *Rhadinea* (Colubridae) from the Caribbean
731 Versant of Guatemala. *Occasional Papers of the Museum of Natural History, University of*
732 *Kansas* 167: 1–9.

733

734 Smith HM, Taylor EH. 1945. An annotated checklist and key to the snakes of Mexico. *United*
735 *States National Museum Bulletin* 187: iv + 1–239.

736

737 Solorzano A. 2004. *Serpientes de Costa Rica - Snakes of Costa Rica*. Costa Rica: Editorial
738 INBio, , 792 pp.

739

740 Stafford PJ. 2004. A new species of *Tantilla* (Serpentes; Colubridae) of the *taeniata* group
741 from Southern Belize. *Journal of Herpetology* 38(1): 43–52. DOI: 10.1670/208-02a.

742

743 Stamatakis A, Hoover P, Rougemont J. 2008. A Rapid Bootstrap Algorithm for the RAxML
744 Web-Servers. *Systematic Biology* 75: 758–771. DOI: 10.1080/10635150802429642

745

746 Starace F. 1998. *Guide des Serpents et Amphisbènes de Guyane*. Guadeloupe, Guyane: IBIS
747 Rouge Editions, 450 pp.

748

749 Stattersfield AJ, Crosby MJ, Long AJ, Wege D. 1998. *Endemic birds areas of the world.*
750 *Priorities for biodiversity conservation*. Cambridge, UK: BirdLife International, 815 pp.

751

752 Stuart LC, Bailey JR. 1941. Three new species of the genus *Rhadinaea* from Guatemala.
753 *Occasional Papers of the Museum of Zoology, University of Michigan* 442: 1–11.

754

- 755 Stuart LC. 1949. A new *Trimetopon* (Ophidia) from Guatemala. *Proceedings of the Biological*
756 *Society of Washington* 62: 165–168.
- 757
- 758 Stuart LC. 1954. Descriptions of some new amphibians and reptiles from Guatemala.
759 *Proceedings of the Biological Society of Washington* 67: 159–178.
- 760
- 761 Van Denburgh J. 1912. Expedition of the California Academy of Sciences to the Galapagos
762 Islands, 1905–1906. IV. The snakes of the Galapagos Islands. *Proceedings of the California*.
763 *Acadademy of. Sciences* (Ser. 4) 1(4): 323–374.
- 764
- 765 Venegas PJ. 2005. Herpetofauna del bosque seco Ecuatorial de Peru: taxonomía, ecología y
766 biogeografía. *Zonas Áridas* 9: 9–26.
- 767
- 768 Venegas PJ, Townsend JH, Koch C, Böhme W.2008. Two new sympatric species of leaf-toed
769 geckos (Gekkonidae: *Phyllodactylus*) from the Balsas region of the upper Marañón Valley,
770 Peru. *Journal of Herpetology* 42(2): 386–396. DOI: 10.1670/07-1341.1.
- 771
- 772 Vidal N, Kindl SG, Wong A, Hedges SB. 2000. Phylogenetic relationships of Xenodontine
773 snakes inferred from 12S and 16S ribosomal RNA sequences. *Molecular Phylogenetics and*
774 *Evolution* 14(3): 389–402. DOI: 10.1006/mpev.1999.0717.
- 775
- 776 Villa JD. 1971. *Crisantophis*, a new genus for *Conophis nevermanni* Dunn. *Journal of*
777 *Herpetology* 5: 173–77. DOI: 10.2307/1562739.
- 778
- 779 Villa JD. 1990. *Hydromorphus* Peters. Central American watersnakes. *Catalogue of American*
780 *Amphibians and Reptiles* 472: 1–2.
- 781
- 782 Werneck FP, Costa GC, Colli GR, Prado DE, Sites Jr JW. 2011. Revisiting the historical
783 distribution of Seasonally Dry Tropical Forests: new insights based on palaeodistribution
784 modelling and palynological evidencegeb. *Global Ecology and Biogeography* 20: 272–288.
785 DOI: 10.1111/j.1466-8238.2010.00596.x.
- 786
- 787 Werner F. 1924. Neue oder wenig bekannte Schlangen aus dem Naturhistorischen
788 Staatsmuseum in Wien. 1. Teil. *Sitzungsberichte der Akademie der Wissenschaften in Wien,*
Abt. I, 133: 29–56.

789

790 Wilson LD. 1976. Variation in the colubrid snake *Tantilla semicincta* (Dumeril, Bibron, and
791 Dumeril), with comments on pattern dimorphism. *Bulletin of the Southern California
792 Academy of Sciences* 75: 42–48.

793

794 Wilson LD. 1987. A resume of the Colubrid snakes of the genus *Tantilla* of South America.
795 *Milwaukee Public Museum Contributions in Biology and Geology* 68: 1–35.

796

797 Wilson LD, Mata-Silva V. 2015. A checklist and key to the snakes of the *Tantilla* clade
798 (Squamata: Colubridae), with comments on distribution and conservation. *Mesoamerican
799 Herpetology* 2(4): 418–498.

800

801 Wilson LD, Mena CE. 1980. Systematics of the *melanocephala* group of the colubrid snake
802 genus *Tantilla*. *San Diego Society of Natural History Memoirs* 11: 5–58.

803

804 Zaher H. 1996. A new genus and species of pseudoboine snake, with a revision of the genus
805 *Clelia* (Serpentes, Xenodontinae). *Bolletino dei Musei Regionale Scienza naturali di Torino*
806 14: 289–337.

807

808 Zaher H. 1999. Hemipenial morphology of the South American xenodontine snakes, with a
809 proposal for a monophyletic Xenodontinae and a reappraisal of colubroid hemipenes. *Bulletin
810 of the American Museum of Natural History* 240:1–168.

811

812 Zaher H, Grazziotin FG, Cadle JE, Murphy RW, Moura-Leite JC. 2009. Molecular phylogeny
813 of advanced snakes (Serpentes, Caenophidia) with an emphasis on South American
814 Xenodontines: a revised classification and descriptions of new taxa. *Papéis Avulsos de
815 Zoologia (São Paulo)* 49(11): 115–153. DOI: 10.1590/S0031-10492009001100001.

816

817 Zaher H, Prudente AL. 2003. Hemipenes of *Siphlophis* (Serpentes, Xenodontinae) and
818 techniques of hemipenial preparation in snakes: A response to Dowling. *Herpetological
819 Review* 34: 302–307.

820

821 APPENDIX. Additional specimens examined.

822

823 *Tantilla capistrata*: PERU: Lambayeque: Chaparrí, 446 m a.s.l. (ZFMK 85028).

824

825 *Tantilla relicta*: USA: Florida: Bushnell (ZFMK 84387).