

Changes in understory species occurrence of a secondary broadleaved forest after mass mortality of oak trees under deer foraging pressure

Hiroki Itô¹

¹Hokkaido Research Center, Forestry and Forest Products Research Institute, Toyohira, Sapporo 305-8687, Japan. Email address: abies.firma@gmail.com

ABSTRACT

The epidemic of mass mortality of oak trees by Japanese oak wilt has affected secondary deciduous broadleaved forests that have been used as coppices in Japan. The dieback of oak trees formed gaps in the crown that would be expected to enhance the regeneration of shade-intolerant pioneer species. However, foraging by sika deer *Cervus nippon* has also affected forest vegetation, and the compound effects of both on forest regeneration should be considered when they simultaneously occur. A field study was conducted in Kyôto City, Japan, to investigate how these compound effects affected the vegetation of the understory layer of these forests. The presence/absence of seedlings and saplings was observed for 200 quadrats sized 5 m × 5 m for each species in 1992, before the mass mortality and deer encroachment, and in 2014 after these effects. A hierarchical Bayesian model was constructed to explain the occurrence, survival, and colonization of each species with their responses to the gaps that were created, expanded, or affected by the mass mortality of *Quercus serrata* trees. The species that occurred most frequently in 1992, *Eurya japonica*, *Quercus glauca*, and *Cleyera japonica*, also had the highest survival probabilities. Deer-unpalatable species such as *Symplocos prunifolia* and *Triadica sebifera* had higher colonization rates in the gaps, while the deer-palatable species *Aucuba japonica* had the smallest survival probability. The gaps thus promoted the colonization of deer-unpalatable plant species such as *Symplocos prunifolia* and *Triadica sebifera*. In the future, such deer-unpalatable species may dominate gaps that were created, expanded, or affected by the mass mortality of oak trees.

Keywords: deer-unpalatable plant species, gap, Japanese oak wilt, sika deer, understory vegetation

INTRODUCTION

Many coppices have been abandoned for socio-economic reasons, such as the replacement of woody fuels with fossil fuels in Europe (Rackham, 2008; Müllerová et al., 2015; Svátek and Matula, 2015). This abandonment parallels that of Japan (Suzuki, 2013), and a considerable number of deciduous oak forests grown from such coppices have suffered from the mass mortality of oak trees (Kuroda et al., 2012; Nakajima and Ishida, 2014). ~~This mass mortality has been~~ caused by Japanese oak wilt (Kuroda et al., 2012).

The direct cause of the mortality is a pathogenic species of fungus *Raffaelea quercivora* Kubono et Shin. Ito, which is carried by the ambrosia beetle *Platypus quercivorus* Murayama (Kubono and Ito, 2002; Kinuura and Kobayashi, 2006). In addition, it has been pointed out that the coppice abandonment is an indirect cause of the epidemic because the ambrosia beetles reproduce more successfully in large oak stems, which are more abundant as abandoned oak stems grow up (Kobayashi and Ueda, 2005). There is ~~an~~ another possible reason why the abandonment indirectly affected the epidemic: in the period when coppices were managed, oak stems that died from the wilt were quickly felled and utilized for fuel or charcoal, and as a result, it prevented outbreaks of the disease (Ida and Takahashi, 2010).

The mass mortality altered the structures of damaged oak forests. The two major deciduous oak species in Japan, *Quercus crispula* Blume and *Quercus serrata* Murray, are vulnerable to the pathogenic fungus. Nakajima and Ishida (2014) showed that $80 \pm 19\%$ (mean $\pm$ standard deviation) stems of *Quercus crispula* died, while $34 \pm 19\%$ stems of *Quercus serrata* died. Naka (1982) studied an old-growth evergreen forest in the Kasugayama Forest Reserve in Nara City, which is located about 40 km south of the study site, and showed that the major gap generator was typhoons, and that the interval was 6.57 years with a tree fall rate for overstory trees of 0.84 trees/ha/year, and that the canopy opening rate was $55.6 \text{ m}^2/\text{ha}/\text{year}$. Though these rates are not directly applicable to deciduous secondary oak forests, the gaps created by the mortality would not be negligible for oak-dominated forests. If *Quercus serrata* covers 30% of the canopy and 30% of them die, $900 \text{ m}^2 / \text{ha}$ of the canopy will become gaps. How such damaged oak forests ~~are~~ regenerated depends on circumstance; sub-canopy trees might grow to canopy trees in some cases (Itô et al., 2009), and dense floor vegetation such as dwarf bamboo might inhibit regeneration in other cases (Itô et al., 2011; Saito and Shibata, 2012). In the latter case, the damaged forests may lack a canopy layer for a long time.

For the last several decades, herbivory by overabundant deer populations has negatively affected forest vegetation in Europe and North America (Rooney, 2001; Côté et al., 2004; Rackham, 2008). This is paralleled in Japan: sika deer (*Cervus nippon* Temminck) is one major inhibitor of forest regeneration (Takatsuki, 2009; Suzuki, 2013; Iijima and Nagaike, 2015). However, little is known about how regeneration proceeds after mass mortality under deer foraging pressure (Obora et al., 2013). Gap formations should improve light conditions on the forest floor and promote the regeneration of many tree species (Suzuki, 2013). However, excessive browsing should inhibit regeneration, except for unpalatable and browse-tolerant species (Shimoda et al., 1994). Therefore, a combination of both factors might promote the regeneration of species that are unpalatable and/or browse-tolerant to deer. This in turn may alter the species composition of forests that regenerate from the mass mortality of oak trees over time.

It is important for the management of secondary forests to predict what tree or shrub species will be recruited in forest stands damaged by the mass mortality of oak trees and deer impact. In a previous study, Itô (2015) described the changes in the canopy/sub-canopy and understory layers of a damaged forest by comparing vegetation before and after mass mortality of oak trees and deer foraging. It was found that regenerating species were limited to the originally abundant species, such as *Quercus glauca* Thunb., *Cleyera japonica* Thunb., and *Eurya japonica* Thunb. var. *japonica*, as well as to species unpalatable to deer, such as *Symplocos prunifolia* Siebold

et Zucc. and *Triadica sebifera* (L.) Small. However, the study only described the changes in species occurrence and failed to estimate the specific probabilities of survival and colonization. In this study, the previous data from the understory layers were reanalyzed using a hierarchical Bayesian model that explicitly incorporates probabilities of ecological processes such as occurrence, survival, and colonization. A hierarchical model incorporating random species effects also makes it possible to estimate those probabilities by species, “borrowing strength from the ensemble” (Kéry and Schaub, 2012). These advantages should be of use in predicting which species will dominate such stands in the future.

MATERIALS AND METHODS

Study site

The field data were collected in the Ginkakuzi-san (or Ginkakuji-san in a different spelling) National Forest located in Kyôto City, Japan (35.029°N, 135.801°E). The yearly average temperature from 1981 to 2010 was 15.9°C and the average precipitation was 1491.3 mm at the Kyôto Local Meteorological Office. Elevation of the forest was 100–290 m above sea level, and the forest was in the warm temperate zone. The national forest was protected for its landscape and erosion control, and most of it was situated in the buffer zone of the UNESCO world heritage historic monuments of ancient Kyôto. The fieldwork was conducted with permission under an agreement between the Kyôto-Ôsaka District Forest Office and the Forestry and Forest Products Research Institute.

In the 1930s, most of the forest was covered with a mix of pines (*Pinus densiflora* Siebold et Zucc.) and broadleaved trees including oak (*Quercus serrata*). After the 1960s, many pine trees had died due to the pine wilt disease. Recently, most of the national forest has been covered with a secondary broadleaved forest consisting of many species such as evergreen oak *Quercus glauca*, evergreen subcanopy species *Symplocos prunifolia*, and deciduous tree species *Ilex macropoda* Miq., although conifers (*Cryptomeria japonica* (L.f.) D.Don and *Chamaecyparis obtusa* (Siebold et Zucc.) Endl.) were planted in a small part of the area (Itô, 2007). In 2005, the mass mortality of oaks was first recognized in the eastern part of Kyôto City, in which the Ginkakuzi-san National Forest is located, and then the damage expanded (Itô, 2015). In addition, damage by sika deer resulting from browsing and bark-stripping have been noticeable over the same period. Deer had been seldom seen in the 1990s, in my observation, but by this time inhabited the forest throughout the year (Itô, 2015).

In 1992, a 0.5 ha (100 × 50 m) plot was established on a south-facing slope in the national forest at an elevation of 140–195 m. The average slope inclination was about 30° and the surface geology was granite. All the stems in the plot were marked and their diameters at breast height (dbh) were measured in 1993, 1996, 1999, 2002, 2005, and 2014. Mainly due to the mass mortality of oak trees *Quercus serrata*, the basal area in the plot decreased from 43.3 m²/ha in 2005 to 39.5 m²/ha in 2014, while the number of stems in the plot increased from 1554 to 1645 (3108 stems/ha to 3290 stems/ha in stem density). There were 36 *Quercus serrata* stems in the plot in 2005, and 21 of them died by 2014. These trees were most likely killed by Japanese oak wilt. This affected the forest structure in the plot via the formation of new canopy gaps or

expanding existing gaps. In some area, the mortality indirectly affected the forest floor via treatments of dead stems by the forest office (cutting down the stems, cutting the fallen stems into pieces, and disinfecting them) (Itô, 2015). On the other hand, none of the evergreen oaks *Quercus glauca* died from the disease, though some were attacked by the ambrosia beetles, being less vulnerable than *Quercus serrata* (Murata et al., 2005, 2009). In the understory layer, *Quercus glauca* and some evergreen shrub species such as *Eurya japonica* and *Cleyera japonica* were frequently observed. Overall changes in the species composition of the site from 1990s to 2010s were reported in Itô (2015).

Data collection

The plot was divided into 200 quadrats sized 5 m × 5 m. In 1992, all woody plant seedling or saplings (age ≥ 1 year and dbh < 3 cm) were tallied by species. The plots were resampled in 2014. In 2014, each quadrat was classified into inside or outside of the gaps, which were formed by the death of oak trees, or had been gaps at least since 2005 and were affected in addition by fallen dead oak trees. Quadrats on the boundary were visually classified; if an open area which was created, expanded, or affected by the mass mortality and occupied most of the quadrat, the quadrat was classified as a gap quadrat. Twenty of 200 quadrats were classified as gaps created or affected by oak death (Fig. 1). The size of the largest gap was approximately 250 m² consisting of 10 adjacent quadrats.




Figure 1. Map showing the study site.

Gap denotes the gap quadrats created, expanded, or affected by the mass mortality of oak trees. Non-gap denotes the rest of the quadrats. Units of the X and Y axes are in meters.

In 1992, 55 species were observed and 58 species in 2014 (Itô, 2015), and 75 species were observed in total, combining data from both years. The maximum number of quadrats in 1992 in which a given species occurred was for *Quercus glauca* and *Eurya japonica*, both of which were observed in 184 of 200 quadrats. The mean and median numbers of quadrats in 1992 in which a given species occurred were 19.5 and 4,

153 respectively, excluding species which were not ~~occurred~~ in 1992 in any quadrats. On the
 154 other hand, the maximum in 2014 was 193 for *Quercus glauca*. The mean and median
 155 were 15.6 and 5, respectively, excluding species which ~~were not occurred~~
 156 quadrats.

157 Data analysis was conducted for 42 of the species that were observed in more than 5
 158 quadrats over both sample years (200 quadrats $\times$ 2 observations).

159 Statistical Modeling

160 A hierarchical Bayesian model was constructed to determine how species were affected
 161 by the recent environmental changes in the forest.

162 The presence/absence (presence = 1, absence = 0) of species i in quadrat j in the
 163 year 1992 (y_{1ij}) and 2014 (y_{2ij}) was assumed to follow the Bernoulli distribution given
 164 the occurrence probability ψ_{1ij} and ψ_{2ij} , as follows,

$$y_{1ij} \sim \text{Bernoulli}(\psi_{1ij})$$

$$y_{2ij} \sim \text{Bernoulli}(\psi_{2ij}).$$

165 To be exact, the “presence/absence” was “detection/nondetection” (Dorazio et al.,
 166 2006; Kéry and Schaub, 2012). It has been pointed out that detection probability should
 167 be considered to correctly estimate population properties such as the occurrence rate
 168 or survival rate, even if the observed objects are plants (Kéry, 2004; Chen et al., 2009,
 169 2013). The present study had, ~~regrettably~~, only one observation for each survey year.
 170 However, the quadrat size was rather small (5 m $\times$ 5 m) and the whole of each quadrat
 171 was explored, so I expected that the detection probability should be near to one and
 172 therefore “detection/nondetection” was regarded as “presence/absence” in this study.
 173 Chen et al. (2009) showed that the detection probability asymptotically approaches one
 174 with larger survey efforts.

The parameter of occurrence in 1992, ψ_{1ij} , was formulated as follows:

$$\text{logit}(\psi_{1ij}) = \beta_o + \varepsilon_{oi} + r_j,$$

175 where β_o denotes an intercept or overall mean of ψ_1 on the logit scale, and ε_{oi} denotes
 176 the random species effect on the intercept of species i . The parameter r_j denotes a
 177 spatially autocorrelated random effect of quadrat j .

The parameter of occurrence in 2014, ψ_{2ij} , was formulated as follows:

$$\psi_{2ij} = y_{1ij}\phi_{ij} + (1 - y_{1ij})\gamma_{ij},$$

178 where parameter ϕ_{ij} denotes the ‘survival’ probability that species i was present in
 179 quadrat j in 1992 and still present in 2014. The parameter γ_{ij} denotes the ‘colonization’
 180 probability that species i was absent in quadrat j in 1992 but present in 2014.

The parameters of survival ϕ_{ij} and colonization γ_{ij} were formulated as follows:

$$\text{logit}(\phi_{ij}) = \beta_s + \varepsilon_{si} + (\beta_{sg} + \varepsilon_{sgi})g_j$$

$$\text{logit}(\gamma_{ij}) = \beta_c + \varepsilon_{ci} + (\beta_{cg} + \varepsilon_{cgi})g_j,$$

where the parameters β_s and β_c are intercepts or overall means of ϕ and γ on the logit scale, respectively. The parameters ε_{si} and ε_{ci} are random species effects on the intercepts, β_{sg} and β_{cg} are coefficients of the gap predictor g_j (0: non-gap quadrats, 1: gap quadrats affected by oak mortality), and ε_{sgi} and ε_{cgi} are random species effects on the coefficients.

Priors of the random species effects were defined hierarchically; hyperparameters, σ_o , σ_s , σ_{sg} , σ_c , and σ_{cg} , scaled the distribution of ε_{oi} , ε_{si} , ε_{sgi} , ε_{ci} , and ε_{cgi} , respectively, as follows:

$$\varepsilon_{oi} \sim \text{Normal}(0, \sigma_o^2)T(-10, 10),$$

where $\text{Normal}(0, \sigma^2)T(-10, 10)$ denote a normal distribution truncated between -10 and 10; the truncation was incorporated to stabilize the logit scale parameters (Kéry and Schaub, 2012). Priors of the parameters β_o , β_{op} , β_{os} , β_s , β_{sg} , β_c , and β_{cg} were defined as $\text{Normal}(0, 10^4)T(-10, 10)$. The prior of the spatial effect r_j was defined as an intrinsic conditional autoregressive model as follows:

$$r_j | r_{-j} \sim \text{Normal} \left(\sum_{k \neq j} \frac{w_{jk} r_j}{w_{j+}}, \frac{\sigma_r^2}{w_{j+}} \right),$$

where r_{-j} denote the values of r except the quadrat j , a variable w_{jk} was defined to be 1 if quadrat j and quadrat k are adjacent, and 0 if not, and w_{j+} was defined to be $\sum_k w_{jk}$. The parameter σ_r^2 denotes a variance of the random effect.

Presence/absence data for 42 species, which were observed in more than 5 quadrats in total over two surveys, 1992 and 2014, were used for the parameter estimation. To estimate the posterior distribution for each parameter, the Markov chain Monte Carlo (MCMC) method was adopted; this simulation method generates Markov chains drawing values from the target posterior distributions if the chains converge to stationary distributions (Gelman et al., 2013). Four parallel chains were generated in this study, and each of them had 13,000 iterations while the first 3,000 iterations were dropped as burn-in. The MCMC sample was taken from the three chains with 10 thinning intervals, so that the sample size was 4,000. OpenBUGS 3.2.3 (Lunn et al., 2009) was used for the computation. The BUGS code is available in List S2. To check the convergence, Gelman-Rubin statistics ($\hat{R}$) were calculated (Gelman and Rubin, 1992; Brooks and Gelman, 1998; Gelman et al., 2013). When the Markov chains successfully converge, the value of $\hat{R}$ becomes nearly one. If the value of $\hat{R}$ is no larger than 1.1, the chains are usually regarded as converged (Kéry and Schaub, 2012).

RESULTS

For each parameter, the values of $\hat{R}$ were no larger than 1.1 for each parameter, so that the Markov chains seemed to reach convergence. However, some random species effects on coefficients of gaps had rather wide posteriors as mentioned later. Those parameters might lack enough information to obtain satisfactory estimates.

216 Posterior mean, median, and 95% CI (Credible Interval) of the overall occurrence
 217 probability in 1992 (probability that a species was present in a quadrat in 1992), β_o ,
 218 was estimated to be -3.25 for the posterior mean, -3.25 for the median, and -3.99 to
 219 -2.54 for the 95% CI (Table 1). The value -3.25 on the logit scale is equivalent to 0.037
 220 ($= 1/(1 + \exp(3.25))$) on the probability scale. Therefore, a species was expected to
 221 occur in 3.7% of quadrats on average in 1992. In the same manner, the 95% CI was
 222 1.8–7.3% on the probability scale.

223 The overall survival probability (probability that a species was present in a quadrat
 224 in 1992 and still present in the same quadrat in 2014), β_s , was estimated to be -2.60 for
 225 the posterior mean, -2.57 for the median, and -4.11 to -1.44 for the 95% CI. The mean
 226 value was equivalent to a probability that a species occurring in 1992 surviving in the
 227 same quadrat in 2014 was expected to be 6.9%, and the 95% CI was 1.6–19.2%.

228 On the other hand, the overall colonization probability (probability that a species was
 229 absent in a quadrat in 1992 but present in the same quadrat in 2014), β_c , was estimated to
 230 be -3.81 for the posterior mean, -3.81 for the median, and -4.45 to -3.18 for the 95% CI.
 231 The value -3.81 was equivalent to 0.022 on the probability scale; so that the probability
 232 that a species which was absent in a quadrat in 1992 had colonized into the quadrat in
 233 2014 was expected to be 2.2%, and the 95% CI was 1.2–4.0%.

234 Posterior mean of coefficients of the gap on the survival β_{sg} and colonization β_{cg}
 235 were -0.17 and 1.51, respectively, and β_{cg} did not include zero in the 95% CI (0.86–
 236 2.08), while β_{sg} included zero in the 90% CI (-1.27–0.81). The mean value 1.51
 237 of β_{cg} meant that gaps improved the average colonization probability 2.2% to 9.1%
 238 ($= 1/(1 + \exp(3.81 - 1.51))$) on average.

Table 1. Parameter estimates (posterior mean, standard deviation (SD), and 2.5%, 5%, 50%, 95%, and 97.5% quantiles) other than random effects.

Positive values of the intercepts (β_o , β_s , and β_c) on the logit scale mean that the corresponding probabilities are larger than 0.5, and vice versa. Positive values of the coefficients of gap (β_{sg} and β_{cg}) mean that the gap increases the corresponding probabilities. Parameters σ_o and below are standard deviations of the corresponding predictors.

	Mean	SD	2.5%	5%	50%	95%	97.5%
β_o	-3.25	0.37	-3.99	-3.88	-3.25	-2.65	-2.54
β_s	-2.60	0.67	-4.11	-3.81	-2.57	-1.61	-1.44
β_{sg}	-0.17	0.65	-1.54	-1.27	-0.14	0.81	1.04
β_c	-3.81	0.32	-4.45	-4.33	-3.81	-3.28	-3.18
β_{cg}	1.51	0.31	0.86	0.98	1.53	1.99	2.08
σ_o	2.27	0.30	1.76	1.82	2.24	2.81	2.95
σ_s	2.89	0.57	1.97	2.07	2.84	3.91	4.13
σ_{sg}	1.51	0.84	0.19	0.31	1.40	3.07	3.48
σ_c	1.89	0.28	1.43	1.48	1.86	2.40	2.51
σ_{cg}	1.35	0.32	0.81	0.88	1.32	1.93	2.07
σ_r	0.44	0.10	0.27	0.29	0.44	0.60	0.65

239 Random species effects on occurrence probability in 1992 (ϵ_o) are shown in Fig. 2.
 240 *Eurya japonica* and *Quercus glauca* had the largest value, followed by *Cleyera japonica*,

241 *Aucuba japonica* Thunb. var. *japonica*, *Ilex crenata* Thunb., *Photinia glabra* (Thunb.)
 242 Maxim., and so on. The posterior mean for *Eurya japonica* and *Quercus glauca* was
 243 5.71, and this meant that the occurrence probability of the species was improved from
 244 an overall mean of 3.7% to 92.1% ($= 1/(1 + \exp(3.25 - 5.71))$). The 95% CI of the
 245 random species effect for *Eurya japonica*, 4.85–6.62, corresponded to 83.2–96.7% on
 246 the occurrence probability for the species. In the same manner, the expected occurrence
 247 probabilities were 59.1% for *Cleyera japonica*, 53.2% for *Aucuba japonica*, 34.8% for
 248 *Ilex crenata*, and 31.6% for *Photinia glabra*. On the other hand, *Carpinus tschonoskii*
 249 Maxim., *Celtis sinensis*, *Zanthoxylum ailanthoides* Siebold et Zucc., and *Triadica*
 250 *sebifera* had the smallest values because these species were not detected in 1992 (Fig. 2).
 251 The posterior mean for *Carpinus tschonoskii* was -3.21 on the logit scale and the expected
 252 occurrence was 0.2% ($= 1/(1 + \exp(3.25 + 3.21))$) on the probability scale. The 95%
 253 CI of the species, -6.08 to -1.11, corresponded to an occurrence probability of 0.0–1.3%
 254 on the probability scale.

255 Random species effects on the intercept of survival from 1992 to 2014 (ϵ_s) are shown
 256 in Fig. 3A. *Quercus glauca* had the largest value, and *Cleyera japonica*, *Eurya japonica*,
 257 and *Camellia japonica* L. followed. The posterior mean of *Quercus glauca* was 5.93,
 258 and the expected survival probability was 96.5% ($= 1/(1 + \exp(2.60 - 5.93))$) without
 259 the gap effects. The expected probabilities were 88.3% for *Cleyera japonica*, 76.7% for
 260 *Eurya japonica*, and 55.7% for *Camellia japonica*. On the other hand, the posterior of
 261 *Aucuba japonica* was less than zero within 95% CI (Fig. 3A). The posterior mean of
 262 the random effect was -3.80 and the expected value of survival probability was 0.2%.
 263 The posterior mean of *Vaccinium bracteatum* Thunb. was the second smallest and the
 264 value was -3.03; the expected survival probability was 0.4%. Species without surviving
 265 individuals had nearly zero means and wider CIs, such as for *Zanthoxylum ailanthoides*.
 266 Random species effects on the survival coefficients of the gap predictor were rather
 267 small as absolute values and included zero in their 90% CI for all species (Fig. 3B).

268 Random species effects on intercepts of colonization from 1992 to 2014 were the
 269 largest in *Quercus glauca*, followed by *Eurya japonica*, *Cleyera japonica*, *Symplocos*
 270 *prunifolia*, *Photinia glabra*, and so on (Fig. 4A). The posterior mean of the random
 271 effect for *Quercus glauca* was 6.09 and the expected colonization probability without
 272 the gap effects was 90.7%. In the same manner, the expected colonization probabilities
 273 for *Cleyera japonica*, *Symplocos prunifolia*, and *Photinia glabra* were 28.3%, 26.7%,
 274 and 14.1%, respectively.

275 Random species effects on the colonization coefficient of the gap predictor were the
 276 largest for *Callicarpa mollis* Siebold et Zucc., followed by *Zanthoxylum ailanthoides*,
 277 *Carpinus tschonoskii*, *Triadica sebifera*, and so on. They were the smallest in *Cleyera*
 278 *japonica*. The posterior mean of the random effect for *Callicarpa mollis* was 2.32, so that
 279 the expected colonization probability for the species in the gap quadrats was estimated
 280 to be 25.9% ($= 1/(1 + \exp(3.81 - 1.51 + 1.07 - 2.32))$), as the posterior mean of the
 281 random species effect on the colonization for the species was -1.07. Some species such
 282 as *Quercus glauca* had wider CI in their posteriors. This might be because there was
 283 little information for such species due to the small number of colonizers within the gaps
 284 (Fig. 4B).

285 Complete estimates of random effects are available in Table S3.




Figure 2. Random species effects on occurrence in 1992 ϵ_0 .

Thin lines denote 95% credible intervals (CI), thick lines denote 90% CI, and circles denote medians. Larger positive values in the posterior indicate that the species will be more likely to be present in 1992 than the overall mean, and negative values indicate that the species will be less likely.




Figure 3. Random species effects on survival.

(A) ϵ_s , on the intercept, (B) ϵ_{sp} and (B) ϵ_{sg} , on the coefficient of the gap. Thin lines denote 95% credible intervals (CI), thick lines denote 90% CI, and circles denote medians. Larger positive values in the posterior indicate that a species that had been present in a quadrat in 1992 will be more likely to be present in the same quadrat in 2014 than the overall mean, and negative values indicate that the species will be less likely.




Figure 4. Random species effects on colonization.

(A) ϵ_c , on the intercept and (B) ϵ_{cg} , on the coefficient of the gap. Thin lines denote 95% credible intervals (CI), thick lines denote 90% CI, and circles denote medians. Larger positive values in the posterior indicate that species that had been absent in a quadrat in 1992 will be more likely to be present in the same quadrat in 2014 than the overall mean, and negative values indicate that the species will be less likely.

DISCUSSION

The mean probability of overall occurrence in 1992 for a single species was expected to be only 3.7% for each quadrat. This reflects the fact that many species occurred only in a small fraction of the quadrats. The most frequent species in 1992 were all evergreen tree or shrub species such as *Eurya japonica*, *Quercus glauca*, *Cleyera japonica*, *Aucuba japonica*, *Ilex crenata*, and *Photinia glabra*; the expected occurrence probabilities of these species were all larger than 30%. Species with large values of random species effects concerning occurrence corresponded to these species as a matter of course (Fig. 2). On the other hand, the least occurring species were deciduous trees or shrubs such as *Carpinus tschonoskii*, *Zanthoxylum ailanthoides*, *Celtis sinensis* Pers., and *Triadica sebifera*; these species were not detected in the quadrats in 1992. The canopy of the forest was almost closed in 1992, so that shade-tolerant evergreen tree or shrub species dominated the understory layer and deciduous early-successional species infrequently occurred.

Overall survival and colonization probability of a species was expected to be 6.9% and 2.2%, respectively, on average. This may seem small, but some species have large random effect values allowing them to survive or colonize at high probabilities. The average colonization probability was estimated to be improved in the gap quadrats. This was in accord with the preceding study showing that gap creation increased floor species richness even under deer pressure (Suzuki, 2013).

The species that had the highest survival probability in the non-gap quadrats was *Quercus glauca* (Fig. 3A); the mean probability of survival was expected to be 96.5%. *Quercus glauca* produces many sprouts (Cai, 2000), and the sprouts may survive when the main stems die. This trait might contribute the high survival probability. *Cleyera japonica*, *Eurya japonica*, and *Camellia japonica* followed *Quercus glauca* in their survival probabilities. These species, including *Quercus glauca*, were all evergreen broadleaved species that could survive under a closed canopy, as shown by their occurrence. In contrast, *Aucuba japonica*, a species of evergreen shrub, had the smallest survival probability at 0.2%. This species was found in 107 of 200 quadrats in 1992, but never found in 2014. *Aucuba japonica* is known to be a preferred food for sika deer and is vulnerable to deer browsing (Hashimoto and Fujiki, 2014). The encroachment of deer likely explains its small survival probability. The posterior mean of the random effect was the second smallest in *Vaccinium bracteatum*. There is relatively little information on the palatability of *Vaccinium bracteatum* for sika deer, and some reports refer to it as a food plant, while others refer to it as unpalatable (Nakajima, 1929; Kabaya, 1988; Takatsuki, 1989; Hashimoto and Fujiki, 2014). The present results suggest that the species is vulnerable to deer impacts. The random species effects on the survival coefficients of the gap predictor were not so clear (Fig 3B). This may due to the small sample size of the gap quadrats to some extent, and due to the small number of light-demanding species occurring in 1992 to some extent.

The colonization probability in the non-gap quadrat was highest in *Quercus glauca*, and evergreen shrub or sub-canopy trees followed (Fig. 4A). Among them, *Quercus glauca* had an especially high colonization probability; the expected value was 90.7%. This might be because numerous seeds of this species are dispersed in the forest due to the abundance of mother trees (Itô, 2007; Itô, 2015), and it could persist at the

seedling stage (Itô, 2009). On the other hand, the four species whose colonization probabilities were largest in the gaps are all shade-intolerant (Shimoda et al., 1994; Shibata and Nakashizuka, 1995). In addition to the four species, *Symplocos prunifolia* is also considered to be a shade-intolerant species though it is an evergreen (Fujii, 1994). However, *Symplocos prunifolia* and *Triadica sebifera* are unpalatable plants for sika deer (Shimoda et al., 1994; Hashimoto and Fujiki, 2014). *Triadica sebifera* is an alien species in Japan. The species is unpalatable for sika deer (Shimoda et al., 1994; Hashimoto and Fujiki, 2014), and as of 2002, it was increasing on Mt. Kasugayama (Maesako et al., 2007). Shimoda et al. (1994) studied the deer effects on pioneer species on Mt. Kasugayama, which was inhabited by many deer; the authors found that pioneer species including *Zanthoxylum ailanthoides* and *Callicarpa mollis* emerged in gaps but rarely survived or matured due to deer foraging pressure.

In the study site, few large stems (height ≥ 50 cm) of *Zanthoxylum ailanthoides*, *Callicarpa mollis*, and *Carpinus tschonoskii* were found, though a greater number of those of *Quercus glauca*, *Cleyera japonica*, and *Eurya japonica* were found in the quadrats that were not affected by the mass oak mortality (Itô, 2015). The latter species are evergreen trees or shrubs, and they had been dominant at least since 1992. In addition to these species, large plants of *Symplocos prunifolia* and *Triadica sebifera* were found in the gap quadrats created, expanded, or affected by the mass mortality (Itô, 2015), and the high colonization rate of these species in the gaps might contribute to their regeneration. Suzuki (2013) pointed out that succession after the abandonment of coppices in which there was gap creation under deer herbivory pressure would make the vegetation be dominated by shade-tolerant and herbivory-tolerant species. In the present study, the two shade-intolerant and deer-unpalatable species frequently colonized into the gap quadrats and grew up. The difference may be due to the abundance of the two species or overall differences in the vegetation.

Obara et al. (2013) studied the effect of deer herbivory on forest regeneration after the mass mortality of oak trees and described that only some fern species that were deer-unpalatable increased under deer herbivory, while many species including trees increased within enclosure. Obara et al. (2013), however, also mentioned that the tree species were suppressed by shrub and grass species even in the enclosures. This suggests the difficulty of forests regenerating from damage caused by the mass mortality of oak trees. A similar issue was also reported in (Itô et al., 2011), in which they studied a forest that had been damaged by mass mortality and whose floor was dominated by dwarf bamboo. In the present study, such shrub or grass species that dominated the shrub layer or floor were not found, but a deer-unpalatable fern, *Hypolepis punctata* (Thunb.) Mett. ex Kuhn, could potentially dominate the floor because it partially colonized the gap quadrats (Itô, 2015). Such deer-unpalatable ferns and/or herbaceous species may suppress regeneration of tree species, in which case they colonize into the gaps and dominate the floor.

CONCLUSION

Gaps that were created, expanded, or affected by the mass mortality of oak trees might be beneficial for the colonization of pioneer species. Shade-intolerant species such as *Callicarpa mollis*, *Zanthoxylum ailanthoides*, *Carpinus tschonoskii*, *Triadica sebifera*,

and *Symplocos prunifolia* were estimated to more frequently colonize the gaps. Among them, deer-unpalatable *Symplocos prunifolia* and *Triadica sebifera* may be more likely to survive or mature under foraging pressure of deer, while deer-palatable species such as *Callicarpa mollis* and *Zanthoxylum ailanthoides* may be unlikely to grow under such pressure. This may change the species composition in regenerating stands.

In the future, deer-unpalatable species such as *Symplocos prunifolia* and *Triadica sebifera* may dominate the understory within the gaps that are created, expanded, or affected by the mass mortality of oak trees rather than the current dominant species such as *Eurya japonica* and *Quercus glauca*, while these current dominant species may retain their dominance within unaffected areas owing to their abundance and shade-tolerance under the current magnitude of deer pressure.

ACKNOWLEDGEMENTS

I thank Dr. K. Hirayama (Kyôto Prefectural University) for cooperating with the fieldwork and the Kyôto-Ôsaka District Forest Office for supporting this work. I also thank Dr. H. Iijima (Yamanashi Forest Research Institute) for reading a previous version of the manuscript and for comments. Computational calculations were conducted on the high performance cluster computing system of AFFIT, Ministry of Agriculture, Forestry and Fisheries, Japan.

REFERENCES

- Brooks, S. P. and Gelman, A. (1998). General methods for monitoring convergence of iterative simulations. *Journal of Computational and Graphical Statistics*, 7:434–455.
- Cai, F. (2000). A study on the structure and dynamics of *Cyclobalanopsis glauca* population at hills around West Lake in Hangzhou. *Scientia Silvae Sinicae*, 36:67–72. (in Chinese with English summary).
- Chen, G., Kéry, M., Plattner, M., Ma, K., and Gardner, B. (2013). Imperfect detection is the rule rather than the exception in plant distribution studies. *Journal of Ecology*, 101:183–191.
- Chen, G., Kéry, M., Zhang, J., and Ma, K. (2009). Factors affecting detection probability in plant distribution studies. *Journal of Ecology*, 97:1383–1389.
- Côté, S. D., Rooney, T. P., Tremblay, J.-P., Dussault, C., and Waller, D. M. (2004). Ecological impacts of deer overabundance. *Annual Review of Ecology and Systematics*, 35:113–147.
- Dorazio, R. M., Royle, J. A., Söderström, B., and Glimskär, A. (2006). Estimating species richness and accumulation by modeling species occurrence and detectability. *Ecology*, 87:842–854.
- Fujii, T. (1994). Stand development process and flowering patterns of a *Symplocos prunifolia* population. *Humans and Nature*, 3:79–83. (in Japanese with English summary).
- Gelman, A., Carlin, J. B., Stern, H. S., Dunson, D. B., Vehtari, A., and Rubin, D. B. (2013). *Bayesian data analysis*. CRC Press, Boca Raton, 3rd edition.
- Gelman, A. and Rubin, D. B. (1992). Inference from iterative simulation using multiple sequences. *Statistical Science*, 7:457–472.

- 417 Hashimoto, Y. and Fujiki, D. (2014). List of food plants and unpalatable plants of sika
418 deer (*Cervus nippon*) in Japan. *Humans and Nature*, 25:133–160. (in Japanese).
- 419 Ida, H. and Takahashi, S. (2010). Mass mortality of oak trees had already occurred at
420 the Edo period. *Journal of Japanese Forest Society*, 92:115–119. (in Japanese with
421 English summary).
- 422 Iijima, H. and Nagaike, T. (2015). Appropriate vegetation indices for measuring the
423 impacts of deer on forest ecosystems. *Ecological Indicators*, 48:457–463.
- 424 Itô, H. (2007). Twelve-years change of a broad-leaved secondary forest in Ginkakuji-san
425 National Forest. *Bulletin of FFPRI*, 6:93–100. (in Japanese with English summary).
- 426 Itô, H. (2009). A comparison of seedling emergence and survival between *Quercus*
427 *glauca* and *Symplocos prunifolia*. *Journal of Forest Research*, 14:245–250.
- 428 Itô, H. (2015). Effects of sika deer (*Cervus nippon*) on the dynamics of a broadleaved
429 secondary forest after mass mortality of oak trees. *Journal of Japanese Forest Society*,
430 97:304–308. (in Japanese with English summary).
- 431 Itô, H., Igarashi, T., and Kinuura, H. (2009). Changes in stand structure after the mass
432 mortality of oak trees in Keihoku area, Kyoto City, Japan. *Journal of Japanese*
433 *Forestry Society*, 91:15–20. (in Japanese with English summary).
- 434 Itô, H., Kinuura, H., and Oku, H. (2011). Stand structure of a broadleaved forest with
435 *Sasa*-type floor after mass mortality by oak wilt disease. *Journal of Japanese Forestry*
436 *Society*, 93:84–87. (in Japanese with English summary).
- 437 Kabaya, H. (1988). Changes of undergrowth of a broadleaved evergreen forest in relation
438 to grazing effects by sika deer. *Bulletin of the Tokyo University Forests*, 78:67–82. (in
439 Japanese with English summary).
- 440 Kéry, M. (2004). Extinction rate estimates for plant populations in revisitation studies:
441 importance of detectability. *Conservation Biology*, 18:570–574.
- 442 Kéry, M. and Schaub, M. (2012). *Bayesian population analysis using WinBUGS: a*
443 *hierarchical perspective*. Academic Press, Waltham.
- 444 Kinuura, H. and Kobayashi, M. (2006). Death of *Quercus crispula* by inoculation with
445 adult *Platypus quercivorus* (Coleoptera: Platypodidae). *Applied Entomology and*
446 *Zoology*, 41:123–128.
- 447 Kobayashi, M. and Ueda, A. (2005). Wilt disease of Fagaceae trees caused by *Platypus*
448 *quercivorus* (murayama) (Coleoptera: Platypodidae) and the associated fungus: aim
449 is to clarify the damage factor. *Journal of Japanese Forest Society*, 87:435–450. (in
450 Japanese with English summary).
- 451 Kubono, T. and Ito, S.-i. (2002). *Raffaelea quercivora* sp. nov. associated with mass mor-
452 tality of Japanese oak, and the ambrosia beetle (*Platypus quercivorus*). *Mycoscience*,
453 43:255–260.
- 454 Kuroda, K., Osumi, K., and Oku, H. (2012). Reestablishing the health of secondary
455 forests “Satoyama” endangered by Japanese oak wilt: A preliminary report. *Journal*
456 *of Agricultural Extension and Rural Development*, 4:192–198.
- 457 Lunn, D., Spiegelhalter, D., Thomas, A., and Best, N. (2009). The bugs project:
458 Evolution, critique, and future directions. *Statistics in Medicine*, 28:3049–3067.
- 459 Maesako, Y., Nanami, S., and Kanzaki, M. (2007). Spatial distribution of two invasive
460 alien species, *Podocarpus nagi* and *Sapium sebiferum*, spreading in a warm-temperate
461 evergreen forest of the Kasugayama Forest Reserve, Japan. *Vegetation Science*,
462 24:103–112.

- 463 Müllerová, J., Hédli, R., and Szabó, P. (2015). Coppice abandonment and its implications
464 for species diversity in forest vegetation. *Forest Ecology and Management*, 343:88–
465 100.
- 466 Murata, M., Matsuda, Y., Yamada, T., and Ito, S. (2009). Differential spread of dis-
467 coloured and non-conductive sapwood among four Fagaceae species inoculated with
468 *Raffaelea quercivora*. *Forest Pathology*, 39:192–199.
- 469 Murata, M., Yamada, T., and Ito, S.-i. (2005). Changes in water status in seedlings of
470 six species in the Fagaceae after inoculation with *Raffaelea quercivora* Kubono et
471 Shin-Ito. *Journal of Forest Research*, 10:251–255.
- 472 Naka, K. (1982). Community dynamics of evergreen broadleaf forests in southwestern
473 Japan. I. Wind damaged trees and canopy gaps in an evergreen oak forest. *Botanical*
474 *Magazine Tokyo*, 95:385–399.
- 475 Nakajima, H. and Ishida, M. (2014). Decline of *Quercus crispula* in abandoned coppice
476 forests caused by secondary succession and Japanese oak wilt disease: Stand dynamics
477 over twenty years. *Forest Ecology and Management*, 334:18–27.
- 478 Nakajima, M. (1929). The feeding experiments of Japanese deer in the university forest
479 in Chiba prefecture. *Bulletin of the Tokyo University Forests*, 8:95–114. (in Japanese).
- 480 Obora, T., Watanabe, H., and Yokoi, S. (2013). The effect of Sika deer browsing damage
481 on regeneration in mass mortality area of Japanese oak trees. *Journal of Japanese*
482 *Society of Revegetation Technology*, 39:260–263. (In Japanese).
- 483 Rackham, O. (2008). Ancient woodlands: Modern threats. *New Phytologist*, 180:571–
484 586.
- 485 Rooney, T. P. (2001). Deer impacts on forest ecosystems: a North American perspective.
486 *Forestry*, 74:202–208.
- 487 Saito, S. and Shibata, M. (2012). The forest structure and tree death rate of forest stands
488 damaged by Japanese oak wilt in Yamagata prefecture. *Journal of Japanese Forest*
489 *Society*, 94:223–228. (in Japanese with English summary).
- 490 Shibata, M. and Nakashizuka, T. (1995). Seed and seedling demography of four co-
491 occurring *Carpinus* species in a temperate deciduous forest. *Ecology*, 76:1099–1108.
- 492 Shimoda, K., Kimura, K., Kanzaki, M., and Yoda, K. (1994). The regeneration of
493 pioneer tree species under browsing pressure of Sika deer in an evergreen oak forest.
494 *Ecological Research*, 9:85–92.
- 495 Suzuki, M. (2013). Succession of abandoned coppice woodlands weakens tolerance of
496 ground-layer vegetation to ungulate herbivory: A test involving a field experiment.
497 *Forest Ecology and Management*, 289:318–324.
- 498 Svátek, M. and Matula, R. (2015). Fine-scale spatial patterns in oak sprouting and
499 mortality in a newly restored coppice. *Forest Ecology and Management*, 348:117–123.
- 500 Takatsuki, S. (1989). Effects of deer on plants and plant communities. *Japanese Journal*
501 *of Ecology*, 39:67–80. (in Japanese with English summary).
- 502 Takatsuki, S. (2009). Effects of sika deer on vegetation in Japan: A review. *Biological*
503 *Conservation*, 142:1922–1929.