

Ant-mediated seed dispersal in a warmed world

Climate change affects communities both directly and indirectly via changes in interspecific interactions. One such interaction that may be altered under climate change is the ant-plant seed dispersal mutualism common in deciduous forests of the eastern US. As climatic warming alters the abundance and activity levels of ants, the potential exists for shifts in rates of ant-mediated seed removal. We used an experimental temperature manipulation at two sites in the eastern US (Harvard Forest in Massachusetts and Duke Forest in North Carolina) to examine the potential impacts of climatic warming on overall rates of seed dispersal (using *Asarum canadense* seeds) as well as species-specific rates of seed dispersal at the Duke Forest site. We also examined the relationship between ant critical thermal maxima (CT_{max}) and the mean seed removal temperature for each ant species. We found that seed removal rates did not change as a result of experimental warming at either study site, nor were there any changes in species-specific rates of seed dispersal. There was, however, a positive relationship between CT_{max} and mean seed removal temperature, whereby species with higher CT_{max} removed more seeds at hotter temperatures. The temperature at which seeds were removed was influenced by experimental warming as well as diurnal and day-to-day fluctuations in temperature. Taken together, our results suggest that while temperature may play a role in regulating seed removal by ants, ant plant seed-dispersal mutualisms may be more robust to climate change than currently assumed.

Ant-mediated seed dispersal in a warmed world

Katharine L. Stuble^{1†}

Courtney M. Patterson¹

Mariano A. Rodriguez-Cabal^{2§}

Relena R. Ribbons¹

Robert R. Dunn³

Nathan J. Sanders^{1‡}

¹ Dept. of Ecology & Evolutionary Biology, University of Tennessee, Knoxville, TN 37996, USA

² Dept. of Zoology, University of British Columbia, Vancouver, B.C. V6T1Z4, Canada

³ Dept. of Biology, North Carolina State University, Raleigh, NC 27695

[†] Current address / Corresponding author: Oklahoma Biological Survey, 111 E Chesapeake St, Norman, OK, 73019 USA

[§]Current address: INIBIOMA, CONICET, Universidad Nacional del Comahue, Bariloche, Río Negro – Argentina, 8400

[‡] Current address: Center for Macroecology, Evolution and Climate, Natural History Museum of Denmark, University of Copenhagen, DK-2100 Copenhagen, Denmark

Abstract

Climate change affects communities both directly and indirectly via changes in interspecific interactions. One such interaction that may be altered under climate change is the ant-plant seed dispersal mutualism common in deciduous forests of the eastern US. As climatic warming alters the abundance and activity levels of ants, the potential exists for shifts in rates of ant-mediated seed removal. We used an experimental temperature manipulation at two sites in the eastern US (Harvard Forest in Massachusetts and Duke Forest in North Carolina) to examine the potential impacts of climatic warming on overall rates of seed dispersal (using *Asarum canadense* seeds) as well as species-specific rates of seed dispersal at the Duke Forest site. We also examined the relationship between ant critical thermal maxima (CT_{max}) and the mean seed removal temperature for each ant species. We found that seed removal rates did not change as a result of experimental warming at either study site, nor were there any changes in species-specific rates of seed dispersal. There was, however, a positive relationship between CT_{max} and mean seed removal temperature, whereby species with higher CT_{max} removed more seeds at hotter temperatures. The temperature at which seeds were removed was influenced by experimental warming as well as diurnal and day-to-day fluctuations in temperature. Taken together, our results suggest that while temperature may play a role in regulating seed removal by ants, ant plant seed-dispersal mutualisms may be more robust to climate change than currently assumed.

Keywords: ants, climate change, myrmecochory, seed dispersal, warming

Introduction

Understanding how organisms will respond to ongoing changes in climate, leading to subsequent changes in key ecological processes, is essential in order to predict the structure and function of ecosystems in the future. For example, the alteration of interspecific interactions is one important mechanism by which climate change may ultimately alter the structure and function of ecosystems ([Tylianakis et al. 2008](#), [van der Putten et al. 2010](#), [Walther 2010](#), [Harley 2011](#)). The majority of studies on the effects of climate change on interspecific interactions have focused on negative interactions, such as competition ([Suttle et al. 2007](#)), predator-prey interactions ([Rothley and Dutton 2006](#), [Barton and Schmitz 2009](#), [Harley 2011](#)), and herbivory ([Bale et al. 2002](#)). Though rare relative to studies that focus on physiological responses ([Norby and Luo 2004](#), [Dillon et al. 2010](#)) or aggregate community-level responses ([Klein et al. 2004](#), [Villalpando et al. 2009](#), [Fitzpatrick et al. 2011](#), [Sheldon et al. 2011](#)), empirical studies of the effects of climate on interactions among species have found that climate change can alter the nature and outcomes of interspecific interactions through a variety of mechanisms such as altered abundance and fitness levels of key species ([Suttle et al. 2007](#)), shifts in phenology ([Both et al. 2009](#)), and species range shifts ([Harley 2011](#)). Each of these mechanisms can decouple interspecific interactions by altering the frequency and intensity of interactions among species.

As with most studies in ecology, work on the effects of climate change on positive interactions is lacking, even though mutualisms play fundamental roles in structuring communities and ecosystems ([Callaway 1995](#), [Stachowicz 2001](#)). Mutualisms including plant-pollinator interactions and mycorrhizal interactions have been altered by climate change ([Parrent et al. 2006](#), [Mommott et al. 2007](#), [Hoover et al. 2012](#)). However, the influence of climate change on other types of positive interactions is not as well studied.

One such mutualism that may be altered by climate change is the ant-plant seed dispersal mutualism, which includes hundreds of ant species and thousands of plant species around the world (Beattie and Hughes 2002; Gove et al 2007; Rico-Gray and Oliviera 2007, Lengyel et al. 2010). In deciduous forests of the United States approximately thirty percent of understory herb species might be ant-dispersed (Beattie and Culver 1981), and a proposed keystone seed-dispersing ant species, *Aphaenogaster rudis*, is responsible for upwards of 90% of ant-mediated seed dispersal (Ness et al. 2009; Zelikova et al. 2008). Such specialization in interactions can make interaction networks more vulnerable to disruption as a result of low levels of functional redundancy within a system ([Aizen et al. 2012](#)). Myrmecorous plant species that rely on a single ant species for seed dispersal may be at increased risk for disruption by ongoing climatic change ([Pelini et al. 2011a](#), [Warren et al. 2011](#)) if that ant species is negatively affected by warming. Inversely, systems in which multiple species are responsible for removing seeds may prove to be more resistant to disruptions because of functional redundancy in the system ([Peterson et al. 1998](#)). However, despite the importance and ubiquity of ant-plant seed-dispersal mutualisms in ecosystems around the world and the importance of temperature in regulating ant foraging, experiments examining the consequences of climatic change on plant-seed dispersal mutualisms are rare (but see Pelini et al. 2011a).

To examine the potential for climate change to alter seed dispersal mutualisms, we experimentally manipulated temperature at two sites in the eastern United States (North Carolina and Massachusetts) and examined overall rates of seed removal as a function of temperature. At the North Carolina site, we also examined species-specific rates of seed dispersal. We test two *a priori* predictions:

Prediction 1: Rates of seed removal by ants would decrease as a result of experimental warming at the southern site where species, including *A. rudis*, are closer to their critical thermal maxima (CT_{max}) ([Deutsch et al. 2008](#), [Diamond et al. 2012a](#), [Diamond et al. 2012b](#), [Huey et al. 2012](#)), a

pattern previously documented in this study system ([Diamond et al. 2012a](#)). Rates of seed removal can be expected to increase at the northern site, where most species, including *A. rudis*, are well below their CT_{max} ([Diamond et al. 2012a](#)) and perhaps even below their thermal optima. *Prediction 2*: Species with higher CT_{max} would remove seeds more readily under warmer conditions as compared to species with lower CT_{max} as these species are more tolerant of higher temperatures and have been found to be more active as temperatures increase (Stuble et al. 2013).

Methods

Site description

This experiment was conducted at Duke Forest (southern site) in North Carolina and Harvard Forest (northern site) in Massachusetts, USA, in order to examine the potential impacts of climate change on seed dispersal mutualisms near the southern extent and northern extent of the ranges of several ant species. The Duke Forest site consists of a closed-canopy oak-hickory (*Quercus* spp., *Carya* spp.) forest with a mean annual temperature of 15.5°C and approximately 1140 mm of precipitation annually. The Harvard Forest site is in a closed-canopy oak-maple (*Quercus* spp.-*Acer* spp.) forest with a mean annual temperature of 7.1°C and 1066 mm of precipitation a year. Though not present in the chambers, several myrmecochorous plant species are common in these forests. These species include *Asarum canadense*, *Hexastylis arifolia*, *Trillium* spp., *Viola rotundifolia*, and *Sanguinaria canadensis*, among others. Seeds of these plant species are typically dispersed by ants in the spring (Thompson 1981).

The two sites share ~30 ant species, with the North Carolina site near the southern range edge and the Massachusetts site near the northern range edge for many of these species ([Pelini et al. 2011b](#)). The most abundant ant species at both sites is the proposed keystone seed disperser *A.*

rudis (Ness et al. 2009) (or at least a species in the *A. rudis* complex). *Crematogaster lineolata*, a behaviorally aggressive species, can be abundant at the southern site in warmer months while *Formica subsericea* and *Camponotus pennsylvanicus* (also behaviorally aggressive) are the two next most abundant ant species at the Harvard Forest site (Stuble et al. 2013b). For the purposes of this study we are combining *A. rudis*, *A. picea*, and *A. carolinensis* into the *A. rudis* complex due to the difficulty of identifying these closely related species in the field.

At each site, there are twelve experimental open-top warming chambers (Fig. 1). Each chamber is 5 m in diameter and 1.2 m tall with a 2-3 cm gap at the bottom to allow ants and other organisms to move in and out. The chambers are approximately 1000 ant body lengths in diameter. The chambers contain about 1 colony per square m, and the average foraging distance of an *A. rudis* colony is ~70cm. Moreover, during the summer of 2012, we watched 72 *A. rudis* workers visiting baits and returning to their colonies. Out of those 72 observations, only 1 worker visited a bait in the chamber and returned to a nest outside the chamber (L. Nichols, unpublished data). So, most of the activity we see is from ants in the chambers. Nine chambers at each site are warmed from 1.5°C to 5.5°C in 0.5°C steps using air warmed by hydronic radiators, while the three control chambers blow air at ambient temperatures into the plots (see Pelini et al. (2011b) for a detailed description of the chambers). Warming treatments have been maintained continuously since January of 2010 and have been successful at maintaining the targeted temperature increases. For 2011, a significantly positive relationship between the target temperature increase and actual temperature increase was maintained ($p < 0.01$, $R^2 = 0.99$). Significant shifts in levels of ant foraging activity at food baits have been documented in the chambers across the temperature treatments, with more thermally tolerant species exhibiting higher levels of foraging activity in warmer chambers than species with lower thermal tolerances (Stuble et al. 2013a). Those results suggest that temperature does mediate foraging behavior. Such a result, that environmental context can mediate foraging behavior, is in line with previous

work in this system (Pelini et al. 2011) and others ([Cerdá et al. 1997](#), [Sanders and Gordon 2000](#), [2003](#), [Gibb and Parr 2010](#)).

Seed removal

To assess the impact of temperature on rates of seed dispersal, we haphazardly positioned one seed cache in each of the 12 chambers at Duke Forest and Harvard Forest. Each cache contained 20 seeds of the myrmecochorous species *Asarum canadense*. The range of *Asarum canadense* extends from New Brunswick, Canada to North Carolina in the southern US ([Cain and Damman 1997](#)). Seeds of *A. canadense* are similar in mass to many other myrmecochorous plant species ([Michaels et al. 1988](#)). Seeds used in the trials at Duke Forest were collected at North Carolina State University's Schenck Forest in Raleigh, North Carolina on May 11, 2011 and those used in the Harvard Forest trials were collected from Mt. Toby in Massachusetts on June 8, 2011. We covered each seed cache with a mesh cage to allow ants to access the seeds while preventing access by rodents. Caches were left out for one hour, after which time the number of seeds remaining in the cache was counted and any remaining seeds were removed from the chamber. Though observing seed removal for an hour limits our ability to account for the fate of all seeds, using this standard timeframe allowed us to compare relative rates of seed removal across treatments. A total of ten trials were conducted at Duke Forest between May 12 and May 25, 2011, with five trials conducted during the day (between 0900 and 1900) and five during the night (between 2100 and 0500). Another five trials were conducted at Harvard Forest between June 16 and June 30, 2011; three during the day and two at night. These dates corresponded with the time periods during which the seeds were naturally released at each site, as opposed to conducting this experiment in the hottest part of the year when the impacts of warming might be expected to be greater, but when any results might be less ecologically relevant.

We calculated the average seed dispersal rate (number of seeds removed in an hour) for each chamber at each site. We used ANCOVA to examine differences in seed dispersal rates as a function of temperature treatment (which we refer to as $\Delta^{\circ}\text{C}$, included as a continuous variable) and site. The number of seeds removed per hour was square root transformed to meet assumptions of normality. All statistics were performed in SAS, version 9.2.

To determine the ant species responsible for removing the seeds, we continuously observed caches of 10 *A. canadense* seeds within the chambers at Duke Forest for one hour, or until all seeds were removed. Four seed removal observations were conducted in each chamber: two during the day and two during the night. Nighttime observations were conducted using red lights, which is typical in studies of ant behavior at night ([Hodgson 1955](#), [Narendra et al. 2010](#)). We recorded the identity of the ant species removing the seeds. When possible, we also followed the seed back to the nest (or under leaf litter in some cases) and noted the distance it had been moved. At the beginning of each observation, we took four ground surface temperature measurements using a handheld infrared thermometer (Raytek® Raynger ST, $\pm 1^{\circ}\text{C}$), one at each corner of the seed cache, which were averaged together. These temperature readings provided us with estimates of ground-surface temperature conditions in the immediate vicinity of the seeds. Ground-surface temperature has been shown to be an important driver of foraging activity in ants ([Whitford and Ettershank 1975](#), [Crist and MacMahon 1991](#)). We calculated the percentage of seeds removed by each species overall, as well as separately for day and night. We also calculated the mean number of seeds removed by each species in each chamber across all trials.

We used linear regressions to examine differences in seed dispersal rates across temperature treatments for each ant species. (We examined several polynomial regressions, but found none of them to be a better fit than simple linear regressions.) Mean numbers of seeds removed were log transformed to meet assumptions of normality for *A. rudis* and *C. lineolata*.

Finally, we calculated the average ground surface temperature (based on temperatures collected with the infrared thermometer) at which each species removed seeds across all treatments and times. We then examined the relationship between the average temperature at which a species removed seeds and the CT_{max} of that species (as calculated by Diamond et al., 2012) across all species observed removing seeds in the system. *Aphaenogaster lamellidens* was excluded from this analysis as it was only observed removing seeds from two seed caches and was an outlier (as indicated by a plot of residuals by predicted values).

Results and Discussion

Seed removal rate did not depend on temperature treatment and did not vary between sites ($F_{2,21} = 0.93$, $p = 0.41$; Fig. 2). This is despite the fact that most of the foragers observed in this study were from colonies within the experimental chambers. At the southern site, where seed dispersal observations were conducted, the mean seed removal distance was 51 cm, and only 2% of observed seeds were removed more than 2 m. The lack of response to experimental warming contrasts with the prediction that, based on the thermal limits of *A. rudis* and its disproportionate role in seed dispersal, seed dispersal rate should decline with increasing temperatures. Regardless of temperature treatment or site, ants removed ~ 23% of seeds per hour (an average of 4.6 seeds out of 20). At the southern site, we observed seven ant species removing seeds across a range of ground surface temperatures from 17°C to 30°C (Table 1). *Aphaenogaster rudis* was the most common seed disperser, removing approximately 45.5% of seeds (Table 2). However, there was no relationship between the rate of seed dispersal by *A. rudis* and temperature treatment (Table 1). With the exception of *C. lineolata*, which showed a marginally significant increase of approximately 0.1 seeds removed per degree of warming, seed removal did not vary systematically with temperature treatment for any ant species (Table 1). This is despite

previously observed shifts in foraging under experimentally warmed conditions ([Pelini et al. 2011a](#), [Stuble et al. 2013a](#)). Pelini et al. (2011) found an approximately 50% decrease in several types of foraging, including seed removal, as a result of 1°C of warming at the southern site, though no change was observed at the northern site. Using the same warming chambers as in this study, Stuble et al. (2013) found species-specific shifts in foraging activity as a result of experimental warming consistent with the thermal tolerances of the foraging species. Further, ant community composition shifts in response to experimental warming, demonstrating the importance of temperature in regulating the ant community ([Diamond et al. 2012a](#)). Despite this, experimental warming apparently does not affect the aspects of the seed-dispersal mutualisms we studied in this system. This begs the question - why isn't this seed-dispersal mutualism disrupted by experimental warming?

One possible answer is that foraging behavior by *Aphaenogaster rudis* may be more tolerant to experimental warming than previously thought. *A. rudis* is a keystone mutualist in this and other systems, responsible for the majority of ant-mediated seed dispersal ([Zelikova et al. 2008](#), [Ness et al. 2009](#), [Canner et al. 2012](#)). The abundance and activity of *A. rudis* declines with elevation (i.e., lower temperatures) at biogeographic scales ([Zelikova et al. 2008](#)). Additionally, the relatively low thermal tolerance of this species accurately predicts its activity relative to other species (Stuble et al. in press). However, when exposed to experimental warming, the abundance ([Pelini et al. 2011a](#)) and foraging activity (Stuble et al. 2013) of *A. rudis* apparently do not decline at either study site. Importantly, the average foraging distance of *Aphaenogaster* spp. was ~70cm at the study site (L. Nichols, unpublished data). Further, based on the proportion of these ants observed foraging into the chambers from outside in observations, we'd predict that only about one of the eighty seeds observed being removed by *A. rudis* was likely to have been removed by a worker originating outside of the chambers. Thus, it is not likely the case that ants

are coming to baits from nests that are outside the chambers. However, even in cases in which individual workers do forage at the experimental baits from colonies outside of the chambers, these individuals are still exposed to the experimental temperature conditions while discovering, foraging at, and recruiting to the seed caches. These results suggest that temperature does not substantially alter this foraging behavior. The apparent tolerance of the foraging activity of this important seed dispersing species to warming may play a major role in promoting the stability of ant-plant seed dispersal in light of global change.

It is important to note that six ant species other than *A. rudis* were observed removing seeds in this study, and they removed > 50% of the seeds. This runs counter to several studies suggesting seed dispersal mutualisms may be highly specialized ([Gove et al. 2007](#), [Ness et al. 2009](#)). Both the foraging activity and abundances of several of these species, including *C. lineolata* and *Formica pallidefulva*, shift with warming ([Pelini et al. 2011a](#), [Diamond et al. 2012a](#), [Stuble et al. 2013a](#)), resulting in an altered community of foragers (Diamond et al. 2012a). By having multiple ant species interacting with myrmecochorous plants, this ant-plant seed dispersal mutualism may be relatively resistant to the effects of warming as some ant species increase in activity and abundance while others decline in abundance with temperature. Previous work on ant foraging and community composition as a result of the experimental warming at these sites suggests quite a bit of variability in species-specific responses to warming which may, in fact, play a role in moderating the effects of climatic warming ([Stuble et al. 2013a](#)).

In addition to the apparent (and of course relative) resistance of the foraging of *A. rudis* to warming, along with the diversity of ants engaging in this mutualism, another factor possibly strengthening the resistance of the ant-plant seed dispersal mutualism may be the timing of ant-mediated seed dispersal within deciduous forests of the eastern US. Ant-dispersed seeds in these forests, including those of *Asarum canadense*, are primarily dispersed in the spring ([Thompson 1981](#)). Temperatures in May in North Carolina and June in Massachusetts at the study sites are

far from the critical thermal maxima of ant species in the system. For example, the critical thermal maximum for *A. rudis* is 38°C and 40°C for populations at the northern and southern sites, respectively ([Diamond et al. 2012b](#)), as opposed to the mean environmental temperatures during the sampling period, which were 20°C at the northern site and 22°C at the southern site. The thermal buffer between CT_{max} and the environmental temperature during the time of year when seeds are dispersed may confer some degree of tolerance on this mutualism. Pelini ([2011a](#)) found that rates of seed removal decreased in a passive experimental warming at the same two sites, despite achieving warming of only 0.3°C above ambient. However, the seed removal trials in Pelini et al.'s experiment were conducted mostly in August when ambient environmental temperatures are hotter than those experienced in the present study. We suggest that the proximity of ants to their upper thermal limits in August may have driven the effects of warming observed in the Pelini et al. ([2011a](#)) study while seed dispersal occurring in the spring when our study was conducted may be less likely to be detrimentally impacted by warming. However, this protection assumes that the peak of *A. rudis* activity and seed set coincide. Phenological shifts in plant reproduction caused by ongoing warming ([Price and Waser 1998](#), [Dahlgren et al. 2007](#), [Inouye 2008](#), [Liu et al. 2011](#), [Wolkovich et al. 2012](#)) have the potential to result in seeds appearing before ants become active ([Warren et al. 2011](#)). Warren et al. (2011) suggest that while both seed release by plants and onset of foraging in ants seem to be driven by temperature, variability in activation temperatures among ant species may result in situations in which early seeding plant species may become decoupled from their foragers in some areas.

Despite the apparent tolerance of the ant-plant seed dispersal mutualism to experimental warming in this study, there was a significant relationship between the ground-surface temperature at which a species removed seeds and the critical thermal maximum of that species ($F_{1,4} = 7.35$, $p = 0.05$, $R^2 = 0.65$, Fig. 3). That is, those species with high thermal tolerances were most active under the warmest temperatures. The positive relationship between CT_{max} and seed

removal temperature suggests that while chronic experimental warming may not affect rates of seed dispersal, temperature does relate to rates of seed removal. This finding incorporates both temperature variability associated with the temperature treatments as well as daily temperature variability and complements other studies that have shown physiological tolerance to be an important predictor of ant activity (Diamond et al, in press).

One important caveat to our study (and to most studies of ant-seed dispersal mutualisms) is that we do not know the ultimate fate of the seeds once they were removed by the ants. It is possible that warming could still alter the dynamics of plant populations by altering rates of germination and seedling survival post-germination ([De Frenne et al. 2012](#)), even in cases in which seed dispersal remains unaffected as temperatures increase. Additionally, some species, including *C. lineolata*, dispersed seeds very short distances (only a few centimeters) while other species, such as *C. castaneus*, often carried seeds several meters. Even slight shifts in dispersal rates among these species may alter plant population dynamics if dispersal distances differ substantially among species. Finally, by observing seed caches for an hour, we fail to collect data on the fate of seeds not removed in that time frame. However, a considerable proportion of the seeds that will be removed by ants are, in fact, removed soon after release from the parent plant ([Turnbill and Culver 1983](#), [Beaumont et al. 2013](#)) and seeds not dispersed by ants are at risk of predation by rodents ([Heithaus 1981](#)). Additionally, as in most studies of ant-seed interactions, we focus on seeds of only a single plant species. Including more seeds from more species would be ideal, but would have been beyond what was possible in this (or in most) studies.

It has become axiomatic that interactions among species are being affected by ongoing climatic change. However, in this study, we found no reduction in overall rates of seed removal as a result of experimental warming. Moreover, there were no differences in species-specific seed removal rates at the southern site. We suggest that the ant-plant seed dispersal mutualism may be

resistant, at least in part, to climatic warming as a result of the diversity of ants active in this mutualism as well as the seasonal timing of this mutualism in the spring when most ant species are far from their upper thermal limits. Importantly, however, if phenological mismatches arise, or if the fate of seeds after dispersal is altered, the consequences of warming on plant populations and communities could emerge in unexpected ways.

Acknowledgements

We thank M. Burt and I. Del Toro for help in the field. A. Ellison and N. Gotelli were integral in developing and improving this project.

Work Cited

- Aizen, M. A., M. Sabatino, and J. M. Tylianakis. 2012. Specialization and rarity predict nonrandom loss of interactions from mutualist networks. *Science* **335**:1486-1489.
- Bale, J. S., G. J. Masters, I. D. Hodkinson, C. Awmack, T. M. Bezemer, V. K. Brown, J. Butterfield, A. Buse, J. C. Coulson, J. Farrar, J. E. G. Good, R. Harrington, S. Hartley, T. H. Jones, R. L. Lindroth, M. Press, C., I. Symrnioudis, A. D. Watt, and J. B. Whittaker. 2002. Herbivory in global change research: direct effects of rising temperature on insect herbivores. *Global Change Biology* **8**:1-16.
- Barton, B. T., and O. J. Schmitz. 2009. Experimental warming transforms multiple predator effects in a grassland food web. *Ecology Letters* **12**:1317-1325.
- Beaumont, K. P., D. A. Mackay, and M. A. Whalen. 2013. Multiphase myrmecochory: the roles of different ant species and effects of fire. *Oecologia* **172**:791-803.
- Cain, M. L., and H. Damman. 1997. Clonal growth and ramet performance in the woodland herb, *Asarum canadense*. *Journal of Ecology* **85**:883-897.
- Callaway, R. M. 1995. Positive interactions among plants. *Botanical Review* **61**:306-349.
- Canner, J. E., R. R. Dunn, I. Giladi, and K. Gross. 2012. Redispersal of seeds by a keystone ant augments the spread of common wildflowers. *Acta Oecologia* **40**:31-39.
- Cerdá, X., J. Retana, and S. Cros. 1997. Thermal disruption of transitive hierarchies in Mediterranean ant communities. *Journal of Animal Ecology* **66**:363-374.
- Crist, T. O., and J. A. MacMahon. 1991. Foraging patterns of *Pogonomyrmex occidentalis* (Hymenoptera: Formicidae) in a shrub-steppe ecosystem: the roles of temperature, trunk trails, and seed resources. *Environmental Entomology* **20**:265-275.
- Dahlgren, J. P., H. von Zeipel, and J. Ehrlén. 2007. Variation in vegetative and flowering phenology in a forest herb caused by environmental heterogeneity. *American Journal of Botany* **94**:1570-1576.
- De Frenne, P., B. J. Graae, J. Brunet, A. Shevtsova, A. De Schrijver, O. Chabrierie, S. A. O. Cousins, G. Decocq, M. Diekmann, M. Hermy, T. Heinken, A. Kolb, C. Nilsson, S. Stanton, and K. Verheyen. 2012. The response of forest plant regeneration to temperature variation along a latitudinal gradient. *Annals of Botany* **109**:1037-1046.
- Deutsch, C. A., J. J. Tewksbury, R. B. Huey, K. S. Sheldon, C. K. Ghalambor, D. C. Haak, and P. R. Martin. 2008. Impacts of climate warming on terrestrial ectotherms across latitude. *Proceedings of the National Academy of Sciences of the USA - Biological Sciences* **105**:6668-6672.
- Diamond, S. E., L. M. Nichols, N. McCoy, C. Hirsch, S. L. Pelini, N. J. Sanders, A. M. Ellison, N. J. Gotelli, and R. R. Dunn. 2012a. A physiological trait-based approach to predicting the responses of species to experimental climate warming. *Ecology* **93**:2305-2312.

347 Diamond, S. E., D. M. Sorger, J. Hulcr, S. L. Pelini, I. Del Toro, C. Hirsch, E. Oberg, and R. R.
 348 Dunn. 2012b. Who likes it hot? A global analysis of the climatic, ecological, and
 349 evolutionary determinants of warming tolerance in ants. *Global Change Biology* **18**:448-
 350 456.

351 Dillon, M. E., G. Wang, and R. B. Huey. 2010. Global metabolic impacts of recent climate
 352 warming. *Nature* **467**:704-706.

353 Fitzpatrick, M. C., N. J. Sanders, S. Ferrier, J. T. Longino, M. D. Weiser, and R. R. Dunn. 2011.
 354 Forecasting the future of biodiversity: a test of single- and multi-species models for ants
 355 in North America. *Ecography* **34**:836-847.

356 Gibb, H., and C. L. Parr. 2010. How does habitat complexity affect foraging success? A test
 357 using functional measures on three continents. *Oecologia* **164**:1061-1073.

358 Gove, A. D., J. D. Majer, and R. R. Dunn. 2007. A keystone ant species promotes seed dispersal
 359 in a "diffuse" mutualism. *Oecologia* **153**:687-697.

360 Harley, C. D. G. 2011. Climate change, keystone predation, and biodiversity loss. *Nature*
 361 **334**:1124-1127.

362 Heithaus, E. R. 1981. Seed predation by rodents on three ant-dispersed plants. *Ecology* **62**:136-
 363 145.

364 Hodgson, E. S. 1955. An ecological study of the behavior of the leaf-cutting ant *Atta cephalotes*.
 365 *Ecology* **36**:293-304.

366 Hoover, S. E. R., J. J. Ladley, A. A. Shchepetkina, M. Tisch, S. P. Gieseg, and J. M. Tylianakis.
 367 2012. Warming, CO₂, and nitrogen deposition interactively affect a plant-pollinator
 368 mutualism. *Ecology Letters* **15**:227-234.

369 Huey, R. B., M. R. Kearney, A. Krockenberger, J. A. M. Holtum, M. Jess, and S. E. Williams.
 370 2012. Predicting organismal vulnerability to climate warming: roles of behaviour,
 371 physiology and adaptation. *Philosophical Transactions of the Royal Society B* **367**:1665-
 372 1679.

373 Inouye, D. W. 2008. Effects of climate change on phenology, frost damage, and floral
 374 abundance of montane wildflowers. *Ecology* **89**:353-362.

375 Klein, J. A., J. Harte, and X.-Q. Zhao. 2004. Experimental warming causes large and rapid
 376 species loss, dampened by simulated grazing on the Tibetan Plateau. *Ecology Letters*
 377 **7**:1170-1179.

378 Liu, Y., P. B. Reich, G. Li, and S. Sun. 2011. Shifting phenology and abundance under
 379 experimental warming alters trophic relationships and plant reproductive capacity.
 380 *Ecology* **92**:1201-1207.

381 Memmott, J., P. G. Craze, N. M. Waser, and M. V. Price. 2007. Global warming and the
 382 disruption of plant-pollinator interactions. *Ecology Letters* **10**:710-717.

383 Michaels, H. J., B. Benner, A. P. Hartgerink, T. D. Lee, and S. Rice. 1988. Seed size variation:
 384 magnitude, distribution, and ecological correlates. *Evolutionary Ecology* **2**:157-166.

385 Narendra, A., S. Reid, and J. M. Hemmi. 2010. The twilight zone: ambient light levels trigger
386 activity in primitive ants. *Proceedings of the Royal Society B: Biological Sciences*
387 **277**:1531-1538.

388 Ness, J. H., D. F. Morin, and I. Giladi. 2009. Uncommon specialization in a mutualism between a
389 temperate herbaceous plant guild and an ant: are *Aphaenogaster* ants keystone
390 mutualists? *Oikos* **118**:1793-1804.

391 Norby, R. J., and Y. Luo. 2004. Evaluating ecosystem responses to rising atmospheric CO₂ and
392 global warming in a multi-factor world. *New Phytologist* **162**:281-283.

393 Parrent, J. L., W. F. Morris, and R. Vilgalys. 2006. CO₂-enrichment and nutrient availability alter
394 ectomycorrhizal fungal communities. *Ecology* **87**:2278-2287.

395 Pelini, S. L., M. Boudreau, N. McCoy, A. M. Ellison, N. J. Gotelli, N. J. Sanders, and R. R. Dunn.
396 2011a. Effects of short-term warming on low and high latitude forest ant communities.
397 *Ecosphere* **2**:1-12.

398 Pelini, S. L., F. P. Bowles, A. M. Ellison, N. J. Gotelli, N. J. Sanders, and R. R. Dunn. 2011b.
399 Heating up the forest: open-top chamber warming manipulation of arthropod
400 communities at Harvard and Duke Forests. *Methods in Ecology and Evolution* **2**:534-
401 540.

402 Peterson, G., C. R. Allen, and C. S. Holling. 1998. Ecological resilience, biodiversity, and scale.
403 *Ecosystems* **1**:6-18.

404 Price, M. V., and N. M. Waser. 1998. Effects of experimental warming on plant reproductive
405 phenology in a subalpine meadow. *Ecology* **79**:1261-1271.

406 Rothley, K. D., and G. Dutton. 2006. Behavioral responses to environmental change alter direct
407 and indirect trait-mediated interactions. *Canadian Journal of Zoology* **84**:1053-1058.

408 Sanders, N. J., and D. M. Gordon. 2000. The effects of interspecific interactions on resource use
409 and behavior in a desert ant. *Oecologia* **125**:436-443.

410 Sanders, N. J., and D. M. Gordon. 2003. Resource-dependent interactions and the organization
411 of desert ant communities. *Ecology* **84**:1024-1031.

412 Sheldon, K. S., S. Yang, and J. J. Tewksbury. 2011. Climate change and community
413 disassembly: impacts of warming on tropical and temperate montane community
414 structure. *Ecology Letters* **14**:1191-1200.

415 Stachowicz, J. J. 2001. Mutualisms, facilitation, and the structure of ecological communities.
416 *BioScience* **51**:235-246.

417 Stuble, K. L., L. K. Kirkman, and C. R. Carroll. 2010. Are red imported fire ants facilitators of
418 native seed dispersal? . *Biological Invasions* **12**:1661-1669.

419 Stuble, K. L., S. L. Pelini, S. E. Diamond, D. A. Fowler, R. R. Dunn, and N. J. Sanders. 2013a.
420 Foraging by forest ants under experimental warming: a test at two sites. *Ecology and*
421 *Evolution* **3**:482-491.

Stuble, K. L., M. A. Rodriguez-Cabal, G. L. McCormick, I. Jurić, R. R. Dunn, and N. J. Sanders. 2013b. Tradeoffs, competition, and coexistence in eastern deciduous forest ant communities. *Oecologia* **171**:981-992.

Suttle, K. B., M. A. Thomsen, and M. E. Power. 2007. Species interactions reverse grassland responses to changing climate. *Science* **315**:640-642.

Thompson, J. N. 1981. Elaiosomes and fleshy fruits: phenology and selection pressures for ant-dispersed seeds. *American Naturalist* **117**:104-108.

Turnbill, C. L., and D. C. Culver. 1983. The timing of seed dispersal in *Viola nuttallii*: attraction of dispersers and avoidance of predators. *Oecologia* **59**:360-365.

Tylianakis, J. M., R. K. Didham, J. Bascompte, and D. A. Wardle. 2008. Global change and species interactions in terrestrial ecosystems. *Ecology Letters* **11**.

van der Putten, W. H., M. Macel, and M. E. Visser. 2010. Predicting species distribution and abundance responses to climate change: why it is essential to include biotic interactions across trophic levels. *Philosophical Transactions of the Royal Society B* **365**:2025-2034.

Villalpando, S. N., R. S. Williams, and R. J. Norby. 2009. Elevated air temperature alters an old-field insect community in a multifactor climate change experiment. *Global Change Biology* **15**:930-942.

Walther, G.-R. 2010. Community and ecosystem responses to recent climate change. *Philosophical Transactions of the Royal Society B* **365**:2019-2024.

Warren, R. J., V. Bahn, and M. A. Bradford. 2011. Temperature cues phenological synchrony in ant-mediated seed dispersal. *Global Change Biology* **17**:2444-2454.

Whitford, W. G., and G. Ettershank. 1975. Factors affecting foraging activity in Chihuahuan Desert Harvester ants. *Environmental Entomology* **4**:689-696.

Wolkovich, E. M., B. I. Cook, J. M. Allen, T. M. Crimmins, J. L. Betancourt, S. E. Travers, S. Pau, J. Regetz, T. J. Davies, N. J. B. Kraft, T. R. Ault, K. Bolmgren, S. J. Mazer, G. J. McCabe, B. J. McGill, C. Parmesan, N. Salamin, M. D. Schwartz, and E. E. Cleland. 2012. Warming experiments underpredict plant phenological responses to climate change. *Nature* **485**:494-497.

Zelikova, T. J., R. R. Dunn, and N. J. Sanders. 2008. Variation in seed dispersal along an elevational gradient in Great Smokey Mountains National Park. *Acta Oecologia* **34**:155-162.

Table 1 (on next page)

Table 1. Test statistics are from linear regressions examining the influence of temperature treatment on the number of seeds removed by each ant species (d.f. = 11 for all species).

Species	<i>F</i>	<i>p</i>
<i>Aphaenogaster lamellidens</i>	2.58	0.58
<i>Aphaenogaster rudis</i>	2.45	0.15
<i>Camponotus castaneus</i>	0.10	0.75
<i>Camponotus pennsylvanicus</i>	0.24	0.64
<i>Crematogaster lineolata</i>	4.14	0.07
<i>Formica pallidefulva</i>	0.34	0.58
<i>Formica subsericea</i>	0.02	0.90

Table 2_(on next page)

Table 2. Percentage of seeds removed by each species overall, during the day, and the night.

Species	Overall percent	Day percent	Night percent
<i>Aphaenogaster lamellidens</i>	8.5	17.2	0.0
<i>Aphaenogaster rudis</i>	45.5	48.3	42.7
<i>Camponotus castaneus</i>	26.7	0.0	52.8
<i>Camponotus pennsylvanicus</i>	2.8	2.3	3.4
<i>Crematogaster lineolata</i>	6.8	12.6	1.1
<i>Formica pallidefulva</i>	2.3	4.6	0.0
<i>Formica subsericea</i>	7.4	14.9	0.0

Figure 1

Figure 1. Warming chamber at Duke Forest.



Figure 2

Figure 2. Number of seeds removed in the course of an hour as a function of temperature treatment. Black dots represent Duke Forest and gray dots represent Harvard Forest.




Figure 3

Figure 3. Temperature at which seeds were removed as a function of a species' critical thermal maximum (CT_{max}).

