

# Neck biomechanics indicate that giant Transylvanian azhdarchid pterosaurs were short-necked arch predators

Darren Naish <sup>Corresp., 1</sup>, Mark P Witton <sup>2</sup>

<sup>1</sup> Ocean and Earth Science, University of Southampton, Southampton, UK

<sup>2</sup> School of Earth & Environmental Sciences, University of Portsmouth, Portsmouth, United Kingdom

Corresponding Author: Darren Naish  
Email address: eotyranus@gmail.com

Azhdarchid pterosaurs include the largest animals to ever take to the skies with some species exceeding 10 metres in wingspan and 220 kg in mass. Associated skeletons show that azhdarchids were long-necked, long-jawed predators that combined a wing planform suited for soaring with limb adaptations indicative of quadrupedal terrestrial foraging. The postcranial proportions of the group have been regarded as uniform overall, irrespective of their overall size, notwithstanding suggestions that minor variation may have been present. Here, we discuss a recently discovered giant azhdarchid neck vertebra referable to *Hatzegopteryx* from the Maastrichtian Sebeş Formation of the Transylvanian Basin, Romania, which shows how some azhdarchids departed markedly from conventional views on their proportions. This vertebra, which we consider a cervical VII, is 240 mm long as preserved and almost as wide. Among azhdarchid cervicals, it is remarkable for the thickness of its cortex (4-6 mm along its ventral wall) and robust proportions. By comparing its dimensions to other giant azhdarchid cervicals and to the more completely known necks of smaller taxa, we argue that *Hatzegopteryx* had a proportionally short, stocky neck highly resistant to torsion and compression. This specimen is one of several hinting at greater disparity within Azhdarchidae than previously considered, but is the first to demonstrate such proportional differences within giant taxa. On the assumption that other aspects of *Hatzegopteryx* functional anatomy were similar to those of other azhdarchids, and with reference to the absence of large terrestrial predators in the Maastrichtian of Transylvania, we suggest that this pterosaur played a dominant predatory role among the unusual palaeofauna of ancient Haţeg.

# Neck biomechanics indicate that giant Transylvanian azhdarchid pterosaurs were short-necked arch predators

Darren Naish<sup>1</sup> and Mark P. Witton<sup>2</sup>

<sup>1</sup>Ocean and Earth Science, University of Southampton, National Oceanography Centre, Southampton, University of Southampton, Southampton SO14 3ZH, UK. <sup>2</sup>School of Earth and Environmental Sciences, University of Portsmouth, Burnaby Building, Burnaby Road, Portsmouth, PO1 3QL, UK.

\*Correspondence to [eotyrannus@gmail.com]

## Abstract

Azhdarchid pterosaurs include the largest animals to ever take to the skies with some species exceeding 10 metres in wingspan and 220 kg in mass. Associated skeletons show that azhdarchids were long-necked, long-jawed predators that combined a wing planform suited for soaring with limb adaptations indicative of quadrupedal terrestrial foraging. The postcranial proportions of the group have been regarded as uniform overall, irrespective of their overall size, notwithstanding suggestions that minor variation may have been present. Here, we discuss a recently discovered giant azhdarchid neck vertebra referable to *Hatzegopteryx* from the Maastrichtian Sebeş Formation of the Transylvanian Basin, Romania, which shows how some azhdarchids departed markedly from conventional views on their proportions. This vertebra, which we consider a cervical VII, is 240 mm long as preserved and almost as wide. Among azhdarchid cervicals, it is remarkable for the thickness of its cortex (4-6 mm along its ventral wall) and robust proportions. By comparing its dimensions to other giant azhdarchid cervicals and to the more completely known necks of smaller taxa, we argue that *Hatzegopteryx* had a proportionally short, stocky neck highly resistant to torsion and compression. This specimen is one of several hinting at greater disparity within Azhdarchidae than previously considered, but is the first to demonstrate such proportional differences within giant taxa. On the assumption that other aspects of *Hatzegopteryx* functional anatomy were similar to those of other azhdarchids, and with reference to the absence of large terrestrial predators in the Maastrichtian of Transylvania, we suggest that this pterosaur played a dominant predatory role among the unusual palaeofauna of ancient Haţeg.

## 32 Introduction

33 Substantial recent interest in the largest known azhdarchid pterosaurs – the Upper Cretaceous  
 34 taxa *Arambourgiania philadelphiae*, *Quetzalcoatlus northropi* and *Hatzegopteryx thambema* –  
 35 has shed much light on their morphology, palaeoecology, and flight capabilities (Witton and  
 36 Naish 2008, 2015; Witton and Habib 2010; Habib 2013). This advanced pterodactyloid clade,  
 37 deeply nested with the morphologically diverse Azhdarchoidea (Nessov 1984; Kellner 2003;  
 38 Unwin 2003; Andres and Meyers 2013), is noted for the proportionally elongate, edentulous  
 39 jaws, remarkably long, cylindrical neck vertebrae and often unusually large size of its  
 40 constituent taxa (Witton and Naish 2008; Witton 2013). Although azhdarchids are comparably  
 41 well represented in the fossil record compared to other pterosaur groups, frustratingly little is  
 42 known of their skeletal anatomy. Partial and complete skeletons of azhdarchids are known and  
 43 are adequately documented (e.g. Godfrey and Currie 2005), but the best represented taxa –  
 44 *Zhejiangopterus linhaiensis* and *Quetzalcoatlus* sp. – remain only preliminarily described (Cai  
 45 and Wei 1994; Lawson 1975; Kellner and Langston 1996). Hypotheses about flight, body mass,  
 46 functional morphology, ecology and lifestyle, all of which remain controversial, are based  
 47 predominantly on knowledge of inadequately described taxa (Witton and Naish 2008, 2015;  
 48 Averianov 2013). Despite these shortfalls of evidence, azhdarchids have been widely assumed  
 49 as uniform in anatomy and ecology (Unwin 2005; Witton and Naish 2008; Witton 2013).

50 Azhdarchids are primarily characterised by their elongate, often tubular neck vertebrae (Nessov  
 51 1984; Kellner 2003; Unwin 2003; Andres and Meyers 2013), and it is a familiar fact of the  
 52 pterosaur literature that these often isolated fossils make up a substantial portion of their fossil  
 53 record. That the giant azhdarchids had the same long necks as their smaller relatives has been  
 54 verified by the discovery of several gigantic vertebrae, including University of Jordan,  
 55 Department of Geology (UJA) specimen VF1: the 620 mm long holotype cervical V of *A.*  
 56 *philadelphiae*. This specimen is argued by some authors to pertain to an animal with a c. 3 m  
 57 long neck (Frey and Martill 1996; Martill et al. 1998), a dimension which would make large  
 58 azhdarchids among the longest-necked animals outside of Sauropoda (Taylor and Wedel 2013)  
 59 and Plesiosauria, despite their necks being formed of only nine vertebrae (Bennett 2014).

60 However, recent discoveries of two proportionally short, isolated azhdarchid cervical vertebrae  
 61 from the Maastrichtian Sebeş Formation (Transylvanian Basin) of western Romania have  
 62 prompted suggestions that some azhdarchids may have been proportionally short necked  
 63 (Vremir 2010; Vremir et al. 2015). The first of these specimens, LPV (FGGUB) R.2395, was  
 64 interpreted as a cervical IV from a small azhdarchid with an estimated 3 m wingspan (Vremir et  
 65 al. 2015). The second represents a gigantic azhdarchid: Transylvanian Museum Society (Cluj-  
 66 Napoca, Romania) specimen EME 315 (Fig. 1). This latter bone is proportionally short and wide,  
 67 of robust construction and bears – for a pterosaur – remarkably thick bone walls. Details of

bone structure and provenance led Vremir (2010) to suggest it may represent a cervical III from *Hatzegopteryx*, a giant azhdarchid described from the middle member of the Densuș-Ciula Formation, Maastrichtian of Vălioara, northern Hațeg basin, deposits contemporary and adjacent to the Sebeș Formation. We discuss the taxonomic identity of the specimen further below. Vremir (2010) concluded that the size and shape of EME 315 is so distinct relative to that of other azhdarchids that it must reflect a departure from expected azhdarchid anatomy and lifestyle.

The concept of short necked azhdarchids is yet to be explored in detail, despite the significance it has for our understanding of azhdarchid palaeoecology and disparity. The functional anatomy of the long, stiffened azhdarchid neck has been the most controversial element in discussions of azhdarchid lifestyles (e.g. Witton and Naish 2013, 2015; Averianov 2013, and references therein), so understanding its variation and biomechanics is paramount to advancing palaeobiological appreciation of the group. Here, we investigate the radical morphological differences between EME 315 and other azhdarchid cervicals from two angles. Firstly, we attempt to estimate the probable neck length of EME 315 and other azhdarchids (both giant and smaller species) to assess possible variation in their proportions and form. Secondly, we assess the bending strength of two giant azhdarchid vertebrae (EME 315 and UJA VF1) to appreciate variation in structural properties and functionality, and relate these to contemporary ideas of azhdarchid behaviour and ecology. It is imperative to these studies that we also investigate the likely identity and vertebral position of EME 315, and this is also discussed below.

## Methods

### *Taxonomic and anatomical identity of EME 315*

EME 315 possesses multiple apomorphies of azhdarchid pterosaur cervical vertebrae, including the characteristic ‘bifid’ neural spine, large, dorsoventrally flattened zygapophyses and a low centrum (e.g. Andres and Ji 2008; Averianov 2010; Buffetaut and Kuang 2010; Vremir et al. 2013). It can thus be referred to Azhdarchidae with confidence. We agree with Vremir (2010) that comparable size, anatomy, and geographical and geological provenance all indicate affinities with *Hatzegopteryx*, a robust giant azhdarchid first described from nearby Vălioara in the Hațeg Basin (Buffetaut et al. 2002, 2003). We draw specific attention to the ventral bone wall of EME 315: at 4-6 mm thick, it is considerably thicker than the 2.6 mm or less reported from most other giant azhdarchids (including the giant *Arambourgiania* holotype cervical – Frey and Martill 1996; Martill et al. 1998) but is comparable to bone walls of the *H. thambema* holotype humerus (Laboratory of Vertebrate Palaeontology, Geological and Geophysical

Faculty, University of Bucharest, Romania) FGGUB R1083 (Buffetaut et al. 2003). A large, elongate cervical vertebra from the Maastrichtian of the French Pyrenees was also described as having thick bone walls of 2-6 mm (Buffetaut et al. 1997) so it is possible that this feature was more widespread in azhdarchids. The spongy internal texture visible at the broken end of EME 315 also recalls the aberrant internal structure of the skull and humerus of the *H. thambema* holotype (Buffetaut et al. 2002). We consider *Hatzegopteryx* and EME 315 to possess a bone construction atypical among pterosaurs, and a close relationship between these specimens likely. The Sebeş Basin material does not overlap with the *H. thambema* holotype, so we, accordingly, provisionally identify the Sebeş Basin vertebra as *Hatzegopteryx* sp. only. This referral of EME 315 to *Hatzegopteryx* is supported by the lack of firm evidence for a second giant azhdarchid in Romania, as well as the fact that, while multiple azhdarchid taxa are known to have been contemporaneous in several Late Cretaceous faunas (Vremir et al. 2013, 2015), we have yet to discover evidence that more than one giant taxon inhabited a given fauna.

Isolated azhdarchid cervicals have typically been regarded as offering little insight to their position within the cervical series, except perhaps for cervical V, which appears distinctly elongate (Frey and Martill 1996; Martill et al. 1998). Recent work on relatively complete azhdarchid cervical skeletons indicates that their vertebrae may show consistent characteristics specific to the position in the cervical series (Pereda-Suberbiola et al. 2003 (*sensu* Kellner 2010); Averianov 2010, 2013) (Fig. 2). Work in this area must be regarded as provisional given that complete azhdarchid necks, or even sufficient material to completely reconstruct entire cervical series, remain few in number. However, we consider known azhdarchid necks of consistent enough form that the likely vertebral position of well-preserved azhdarchid cervicals, such as EME 315, can be determined with some degree of confidence.

Vremir (2010) considered EME 315 as a cervical III, but we consider this unlikely. The neural spines of cervical III in *Azhdarcho lancicollis* (Zoological Institute of the Russian Academy of Sciences, St. Petersburg, Russia, ZIN PH 131/44) and *Quetzalcoatlus* sp. (Texas Memorial Museum, Austin, USA, TMM 41544.16) extend for the length of the entire centrum and lack any obvious reduction in height at mid-length (Fig. 2a; Howse 1986; Averianov 2010), a significant contrast to the bifid neural spine of EME 315. Indeed, Howse (1986) reported that the *Quetzalcoatlus* cervical III neural spine is at its highest point mid-way along its length, a marked contrast to the condition in EME 315. The proportions of cervical III cotyles, which are approximately twice as wide as tall and subequal in height to the neural arch, also contrast with EME 315, as does the continuous tapering of cervical III zygapophyses when viewed in dorsal aspect. Cervical IIIs also seem generally longer-bodied than the proportionally short EME 315. We find greater similarity with other azhdarchid cervicals (below) and thus disagree with a cervical III identity for EME 315.

Azhdarchid cervicals IV and V can be up to eight times longer than wide (Lawson 1975; Howse 1986; Frey and Martill 1996). Their neural spines comprise low anterior and posterior ridges with a mid-length so reduced that they are confluent with the vertebral corpus, sometimes being represented by a faint, narrow ridge at best (Fig. 2d-f). EME 315 is not elongate relative to its width (Fig. 1E) and, though possessing a bifid neural spine, the breadth of the preserved neural spine bases suggests they were robust, tall structures. Azhdarchid cervical VIs seem similar to fourth and fifth elements, but have a proportionally tall posterior neural spine (Fig. 2g-h). EME 315 contrasts with most or all of these conditions, and thus likely pertains to a posterior section of the neck – that is, to cervicals VII or VIII.

Strong similarity occurs between EME 315 and cervicals VII and VIII of *Azhdarcho lancicollis* (ZIN PH 138/44 and 137/44, respectively (Averianov 2010, 2013), Fig. 2i-k), with the most notable similarity pertaining to cervical VII. The cotyle heights of these vertebrae are characteristically shallower than their neural arches, and four times wider than high (Averianov 2010). The cotyle width:height ratio of EME 315 approximates this at ca. 3.7. Both EME 315 and *Azhdarcho* cervical VII possess hypapophyses, a contrast to cervical VIII of *Azhdarcho* where a hypapophysis is absent (Averianov 2010). Reconstructed length:width ratios of EME 315 and the posterior cervicals of *Azhdarcho* are similar (1.36 in *Azhdarcho* cervical VII, 1.06 in cervical VIII, versus 1.25 in EME 315; based on a reconstructed EME 315 length and width of 300 mm and 240 mm, respectively), as are the presences of pneumatic foramina dorsal to the neural canal. The relatively splayed prezygapophyses of cervicals VII and VIII in *Azhdarcho* also correspond well with EME 315, although they are much smaller in *Azhdarcho* cervical VIII. The articular faces in the latter are joined to the vertebral body via a constricted bony shaft, whereas the zygapophyses of EME 315 and *Azhdarcho* cervical VII are more massive overall. Cervical VII in *Azhdarcho* and EME 315 are also similar in having a tapered ‘waist’ mid-way along the length of the centrum. This feature is absent in cervical VIII of *Azhdarcho* which has, in contrast, subparallel lateral margins. The pneumatic foramina are larger than the neural canal in *Azhdarcho*’s cervical VII, which contrasts with the condition in EME 315 and cervical VIII of *Azhdarcho*. EME 315 also lacks pneumatic foramina on the lateral surface of the centrum, in contrast to *Azhdarcho*’s cervical VIII where they are present. The neural spines on the posterior cervicals of *Azhdarcho* are unknown, but those of the posteriormost cervicals of *Phosphatodraco mauritanicus* are proportionally tall and anteroposteriorly restricted (Fig. 2J; Pereda-Suberbiola et al. 2003). This condition matches the one that appears to have been present in EME 315.

EME 315 thus possesses a combination of anatomical traits that are a good match for the posterior cervical vertebrae of at least two other azhdarchid taxa, and it differs markedly from the middle or anterior neck vertebrae of any taxon. We note particular similarity with cervical VII of *Azhdarcho* and hence provisionally consider a seventh cervical position most likely for

EME 315, the caveat being that additional discoveries of azhdarchid posterior cervical vertebrae are needed to bolster our identification.

# *Size of the EME 315 individual*

We refrain from providing a specific wingspan estimate for the EME 315 individual because the relationships between wingspans and cervical vertebrae are not reliably predicted using existing data. Disagreements over the wingspan of the individual represented by the *Arambourgiania* holotype cervical (stated as having a wingspan of 7-8 m wingspan by Pereda-Suberbiola et al. 2003 and yet argued as 10 m or more by others – Frey et al. 1996; Steel et al. 1997; Martill et al. 1998) demonstrate the uncertainty surrounding size estimates of the largest pterosaurs known only from vertebral remains. Vremir (2010) indicated that the great width of EME 315 suggested a similarly expanded postcervical column and perhaps a much larger overall size than that of other giant azhdarchids. This interpretation is questionable as the cervical and anteriormost dorsal vertebrae of giant pterodactyloids are wider and more massive than the rest of the axial column (Bennett 2001; Kellner et al. 2013). A lack of study on proportional scaling of the pterosaur axial column further precludes reliable predictions of the dimensions of the dorsal column belonging to the animal represented by EME 315.

Nevertheless, it is possible to provide a qualified assessment of the general size represented by this vertebra. EME 315 is the most robust pterosaur cervical yet reported and likely conforms to approximate size predictions for the *Hatzegopteryx* holotype, estimated to have a 10-12 m wingspan from the FGGUB R1083 humerus (Buffetaut et al. 2003; Witton and Habib 2010). The size of pterodactyloid cervical condyles and cotyles appears to be relatively uniform along the cervical series (e.g. *Anhanguera* (Wellnhofer 1991a); *Quetzalcoatlus* sp. (Witton and Naish 2008); *Azhdarcho* (2010)), allowing us to assume that the 150 mm wide cotyle of EME 315 is similar to the condylar and cotylar dimensions present along the preceding part of the neck. In the reconstructed neck of *Azhdarcho*, and in completely known necks of *Anhanguera*, atlas cotyle width (assumed to correspond to the dimensions of the occipital condyle) is 30-40% of condyle and cotyle width in the remainder of the neck: the 55 mm wide occipital condyle of the *H. thambema* skull therefore corresponds to the 150 mm wide cotyle of EME 315. The unprecedented width and robust construction of EME 315 also corresponds with the unusually broad skull of *H. thambema*, estimated to span 500 mm across the quadrates (Buffetaut et al. 2003). We take these rough comparisons to indicate that EME 315 probably represents an animal at the upper known limit of pterosaur size.

# *Neck length estimate*

Incredibly long necks incorporating elongate, tubular mid-cervical vertebrae are a well-known feature of Azhdarchidae (e.g. Nessel 1984; Frey and Martill 1996; Kellner 2003; Unwin 2003; Witton and Naish 2008; Averianov 2013). However, published attempts to estimate the length of giant azhdarchid necks are rare and presently limited to isometric scaling of *Quetzalcoatlus* bones to the same linear proportions as the *Arambourgiania* holotype (Frey and Martill 1996; Steel et al. 1997). Subsequent discussions of neck length in giant azhdarchids (e.g. Martill 1997; Taylor and Wedel 2013) have relied on these figures. However, some pterosaur necks, like those of virtually all long-necked tetrapods, are known to scale with positive allometry against body size (Wellnhofer 1970): this calls into question the assumption that azhdarchid necks scaled isometrically and the accuracy of these predicted values.

To estimate and compare the lengths of cervicals III-VII for the EME 315 individual and other azhdarchids, we compiled vertebral length data from six azhdarchid necks: four associated and complete cervical series – representing three skeletons of *Zhejiangopterus linhaiensis* (Cai and Wei 1994), and the holotype of *Phosphatodraco mauritanicus* (Pereda-Suberbiola et al. 2003, as interpreted by Kellner 2010) – in addition to reconstructed, composite skeletons of *Azhdarcho* and *Quetzalcoatlus* sp. (Steel 1997; Averianov 2013) (Table 1). Our sample represents animals with wingspans ranging from 2.5-4.6 m and cervical III-VII lengths of 326–1495 mm. Regression analyses of these data provided reliable relationships between azhdarchid cervical vertebrae and cervical III-VII length (Fig. 3). Surprisingly, we find that azhdarchid necks scale rather differently to other long necked tetrapods. In most long necked animals – examples include sauropods, giraffes, plesiosaurs and tanystropheids (Tschanz 1988; O’Keefe and Hiller 2006; Parrish 2006) – extreme neck length is often associated with a disproportionate increase in the size of cervical vertebrae (i.e. larger animals have disproportionately elongate neck bones against neck length). However, azhdarchid cervical vertebrae seem to largely retain their proportions with increasing neck length, scaling with only slight deviations from isometry. Cervicals III and VII show slightly negative scaling exponents of 0.88 and 0.78 (respectively), while cervicals IV – VI show exponents within 0.9 – 1.11 (Fig. 3). This ‘conservative’ approach to scaling is discussed more below.

### *Vertebral strength analysis*

Azhdarchid cervicals are essentially hollow tubes with near-circular or elliptical cross sections (Fig. 4), and are thus conducive to beam loading calculations to ascertain their strength. We modelled bending load capacity for both UJA VF1 (the holotype vertebra of *Arambourgiania*) and EME 315 based on their minimal central diameters, and using both their preserved and estimated total lengths (Table 2). Cortical thicknesses were measured from broken mid-shaft sections of each bone. To enhance comparability between these vertebrae, we also modelled a



hypothetical *Hatzegopteryx* cervical V based on length projections from our azhdarchid neck dataset and the centrum dimensions of EME 315: we estimate this bone's length as 413 mm. This also provides a minimum estimate of neck strength because, as noted above, cervical V is the longest bone in the azhdarchid neck and thus the most susceptible to distortion under loading. Vertebral sections were modelled as consistent along the vertebral length and internal supporting structures were not factored into our equations. Because the vertebrae in question are elliptical in cross-section, we modelled both dorsoventral and lateral bending resistance. To calculate second moment of area ( $I$ , required to calculate section modulus for stress calculations) for each vertebral axis, we used:

$$1. I = \pi/4(R_1R_2^3 - R_3R_4^3)$$

Where  $R_1$  and  $R_2$  represents the total bone radii in perpendicular x and y axes (relative to loading regime), and  $R_3$  and  $R_4$  represent radii of the internal bone cavity. Bone stress was modelled using cantilever-style loading, where one end of the bone is fixed and the total length of the bone equals the moment arm. Stress values reflect those experienced at the supported end of the bone. Vertebrate bones are rarely loaded as true cantilevers in life but such a reductionist approach provides a quantified means of comparing bone structure and robustness (Witton and Habib 2010). We calculated stresses ( $\sigma$ , Mpa) experienced at the supported end of the vertebrae during cantilevered loading:

$$2. \sigma = WL/Z$$

Where  $L$  is bone length (mm),  $W$  (N) is the weight loaded onto the bone and  $Z$  is section modulus (second moment of area/distance to neutral surface of the vertebra). Calculating bone strength requires some assumptions about the Young's modulus of pterosaur bone. We follow Palmer and Dyke (2009) in using 22 GPa – a value agreeing with several avian long bones – which seems a reasonable proxy for pterosaur bones. Following Currey (2004) and Palmer and Dyke (2010), we used the relationship between Young's modulus and yield stress in tension of 162 MPa. We modelled a range of values reflecting different upper limits for giant pterosaur body mass (180–250 kg) for  $W$  to demonstrate the sensitivity of our results and calculate Relative Failure Force (RFF; Witton and Habib 2010) for each model. RFF is bone failure force, in bending, divided by total body weights. Although pterosaur axial elements were unlikely to ever bear a full loading of body mass in life, it provides a useful proxy by which we might compare the results here with those of other studies (e.g. Witton and Habib 2010) and to compare strengths of pterosaur vertebra against their respective body masses.

## Results and discussion

# 281 Neck length of EME 315 and other azhdarchid pterosaurs

282 The results of our neck length estimates are summarised in Fig. 5. Our dataset shows a  
283 reasonable ( $r^2=0.973$ ) relationship between the length of cervical VII and the combined lengths  
284 of cervicals III-VII:

$$285 \quad 3. \text{CIII-VII} = 17.283\text{CVII}^{0.7835}$$

286 where *CIII-VII* represents the length of cervicals III-VII (mm), and *CVII* represents the length of  
287 cervical VII (mm). Assuming EME 315 is a seventh cervical, its preserved length (240 mm)  
288 predicts a cervical III-VII length of only 1266 mm, while the estimated total length (300 mm)  
289 projects cervical III-VII values of 1508 mm. These values must be considered low given the size  
290 of EME 315 and its indications of body size similar to that of the *H. thambema* holotype. Using  
291 the estimated 770 mm length (Frey and Martill 1996), we modelled the cervical III-VII length of  
292 *Arambourgiania* at 2652 mm, a value shorter than estimates based on strictly isometric scaling  
293 (2817 mm; Steel 1997) but still 75% longer than that predicted for the EME 315 azhdarchid. This  
294 discrepancy is further borne out in our estimate of 412 mm for a *Hatzegopteryx* cervical V –  
295 almost half the estimated length of *Arambourgiania* cervical V. Indeed, predicted cervical  
296 values of EME 315 match those measured from the reconstructed neck of the 4.6 m wingspan  
297 *Quetzalcoatlus* sp. (Steel 1997): its estimated cervical V length and neck length are near  
298 identical to values measured from *Q. sp.* (410 mm and 1495 mm, respectively) despite this  
299 taxon being substantially smaller (Fig. 6).

300 These calculations converge in establishing that *Hatzegopteryx* had a proportionally short neck  
301 (Vremir 2010) c. 50-60% of the length expected for a ‘typical’ giant azhdarchid like  
302 *Arambourgiania*. Our estimates indicate that giant azhdarchids included both *Hatzegopteryx*-  
303 like forms with short, wide necks, and *Arambourgiania*-like species with long, gracile necks. The  
304 former befits an animal with the unusually robust cranial anatomy known for *H. thambema* and  
305 is consistent with the view that this pterosaur was robust overall (Buffetaut et al. 2002, 2003).  
306 As noted above, short necks have been postulated for a much smaller Romanian azhdarchid  
307 known from a likely cervical IV, LPV (FGGUB) R.2395 (Vremir et al. 2015). This neck of this  
308 animal, considered to have a 3-4 m wingspan, was estimated at 352–419 mm using an earlier  
309 version of the data presented above: we revise this estimate upwards here to 460 mm.  
310 Nevertheless, this value is still shorter than that measured from smaller azhdarchids (e.g. the  
311 2.5 m wingspan *Zhejiangopterus linhaiensis*, 502 mm measured neck length) and suggests that  
312 short necks may not be restricted to giant taxa (Vremir et al. 2015). Overall, these data suggest  
313 that there is more variation in neck proportions and robustness within Azhdarchidae than  
314 previously anticipated: the concept of the clade as one with a uniformly long-necked  
315 morphotype (e.g. Witton and Naish 2008) now warrants significant reappraisal.

316

# 317 Neck biomechanics in giant azhdarchids

318 EME 315 represents an anatomical extreme among pterosaur neck vertebrae: its size, bone wall  
319 thickness and massiveness are unprecedented among other flying reptile remains. Its structural  
320 properties, and utility within a possibly shorter variant of the azhdarchid neck, are therefore  
321 significant not only to our understanding of azhdarchid palaeobiology as a whole, but in that  
322 they represent a hitherto unreported morphological class of pterosaur anatomy.

323 Our strength analysis (Table 2) shows that *Hatzegopteryx* neck vertebrae are considerably  
324 stronger than those of *Arambourgiania*. Even at the lowest loading threshold, and in its  
325 strongest bending plane (sagittal), the holotype *Arambourgiania* cervical does not withstand  
326 the strain of one bodyweight. At most, the UJA VF1 vertebrae has RFFs of 0.57 (1765 N loading  
327 in sagittal plane), this decreasing to 0.38 in 2452 N coronal loading. *Hatzegopteryx*, however,  
328 shows consistent capacity for the withstanding of high stresses. The (reconstructed) 300 mm  
329 long EME 315 model has an RFF of 10.04 when loaded with 1765 N in the coronal plane, and  
330 maintains high RFFs (5.57) even when loaded by 2452 N on its weakest axis. The longer (412  
331 mm) hypothetical *Hatzegopteryx* cervical IV is also consistently strong in all tests, able to  
332 withstand 4.05 - 7.3 RFFs in various loading regimes.

333 These findings confirm predictions that giant azhdarchid vertebrae are not functionally uniform  
334 (Vremir 2010). The detailed anatomy of giant azhdarchid cervicals provide insights into the  
335 contrasting figures generated by our bone strength analysis. *Arambourgiania* cervical V can be  
336 viewed as a giant variant on a 'typical' azhdarchid cervical, being a thin-walled (bone wall  
337 thickness 2.6 mm), elongate tube supported internally by a network of bony trabeculae (Frey  
338 and Martill 1996; Martill et al. 1998). It mainly differs from other azhdarchid cervicals in bearing  
339 a mid-centrum section which is taller than wide (55 mm tall vs. 48 mm wide). As is well  
340 documented for other long pterosaur bones, this form is ideally suited to maximising stiffness,  
341 and thus resisting bending and torsion over long dimensions and within constrained loading  
342 regimes. The ratio of bone shaft thickness to wall thickness (bone radius/bone thickness, R/t) in  
343 UJA VF1 is 9.9, a value greater than recorded from other tetrapods but comparable to those  
344 measured from large pterosaur wing bones (Currey 2002; Fastnacht 2005).

345 Frey and Martill (1996) suggested that the unusually tall cross section of *Arambourgiania* likely  
346 improved its resistance to dorsoventral loading, and this is corroborated by our bending  
347 analysis. Dorsoventral expansion of a cervical vertebra is an efficient means to increase vertical  
348 bending strength without incurring additional mass (Frey and Martill 1996), and we might  
349 predict this to be an evolutionary response to an increase in the weight of the neck and head in  
350 large azhdarchid pterosaurs. Even accounting for the 'conservative' scaling of pterosaur necks

(Fig. 3), mass compounds exponentially against length, and giant pterosaurs would thus have experienced proportionally higher loading on their neck skeleton than similarly proportioned smaller species. We predict that the size and weight of large azhdarchid heads explains their relatively 'conservative' cervical scaling compared to long-necked species with small heads, as the heads of the latter impart relatively low structural demands on the supporting neck and permit development of proportionally longer neck components even at large size. In having heads which are predicted to be several metres long (Buffetaut et al. 2003; Witton 2013), giant azhdarchids would have experienced much greater structural demand on their cervical skeletons, even accounting for cranial pneumaticity, and this almost certainly lessened the potential for pronounced cervical allometry.

As with most pterosaur bones, the greatest risk of structural failure to UJA VF1 is buckling: this can be caused by high compressive loads along the long axis of the vertebra or large bending moments. This may explain why the R/t of the *Arambourgiania* cervical is not as high as those measured from other long pterosaur bones (Fastnacht 2005 reports an R/t of 20 for some pterosaur bones), as lowering R/t is one way to increase buckling strength.

The structural characteristics of EME 315 frequently contrast with this configuration. As noted above, the vertebra is proportionally short overall, and although its mid-centrum section has an elliptical shape typical for an azhdarchid, it is broader than other azhdarchid centra in all respects, being 74 mm tall by 115 mm wide. The large second moment of area created by the expanded centrum can be seen as being particularly significant as goes resisting bending through experimental modelling of a vertebra with the *Hatzegopteryx* section profile and the 770 mm length predicted for *Arambourgiania* cervical V. Even when loaded at 2452 N, this hypothetical vertebra still produces high (over 2.17) RFF scores. By contrast, the smaller, thinner-walled section of *Arambourgiania* only achieves an RFF of 1.47 when shortened to 300 mm (the predicted complete length of EME 315) and modelled with the lightest loading in our experiments.

The EME 315 bone wall is relatively thick (4-6 mm) which means that – despite the size of the centrum – it has an R/t comparable to that of *Arambourgiania* at 9.45. This is noteworthy, as its larger size hypothetically permits a much higher R/t, which would be advantageous to decreasing mass and increasing performance against bending (see Currey 2002 for discussion). However, it may be that the thicker cortices of this bone enhanced buckling strength without drastically altering bending strength (Currey 2002) or that its cross-sectional proportions are sufficient to provide high bending resistance alone. Such thick bone walls are not without precedent in pterosaurs – they appear in certain dsungaripterid limb bones (Fastnacht 2005), a partial vertebra from another European azhdarchid (Buffetaut et al. 1997) and the *Hatzegopteryx* type material (Buffetaut et al. 2002, 2003). Buckling resistance, as well as

increased performance against compressive loading, has been posited as an explanation for this phenomenon (Fastnacht 2005).

Several other structural features are of interest for EME 315. Well-preserved endosteal regions of EME 315 show that a system of camellate bone, rather than the trabeculae seen in *Arambourgiania* (Martill et al. 1998), occupied at least the ventral part of the centrum's interior. Such tissues seem pervasive throughout *Hatzegopteryx* bones, also being present in the jaw and humerus. We interpret these features as evidencing further resistance to buckling elsewhere in the skeleton. Furthermore, the already large mid-length centrum of EME 315 is considerably expanded at its anterior and posterior ends. This allows for broadened cotyle/condyle articulations and a greater capacity to distribute high stresses between vertebral joints, and their relatively wide, shallow profile is ideally shaped to resist torsion.

Assuming that the general characteristics and proportions of these giant azhdarchid neck vertebrae apply to their entire cervical series (which seems reasonable, given the profiles of other pterosaur vertebrae), we predict that at least two structural configurations existed among these giant forms. Selection pressures on *Arambourgiania* seem to have prioritised mass reduction and stiffness, which are ideal for elongating bones at the expense of loading capacity. We predict that the anterior cervical skeleton and crania of *Arambourgiania* were relatively slender and lightweight, with a gracile skull more akin to that of *Quetzalcoatlus* than the proportionally broad or deep skulls of *Hatzegopteryx* or the unnamed Texas Memorial Museum specimen 42489-2. EME 315 seems contrarily adapted: its cross-sectional proportions, massive features and thick bone walls are not advantageous for producing a long, lightweight neck skeleton (at least within the context of pterosaur anatomy), but better suited to resisting high bending and compressive stresses. Assuming the other neck bones of the EME 315 individual were similarly adapted, *Hatzegopteryx* must have possessed a significantly stronger neck skeleton than *Arambourgiania*, and perhaps the strongest neck of any known pterosaur. Our stress analysis accords with observations that the very large jaw bones of *Hatzegopteryx* indicate a very wide (0.5 m), and thus potentially relatively large and heavy, skull (Buffetaut et al. 2002, 2003).

#### *Supporting and utilising the azhdarchid neck skeleton*

The robustness and apparent strength of EME 315 raises questions about the function of the *Hatzegopteryx* neck, particularly with respect to how it may have performed in tasks other than supporting a large skull. Investigating this requires some appreciation of pterosaur neck musculature. Pterosaur cervical myology has not featured prominently in technical discussions of this group, but artistic representations of azhdarchids – many of them overseen by pterosaur

researchers – frequently show an extremely reduced cervical musculature relative to the typical tetrapod condition. We assume that these reconstructions were produced following observation of mid-series vertebrae, which are very long, have reduced processes, and have indications of limited arthrological range (Averianov 2013).

However, azhdarchid fossils – including the specimens discussed here – show that the assumption of a paltry, reduced neck musculature represents an oversimplification and is inconsistent with anatomical data from other animals. Our arguments can be summarised as follows: 1) azhdarchid skeletal anatomy suggests that certain muscle groups related to neck function were indeed minimised, but that many aspects of axial, skull and pectoral skeletal anatomy show potential for large muscle attachments; 2) comparisons made between azhdarchid neck skeletons and those of extant animals suggest they are not as atypical as often assumed, and that reptilian cervical musculature correlates well with large muscle attachment sites on azhdarchid cervicals; and 3), that various aspects of azhdarchid anatomy counter proposals of a reduced degree of soft-tissue neck support. We will briefly explore these points here to further elaborate on the functional capacity of giant azhdarchid necks.

Our most general observation is that complete, associated azhdarchid neck skeletons show that they are not solely composed of simple, stiff-jointed, near-featureless tubes. As outlined in Fig. 2, cervicals III, VI, VII and (probably) VIII possess relatively prominent neural spines, indicating differential development of epaxial musculature (Witton and Naish 2008). The ‘tubular’ morphology often ascribed to their neck skeletons only really applies to cervicals IV and V. Averianov (2013) demonstrated that intervertebral cervical articulations are variable along the neck, those of the posterior vertebrae being less restrictive than those of the anterior- and mid-sections. In these respects, azhdarchid necks are comparable to those of other amniotes. X-rays of living animals show that the middle section of the cervical series is often relatively immobile, and that the majority of movement in the neck is achieved via movement at either end of the cervical series (Vidal et al. 1986; Graf et al. 1992, 1995; Taylor et al. 2009). Relatively long-necked mammals (examples include horses, deer, giraffes and camels), as well as extinct long-necked reptiles such as tanystropheids, possess reduced processes and relative immobility associated with their mid-length cervical vertebrae (Fig. 7; Goldfinger 2004; Renesto 2005). Azhdarchid neck skeletons are thus typical in that greater complexity and robustness was present at the extreme ends of their cervical skeleton, as well as in neighbouring cranial or torso skeletal elements; this was surely associated with the anchoring of powerful neck musculature and large ligaments at the base and anterior end of the neck. These are optimal positions from which to support and operate long necks. In view of this, the elongate and tubular, relatively immobile mid-series vertebrae of azhdarchids should be viewed as a pronounced development of a skeletal adaptation common across tetrapods, not as an unusual or unprecedented anatomical configuration.

Azhdarchid skeletons show ample attachment sites for neck musculature. For example, the occiput of *Hatzegopteryx* shows obvious signs of substantial soft-tissue attachment: the nuchal line is well developed and long, and its dorsolateral edges are deeply dished and marked with vertical scarring (Buffetaut et al. 2002, 2003). Comparison with extant reptile anatomy (Herrel and de Vree 1999; Cleuren and de Vree 2000; Tsuihiji 2005, 2010; Snively and Russell 2007; Snively et al. 2013) suggests that these features reflect large insertion areas for transversospinalis musculature (specifically m. transversospinalis capiti and the m. epistropheo-capitis/splenius group), cervical musculature devoted to head and neck extension and lateral flexion. The large neural spines on posterior azhdarchid cervicals and anterior thoracic vertebrae provide potential origin sites for m. transversospinalis capiti, while the long neural spine of cervical III likely anchored m. epistropheo-capitis. The opisthotic process of *Hatzegopteryx* is poorly known but was evidently large and robust and likely facilitated attachment of large neck extensors and lateral flexors (m. semispinalis capitis/spinocapitis posticus). Similarly, the broken basioccipital tuberosities of *Hatzegopteryx* are long even as preserved: neck and head flexors anchoring to these (m. longissimus capitis profundus, m. rectus capitis ventralis) would have had high mechanical advantage. The length and size of these occipital features suggest that large muscles with augmented lever arms were anchored to the azhdarchid skull. Witmer et al. (2003) and Habib and Godfrey (2010) made similar observations about the occipital regions of other pterodactyloids: at least the anterior neck skeleton of pterosaurs was likely strongly muscled.

At the other extreme of the axial column, the azhdarchid scapulocoracoid suggests that their superficial neck musculature may have been well developed. Their scapulae are large and dorsoventrally expanded compared to those of other pterosaurs (e.g. Elgin and Frey 2011), permitting broad insertions of m. levator scapulae and m. serratus (Bennett (2003) shows their likely origin in other pterosaurs). These muscles originate on the anterior cervicals in modern reptiles and can function as neck elevators and retractors if the scapulae are immobile. Azhdarchid scapulocoracoids articulated tightly with the dorsal vertebrae and sternum (Frey et al. 2003) and were buried within deep flight musculature, so were likely capable of little, if any, motion. Contraction of cervical-pectoral muscle groups would thus likely elevate the neck, and asymmetric contraction of these muscles would move the neck laterally. These muscles (or homologues thereof) are particularly large in long-necked, large-headed mammals such as horses and deer (Goldfinger 2004), and we propose that the enlarged pectoral skeleton of azhdarchids may indicate similar enhancement of the posterior neck musculature.

Comparison with the anatomy of modern reptiles suggest that both m. levator scapulae and m. serratus, as well as muscles operating within the cervical series, anchored to the lateral faces of the neural arch and zygapophyses in azhdarchids (Herrel and de Vree 1999; Cleuren and de Vree 2000; Snively and Russell 2007). This is important for consideration of azhdarchid

palaeobiology, it being a clear indication that neural spine height is not the only indicator of neck muscle size. Reptilian cervical extensor musculature, such as *m. longissimus cervicis* and *m. transversospinalis cervicis*, originate and insert on cervical zygapophyses as well as the vertebral corpus (Herrel and de Vree 1999; Snively and Russell 2007). Other muscles, including those superficial muscles outlined above and *m. longus colli ventralis*, originate on cervical ribs, diapophyses and transverse processes (Herrel and de Vree 1999; Cleuren and de Vree 2000; Snively and Russell 2007). These structures are reduced in azhdarchids, but not absent. Juvenile specimens show that vestigial cervical ribs occur on the ventral surfaces of their prezygapophyses (Godfrey and Currie 2005), fusing to the zygapophyses in older animals to form the ventral face of the prezygapophysis (Unwin 2003). In well preserved specimens, fused cervical ribs form prominent ventral prezygapophyseal tubercles (Company et al. 1999; Vremir et al. 2015). The retention of cervical ribs in tubercle form may indicate that these structures maintained a functional role, perhaps persisting as attachment sites for muscles ancestrally anchored to non-reduced cervical ribs, diapophyses or transverse processes. We refrain from making more specific comment on this issue until an improved understanding of pterosaur cervical musculature is achieved, but note that well-preserved azhdarchid zygapophyses have complicated morphologies with crests, prominences, concave facets and well-defined edges (e.g. Frey and Martill 1996; Company et al. 1999; Averianov 2010; Vremir et al. 2013, 2015): anatomy expected of structures that act as anchorage sites for prominent musculature. The atypically elongate, broad zygapophyses of azhdarchids (e.g. Howse 1986; Unwin 2003; Kellner 2003; Witton and Naish 2008) can be viewed with new significance if, as proposed here, they accommodated muscle attachment.

Finally, the idea that azhdarchids had thinly muscled necks is at odds with their cranial proportions, which are among the most extreme of any animal. The skulls of azhdarchids are proportionally huge (Fig. 6; Cai and Wei 1994; Kellner and Langston 1996; Witton 2013) and – even accounting for their high degree of pneumaticity – would have subjected their neck tissues to high amounts of strain and stress. A well-developed system of ligaments and epaxial musculature was likely needed for cranial movement and support.

We thus propose that hypotheses of highly reduced neck muscles in azhdarchids are likely erroneous. The reduction of some axial structures – in particular the neural spines of the mid-series cervicals and small size of cervical rib homologues – suggest that some muscle groups were likely reduced, but other areas for muscle attachment were prominent enough to indicate that their necks were neither weak nor underpowered. Indeed, several of their likely attachment sites must be viewed as expanded compared to those of other pterosaurs, and with effective mechanical advantage for operating the head and neck.



Our hypotheses regarding azhdarchid neck musculature allow us to make some provisional, general comments on the vertebral myology of giant forms. We note that areas likely to anchor muscle – such as neural spines and zygapophyses – of EME 315 are proportionally expanded. The bifid neural spine of EME 315 is broken at the base of each process, but the broken surfaces are sufficiently broad and elongate (Fig. 1) to suggest that the spines were broad, long and perhaps tall when complete. The geometry of the zygapophyses are complex. Low crests and prominent edges extend from the vertebral corpus towards their articular surfaces, and their lateral and medial faces show complex concavities and edges: we posit that these mark muscle scarring. The ventrolateral surfaces of the EME 315 corpus are also notably concave and meet the ventral face along a defined, sweeping edge. These features suggest that EME 315 was well-muscled in life. This seems appropriate given the size of the *Hatzegopteryx* skull, and those features indicating large muscle insertions on its occipital face.

The holotype cervical of *Arambourgiania* may also show some evidence of muscle scarring: a sagittal crest on its anterior ventral surface and two low crests on the dorsal surface of the prezygapophyses. These latter features are topographically similar, though less defined, to crests seen on EME 315 and other azhdarchid vertebrae. However, the overall potential area for muscle attachment in this giant vertebra is much lower than it is in EME 315. The broken section of the anterior surface of the neural spine is smaller than that seen in EME 315, indicating a shallower neural spine overall. The zygapophyses are also shorter and more gracile. These differences might be partly explained by the different likely positions of EME 315 and UJA VF1 within the cervical skeleton (a cervical V is expected to have lesser muscle attachment than preceding or following vertebrae) but better known azhdarchid necks suggest that generalities of morphology will be common in other, adjacent vertebrae along the column (Fig. 5). We therefore conclude that *Arambourgiania* likely had a relatively lightly muscled neck relative to that of *Hatzegopteryx*. This is in keeping with the reduced strength of UJA VF1 predicted in our testing.

# *Disparity and ecological diversity in giant azhdarchids*

EME 315 and the other *Hatzegopteryx* material provides the strongest evidence yet that azhdarchids were not anatomically uniform (Vremir et al. 2013; Witton 2013). Understanding the overall form of azhdarchids is hampered by a lack of associated material, but fragmentary specimens indicate that azhdarchids were variable in at least three major anatomical respects (Figs. 5 and 8). The first is neck type, since some taxa had relatively short (though perhaps not shorter than expected for other pterodactyloids), robust necks (such as *Hatzegopteryx*; R2395), and others had much longer, more gracile and mechanically weaker necks (e.g. *Quetzalcoatlus* sp., *Arambourgiania*). The second is cranial morphotype: this also comprises robust forms, with

relatively short skulls and proportionally broad jaws (e.g. the possible azhdarchid *Bakonydraco*; Javelina Formation specimen TMM 42489-2), and gracile forms with elongate rostra and slender jaws (*Quetzalcoatlus* sp.; *Zhejiangopterus*; *Alanqa*). Some azhdarchids also appear to have relatively slender rostra, as indicated by the concave dorsal skull margin of *Azhdarcho* (Fig. 8A, Averianov 2010). A third category concerns the wing skeletons: we note that the relatively abbreviated metacarpal IV and proximal wing phalanx of the diminutive azhdarchid *Montanazhdarcho minor* contrasts markedly with the elongate distal forelimb elements of *Quetzalcoatlus* sp. and *Zhejiangopterus* (McGowan et al. 2002). It has been speculated that azhdarchids might be roughly grouped into ‘robust’ and ‘gracile’ forms based on these differences (Witton 2013). It certainly seems appropriate to consider forms like *Hatzegopteryx* ‘robust’ and others – e.g., *Quetzalcoatlus* and *Zhejiangopterus* – ‘gracile’, but some taxa show ‘mixed’ anatomies (e.g. *Montanazhdarcho* has proportionally stocky wing bones, but elongate neck bones (McGowan et al. 2002)), suggesting these categories must be considered loose. Azhdarchid body plans may have been rather more varied than imagined previously.

Our assessment of vertebral mechanics in *Hatzegopteryx* and *Arambourgiania* suggests that azhdarchid necks had drastically different functional capabilities. We presume that cranial and cervical disparity reflects distinct foraging habits and prey preferences, with robust azhdarchids tackling relatively larger prey than their gracile counterparts. The stout, thick-walled cervicals of *Hatzegopteryx*, as well as its generally reinforced bones and wide jaws (Buffetaut et al. 2002, 2003), seem better suited to tackling larger, more powerful prey, or for using greater force and violence when obtaining food, than azhdarchid species with thin-walled bones, long, gracile necks and narrow skulls. Undescribed fossils likely referable to *Hatzegopteryx* (including additional skull and limb elements that cannot be described here) show that robust construction was consistent across its body. The high resistance to bending stresses and indications of large cervical muscles in *Hatzegopteryx* are consistent with this concept, as are the inverse findings for *Arambourgiania*.

Modern studies on azhdarchid foraging behaviour suggest that they were terrestrially-foraging generalists (Witton and Naish 2008, 2015; Carroll et al. 2013; Witton, 2017). What little is known of giant azhdarchid anatomy is similar enough to that of the smaller, better known azhdarchids to assume that they also foraged terrestrially, albeit perhaps with a greater emphasis on carnivory. We propose that gracile giants like *Arambourgiania* consumed relatively small prey such as early juvenile and hatchling dinosaurs, large eggs and other diminutive components of Cretaceous terrestrial ecosystems (Fig. 9A). This is in keeping with proposals that some giants occupied ‘middle tier’ predatory niches in some Cretaceous ecosystems (Witton and Naish 2015). *Hatzegopteryx*, however, shows potential for tackling much larger prey items, perhaps even killing animals too large to ingest whole (modern azhdarchid analogues, such as storks, are capable of attacking large animals, and killing human children,

with their azhdarchid-like beaks: see Witton and Naish 2015 for discussion). *Hatzegopteryx* is the largest terrestrial predator known in Maastrichtian eastern Europe by some margin (Witton and Naish 2015): its size, robust anatomy, and the deficit of other large carnivores in well-sampled European deposits implies that it may have been an arch predator in its community (Fig. 9B). The idea that a pterosaur may have played such an important role in a terrestrial Cretaceous ecosystem is far removed from previous interpretations of azhdarchids and pterosaurs generally, and perhaps a clear sign of how far pterosaur studies have progressed in recent decades (see Wellnhofer 1991b; Hone 2012 and Witton 2013 for overviews of pterosaur research).

Finally, the growing evidence for distinct bauplans within Azhdarchidae complicates assessments of pterosaur disparity at the close of the Cretaceous and ideas surrounding the extinction of the group. Azhdarchids dominate pterosaur faunas in the Maastrichtian, with only two localities recording non-azhdarchid pterosaurs from this time (Price 1953; Longrich et al. 2016). Assumptions that azhdarchids were morphologically uniform have led to proposals that Maastrichtian pterosaurs were ecologically constrained at the end of the Cretaceous, and that their extinction represents the unspectacular end of a long, gradual pterosaurian decline (Unwin 2005; Witton 2013). The identification of clear distinctions in form and function within Azhdarchidae, along with the recent potential identification of the first small-bodied azhdarchid species from Campanian sediments (Martin-Silverstone et al. 2016), argues that latest Cretaceous pterosaurs were not as ecologically homogenous as previously thought, and that their extinction may have coincided with their exploitation of niches previously unused in pterosaur evolution. Pterosaur extinction in the K/Pg event may thus have been more significant than traditionally considered.

## Acknowledgements

This work reflects discussion with a number of individuals on azhdarchid pterosaur anatomy, animal scaling and functional anatomy. We are grateful to S. Brusatte, E. Buffetaut, G. Dyke, M. Habib, D. Hone, A. Kellner, C. Palmer, M. Norell, D. Unwin, M. Vremir and M. Wedel for their assistance in helping us develop the ideas outlined in this paper. We are grateful to three referees (Eric Snively, David Hone and Elizabeth Martin-Silverstone) who provided very useful and constructive comments on an earlier version of this paper.

## References

- 637 Andres, B., and Myers, T. S. 2013. Lone star pterosaurs. *Earth and Environmental Science*  
638 *Transactions of the Royal Society of Edinburgh*, 103, 383-398.
- 639 Andres, B. and Qiang, J. 2008. A new pterosaur from the Liaoning Province of China, the  
640 phylogeny of the Pterodactyloidea, and convergence in their cervical vertebrae.  
641 *Palaeontology*, 51, 453-469.
- 642 Averianov, A. O. 2013. Reconstruction of the neck of *Azhdarcho lancicollis* and lifestyle of  
643 azhdarchids (Pterosauria, Azhdarchidae). *Paleontological Journal*, 47, 203-209.
- 644 Averianov, A. O. 2010. The osteology of *Azhdarcho lancicollis* Nessel, 1984 (Pterosauria,  
645 Azhdarchidae) from the late Cretaceous of Uzbekistan. *Proceedings of the Zoological*  
646 *Institute RAS*, 314, 264-317.
- 647 Bennett, S. C. 2001. The osteology and functional morphology of the Late Cretaceous pterosaur  
648 *Pteranodon* Part I. General description of osteology. *Palaeontographica Abteilung A*, 1-112.
- 649 Bennett, S. C. 2003. Morphological evolution of the pectoral girdle in pterosaurs: myology and  
650 function. In Buffetaut, E. and Mazin, J. M. (eds) *Evolution and Palaeobiology of Pterosaurs*,  
651 *Geological Society Special Publications*, 217, 191-215.
- 652 Bennett, S. C. 2014. A new specimen of the pterosaur *Scaphognathus crassirostris*, with  
653 comments on constraint of cervical vertebrae number in pterosaurs. *Neues Jahrbuch für*  
654 *Geologie und Paläontologie-Abhandlungen*, 271, 327-348.
- 655 Buffetaut, E. and Kuang, X. 2010. The *tuba vertebralis* in dsungaripterid pterosaurs. *Acta*  
656 *Geoscientia Sinica*, 31, 10–11.
- 657 Buffetaut, E., Laurent, Y., Le Loeuff, J. and Bilotte, M. 1997. A terminal Cretaceous giant  
658 pterosaur from the French Pyrenees. *Geological Magazine*, 134, 553-556.
- 659 Buffetaut, E., Grigorescu, D. and Csiki, Z. 2002. A new giant pterosaur with a robust skull from  
660 the latest Cretaceous of Romania. *Naturwissenschaften*, 89, 180-184.
- 661 Buffetaut, E., Grigorescu, D. and Csiki, Z. 2003. Giant azhdarchid pterosaurs from the terminal  
662 Cretaceous of Transylvania (western Romania). In Buffetaut, E. and Mazin, J. M. (eds)  
663 *Evolution and Palaeobiology of Pterosaurs*, *Geological Society, London, Special Publications*,  
664 217, 91-104.
- 665 Cai, Z., and Wei, F. 1993. On a new pterosaur (*Zhejiangopterus linhaiensis* gen. et sp. Nov.) from  
666 Upper Cretaceous in Linhai, Zhejiang, China. *Vertebrata Pal Asiatica*, 32, 181-194.

- 667 Carroll, N.R., Poust, A.W. and Varricchio, D.J. 2013. A third azhdarchid pterosaur from the Two  
668 Medicine Formation (Campanian) of Montana. In Sayão, J.M., Costa, F.R., Bantim, R.A.M.,  
669 and Kellner, A.W.A. (eds) *International Symposium on Pterosaurs, Rio Ptero 2013, Short*  
670 *Communications*, 40–42. Universidad Federal do Rio de Janeiro, Rio de Janeiro.
- 671 Cleuren, J. and de Vree, F. 2000. Feeding in crocodilians. In Schwenk, K. (ed) *Feeding: form,*  
672 *function, and evolution in tetrapod vertebrates*. Elsevier Academic Press, 337-358.
- 673 Company, J., Ruiz-Omeñaca, J. I. and Pereda Suberbiola, X. 1999. A long-necked pterosaur  
674 (Pterodactyloidea, Azhdarchidae) from the Upper Cretaceous of Valencia, Spain. *Geologie en*  
675 *Mijnbouw*, 78, 319-333.
- 676 Currey, J. D. 2002. *Bones: structure and mechanics*. Princeton University Press.
- 677 Currey, J. D. 2004. Tensile yield in compact bone is determined by strain, post-yield behaviour  
678 by mineral content. *Journal of biomechanics*, 37, 549-556.
- 679 Elgin, R. A., and Frey, E. 2011. A new azhdarchoid pterosaur from the Cenomanian (Late  
680 Cretaceous) of Lebanon. *Swiss Journal of Geosciences*, 104, 21-33.
- 681 Fastnacht, M. 2005. The first dsungaripterid pterosaur from the Kimmeridgian of Germany and  
682 the biomechanics of pterosaur long bones. *Acta Palaeontologica Polonica*, 50, 273–288.
- 683 Frey, E., and Martill, D. M. 1996. A reappraisal of *Arambourgiania* (Pterosauria,  
684 Pterodactyloidea): One of the world’s largest flying animals. *Neues Jahrbuch fur Geologie*  
685 *und Palaontologie-Abhandlungen*, 199, 221-248.
- 686 Frey, E., Buchy, M. C. and Martill, D. M. 2003. Middle- and bottom- decker Cretaceous  
687 pterosaurs: unique designs in active flying vertebrates. In Buffetaut, E. and Mazin, J. M. (eds)  
688 *Evolution and Palaeobiology of Pterosaurs, Geological Society Special Publications*, 217, 267-  
689 274.
- 690 Graf, W., de Waele, C. and Vidal, P. P. 1992. Skeletal geometry in vertebrates and its relation to  
691 the vestibular end organs. In Berthoz, A., Graf, G. and Vidal, P. P. (eds.) *The Head-Neck*  
692 *Sensory Motor System*. Oxford University Press (New York and Oxford), pp. 129–134.
- 693 Graf, W., de Waele, C. and Vidal, P. P. 1995. Functional anatomy of the head-neck movement  
694 system of quadrupedal and bipedal mammals. *Journal of Anatomy*, 186, 55–74.
- 695 Goldfinger, E. 2004. *Animal Anatomy for Artists: The Elements of Form*. Oxford University Press.
- 696 Habib, M. 2013. Constraining the air giants: limits on size in flying animals as an example of  
697 constraint-based biomechanical theories of form. *Biological Theory*, 8, 245-252.

- 698 Habib, M. and Godfrey, S. 2010. On the hypertrophied opisthotic processes in *Dsungaripterus*  
699 *weii* Young (Pterodactyloidea, Pterosauria). *Acta Geoscientica Sinica*, 31, 26.
- 700 Herrel, A., and Vree, F. D. 1999. The cervical musculature in helodermatid lizards. *Belgian*  
701 *Journal of Zoology*, 129, 175-186.
- 702 Hone, D.W.E. 2012. Pterosaur research: recent advances and a future revolution. *Acta*  
703 *Geologica Sinica*, 86, 1366-1376.
- 704 Howse, S. C. B. 1986. On the cervical vertebrae of the Pterodactyloidea (Reptilia: Archosauria).  
705 *Zoological Journal of the Linnaean Society*, 88, 307-328.
- 706 Godfrey, S. J. and Currie, P. J. 2005. Pterosaurs. In Currie, P.J., and Koppelhus, E.B. (eds)  
707 *Dinosaur Provincial Park: A Spectacular Ancient Ecosystem Revealed*. Indiana University  
708 Press, Bloomington and Indianapolis, 292-311.
- 709 Kellner A. W. A. 2003. Pterosaur phylogeny and comments on the evolutionary history of the  
710 group. In Buffetaut, E. and Mazin, J. M. (eds) *Evolution and Palaeobiology of Pterosaurs*,  
711 *Geological Society Special Publications*, 217, 105–137.
- 712 Kellner, A. W. 2010. Comments on the Pteranodontidae (Pterosauria, Pterodactyloidea) with  
713 the description of two new species. *Anais da Academia Brasileira de Ciências*, 82, 1063-1084.
- 714 Kellner, A. W. A. and Langston Jr, W. 1996. Cranial remains of *Quetzalcoatlus* (Pterosauria,  
715 Azhdarchidae) from Late Cretaceous sediments of Big Bend National Park, Texas. *Journal of*  
716 *Vertebrate Paleontology*, 16, 222-231.
- 717 Kellner, A. W., Campos, D. A., Sayao, J. M., Saraiva, A. A., Rodrigues, T., Oliveira, G., Cruz, L. A.,  
718 Costa, F. R., Silva, H. P. and Ferreira, J. S. 2013. The largest flying reptile from Gondwana: a  
719 new specimen of *Tropeognathus* cf. *T. mesembrinus* Wellnhofer, 1987 (Pterodactyloidea,  
720 Anhangueridae) and other large pterosaurs from the Romualdo Formation, Lower  
721 Cretaceous, Brazil. *Anais da Academia Brasileira de Ciências*, 85, 113-135.
- 722 Lawson, D. A. 1975. Pterosaur from the Latest Cretaceous of West Texas: Discovery of the  
723 largest flying creature. *Science*, 185, 947–948.
- 724 Longrich, N., Martill, D. M. and Andres, B. 2016. Late Maastrichtian pterosaurs from the Tethys  
725 Seaway provide evidence for mass extinction of pterosaurs at the Cretaceous-Palaeogene  
726 boundary. *SVPCA and SPPC 2016, Liverpool, Abstract Book*, 22.
- 727 Martill, D. M., Frey, E., Sadaqah, R. M. and Khoury, H. N. 1998. Discovery of the holotype of the  
728 giant pterosaur *Titanopteryx philadelphiae* Arambourg 1959, and the status of

- 729     *Arambourgiania* and *Quetzalcoatlus*. *Neues Jahrbuch fur Geologie und Palaontologie-*  
730     *Abhandlungen*, 207, 57-76.
- 731     Martin-Silverstone, E., Witton, M. P., Arbour, V. M., and Currie, P. J. 2016. A small azhdarchoid  
732     pterosaur from the latest Cretaceous, the age of flying giants. *Royal Society Open Science*, 3,  
733     160333.
- 734     McGowan, M. R., Padian, K., de Sosa, M. A. and Harmon, R. J. 2002. Description of  
735     *Montanazhdarcho minor*, an azhdarchid pterosaur from the Two Medicine Formation  
736     (Campanian) of Montana. *PaleoBios* 22, 1-9.
- 737     Nesov L. A. 1984. [Pterosaurs and birds from the Late Cretaceous of Middle Asia]. *Paleontol.*  
738     *Zhurnal*, 1, 47–57. [In Russian]].
- 739     O’Keefe, F. R. and Hiller, N. 2006. Morphologic and ontogenetic patterns in elasmosaur neck  
740     length, with comments on the taxonomic utility of neck length variables. *Paludicola*, 5, 206–  
741     229.
- 742     Parrish, J. M. 2006. The origins of high browsing and the effects of phylogeny and scaling on  
743     neck length in sauropodomorphs. In Carrano, M. T., Gaudin, T. J., Blob, R. W. and Wible, J. R.  
744     (eds). *Amniote paleobiology: perspectives on the evolution of mammals, birds, and reptiles*.  
745     University of Chicago Press, 201-224.
- 746     Palmer, C., and Dyke, G. J. 2009. Biomechanics of the unique pterosaur pteroid. *Proceedings of*  
747     *the Royal Society of London B: Biological Sciences*, rspb20091899.
- 748     Price, L. I. 1953. A presença de Pterosáuria no Cretáceo superior do Estado da Paraíba. *Divisão*  
749     *de Geologia e Mineralogia Notas Preliminares e Estudos*, 71, 1-10.
- 750     Renesto, S. 2005. A new specimen of *Tanystropheus* (Reptilia Protorosauria) from the Middle  
751     Triassic of Switzerland and the ecology of the genus. *Rivista Italiana di Paleontologia e*  
752     *Stratigrafia (Research In Paleontology and Stratigraphy)*, 111, 377-394.
- 753     Rieppel, O., Jiang, D. Y., Fraser, N. C., Hao, W. C., Motani, R., Sun, Y. L., and Sun, Z. Y. 2010.  
754     *Tanystropheus* cf. *T. longobardicus* from the early Late Triassic of Guizhou Province,  
755     southwestern China. *Journal of Vertebrate Paleontology*, 30, 1082-1089.
- 756     Snively, E., and Russell, A. P. 2007. Functional variation of neck muscles and their relation to  
757     feeding style in Tyrannosauridae and other large theropod dinosaurs. *The Anatomical*  
758     *Record*, 290, 934-957.

Snively, E., Russell, A. P., Powell, G. L., Theodor, J. M., and Ryan, M. J. 2014. The role of the neck in the feeding behaviour of the Tyrannosauridae: inference based on kinematics and muscle function of extant avians. *Journal of Zoology*, 292, 290-303.

Steel, L., Martill, D. M., Kirk, J., Anders, A., Loveridge, R. F., Frey, E. and Martin, J. G. 1997. *Arambourgiania philadelphiae*: giant wings in small halls. *The Geological Curator*, 6, 305-313

Suberbiola, X. P., Bardet, N., Jouve, S., Iarochène, M., Bouya, B. and Amaghazaz, M. 2003. A new azhdarchid pterosaur from the Late Cretaceous phosphates of Morocco. In Buffetaut, E. and Mazin, J. M. (eds) *Evolution and Palaeobiology of Pterosaurs, Geological Society Special Publications*, 217, 79-90.

Taylor, M. P., and Wedel, M. J. 2013. Why sauropods had long necks; and why giraffes have short necks. *PeerJ*, 1, e36.

Taylor, M. P., Wedel, M. J., and Naish, D. 2009. Head and neck posture in sauropod dinosaurs inferred from extant animals. *Acta Palaeontologica Polonica*, 54, 213-220.

Tschanz, K. A. R. L. 1988. Allometry and heterochrony in the growth of the neck of Triassic prolacertiform reptiles. *Palaeontology*, 31, 997-1011.

Tsuihiji, T. 2005. Homologies of the transversospinalis muscles in the anterior presacral region of Sauria (crown Diapsida). *Journal of Morphology*, 263, 151-178.

Tsuihiji, T. 2010. Reconstructions of the axial muscle insertions in the occipital region of dinosaurs: evaluations of past hypotheses on Marginocephalia and Tyrannosauridae using the Extant Phylogenetic Bracket approach. *The Anatomical Record*, 293, 1360-1386.

Unwin D. M. 2003. On the phylogeny and evolutionary history of pterosaurs. In Buffetaut, E. and Mazin, J. M. (eds) *Evolution and Palaeobiology of Pterosaurs, Geological Society Special Publications*, 217, 139–190.

Unwin, D. M. 2005. *The Pterosaurs from Deep Time*. Pi Press, New York.

Vidal, P. P., Graf, W. and Berthoz, A. 1986. The orientation of the cervical vertebral column in unrestrained awake animals. *Experimental Brain Research*, 61, 549-559.

Vremir, M. 2010. New faunal elements from the Late Cretaceous (Maastrichtian) continental deposits of Sebeş area (Transylvania). *Acta Musei Sabesiensis*, 2, 635-684.

Vremir, M., Kellner, A. W. A., Naish, D. and Dyke, G. J. 2013. A new azhdarchid pterosaur from the Late Cretaceous of the Transylvanian Basin, Romania: implications for azhdarchid diversity and distribution. *PLoS ONE* 8, e54268.



790 Vremir, M., Witton, M., Naish, D., Dyke, G., Brusatte, S. L., Norell, M. and Totoianu, R. 2015. A  
 791 Medium-Sized Robust-Necked Azhdarchid Pterosaur (Pterodactyloidea: Azhdarchidae) from  
 792 the Maastrichtian of Pui (Hațeg Basin, Transylvania, Romania). *American Museum Novitates*,  
 793 3827, 1-16.

794 Wellnhofer, P. 1970. Die Pterodactyloidea (Pterosauria) der Oberjura-Plattenkalke  
 795 Süddeutschlands. *Bayerische Akademie der Wissenschaften, Mathematisch-  
 796 Wissenschaftlichen Klasse, Abhandlungen*, 141, 1-133.

797 Wellnhofer, P. 1991a. Weitere Pterosaurierfunde aus der Santana-Formation (Apt) der Chapada  
 798 do Araripe, Brasilien. *Palaeontographica* 215, 43-101.

799 Wellnhofer, P. 1991b. *The Illustrated Encyclopedia of Pterosaurs*. Salamander Books.

800 Witmer, L. M., Chatterjee, S., Franzosa, J., and Rowe, T. 2003. Neuroanatomy of flying reptiles  
 801 and implications for flight, posture and behaviour. *Nature*, 425, 950-953.

802 Witton, M. P. 2013. *Pterosaurs: Natural History, Evolution, Anatomy*. Princeton University Press.

803 Witton, M. P. 2017. Pterosaurs in Mesozoic ecosystems: a review of fossil evidence. In Hone, D.  
 804 W. E., Martill, D. M. and Witton, M. P. (eds) *New Perspectives on Pterosaur Palaeobiology*.  
 805 *Geological Society Special Publications*. In press.

806 Witton, M. P. and Habib, M. B. 2010. On the size and flight diversity of giant pterosaurs, the use  
 807 of birds as pterosaur analogues and comments on pterosaur flightlessness. *PLoS ONE*, 5,  
 808 e13982.

809 Witton, M. P. and Naish, D. 2008. A reappraisal of azhdarchid pterosaur functional morphology  
 810 and paleoecology. *PLoS ONE*, 3, e2271.

811 Witton, M. P., and Naish, D. (2013). Azhdarchid pterosaurs: water-trawling pelican mimics or  
 812 “terrestrial stalkers”? *Acta Palaeontologica Polonica*, 60, 651-660.

# Figure captions

Figure 1. Giant azhdarchid cervical vertebra referred to *Hatzegopteryx* sp. A-D, line drawings of EME 315 in anterior (A), right lateral (B), ventral (C) and dorsal (D) views; E, proportions of EME 315 compared to other azhdarchid cervicals: note atypical combination of length/width ratio (l:w) and length compared to other azhdarchid cervicals, and especially against the only other known giant cervical, *Arambourgiania* (UJA RF1). Light shading indicates damage; dark shading indicates filler. Abbreviations: co, cotyle; hy, hypapophysis; nc, neural canal; ns; neural spine (anterior region); nsp, neural spine (posterior region); pnf, pneumatic foramen; prz, prezygapophysis; poz, postzygapophysis; vprzt, ventral prezygapophyseal tubercle (fused cervical rib). Scale bar is 100 mm.

Figure 2. Characteristics of azhdarchid vertebrae across their cervical series, demonstrated by several azhdarchid taxa. A, *Azhdarcho lancicollis* cervical III (ZIN PH 131/44), left lateral aspect; B-C, *Quetzalcoatlus* sp. cervical III (TMM 41544.16) in dorsal (B) and left lateral (C) aspect; D, *A. lancicollis* cervical IV (ZIN PH 144/44), left lateral aspect; E, *Q.* sp. cervical V (TMM 41455.15), left lateral aspect; F, *Arambourgiania philadelphiae* cervical V (UJA VF1), dorsal aspect; G-H, *A. lancicollis* cervical VI (ZIN PH 147/44) in left lateral (G) and posterior (H) aspect (note especially large neural spine); I, *A. lancicollis* cervical VII (ZIN PH 138/44), dorsal aspect; J, *Phosphatodraco* cervical VII (OCP DEK/GE 111), left lateral aspect; K, *A. lancicollis* cervical VIII (ZIN PH 137/44), dorsal aspect. Abbreviations as for Figure 2, also with con; condyle; ex, exapophysis; ns, neural spine. A, D, G-H, and K after Averianov (2010); F, after Frey and Martill (1996); J, after Pereda-Suberbiola et al. 2003.

Figure 3. Relationships between azhdarchid cervical vertebrae to cervical III-VII length.

Figure 4. Metrics and cross sections used in estimates of bending strength analysis. A, EME 315 in dorsal view showing line of modelled section (dotted line) and projected 300 mm length; B, UJA VF1 in dorsal view showing line of section and projected 770 mm length (Frey and Martill 1996); C, cross section and dimensions of EME 315; D, cross section of UJA VF1. Note difference in shape and bone wall thicknesses in C and D.

Figure 5. Measured and estimated azhdarchid pterosaur neck lengths against approximate wingspans.

Figure 6. Speculative skeletal reconstructions of *Hatzegopteryx* sp. and *Arambourgiania philadelphiae* (estimated wingspans  $\geq 10$  m – Martill and Frey 1996; Buffetaut et al. 2003) to show discrepancy in neck length alongside a ‘typical’ azhdarchid body plan. A, *Hatzegopteryx* skeleton in lateral aspect; B, dorsal view of EME 315 and FGGUB R1083 jaw elements, proportionate to actual size, suggesting *Hatzegopteryx* bore a wide, as well as relatively short,

neck construction (soft-tissue outline in black). Jaw width after Buffetaut et al. (2003); C, reconstructed *Arambourgiania philadelphiae* cervicals III-VII in lateral aspect; D, 4.6 m wingspan *Q. sp.* skeleton in lateral aspect; E, *Q. sp.* cervical vertebrae III-V and skull in dorsal view; Note how the neck length of *Hatzegopteryx* is similar to this much smaller pterosaur. *H. thambema* holotype (FGGUB R1083) and undescribed referred elements are shown in A; known elements of *A. philadelphiae* (UJA JF1) indicated in white shading in C. Scale bar represents 1 m.

Figure 7. Azhdarchid craniocervical skeleton compared to those of some other tetrapods. A, *Tanystropheus cf. longobardicus*; B, reconstruction of *Zhejiangopterus linhaiensis* cervical skeleton, vertebral morphology adapted from Averianov (2010); C, *Giraffa camelopardalis*; D, *Camelus dromedarius*; E, *Odocoileus virginianus*. Note that the mid-series vertebrae of all taxa – even those with highly complex, strongly-muscled neck skeletons – have reduced features compared to those at the posterior and anterior: the fact that azhdarchid mid-series cervicals have reduced features does not necessarily reflect underdeveloped cervical soft-tissues. A, reconstructed from fossils illustrated by Rieppel et al. (2010); B, reconstructed from Cai and Wei (1994) and Averianov (2010); C-E after Goldfinger (2004). Images not to scale.



Figure 8. Azhdarchid disparity in cranial and limb anatomy. A, ZIN PH 112/44, rostral fragment of *Azhdarcho lancicollis* showing concave dorsal skull margin (after Averianov 2010); B, anterior skull and mandible of TMM 42489-2, unnamed azhdarchid from the Javelina Formation, USA; C, restored skull of *Quetzalcoatlus* sp. (based on Kellner and Langston 1996); D, skull of *Zhejiangopterus linhaiensis* (based on Cai and Wei 1994); E, MOR 69I, *Montanazhdarcho minor* holotype pectoral girdle and left forelimb (note stunted metacarpal IV); F, M1323 postcrania of *Z. linhaiensis*. Abbreviations: car, carpals; cer, cervical vertebrae; cor, coracoid; fem, femur; hum, humerus; mcIV, metacarpal IV; pt, pteroid; rad, radius; tib, tibia; ul, ulna; wpl, wing phalanx I. Scale bars represent 100 mm, except for A (10 mm).

Figure 9. Diversity in predicted life appearance and ecologies for giant azhdarchid pterosaurs. A, two giant, long-necked azhdarchids – the Maastrichtian species *Arambourgiania philadelphiae* – argue over a small theropod; B, the similarly sized but more powerful Maastrichtian, Transylvanian giant azhdarchid pterosaur *Hatzegopteryx* sp. preys on the rhabdodontid iguanodontian *Zalmoxes*. Because large predatory theropods are unknown on Late Cretaceous Hațeg Island, giant azhdarchids may have played a key role as terrestrial predators in this community.

# Figure 1

Giant azhdarchid cervical vertebra referred to *Hatzegopteryx* sp.


Figure 1. Giant azhdarchid cervical vertebra referred to *Hatzegopteryx* sp. A-D, line drawings of EME 315 in anterior (A), right lateral (B), ventral (C) and dorsal (D) views; E, proportions of EME 315 compared to other azhdarchid cervicals: note atypical combination of length/width ratio (l:w) and length compared to other azhdarchid cervicals, and especially against the only other known giant cervical, *Arambourgiania* (UJA RF1). Light shading indicates damage; dark shading indicates filler. Abbreviations: co, cotyle; hy, hypapophysis; nc, neural canal; nsa; neural spine (anterior region); nsp, neural spine (posterior region); pnf, pneumatic foramen; prz, prezygapophysis; poz, postzygapophysis; vprzt, ventral prezygapophyseal tubercle (fused cervical rib). Scale bar is 100 mm.



# Figure 2

Characteristics of azhdarchid vertebrae across their cervical series, demonstrated by several azhdarchid taxa.


Figure 2. Characteristics of azhdarchid vertebrae across their cervical series, demonstrated by several azhdarchid taxa. A, *Azhdarcho lancicollis* cervical III (ZIN PH 131/44), left lateral aspect; B-C, *Quetzalcoatlus* sp. cervical III (TMM 41544.16) in dorsal (B) and left lateral (C) aspect; D, *A. lancicollis* cervical IV (ZIN PH 144/44), left lateral aspect; E, *Q.* sp. cervical V (TMM 41455.15), left lateral aspect; F, *Arambourgiania philadelphiae* cervical V (UJA VF1), dorsal aspect; G-H, *A. lancicollis* cervical VI (ZIN PH 147/44) in left lateral (G) and posterior (H) aspect (note especially large neural spine); I, *A. lancicollis* cervical VII (ZIN PH 138/44), dorsal aspect; J, *Phosphatodraco* cervical VII (OCP DEK/GE 111), left lateral aspect; K, *A. lancicollis* cervical VIII (ZIN PH 137/44), dorsal aspect. Abbreviations as for Figure 2, also with con; condyle; ex, exapophysis; ns, neural spine. A, D, G-H, and K after Averianov (2010); F, after Frey and Martill (1996); J, after Pereda-Suberbiola et al. 2003.



# Figure 3

Relationships between azhdarchid cervical vertebrae to cervical III-VII length.

Figure 3. Relationships between azhdarchid cervical vertebrae to cervical III-VII length.






# Figure 4

Metrics and cross sections used in estimates of bending strength analysis.


Figure 4. Metrics and cross sections used in estimates of bending strength analysis. A, EME 315 in dorsal view showing line of modelled section (dotted line) and projected 300 mm length; B, UJA VF1 in dorsal view showing line of section and projected 770 mm length (Frey and Martill 1996); C, cross section and dimensions of EME 315; D, cross section of UJA VF1. Note difference in shape and bone wall thicknesses in C and D.



# Figure 5

Measured and estimated azhdarchid pterosaur neck lengths against approximate wingspans.

Figure 5. Measured and estimated azhdarchid pterosaur neck lengths against approximate wingspans.



# Figure 6

Speculative skeletal reconstructions of *Hatzegopteryx* sp. and *Arambourgiania philadelphiae* to show discrepancy in neck length alongside a 'typical' azhdarchid body plan.


Figure 6. Speculative skeletal reconstructions of *Hatzegopteryx* sp. and *Arambourgiania philadelphiae* (estimated wingspans  $\geq 10$  m – Martill and Frey 1996; Buffetaut et al. 2003) to show discrepancy in neck length alongside a 'typical' azhdarchid body plan. A, *Hatzegopteryx* skeleton in lateral aspect; B, dorsal view of EME 315 and FGGUB R1083 jaw elements, proportionate to actual size, suggesting *Hatzegopteryx* bore a wide, as well as relatively short, neck construction (soft-tissue outline in black). Jaw width after Buffetaut et al. (2003); C, reconstructed *Arambourgiania philadelphiae* cervicals III-VII in lateral aspect; D, 4.6 m wingspan *Q.* sp. skeleton in lateral aspect; E, *Q.* sp. cervical vertebrae III-V and skull in dorsal view; Note how the neck length of *Hatzegopteryx* is similar to this much smaller pterosaur. *H. thambema* holotype (FGGUB R1083) and undescribed referred elements are shown in A; known elements of *A. philadelphiae* (UJA JF1) indicated in white shading in C. Scale bar represents 1 m.



# Figure 7

Azhdarchid craniocervical skeleton compared to those of some other tetrapods.


Figure 7. Azhdarchid craniocervical skeleton compared to those of some other tetrapods. A, *Tanystropheus* cf. *longobardicus*; B, reconstruction of *Zhejiangopterus linhaiensis* cervical skeleton, vertebral morphology adapted from Averianov (2010); C, *Giraffa camelopardalis*; D, *Camelus dromedarius*; E, *Odocoileus virginianus*. Note that the mid-series vertebrae of all taxa – even those with highly complex, strongly-muscled neck skeletons – have reduced features compared to those at the posterior and anterior: the fact that azhdarchid mid-series cervicals have reduced features does not necessarily reflect underdeveloped cervical soft-tissues. A, reconstructed from fossils illustrated by Rieppel et al. (2010); B, reconstructed from Cai and Wei (1994) and Averianov (2010); C-E after Goldfinger (2004). Images not to scale.



# Figure 8

Azhdarchid disparity in cranial and limb anatomy.

Figure 8. Azhdarchid disparity in cranial and limb anatomy. A, ZIN PH 112/44, rostral fragment of *Azhdarcho lancicollis* showing concave dorsal skull margin (after Averianov 2010); B, anterior skull and mandible of TMM 42489-2, unnamed azhdarchid from the Javelina Formation, USA; C, restored skull of *Quetzalcoatlus* sp. (based on Kellner and Langston 1996); D, skull of *Zhejiangopterus linhaiensis* (based on Cai and Wei 1994); E, MOR 69I, *Montanazhdarcho minor* holotype pectoral girdle and left forelimb (note stunted metacarpal IV); F, M1323 postcrania of *Z. linhaiensis*. Abbreviations: car, carpals; cer, cervical vertebrae; cor, coracoid; fem, femur; hum, humerus; mclV, metacarpal IV; pt, pteroid; rad, radius; tib, tibia; ul, ulna; wpl, wing phalanx I. Scale bars represent 100 mm, except for A (10 mm).



# Figure 9

Diversity in predicted life appearance and ecologies for giant azhdarchid pterosaurs.

Figure 9. Diversity in predicted life appearance and ecologies for giant azhdarchid pterosaurs. A, two giant, long-necked azhdarchids – the Maastrichtian species *Arambourgiania philadelphiae* - argue over a small theropod; B, the similarly sized but more powerful Maastrichtian, Transylvanian giant azhdarchid pterosaur *Hatzegopteryx* sp. preys on the rhabdodontid iguanodontian *Zalmoxes*. Because large predatory theropods are unknown on Late Cretaceous Hațeg Island, giant azhdarchids may have played a key role as terrestrial predators in this community.







**Table 1**(on next page)

Azhdarchid cervical vertebrae data used in neck length estimates.

	Lengths (mm)					
Taxon	<i>Zhejiangopterus linhaiensis</i>			<i>Phosphatodraco mauritanicus</i>	<i>Azhdarcho lancicollis</i>	<i>Quetzalcoatlus</i> sp.
Reference	Cai and Wei 1994			Pereda-Suberbiola et al. 2003; Kellner 2009	Averianov 2013	Steel et al. 1997
Specimen number	M1323	M13234	M1328	OCP/DEK GE 111	Reconstruction	Reconstruction
Cervical number						
III	36	50	57	110	57.2	170.0
IV	114	82	92	190	78.1	265.0
V	142	84	98	225	156.2	410.0
VI	120	72	81	190	102.3	380.0
VII	90	38	56	150	60.0	270.0
CIII-VII neck length	502	326	384	865	453.75	1495.00
Proportion of CV/neck length	0.283	0.258	0.255	0.260	0.34	0.27
Proportion of CVII/neck length	0.179	0.117	0.146	0.173	0.13	0.18

1

2 Table 1. Azhdarchid cervical vertebrae data used in neck length estimates.

# **Table 2**(on next page)

Giant azhdarchid cervical vertebra bending strength compared.

Taxonomic ID	Specimen	Body mass (kg)	W (N)	Vertebral length (mm)	Centrum width radius (mm)	Centrum height radius (mm)	Cortical thickness (mm)	R/t	Section modulus		Maximum stress (Mpa)		RFF	
									Sagittal bending	Coronal bending	Sagittal bending	Coronal bending	Sagittal bending	Coronal bending
<i>Hatzegopteryx</i> sp.	EME 315	250	2,452	240	57.5	37	5	9.45	25307	32830	23.25	17.92	6.97	9.04
<i>Hatzegopteryx</i> sp.	EME 315 (reconstructed)	250	2,452	300	57.5	37	5	9.45	25307	32830	29.06	22.40	5.57	7.23
<i>Hatzegopteryx</i> sp.	EME 315 (hypothetical CV)	250	2,452	412.7	57.5	37	5	9.45	25307	32830	39.98	30.82	4.05	5.26
<i>Arambourgiania philadelphiae</i>	UJA VF1	250	2,452	620	24	27.5	2.6	9.90	4819	4455	315.40	341.20	0.51	0.47
<i>Arambourgiania philadelphiae</i>	UJA VF1 (reconstructed)	250	2,452	770	24	27.5	2.6	9.90	4819	4455	391.71	423.75	0.41	0.38
<i>Hatzegopteryx</i> sp.	EME 315	200	1,961	240	57.5	37	5	9.45	25307	32830	18.60	14.34	8.71	11.30
<i>Hatzegopteryx</i> sp.	EME 315 (reconstructed)	200	1,961	300	57.5	37	5	9.45	25307	32830	23.25	17.92	6.97	9.04
<i>Hatzegopteryx</i> sp.	EME 315 (hypothetical CV)	200	1,961	412.7	57.5	37	5	9.45	25307	32830	31.98	24.66	5.07	6.57
<i>Arambourgiania philadelphiae</i>	UJA VF1	200	1,961	620	24	27.5	2.6	9.90	4819	4455	252.32	272.96	0.64	0.59
<i>Arambourgiania philadelphiae</i>	UJA VF1 (reconstructed)	200	1,961	770	24	27.5	2.6	9.90	4819	4455	313.36	339.00	0.52	0.48
<i>Hatzegopteryx</i> sp.	EME 315	180	1,765	240	57.5	37	5	9.45	25307	32830	16.74	12.90	9.68	12.55
<i>Hatzegopteryx</i> sp.	EME 315 (reconstructed)	180	1,765	300	57.5	37	5	9.45	25307	32830	20.93	16.13	7.74	10.04
<i>Hatzegopteryx</i> sp.	EME 315 (hypothetical CV)	180	1,765	412.7	57.5	37	5	9.45	25307	32830	28.79	22.19	5.63	7.30
<i>Arambourgiania philadelphiae</i>	UJA VF1	180	1,765	620	24	27.5	2.6	9.90	4819	4455	227.09	245.67	0.71	0.66
<i>Arambourgiania philadelphiae</i>	UJA VF1 (reconstructed)	180	1,765	770	24	27.5	2.6	9.90	4819	4455	282.03	305.10	0.57	0.53

1

2 Table 2. Giant azhdarchid cervical vertebra bending strength compared.