

Exploration of immunoglobulin transcriptomes from mice immunized with three-finger toxins and phospholipases A₂ from the Central American coral snake, *Micrurus nigrocinctus*

Andreas H Laustsen^{Corresp., 1}, Mikael Engmark^{1,2}, Christopher Clouser^{3,4}, Sonia Timberlake⁵, Francois Vigneault^{3,4}, José María Gutiérrez⁶, Bruno Lomonte⁶

¹ Department of Biotechnology and Biomedicine, Technical University of Denmark, Kgs. Lyngby, Denmark

² Department of Bio and Health Informatics, Technical University of Denmark, Kgs. Lyngby, Denmark

³ Juno Therapeutics, Seattle, Washington, United States of America

⁴ AbViro, Boston, Massachusetts, United States of America

⁵ Finch Therapeutics, Somerville, Massachusetts, United States of America

⁶ Instituto Clodomiro Picado, Universidad de Costa Rica, San José, Costa Rica

Corresponding Author: Andreas H Laustsen

Email address: ahol@bio.dtu.dk

Snakebite envenomings represent a neglected public health issue in many parts of the rural tropical world. Animal-derived antivenoms have existed for more than a hundred years and are effective in neutralizing snake venom toxins when timely administered. However, the low immunogenicity of many small but potent snake venom toxins represents a challenge for obtaining a balanced immune response against the medically relevant components of the venom. Here, we employ high-throughput sequencing of the immunoglobulin (Ig) transcriptome of mice immunized with a three-finger toxin and a phospholipase A₂ from the venom of the Central American coral snake, *Micrurus nigrocinctus*. Although exploratory in nature, our indicate results showed that only low frequencies of mRNA encoding IgG isotypes, the most relevant isotype for therapeutic purposes, were present in splenocytes of five mice immunized with 6 doses of the two types of toxins over 90 days. Furthermore, analysis of Ig heavy chain transcripts showed that no particular combination of variable (V) and joining (J) gene segments had been selected in the immunization process, as would be expected after a strong humoral immune response to a single antigen. Combined with the titration of toxin-specific antibodies in the sera of immunized mice, these data support the low immunogenicity of three-finger toxins and phospholipases A₂ found in *M. nigrocinctus* venoms, and highlight the need for future studies analyzing the complexity of antibody responses to toxins at the molecular level.

Exploration of immunoglobulin transcriptomes from mice immunized with three-
finger toxins and phospholipases A₂ from the Central American coral snake,
Micrurus nigrocinctus

Andreas H. Laustsen^{1*}, Mikael Engmark^{1,2}, Christopher Clouser^{3,4}, Sonia Timberlake⁵, Francois
Vigneault^{3,4}, José María Gutiérrez⁶, Bruno Lomonte⁶

¹ Department of Biotechnology and Biomedicine, Technical University of Denmark, Denmark

² Department of Bio and Health Informatics, Technical University of Denmark, Denmark

³ Juno Therapeutics, Seattle (WA), USA

⁴ AbVitro, Boston (MA), USA

⁵ Finch Therapeutics, Somerville (MA), USA

⁶ Instituto Clodomiro Picado, Facultad de Microbiología, Universidad de Costa Rica,
San José, Costa Rica

Running title: Antibody isotyping of immune response against coral snake toxins

*** Address correspondence to:**

Dr. Andreas H. Laustsen (ahola@bio.dtu.dk), Department of Biotechnology and Biomedicine,
Technical University of Denmark, Denmark

Abstract

Snakebite envenomings represent a neglected public health issue in many parts of the rural tropical world. Animal-derived antivenoms have existed for more than a hundred years and are effective in neutralizing snake venom toxins when timely administered. However, the low immunogenicity of many small but potent snake venom toxins represents a challenge for obtaining a balanced immune response against the medically relevant components of the venom. Here, we employ high-throughput sequencing of the immunoglobulin (Ig) transcriptome of mice immunized with a three-finger toxin and a phospholipase A₂ from the venom of the Central American coral snake, *Micrurus nigrocinctus*. Although exploratory in nature, our indicate results showed that only low frequencies of mRNA encoding IgG isotypes, the most relevant isotype for therapeutic purposes, were present in splenocytes of five mice immunized with 6 doses of the two types of toxins over 90 days. Furthermore, analysis of Ig heavy chain transcripts showed that no particular combination of variable (V) and joining (J) gene segments had been selected in the immunization process, as would be expected after a strong humoral immune response to a single antigen. Combined with the titration of toxin-specific antibodies in the sera of immunized mice, these data support the low immunogenicity of three-finger toxins and phospholipases A₂ found in *M. nigrocinctus* venoms, and highlight the need for future studies analyzing the complexity of antibody responses to toxins at the molecular level.

(232 words)

1. Introduction

Snakebite envenomings represent a major public health concern in tropical regions of the world (Williams *et al.*, 2011). Despite emerging discoveries that may one day pave the way for novel biotechnology-based antivenoms (reviewed by Laustsen *et al.*, 2016a; 2016b), animal serum-derived antivenoms remain the cornerstone of snakebite envenoming treatment (Gutiérrez *et al.*, 2011). Production of antivenom is challenged by a large variation in immunogenicity of many key snake venom toxins resulting in unpredictable immune responses in production animals (Cook *et al.*, 2010; Guidolin *et al.*, 2010). It has been shown that many of the immunogenic venom components are in fact not important for toxicity (Antúnez *et al.*, 2010; Gutiérrez *et al.*, 2009; Laustsen *et al.*, 2015), and conversely, that some highly toxic venom components, such as α -neurotoxins (both short and long neurotoxins), phospholipases A₂, and P-I snake venom metalloproteinases may be poorly immunogenic (Schottler, 1951; Gutiérrez *et al.*, 2009; Chotwiwatthanakun *et al.*, 2001; Ownby & Colberg, 1990; Judge *et al.*, 2006; Wong *et al.*, 2016; Tan *et al.*, 2016; Leong *et al.*, 2015; Tan *et al.*, 2015). Combined, this creates a challenge for antivenom production, since the goal of obtaining an antivenom with a strong, yet balanced response against all the medically relevant toxins becomes a complex endeavor.

Coral snakes (genera *Micrurus*, *Leptomicrurus*, and *Micruroides*) are, together with the sea snake *Hydrophis* (*Pelamis*) *platura*, the representatives of the snake family Elapidae in the Americas, comprising approximately 85 species (Campbell & Lamar, 2004; the Reptile Database - www.reptile-database.org). Although *Micrurus* species are only responsible for about 1-2% of snakebite cases in this continent, roughly corresponding to 750 to 1000 cases per year, envenomings by these snakes can be fatal if not treated properly and timely (Warrell, 2004; Gutiérrez, 2014; Bucarechi *et al.*, 2016). Envenomings resulting from coral snakebites are

predominantly associated with descending neuromuscular paralysis, which may end in respiratory arrest ([Warrell, 2004](#); [Bucarechi et al., 2016](#)).

Production of antivenoms against *Micrurus* snakes is particularly challenging, as (a) it is very difficult to maintain coral snakes in captivity ([Chacón et al., 2012](#)); (b) the majority of *Micrurus* species provide a very low yield of venom, implying that the collection of the quantities of venom required for horse immunization and quality control testing demands the ‘milking’ of many specimens ([Chacón et al., 2012](#); [Bolaños, 1972](#)); and (c) there is a variable extent of immunological cross-recognition between venoms from coral snakes of different species; hence, antivenoms raised against some species are not always effective in the neutralization of venoms of other species ([Bolaños, Cerdas & Abalos, 1978](#); [Tanaka et al., 2016](#)). As a result, only a few laboratories manufacture *Micrurus* antivenoms, and several countries where these snakes inhabit completely lack this therapeutic resource, e.g. Venezuela, Ecuador, Peru, Bolivia, the Guyanas, and Paraguay, which severely limits the clinical management of these accidents.

Knowledge on the composition of the venoms of *Micrurus* species has increased steadily over the last years, as a consequence of proteomic characterizations (reviewed by [Lomonte et al., 2016b](#)). Two main venom phenotype patterns have been identified, i.e. venoms rich in neurotoxins of the three-finger toxin (3FTx) family, and venoms rich in phospholipases A₂ (PLA₂s) ([Fernández et al., 2015](#)). In addition to these two main protein families, other minor components of these venoms include L-amino acid oxidases, serine proteinases, metalloproteinases, nerve growth factor, C-type lectin-like proteins, Kunitz-type inhibitors, among others ([Fernández et al., 2011, 2015](#); [Corrêa-Netto et al., 2011](#); [Lomonte et al., 2016a](#); [Sanz et al., 2016](#); [Rey-Suárez et al., 2011, 2016](#)). In some cases, the toxins playing the main role

in overall toxicity have been identified, these being 3FTxs and PLA₂s (Rey-Suárez *et al.*, 2012; Vergara *et al.*, 2014; Fernández *et al.*, 2015; Castro *et al.*, 2015; Ramos *et al.*, 2016).

The limited immunogenicity of the highly toxic PLA₂s and 3FTxs (Fernández *et al.*, 2011, Rosso *et al.*, 1996; Alape-Girón *et al.*, 1996) represents another difficulty in production of *Micrurus* antivenom, since it thwarts the goal of raising a balanced immune response against these medically relevant toxins. In order to further explore how these toxins interact with the mammalian immune system, we chose a mouse model and employed an NGS approach using the AbSeq™ technology developed by AbVitro (now Juno Therapeutics, <https://www.junotherapeutics.com>), based on Illumina sequencing (Fig. 1). The methodology was utilized to sequence immunoglobulin (Ig) encoding mRNA transcripts from splenic B-lymphocytes in mice subjected to immunization with either a 3FTx or a PLA₂ toxin from the venom of *M. nigrocinctus* (Central American coral snake). By this approach, the transcription levels of different immunoglobulin isotypes and dominant clones of B-lymphocytes with a particular usage of V (variable) and J (joining) gene segments can be determined for Ig heavy chain transcripts. This methodology has previously been employed for investigating B-cell populations in autoimmune (Stern *et al.*, 2014) or infectious diseases (Tsioris *et al.*, 2015; Di Niro *et al.*, 2015), for example. By employing the AbSeq™ high-throughput approach, we explore, for the first time, the Ig transcriptome including VJ usage patterns in individual animals subjected to immunization with two relevant toxin classes of elapid snakes. Although exploratory in its nature and somewhat limited by a small sample size, this study thus provides novel insight into the humoral response of mice immunized with 3FTx or PLA₂ toxins and highlights important challenges of raising antibodies against poorly immunogenic toxins.




Figure 1: Schematic overview of the experimental strategy. Following extraction from the snakes, venom is fractionated by HPLC, and the fractions of interest are used for immunization of rodents. Upon completion of the immunization protocol, the rodents are sacrificed and RNA is extracted and subjected to the AbSeq protocol (see Materials and Methods), whereby the RNA is reverse transcribed and barcoded, allowing for correct pairing of V_H and V_L chains after DNA sequencing.

2. Materials and Methods

2.1 Snake venom and toxins

Venom from *M. nigrocinctus* was obtained from a pool of more than 50 adult specimens collected in the Central Pacific region of Costa Rica, kept at the serpentarium of Instituto Clodomiro Picado, Universidad de Costa Rica. The snakes were not collected for this study, but belong to Instituto Clodomiro Picado, where their venoms are routinely used for the production of antivenom, wherefore a field permit was not necessary. Venom extraction is performed every four months (Chacón et al., 2012). The venom was lyophilized and stored at -20°C.

Fractionation of the venom was performed by RP-HPLC on a C₁₈ column (4.6 x 250 mm, 5 µm particle diameter; Supelco) as previously described (Fernández et al., 2011). In brief, 2 mg of venom dissolved in 200 µL of water containing 0.1% trifluoroacetic acid (TFA; solution A) were separated at 1 mL/min in an Agilent 1200 chromatograph monitored at 215 nm, applying a gradient towards solution B (acetonitrile, containing 0.1% TFA): 0% B for 5 min, 0–15% B over 10 min, 15–45% B over 60 min, 45–70% B over 10 min, and 70% B over 9 min. Fractions of interest (the major components of the venom, belonging to the 3FTx (lethal) and PLA₂ (myotoxic) protein families) were collected manually, dried in a vacuum centrifuge, and identified by trypsin digestion followed by MALDI-TOF/TOF mass spectrometry (Fernández et al., 2011). Proteins were redissolved in water and their concentrations were estimated on the basis of their absorbance at 280 nm, using a NanoDrop (Thermo) instrument.

2.2 Immunization of mice

Three female CD-1 mice (16-18 g) from Instituto Clodomiro Picado were immunized with a three-finger toxin (3FTx), and two with a phospholipase A₂ (PLA₂), respectively. These correspond to fractions #3 (~P80548) and #30 (~P81166/P81167) described in the previous proteomic characterization of this venom (Fernández et al., 2011). All toxin doses were injected

by the intraperitoneal route. The priming dose was 1 μg emulsified in Freund's complete adjuvant, followed by five booster doses injected in physiological saline without adjuvant, at days 15 (1 μg), 43 (2 μg), 63 (4 μg), and 83 (6 μg for the 3FTx and 8 μg for the PLA₂). At day 90, after obtaining a blood sample for monitoring of the antibody response by enzyme-immunoassay, mice were euthanized by CO₂ inhalation. Their spleens were immediately removed, cut in small pieces, and disaggregated over a stainless steel mesh to obtain splenocytes. These cell suspensions were aliquoted in RNeasy lysis solution (Qiagen) and shipped within 24 h to AbViro, at room temperature, for subsequent molecular studies. The use of animals for these experiments followed the ethical guidelines of the *Comité Institucional para el Uso y Cuido de Animales* (CICUA), Universidad de Costa Rica, with the approval number 82-08.

2.3 Enzyme-immunoassay (ELISA)

In order to evaluate the individual antibody responses of the mice, wells in MaxiSorp 96-well plates (NUNC, Roskilde, Denmark) were coated overnight with 1 μg of either 3FTx or PLA₂, dissolved in 100 μL PBS (0.12 M NaCl, 0.04 M sodium phosphate, pH 7.2). Wells were washed five times with PBS and blocked by adding 100 μL PBS containing 2% (w:v) bovine serum albumin (BSA, Sigma), and incubated at room temperature for 1 h. Plates were then washed five times with PBS. Serial dilutions of serum from each mouse were prepared in PBS + 2% BSA and 100 μL was added to each well, in triplicates, and incubated overnight at 4°C. Normal mouse serum, run simultaneously under identical conditions was used as a control for background. Plates were then washed five times with PBS, followed by the addition of 100 μL of a 1:3000 dilution of anti-mouse IgG (whole molecule) antibodies conjugated to alkaline phosphatase, in PBS + 1% BSA. The plates were incubated for 2 h, and then washed five times with FALC

buffer (0.05 M Tris, 0.15 M NaCl, 20 μ M ZnCl₂, 1 mM MgCl₂, pH 7.4). Development of color was attained by addition of 100 μ L *p*-nitrophenyl phosphate (1 mg/mL in 9.7% v/v diethanolamine buffer, pH 9.8) and absorbances at 405 nm were recorded (Multiskan FC, Thermo Scientific).

2.4 Assessment of mRNA quality

Assessment of RNA quality was performed using Agilent's TapeStation according to the manufacturers protocol and algorithm to calculate RIN^e scores (<http://www.agilent.com/cs/library/technicaloverviews/public/5990-9613EN.pdf>).

2.5 Library preparation and high-throughput sequencing of B-cell receptors

The method for high-throughput sequencing of the B-cell repertoire was performed as described elsewhere (*Di Niro et al., 2015; Tsioris et al., 2015*). Briefly, RNA was reverse-transcribed into cDNA using a biotinylated oligo dT primer. An adaptor sequence was added to the 3' end of all cDNA, which contains the Illumina P7 universal priming site and a 17-nucleotide unique molecular identifier (UMI). Products were purified using streptavidin-coated magnetic beads followed by a primary PCR reaction using a pool of primers targeting the IGHA, IGHD, IGHE, IGHG, IGHM, IGKC and IGLC regions, as well as a sample-indexed Illumina P7C7 primer. The immunoglobulin-specific primers contained tails corresponding to the Illumina P5 sequence. PCR products were then purified using AMPure XP beads. A secondary PCR was then performed to add the Illumina C5 clustering sequence to the end of the molecule containing the constant region. The number of secondary PCR cycles was tailored to each sample to avoid entering plateau phase, as judged by a prior quantitative PCR analysis. The final products

(reverse-transcribed UTR + VDJ + partial - Cexon segments of the transcripts of the immunoglobulin chains, plus molecular barcode, Illumina multiplexing barcode, and Illumina sequencing adapters) were purified, quantified with Agilent TapeStation and pooled in equimolar proportions, followed by high-throughput paired-end sequencing on the Illumina MiSeq platform. For sequencing, the Illumina 600 cycle kit was used with the modifications that 325 cycles was used for read 1, 6 cycles for the index reads, 300 cycles for read 2 and a 10% PhiX spike-in to increase sequence diversity.

2.6 VJ repertoire sequencing data analysis

MiSeq reads were demultiplexed using Illumina software, and processed with the pRESTO toolsuite ([Vander Heiden et al., 2014](#)) as following: Positions with less than Phred quality 5 were masked with Ns. Isotype-specific primers and molecular barcodes (UIDs or UMIs) were identified in the amplicon and trimmed, using pRESTO MaskPrimers-cut. A read 1 and read 2 consensus sequence was generated separately for each mRNA from reads grouped by unique molecular identifier, which are PCR replicates arising from the same original mRNA molecule of origin. UMI read groups were aligned with MUSCLE ([Edgar, 2004](#)), and pRESTO was used to BuildConsensus, requiring $\geq 60\%$ of called PCR primer sequences agree for the read group, maximum nucleotide diversity of 0.1, using majority rule on indel positions, and masking alignment columns with low posterior (consensus) quality. Paired end consensus sequences were then stitched in two rounds. First, ungapped alignment of each read pair's consensus sequence termini was optimized using a Z-score approximation and scored with a binomial p -value as implemented in pRESTO AssemblePairs-align. For read pairs failing to stitch this way, stitching was attempted using the human BCR germline V exons to scaffold each read prior to stitching or

gapped read-joining, using pRESTO's AssemblePairs-reference. Positions with posterior consensus quality less than Phred 5 were masked again with Ns. Each mRNA was annotated for V, D, J germline gene of origin, productivity, and CDR3 region using igblastn (Ye et al, 2013). Igblast output was parsed and analyzed with custom scripts as described below (proprietary to AbVitro (now Juno Therapeutics)) and visualized with R (R Core Team, 2014). Briefly, clones were defined using a conservative approach, grouping mRNAs from the same V and J germline gene of origin and having the same isotype and CDR3 nucleotide sequence. Non-productive rearrangements (as predicted by Igblast) were excluded from the analysis. Within each sample, clones were ranked by abundance (mRNAs/clone) and plotted to highlight the most expanded binding motifs. To examine preferences for V-J usage common across mice exposed to the same toxin, clones were grouped into larger bins that each encompass a single V gene-J gene combination, but all isotypes and CDR3s.

2.7 In silico epitope predictions

The sequences of selected toxin components in the two venom fractions used for immunization were obtained from the UniProtKB database (<http://www.uniprot.org>) and linear parts of B-cell epitopes were predicted using the Bepi-Pred 1.0 server (Larsen et al., 2006) using 0.9 as Threshold to obtain a sensitivity of 0.25 and Specificity of 0.91. As no experimental structures were available, homology models were build based CPHmodels 3.2 (Nielsen et al., 2010) and the pdb-files were submitted to DiscoTope 2.0 (Kringelum et al., 2012) using 0.5 as Threshold to obtain a sensitivity of 0.23 and Specificity of 0.90.

3. Results and Discussion

Three-finger toxins (3FTx) and phospholipase A₂s (PLA₂) are the two most abundant toxin families in the venom of *M. nigrocinctus* (Fernández et al., 2011), and generally they are the two snake toxin families which have been most investigated (Laustsen et al., 2016a). In the venom of *M. nigrocinctus* these toxins cause neuromuscular paralysis, owing to a combination of pre- and post-synaptic actions, and myotoxicity, providing the venom with its high toxicity (Rosso et al., 1996; Alape-Girón et al., 1996). In previous studies it was observed that 3FTxs and PLA₂s were recognized more weakly than larger proteins from this venom, by a therapeutic equine antivenom (Fernández et al., 2011). Despite their low immunogenicity, it was possible to raise an antibody response against both toxins in four out of five mice, although high variation in the antibody titer was observed (Fig. 2). Mice immunized with PLA₂ had a higher antibody titer than mice immunized with the 3FTx, in agreement with the higher molecular mass of the former.





Figure 2: ELISA titrations of serum antibodies against *M. nigrocinctus* PLA₂ or 3FTx in mice following a 90-day immunization protocol. Two mice were immunized with PLA₂, and three were immunized with 3FTx. Plates were coated with either PLA₂ or 3FTx, and antibodies were detected as described in Materials and Methods.

255

Assessment of the mRNA from harvested mouse splenocytes indicated that it was of sufficient quality to proceed to sequencing (RIN^e scores between 5.2 and 6.4). A high-throughput sequencing approach (AbSeq™) was employed to investigate transcription levels of Ig isotypes and the usage of V and J gene segments for heavy chain assembly in mice that were immunized with a 3FTx or a PLA₂. Investigation of the 50 most frequent VJ combinations for the immunized mice did not, however, result in identification of a dominant combination, as the VJ usage was found to be similar across all samples (Fig. 3). This finding suggests that the generated antibody responses might be diverse and that multiple specific antibodies with low abundance are generated in each mouse.



265
266

Figure 3. Comparison of the relative abundance of mRNA for the 50 most abundant VJ combinations for the mouse 3FTx-3 and mouse PLA₂-2 showing VJ usage to be similar across samples. Similar VJ usage patterns were observed for other pairs of immunized mice.

Looking at the sequences of all mRNA transcripts encoding heavy chain variable domain (V_H) clones across each sample, we were able to find shared V_H clones with similar relative abundances in either the PLA₂-immunized or the 3FTx-immunized mice (Fig. 4). In comparison, almost no V_H clones were shared between mice immunized with different toxins (Fig. 5). This implies that the immunization procedure did indeed elicit specific, but different responses dependent on whether PLA₂s or 3FTxs were employed for immunization. The relatively high number of V_H clones found in both of the PLA₂-immunized mice (Fig. 4A) compared to lower number of V_H clones found across the three 3FTx-immunized mice (Fig. 4B-D) further indicate that immunization with PLA₂s is more prone to give rise to antibodies transcribed in similar quantities. Also, an intermediate number of similar V_H clones was found in both the PLA₂-1 and 3FTx-3 samples (Fig. 5E), even though the correlation in relative abundance was not equally pronounced. This is likely explained by the fact that the majority of V_H clones found in both PLA₂-immunized mice are not expected to be specific towards the toxins, but instead are likely to be directed against other (environmental) antigens that the mice have encountered throughout their life.





Figure 4: Relative abundance of unique V_H clone transcripts compared between samples. A large group of V_H transcripts are found in similar abundance in different mice immunized with the same toxin. **A)** Comparison between mouse PLA_2-1 and PLA_2-2 , **B)** Comparison between mouse $3FTx-1$ and $3FTx-2$, **C)** Comparison between mouse $3FTx-1$ and $3FTx-3$, **D)** Comparison between mouse $3FTx-2$ and $3FTx-3$.




294

295 **Figure 5:** Relative abundance of unique V_H clone transcripts compared between samples. Only
 296 few V_H transcripts are found in similar abundance in more than one mouse, when mice
 297 immunized with different toxins are compared. **A)** Comparison between mouse PLA₂-1 and
 298 3FTx-1, **B)** Comparison between mouse PLA₂-2 and 3FTx-1, **C)** Comparison between mouse
 299 PLA₂-1 and 3FTx-2, **D)** Comparison between mouse PLA₂-2 and 3FTx-2, **E)** Comparison
 300 between mouse PLA₂-1 and 3FTx-3, **F)** Comparison between mouse PLA₂-2 and 3FTx-3.

301

302 The AbSeq™ antibody sequencing methodology is capable of determining the Ig isotype of the
 303 identified V_H clones. The coloring of the V_H clones in [Fig. 4](#) and [Fig. 5](#) reveals that a large
 304 number of the most abundant V_H clones present in the mice are of the IgA isotype, which was
 305 confirmed by further investigation of all mRNA transcripts from the splenocytes ([Fig. 6](#)). This is
 306 surprising, as IgG is known to be the dominant immunoglobulin class in mouse blood after the
 307 response to T-dependent protein antigens. The observation could possibly be explained by
 308 differences in expression levels due to different translation rates and half-lives of mRNA
 309 transcripts encoding different immunoglobulin isotypes. All approved antibody-based therapies
 310 on the market are based on IgGs ([Walsh, 2014](#)), which are also the desired isotype for
 311 antivenoms. In immunized horses for antivenom production, two isotypes of IgG are largely
 312 responsible for the neutralization of toxic effects in the case of viperid snake venoms ([Fernandes](#)
 313 [et al., 2000](#)). However, analysis of the transcripts obtained from the immunized mice revealed
 314 only a low percentage of IgG transcripts, as compared to the transcripts for other Igs ([Figs. 4-6](#)).
 315 With the acknowledgement of the small sample size employed in this study, this finding may
 316 indicate a difficulty in raising potent IgG antibodies against both 3FTxs and PLA₂s in rodents. If
 317 similar difficulty is present in horses, this may therefore have implications for antivenom
 318 production. Our results may further suggest that the immune response is slightly lower against
 319 3FTxs than for PLA₂s based on the lower abundance of IgG transcripts in mice immunized with
 320 3FTx ([Fig. 6](#)). Taken together with the results from the ELISA assay ([Fig. 2](#)) and the observation
 321 that immunization with PLA₂s is more prone to give rise to similar Ig transcripts ([Fig. 4A](#) vs.
 322 [Fig. 4B-D](#)), we suggest that the PLA₂ toxins may possibly be slightly more immunogenic than
 323 the 3FTx, although neither toxin seems to have high immunogenicity. The underlying reason for

this could possibly be due to the smaller molecular size of 3FTx compared to PLA₂s, or that PLA₂s may contain distinct epitopes better capable of eliciting an adaptive immune response than 3FTxs. Predictions of possible B-cell epitopes using the protein sequences and BepiPred 1.0 ([Larsen et al. 2006](#)) or DiscoTope 2.0 ([Kringelum et al. 2012](#)) with structural models of the investigated toxins as input do not indicate a major difference in the bare number of possible epitopes. However, the suggested difference in immunogenicity is further indicated by the fact that only two IgG-encoding mRNA transcripts are found in the top 20 most abundant Ig-encoding mRNA transcripts for only one out of three of the 3FTx-immunized mice. In comparison, six and nine of the top 20 mRNA transcripts for mice immunized with PLA₂s encode the IgG isotype ([Fig. 7](#)). It would be interesting to assess whether the immune response of horses against these elapid venom toxins is also characterized by a low proportion of IgG – a finding that would have evident implications for antivenom manufacture, as IgG has been shown to be the antibody isotype of therapeutic value ([Fernandes et al., 2000](#)). However, this is beyond the scope of this exploratory study.



339

340 **Figure 6:** Overview of total mRNA transcripts encoding different immunoglobulin isotypes from
 341 the immunized mice (normalized). Numbers above each bar represents the mRNA count in each
 342 sample.

343




Figure 7: The 20 most abundant V_H clone transcripts and their corresponding isotypes in each immunized mouse based on their fraction of total immunoglobulin mRNA.

4. Concluding remarks and outlook

In addition to demonstrating the utility of high-throughput sequencing technology, AbSeq™, for investigation of immune responses in animals immunized with snake venom toxins, the exploratory findings presented here may indicate possible difficulties in obtaining an IgG response against the medically important toxins of the 3FTx and PLA₂ families from *M. nigrocinctus*. Given that these proteins play key toxic roles in envenomings by elapid snakes, this underlines a drawback of current antivenom production based on immunized animal serum, since IgG has been shown to be the antibody isotype of therapeutic value (Fernandes et al., 2000). Although based on a small exploratory sample size, these findings therefore contribute to the understanding of snake toxin immunogenicity and indicate the possible difficulty in obtaining balanced immune responses in animals during the immunization process.

360 **Acknowledgments**

361 We thank Mikael Rørdam Andersen from the Technical University of Denmark for
 362 fruitful scientific discussion.

References

- Alape-Girón A, Stiles B, Schmidt J, Girón-Cortés M, Thelestam M, Jörnvall H, Bergman T. 1996. Characterization of multiple nicotinic acetylcholine receptor-binding proteins and phospholipases A₂ from the venom of the coral snake *Micrurus nigrocinctus nigrocinctus*. *FEBS Letters* 380:29–32.
- Antúnez J, Fernández J, Lomonte B, Angulo Y, Sanz L, Pérez A, Calvete JJ, Gutiérrez JM. 2010. Antivenomics of *Atropoides mexicanus* and *Atropoides picadoi* snake venoms: Relationship to the neutralization of toxic and enzymatic activities. *Journal of Venom Research* 1:8–17.
- Bolaños R, Cerdas L, Abalos JW. 1978. Venoms of coral snakes (*Micrurus* spp.): Report on a multivalent antivenin for the Americas. *Bulletin of the Pan American Health Organization* 12:23–27.
- Bolaños-Herrera R. 1972. Toxicity of Costa Rican snake venoms for the white mouse. *The American Journal of Tropical Medicine and Hygiene* 21:360–363.
- Bucarechi F, De Capitani EM, Vieira RJ, Rodrigues CK, Zannin M, Da Silva Jr. NJ, Casais-e-Silva LL, Hyslop S. 2016. Coral snake bites (*Micrurus* spp.) in Brazil: A review of literature reports. *Clinical Toxicology* 54:222–234.
- Campbell JA, Lamar WW. 2004. *The Venomous Reptiles of the Western Hemisphere. Vol. II*. Ithaca, New York: Comstock Publishing Associates, Cornell University Press.
- Castro KL, Duarte CG, Ramos HR, de Avila RAM, Schneider FS, Oliveira D, Freitas CF, Kalapothakis E, Ho PL, Chávez-Olortegui C. 2015. Identification and characterization of B-cell epitopes of 3FTx and PLA₂ toxins from *Micrurus corallinus* snake venom. *Toxicon* 93:51–60.

387 Chacón D, Arias J, Solano G, Bonilla F, Gómez A. 2012. Maintaining Coral Snakes (*Micrurus*
388 *nigrocinctus*, Serpentes: Elapidae) for venom production on an alternative fish-based diet.
389 *Toxicon* 60:249–253.

390 Chotwiwatthanakun C, Ronachai P, Akesowan S, Sriprapat S, Ratanabanangkoon K. 2001.
391 Production of potent polyvalent antivenom against three elapid venoms using a low dose,
392 low volume, multi-site immunization protocol. *Toxicon* 39:1487–1494.

393 Cook DAN, Owen T, Wagstaff SC, Kinne J, Wernery U, Harrison RA. 2010. Analysis of
394 camelid IgG for antivenom development: serological responses of venom-immunised
395 camels to prepare either monospecific or polyspecific antivenoms for West Africa. *Toxicon*
396 56:363–372.

397 Corrêa-Netto C, Junqueira-de-Azevedo IL, Silva DA, Ho PL, Leitão-de-Araújo M, Alves ML,
398 Sanz L, Foguel D, Zingali RB, Calvete JJ. 2011. Snake venomomics and venom gland
399 transcriptomic analysis of Brazilian coral snakes, *Micrurus altirostris* and *M. corallinus*.
400 *Journal of Proteomics* 74:1795–1809.

401 Di Niro R, Lee SJ, Vander Heiden JA, Elsner RA, Trivedi N, Bannock JM, Gupta NT, Kleinstein
402 SH, Vigneault F, Gilbert TJ, Meffre E. 2015. *Salmonella* infection drives promiscuous B
403 cell activation followed by extrafollicular affinity maturation. *Immunity* 43:120–131.

404 Edgar RC. 2004. MUSCLE: multiple sequence alignment with high accuracy and high
405 throughput. *Nucleic Acids Research* 32:1792–1797.

406 Fernández J, Alape-Girón A, Angulo Y, Sanz L, Gutiérrez JM, Calvete JJ, Lomonte B. 2011.
407 Venomic and antivenomic analyses of the Central American coral snake, *Micrurus*
408 *nigrocinctus* (Elapidae). *Journal of Proteome Research* 10:1816–1827.

409 Fernandes I, Lima EX, Takehara HA, Moura-da-Silva AM, Tanjoni I, Gutiérrez JM. 2000. Horse
410 IgG isotypes and cross-neutralization of two snake antivenoms produced in Brazil and
411 Costa Rica. *Toxicon* 38:633–44.

412 Fernández J, Vargas N, Pla D, Sasa M, Rey-Suárez P, Sanz L, Gutiérrez JM, Calvete JJ,
413 Lomonte B. 2015. Snake venomomics of *Micrurus alleni* and *Micrurus mosquitensis* from the
414 Caribbean region of Costa Rica reveals two divergent compositional patterns in New
415 World elapids. *Toxicon* 107:217–233.

416 Guidolin RG, Marcelino RM, Gondo HH, Morais JF, Ferreira RA, Silva CL, Kipnis TL, Silva
417 JA, Fafetine J, da Silva WD. 2010. Polyvalent horse F(ab')₂ snake antivenom: development
418 of process to produce polyvalent horse F(ab')₂ antibodies anti-african snake venom. *African*
419 *Journal of Biotechnology* 9:2446–2455.

420 Gutiérrez JM, Lomonte B, Aird SD, da Silva Jr. NJ, 2016. Mecanismo de ação dos venenos de
421 cobras corais. In: As Cobras Corais do Brasil: Biologia, Taxonomia, Venenos e
422 Envenenamientos (da Silva Jr, N.J., Ed.), 415 pp. Editora PUC Goiás, GO, Brazil.

423 Gutiérrez JM, León G, Lomonte B, Angulo Y. 2011. Antivenoms for snakebite envenomings.
424 *Inflammation & Allergy Drug Targets* 10:369–380.

425 Gutiérrez JM, Sanz L, Flores-Díaz M, Figueroa L, Madrigal M, Herrera M, Villalta M, León G,
426 Estrada R, Borges A, Alape-Girón A, Calvete JJ. 2009. Impact of regional variation in
427 *Bothrops asper* venom on the design of antivenoms: Integrating antivenomics and
428 neutralization approaches. *Journal of Proteome Research* 9:564–577.

429 Judge RK, Henry PJ, Mirtschin P, Jelinek G, Wilce JA. 2006. Toxins not neutralized by brown
430 snake antivenom. *Toxicology and Applied Pharmacology* 213:117–125.

431 Kringelum JV, Lundegaard C, Lund O, Nielsen M. 2012. Reliable B cell epitope predictions:

impacts of method development and improved benchmarking. *PLoS Computational Biology* 8:e1002829.

Larsen JE, Lund O, Nielsen M. 2006. Improved method for predicting linear B-cell epitopes. *Immunome Research* 2:1.

Laustsen AH, Lomonte B, Lohse B, Fernández J, Gutiérrez JM. 2015. Unveiling the nature of black mamba (*Dendroaspis polylepis*) venom through venomomics and antivenom immunoprofiling: identification of key toxin targets for antivenom development. *Journal of Proteomics* 119:126–142.

Laustsen AH, Engmark M, Milbo C, Johannesen J, Lomonte B, Gutiérrez JM, Lohse B. 2016. From Fangs to Pharmacology: The Future of Snakebite Envenoming Therapy. *Current Pharmaceutical Design* 22: 5270–5293.

Laustsen AH, Solà M, Jappe EC, Oscoz S, Lauridsen LP, Engmark M. 2016. Biotechnological trends in spider and scorpion antivenom development. *Toxins* 8:1–33.

Leong PK, Fung SY, Tan CH, Sim SM, Tan NH. 2015. Immunological cross-reactivity and neutralization of the principal toxins of *Naja sumatrana* and related cobra venoms by a Thai polyvalent antivenom (Neuro Polyvalent Snake Antivenom). *Acta Tropica* 149:86–93.

Lomonte B, Sasa M, Rey-Suárez P, Bryan W, Gutiérrez JM. 2016a. Venom of the coral snake *Micrurus clarki*: proteomic profile, toxicity, immunological cross-neutralization, and characterization of a three-finger toxin. *Toxins* 8:138.

Lomonte B, Rey-Suárez P, Fernández J, Sasa M, Pla D, Vargas N, Bénard-Valle M, Sanz L, Corrêa-Netto C, Núñez V, Alape-Girón A, Alagón A, Gutiérrez JM, Calvete JJ, 2016b. Venoms of *Micrurus* coral snakes: evolutionary trends in compositional patterns emerging from proteomic analyses. *Toxicon* (submitted).

455 Mardis ER. 2008. Next-generation DNA sequencing methods. *Annual Review of Genomics and*
 456 *Human Genetics* 9:387–402.

457 Nielsen M, Lundegaard C, Lund O, Petersen TN. 2010. CPHmodels-3.0—remote homology
 458 modeling using structure-guided sequence profiles. *Nucleic Acids Research* gkq535.

459 Ownby C, Colberg T. 1990. Comparison of the immunogenicity and antigenic composition of
 460 several venoms of snakes in the family Crotalidae. *Toxicon* 1990 28:189–199.

461 R Core Team (2014). R: A language and environment for statistical computing. R Foundation for
 462 Statistical Computing, Vienna, Austria. URL <http://www.R-project.org/>.

463 Ramos HR, Junqueira-de-Azevedo ILM, Novo JB, Castro K, Duarte CG, Machado-de-Avila RA,
 464 Chavez-Olortegui C, Ho PL. 2016. A heterologous multiepitope DNA prime/recombinant
 465 protein boost immunisation strategy for the development of an antiserum against *Micrurus*
 466 *corallinus* (coral snake) venom. *PLoS Neglected Tropical Diseases* 10:e0004484.

467 Rey-Suárez P, Núñez V, Gutiérrez JM, Lomonte B. 2011. Proteomic and biological
 468 characterization of the venom of the redbellied coral snake, *Micrurus mipartitus* (Elapidae),
 469 from Colombia and Costa Rica. *Journal of Proteomics* 75:655–667.

470 Rey-Suárez P, Stuardo-Florian R, Rostelato-Ferreira S, Saldarriaga M, Núñez V, Rodrigues-
 471 Simioni L, Lomonte B. 2012. Mipartoxin-I, a novel three-finger toxin, is the major
 472 neurotoxic component in the venom of the redbellied coral snake *Micrurus mipartitus*
 473 (Elapidae). *Toxicon* 60:851–863.

474 Rey-Suárez P, Núñez V, Fernández J, Lomonte B. 2016. Integrative characterization of the
 475 venom of the coral snake *Micrurus dumerilii* (Elapidae) from Colombia: proteome,
 476 toxicity, and cross-neutralization by antivenom. *Journal of Proteomics* 136:262–273.

477 Rosso JP, Vargas-Rosso O, Gutiérrez JM, Rochat H, Bougis PE. 1996. Characterization of α -
478 neurotoxin and phospholipase A₂ activities from *Micrurus* venoms. *European Journal of*
479 *Biochemistry* 238:231–239.

480 Sanz L, Pla D, Pérez A, Rodríguez Y, Zavaleta-Martínez A, Salas M, Lomonte B, Calvete JJ.
481 2016. Venomic analysis of the poorly studied desert coral snake, *Micrurus tschudii*
482 *tschudii*, supports the 3FTx/PLA₂ dichotomy across *Micrurus* venoms. *Toxins* 8:178.

483 Schottler WH. 1951. Antigen-antibody relations in the present antivenin production of Brazil.
484 *American Journal of Tropical Medicine and Hygiene* 31:500–509.

485 Stern JN, Yaari G, Vander Heiden JA, Church G, Donahue WF, Hintzen RQ, Huttner AJ, Laman
486 JD, Nagra RM, Nylander A, Pitt D. 2014. B cells populating the multiple sclerosis brain
487 mature in the draining cervical lymph nodes. *Science Translational Medicine* 6:248ra107.

488 Tan KY, Tan CH, Fung SY, Tan NH. 2016. Neutralization of the Principal Toxins from the
489 Venoms of Thai *Naja kaouthia* and Malaysian *Hydrophis schistosus*: Insights into Toxin-
490 Specific Neutralization by Two Different Antivenoms. *Toxins* 8:1–17.

491 Tan CH, Tan KY, Lim SE, Tan NH. 2015. Venomics of the beaked sea snake, *Hydrophis*
492 *schistosus*: A minimalist toxin arsenal and its cross-neutralization by heterologous
493 antivenoms. *Journal of Proteomics* 126:121–30.

494 Tanaka G, Furtado MFD, Portaro FCV, Sant'Anna OA, Tambourgi DV. 2010. Diversity of
495 *Micrurus* snake species related to their venom toxic effects and the prospective of
496 antivenom neutralization. *PLoS Neglected Tropical Diseases* 4:622.

497 Tanaka GD, Sant'Anna OA, Marcelino JR, da Luz ACL, da Rocha MMT, Tambourgi DV. 2016.
498 *Micrurus* snake species: venom immunogenicity, antiserum cross-reactivity and
499 neutralization potential. *Toxicon* 117:59–68.

500 Tsioris K, Gupta NT, Ogunniyi AO, Zimnisky RM, Qian F, Yao Y, Wang X, Stern JN, Chari R,
501 Briggs AW, Clouser CR. 2015. Neutralizing antibodies against West Nile virus identified
502 directly from human B cells by single-cell analysis and next generation sequencing.
503 *Integrative Biology* 7:1587–1597.

504 Vander Heiden JA, Yaari G, Uduman M, Stern JN, O'Connor KC, Hafler DA, Vigneault F,
505 Kleinstein SH. 2014. pRESTO: a toolkit for processing high-throughput sequencing raw
506 reads of lymphocyte receptor repertoires. *Bioinformatics* 30:1930–1932

507 Vergara I, Pedraza-Escalona M, Paniagua D, Restano-Cassulini R, Zamudio F, Batista CV,
508 Possani L., Alagón A. 2014. Eastern coral snake *Micrurus fulvius* venom toxicity in mice is
509 mainly determined by neurotoxic phospholipases A₂. *Journal of Proteomics* 105:295–306.

510 Walsh G. 2014. Biopharmaceutical benchmarks 2014. *Nature biotechnology* 32:992–1000.

511 Warrell DA. 2010. Snake bite. *Lancet* 375:77–88.

512 Warrell DA. 2004. Snakebites in Central and South America: Epidemiology, clinical features
513 and clinical management. In: Campbell JA, Lamar WW, Eds. *The Venomous Reptiles of the*
514 *Western Hemisphere. Vol. II.* Ithaca, New York: Comstock Publishing Associates, Cornell
515 University Press, 709–761.

516 Williams DJ, Gutiérrez JM, Calvete JJ, Wüster W, Ratanabanangkoon K, Paiva O, Brown NI,
517 Casewell NR, Harrison RA, Rowley PD, O'Shea M. 2011. Ending the drought: new
518 strategies for improving the flow of affordable, effective antivenoms in Asia and Africa.
519 *Journal of Proteomics* 74:1735–1767.

520 Wong KY, Tan CH, Tan NH. 2016. Venom and purified toxins of the spectacled cobra (*Naja*
521 *naja*) from Pakistan: Insights into toxicity and antivenom neutralization. *The American*
522 *Journal of Tropical Medicine and Hygiene* 94:1392–1399.

523 Ye J, Ma N, Madden TL, Ostell JM. 2013. IgBLAST: an immunoglobulin variable domain
524 sequence analysis tool. *Nucleic Acids Research* 41:W34-W40.

525