

A new glimpse on Mesozoic zoo-plankton - 150 million years old lobster larvae (#13428)

1

First submission

Please read the **Important notes** below, and the **Review guidance** on the next page. When ready [submit online](#). The manuscript starts on page 3.

Important notes

Editor and deadline

Kenneth De Baets / 8 Oct 2016

Files

Please visit the overview page to [download and review](#) the files not included in this review pdf.

Declarations

No notable declarations are present



Please in full read before you begin

How to review

When ready [submit your review online](#). The review form is divided into 5 sections. Please consider these when composing your review:

- 1. BASIC REPORTING**
- 2. EXPERIMENTAL DESIGN**
- 3. VALIDITY OF THE FINDINGS**
4. General comments
5. Confidential notes to the editor

 You can also annotate this **pdf** and upload it as part of your review

To finish, enter your editorial recommendation (accept, revise or reject) and submit.

BASIC REPORTING

-  Clear, unambiguous, professional English language used throughout.
-  Intro & background to show context. Literature well referenced & relevant.
-  Structure conforms to [PeerJ standard](#), discipline norm, or improved for clarity.
-  Figures are relevant, high quality, well labelled & described.
-  Raw data supplied (See [PeerJ policy](#)).

EXPERIMENTAL DESIGN

-  Original primary research within [Scope of the journal](#).
-  Research question well defined, relevant & meaningful. It is stated how research fills an identified knowledge gap.
-  Rigorous investigation performed to a high technical & ethical standard.
-  Methods described with sufficient detail & information to replicate.

VALIDITY OF THE FINDINGS

-  Impact and novelty not assessed. Negative/inconclusive results accepted. *Meaningful* replication encouraged where rationale & benefit to literature is clearly stated.
-  Data is robust, statistically sound, & controlled.
-  Conclusion well stated, linked to original research question & limited to supporting results.
-  Speculation is welcome, but should be identified as such.

The above is the editorial criteria summary. To view in full visit <https://peerj.com/about/editorial-criteria/>

A new glimpse on Mesozoic zoo-ankton - 150 million years old lobster larvae

Joachim T Haug^{Corresp., 1, 2}, Carolin Haug^{1, 2}

¹ Department of Biology, Ludwig-Maximilians-Universität München, Planegg-Martinsried, Germany

² GeoBio-Center, Ludwig-Maximilians-Universität München, Germany

Corresponding Author: Joachim T Haug

Email address: joachim.haug@palaeo-evo-devo.info

Larvae of malacostracan crustaceans represent a large fraction of modern day zoo-plankton. Plankton is not only a major part of the modern marine ecosystem, but must have played an important role in the ecosystems of the past as well. Unfortunately, our knowledge about plankton composition of the past is still quite limited. As part of the fossil plankton, malacostracan larvae are still a rarity in the fossil record; many types of malacostracan larvae dominating the modern plankton have so far not been found as fossils. Here we report a new type of fossil malacostracan larva, found in the 150 million years old lithographic limestones of southern Germany (Solnhofen Lithographic Limestones). The three rather incomplete specimens mainly preserve the telson. A pronounced middle spine on the posterior edge of these specimens indicates that they are either larval forms of a clawed lobster or of a thalassinidean lobster, or of a closer relative to one of the two groups. The tergo-pleura are drawn out into distinct spines in one specimen, further supporting the interpretation as a larva of a clawed lobster or an early relative. The telson morphology also shows adaptations to a prolonged planktic life style, the latero-posterior edges are drawn out into distinct spines. Similar adaptations are known in larvae of the modern homarid lobster *Nephrops norvegicus*, not necessarily indicating a closer relationship, but convergent life styles. The new finds provide an important new insight into the composition of Mesozoic zoo-plankton and demonstrate the preservation potential of lithographic limestones.

A new glimpse on Mesozoic zoo-plankton – 150 million years old lobster larvae

Joachim T. Haug & Carolin Haug

LMU Munich, Biocenter, Department of Biology II and GeoBio-Center, Großhaderner Str. 2, 82152 Planegg-Martinsried, Germany.

Abstract

Larvae of malacostracan crustaceans represent a large fraction of modern day zoo-plankton. Plankton is not only a major part of the modern marine ecosystem, but must have played an important role in the ecosystems of the past as well. Unfortunately, our knowledge about plankton composition of the past is still quite limited. As part of the fossil plankton, malacostracan larvae are still a rarity in the fossil record; many types of malacostracan larvae dominating the modern plankton have so far not been found as fossils. Here we report a new type of fossil malacostracan larva, found in the 150 million years old lithographic limestones of southern Germany (Solnhofen Lithographic Limestones). The three rather incomplete specimens mainly preserve the telson. A pronounced middle spine on the posterior edge of these specimens indicates that they are either larval forms of a clawed lobster or of a thalassinidean lobster, or of a closer relative to one of the two groups. The tergo-pleura are drawn out into distinct spines in one specimen, further supporting the interpretation as a larva of a clawed lobster or an early relative. The telson morphology also shows adaptations to a prolonged planktic life style, the latero-posterior edges are drawn out into distinct spines. Similar adaptations are known in larvae of the modern homarid lobster *Nephrops norvegicus*, not necessarily indicating a closer relationship, but convergent life styles. The new finds provide an important new insight into the composition of Mesozoic zoo-plankton and demonstrate the preservation potential of lithographic limestones.

Key words: Homarida, Nephropida, Zoea, Solnhofen, Upper Jurassic

Introduction

Plankton describes the entirety of organisms floating in the water column without the ability to actively swim against the current. Plankton has been recognised as an important basis for marine ecosystems in modern seas. Our understanding of changes of composition of plankton are therefore important in modern conservation biology and ecology, but are also of economic importance.

For long-time comparisons of changes in plankton composition, data of fossil plankton is of major interest. Plankton must have been similarly important in the past times as it is today. Unfortunately, our knowledge of plankton in the past is often limited to very specific groups of organisms (examples in, e.g. Lipps 1969; Tappan & Loeblich 1973; Leckie 2009), while such groups dominating the modern plankton often have a very scarce or absent fossil record (e.g. Signor & Vermeij 1994; Rigby & Molsom 2000; Perrier et al. 2015). It is nevertheless already possible to recognise that plankton was composed quite differently in the past (e.g. Nützel & Fryda 2003; Servais et al. 2015).

In modern seas crustaceans are a major part of the zooplankton. Among these we need to especially mention copepod crustaceans and larval stages of malacostracan crustaceans. Copepods are mainly a part of the microplankton, while decapod larvae additionally include forms of meso- and macroplankton.

So how about the fossil record of these groups? Fossil copepods are extremely rare. They are limited to about only half a dozen instances, often only very incomplete or indirect findings (Bennicke 1998; Cressey & Boxshall 1989; Cressey & Patterson 1973; Selden et al. 2010; Palmer 1960; Radwanska & Radwanski 2005; Radwanska & Poiriot 2010). Also malacostracan larvae are still rare in the fossil record, but new forms have have been identified repetitively in recent years

(e.g. Haug et al. 2008, 2011a, 2014, 2015a, b, c), currently limited to the Mesozoic. Despite the rarity, each of these finds marks another important data point for our understanding of plankton in the past.

Due to preservation biases, especially large larval forms appear to be more commonly found as fossils, not necessarily representing the original composition of the fauna, or the true diversity. The most commonly found fossil malacostracan larvae are consequently the super-sized larvae of spiny lobsters and slipper lobsters (with up to 150 mm in the extant fauna). These are represented by at least a dozen different forms (Polz 1984, 1995, 1996; Haug et al. 2013a; Haug & Haug 2016), some known from thousands of individuals (Polz 1971, 1972, 1973) and occurring in at least three lagerstätten (Polz 1984; Pasini & Garassino 2009; Tanaka et al. 2009; Haug et al. 2011). Other more uncommon fossils are also giant larval forms like those of the raptorial mantis shrimps (Haug et al. 2008, 2009, 2010, 2014, 2015b) or the polychelidan lobsters (today only represented by a relic group, mostly deep sea; Haug et al. 2015c; Eiler et al. 2016). Groups of larvae dominating the modern plankton, like larvae of true and false crabs (*Brachyura* and *Anomala*, the latter also called *Anomura*) are very rare on the other hand (Luque 2015; Haug et al. 2015a; Hyžný et al. 2016), as their larvae are significantly smaller and more unlikely to be preserved.

Also, brachyuran and anomalan crustaceans are “still on their way” in the Mesozoic, only diversifying in the later Mesozoic (see discussion in Haug et al. 2015a). It is therefore not to be expected that their larvae were as abundant as in modern oceans. Instead the lobster-like crustaceans, especially polychelidan and clawed lobsters (and their relatives) were dominating parts of the faunas as adults, especially in the mid-Mesozoic, hence the Jurassic. We should therefore expect that larval forms of these groups have represented important parts of the Jurassic plankton. Yet, so far no definite fossils of such planktic larvae have been identified. Possible late larvae of Jurassic polychelidans are late transitory stages, not the truly planktic ones (Eiler & Haug 2016). Some fragmentary specimens have been discussed as possible remains of small malacostracans (Haug et al. 2011a, 2014) and might well be parts of larvae of clawed lobsters, but due to the limitations of preservation this must remain speculative.

Here we can report the first definite finds of zoea-type larvae of clawed lobsters, hence truly planktic larval forms of this group. We discuss the impact on this new finding and how it should influence our strategies for finding further material of fossil malacostracan larvae.

Material and Methods

Material

Three specimens were available for this study (Fig. 1). All originate from the private collection of Roger Frattigiani, Laichingen, and are now deposited in the Staatliches Museum für Naturkunde Stuttgart (SMNS). **NUMBERS WILL BE ADDED LATER** Specimens were originally found in the Birkhof quarry in the Blumenberg area near Eichstätt (Solnhofen Lithographic Limestones). For comparison an extant albuneid zoea from the collections of the Muséum national d'Histoire naturelle Paris was documented.

Documentation method

All specimens were documented on a Keyence BZ-9000 inverse epifluorescence microscope, exploiting the autofluorescence of the fossils (Haug et al. 2011b). Blue-green fluorescence (GFP) was used instead of the commonly used UV fluorescence (e.g., Tischlinger & Arratia 2013). UV fluorescence is often unfortunate due to dust, which shows a very strong fluorescence and is in many cases not removable from the fossils. This is especially true for the comparably high magnifications as applied here (4x objective lens, resulting in 40x magnification). Due to the magnification, depth of field and field of view were limited. To overcome these limitations several stacks of images for several adjacent image details were recorded (see details Haug et al. 2008). Additionally, in some areas where the fluorescence capabilities differed strongly, two stacks were

recorded, each with a different exposure time (Haug et al. 2013b).

Image processing

Stacks of images were fused to sharp images with CombineZP. Fused images were stitched to panoramas using the photomerge function of Adobe Photoshop CS3 or Elements 11. Images of different exposure times were combined into a single evenly illuminated image following the procedure described in Haug et al. (2013b).

Description and interpretation

Specimen 1

Specimen 1 is most complete, but still largely represents fragmentary remains (Fig. 2). The overall colour and texture of the surface already clearly indicate that these fragments are the remains of a crustacean. This is also in concordance with the preserved structures.

Most anteriorly a shield structure is apparent. It appears to be embedded in a dorso-lateral orientation. The anterior rim is drawn out into a distinct but stout rostrum. Along the edge at least two spines are apparent (Fig. 2A). Close to the shield an elongate structure is preserved composed of five elements. Further distal elements are narrower than proximal ones. The structure most likely represents the flagellum of an antennula or antenna. Close to the posterior of the shield a piece of rectangular outline is apparent, most likely representing an isolated element of one the trunk appendages (posterior thoracopods, "pereiopods").

The posterior trunk (pleon) is incompletely preserved and an isolated piece is interpreted as the tergite of pleon segment 3. It is domed, the latero-posterior edges are drawn out into distinct spines. Medially along the posterior rim a posteriorly pointing spine is apparent.

The next posterior preserved piece resembles the tergite of pleon segment 3 in overall morphology and is interpreted as the tergite of pleon segment 5. It is slightly larger than the tergite of pleon segment 3, also the spines are more pronounced.

Articulated to pleon segment 5 is an elongated part posteriorly extending into a more or less triangular structure. This is interpreted as a compound part of pleon segment six and the telson, hence a pleotelson. The posterior part of the pleon segment is partly twisted, and folded onto itself. Still a principle triangular outline (in dorsal view) of the telson is apparent. The posterior rim bears a prominent median spine (Fig. 2B). Left and right to it numerous hair-like structures are apparent. It remains unclear whether these are jointed (true setae) or not (trichomes). The latero-posterior corners are drawn out into elongate distally tapering spines. Close to the base of each large spine, on its median side, slightly laterally from the hair-like structures is a smaller spine, about the same length as the hair-like structures but more massive.

Specimen 2

In comparison to specimen 1, specimen 2 is clearly identified as an isolated telson (Fig. 3A-D). It strongly resembles the posterior part of the pleotelson of specimen 1, but is not twisted and therefore provides additional structural information. The overall size is similar to that of specimen 1. The outline is strongly triangular in dorsal view. The anterior edges, most likely marking the transition to pleon segment 6 are marked by a pair of laterally extending small spines (Fig. 3A).

Postero-lateral edges are drawn out into massive spines, forming a shallow angle. The number of hair-like structures along the posterior rim is 13 per side (Fig. 3D). They are all roughly the same length and the distances between them appear evenly distributed.

Remains of the uropods appear to be preserved left and right to the telson (Fig. 3B, C). Most likely these remains represent the outer, stronger sclerotised edges of the exopods. One small spine appears to be preserved close to the distal end on the posterior surface of the exopod.

Specimen 3

This specimen strongly resembles specimen 2 and is therefore also interpreted as an isolated telson (Fig. 3E-G). Size, principle morphology and number of structures are all similar to specimen 2. Yet, it is not as complete; for example, the posterior edge right to the median spine is broken. It differs from specimen 2 only in the angle of the posterior spines. These form a much narrower angle, as they point less far laterally, but more posteriorly.

Discussion

Systematic interpretation

The specimens are considered conspecific, but differ in the angle between the postero-lateral spines. As all specimens have a similar overall size of the telson region (Fig. 1), it seems unlikely that this difference is a ontogenetic one indicating the presence of several instars. In comparison to modern forms it seems most likely that the spines originally had a certain flexibility and that the difference in angle reflects a preservational difference. We therefore see no possibility to diagnose different forms and see conspecificity as the most parsimonious explanation.

The overall morphology of the specimens indicates that they represent larval malacostracans. An important character in this aspect is the pronounced middle spine of the posterior edge of the telson. In adult malacostracans the telson is often elongate triangular in dorsal view, but with the tip pointing posteriorly, or rectangular to square-shaped in dorsal view. In many larval forms, for example, in decapods, the telson appears forked with a pronounced median indent or, similar to the adults, rectangular, with an evenly armed posterior edge (Martin et al. 2014). Forward pointing triangular to trapezoid/trapezium telson shapes with a pronounced median spine occur in modern forms only in larvae of nephropid or thalassinidean lobsters (Fig. 4). Thalassinidean lobster larvae seem to lack tergo-pleura drawn out into posteriorly pointing spines on the pleon (Dos Santos & González-Gordillo 2004; Pohle & Santana 2014). As such spines are present in one of the fossils and in modern nephropid lobsters (Jorgensen 1925; Wear 1976; Smith 1987; Goy 2014), the fossil larvae most likely represent larvae of clawed lobsters, i.e., nephropids or now extinct relatives of them.

The difficulties with 'clawed lobsters'

Modern clawed lobsters comprise the true lobsters and the reef lobsters, yet quite a number of fossil forms also in principle resembles clawed lobsters, such as erymid or glypheid lobsters (Garassino & Schweigert 2006). The exact relationship of these groups remains still partly unclear. As also (some?) thalassinidean lobster larvae possess a pronounced median spine on the posterior edge of the telson it is possible that this feature characterises a larger group including erymid and glypheid lobsters. Therefore, we can currently not further narrow down the systematic interpretation of the here described larvae. They may represent larval forms of nephropid, erymid or glypheid lobsters or a form closely related to them.

Functional comparison

The telson of the here described larvae clearly shows adaptations to a prolonged life in the pelagic realm. The overall size of the specimens is relatively large for a pelagic larvae (although not as much as in polychelid or achelatan lobsters), and the telson bears pronounced latero-posterior spines.

This distantly resembles the telson of larvae of *Nephrops norvegicus* (scampi). Here also the latero-posterior corners of the telson are strongly drawn out into spines, even more so than in the fossils (Fig. 4). As other nephropids, *N. norvegicus* larvae also possess a median spine on the posterior edge. The telson of *N. norvegicus* larvae is also triangular in dorsal view, yet not as pronounced as in other nephropids.

The overall morphology of the telson of the fossils additionally shows an overall similarity to the larvae of certain false sand crabs (Albuneidae; e.g. Harvey et al. 2014), besides the fact that

these lack the median spine (Figs. 4, 5). Still the overall shape is triangular, the corners are drawn out into spines and the posterior edge is armed. We can assume that the specialised telson of the fossil larvae provided additional bouyancy for the rather large larvae, similar to albuneid larvae.

Significance

The fossil record of arthropod zoo-plankton appears to be very incomplete (Perrier et al. 2015). In general, the overall reconstruction of fossil zoo-plankton seems based largely on estimations, larvae are in such approaches rarely treated in detail (Rigby & Milsom 2000), and, although the fossil record of malacostracan larvae is growing, it appears to be generally regarded as virtually absent (see recent review of Perrier et al. 2015)

Yet, arthropods do play an important role in the modern plankton, especially the larval stages of malacostracans. As pointed out above, clawed lobsters and their relatives, i.e. nephropid, erymid and glypheid lobsters represent an important part of the marine benthic fauna in the Mesozoic. We should therefore expect that their larvae are a major share of the plankton of that time. Finding such larvae is thus important for corroborating this assumption.

As discussed above, the here described specimens show adaptations for prolonged life in the plankton and therefore will represent the upper threshold of size for such larvae. Other larvae of clawed lobsters will be significantly smaller. The larvae furthermore most likely represent only a single specialised form of a wider range of different types of larvae. It has been demonstrated that zoea-type larvae of achelatan lobsters were morphologically more diverse than the larvae of modern forms (Haug et al. 2013a). We can expect that clawed lobster larvae also were morphologically more diverse, possibly similarly diverse to larvae of modern meiruran forms.

The fossils demonstrate that it is possible to find such important components of the plankton and also give an important hint what to look for. Haug et al. (2011a) suggested that some incomplete remains represent isolated shields of larvae, as these might have had a higher preservation potential. The fossils described here show that quite the other end of such a larva, the telson, might also have a high preservation potential. Focussed search for such remains should provide additional insights into the plankton composition of the past.

Acknowledgements

We would like to thank Roger Frattigani, Laichingen for providing the specimens. Gideon T. Haug, Neuried photographed the specimens, for which we are very grateful. Laure Corbari, MNHN Paris, kindly provided access to the extant crustacean collections. J. Matthias Starck, München is thanked for support and discussions. JTH and CH were kindly supported by the German Research Foundation (DFG HA 6300/3-1, DFG HA 7066/3-1). The research visits at MNHN Paris of CH and JTH have been made possible by grants from the European Commission's (FP 6) Integrated Infrastructure Initiative programme SYNTHESYS (FR-TAF-5175, FR-TAF-5181). We thank all people providing free and low-cost software.

References


- Bennike, O. (1998). Fossil egg sacs of *Diaptomus* (Crustaceae: Copepoda) in Late Quaternary lake sediments. *Journal of Paleolimnology*, 19(1), 77-79.
- Cressey, R., & Boxshall, G. (1989). *Kabatarina pattersoni*, a fossil parasitic copepod (Dichelesthidae) from a Lower Cretaceous fish. *Micropaleontology*, 35, 150-167.
- Cressey, R., & Patterson, C. (1973). Fossil parasitic copepods from a Lower Cretaceous fish. *Science*, 180(4092), 1283-1285.

- 261 Dos Santos, A., & González-Gordillo, J. I. (2004). Illustrated keys for the identification of the
262 Pleocyemata (Crustacea: Decapoda) zoeal stages, from the coastal region of south-western Europe.
263 Journal of the Marine Biological Association of the UK, 84(01), 205-227.
264
- 265 Eiler, S.M., & Haug, J.T. (2016). Larval development of fossil polychelidan crustaceans,
266 exemplified by the 150 million years old species *Palaeopentacheles rübenbacheri*. Neues
267 Jahrbuch für Geologie und Paläontologie - Abhandlungen 279(3), 295-310.
268
- 269 Eiler, S.M., Haug, C., & Haug, J.T. (2016). Detailed description of a giant polychelidan eryoneicus-
270 type larva with modern imaging techniques (Eucrustacea, Decapoda, Polychelida). Spixiana 39(1),
271 39-60.
272
- 273 Garassino, A., & Schweigert, G. (2006). The upper Jurassic Solnhofen decapod crustacean fauna:
274 review of the types from old descriptions. (infraorders Astacidea, Thalassinidea, and Palinura).
275 Memorie della Società Italiana di Scienze Naturali e del Museo Civico di Storia naturale in Milano,
276 34 (1), 1-64.
277
- 278 Goy, J. 2014. Astacidea. In: Martin, J. W., Olesen, J., & Høeg, J. T. (eds.), Atlas of Crustacean
279 Larvae, 256-262. The Johns Hopkins University Press, Baltimore.
280
- 281 Harvey A., Boyko, C.B., McLaughlin, P., & Martin J.W. 2014. Martin, J. W., Olesen, J., & Høeg,
282 J. T. (eds.), Atlas of Crustacean Larvae, 283-294. The Johns Hopkins University Press, Baltimore.
283
- 284 Haug, C., Haug, J. T., Waloszek, D., Maas, A., Frattigiani, R., & Liebau, S. 2009. New methods to
285 document fossils from lithographic limestones of southern Germany and Lebanon. Palaeontologia
286 Electronica 12(3); 6T; 12p.
287
- 288 Haug, C., Shannon, K. R., Nyborg, T., & Vega, F. J. (2013b). Isolated mantis shrimp dactyli from
289 the Pliocene of North Carolina and their bearing on the history of Stomatopoda. Bolétin de la
290 Sociedad Geológica Mexicana, 65(2), 273-284.
291
- 292 Haug, C., Wiethase, J. H., & Haug, J. T. 2015b. New records of Mesozoic mantis shrimp larvae and
293 their implications on modern larval traits in stomatopods. Palaeodiversity 8, 121-133.
294
- 295 Haug, J. T., Haug, C., & Ehrlich, M. 2008. First fossil stomatopod larva (Arthropoda: Crustacea) and
296 a new way of documenting Solnhofen fossils (Upper Jurassic, Southern Germany). Palaeodiversity
297 1, 103-109.
298
- 299 Haug, J. T., Ahyong, S., & Haug, C. 2014. Fossil malacostracan larvae. In: Martin, J. W., Olesen, J.,
300 & Høeg, J. T. (eds.), Atlas of Crustacean Larvae, 176-179. The Johns Hopkins University Press,
301 Baltimore.
302
- 303 Haug, J. T., Audo, D., Charbonnier, S., & Haug, C. 2013a. Diversity of developmental patterns in
304 achelate lobsters—today and in the Mesozoic. Development Genes and Evolution 223, 363-373.
305
- 306 Haug, J. T., Martin, J. W., & Haug, C. 2015a. A 150-million-year-old crab larva and its implications
307 for the early rise of brachyuran crabs. Nature Communications 6, art. 6417.
308
- 309 Haug, J. T., Audo, D., Haug, C., Abi Saad, P., Petit, G., & Charbonnier, S. 2015c. Unique occurrence
310 of polychelidan lobster larvae in the fossil record and its evolutionary implications. Gondwana
311 Research 28, 869-874.
312

- 313 Haug, J. T., Haug, C., Kutschera, V., Mayer, G., Maas, A., Liebau, S., Castellani, C., Wolfram, U.,
314 Clarkson, E. N. K. & Waloszek, D. 2011b. Autofluorescence imaging, an excellent tool for
315 comparative morphology. *Journal of Microscopy* 244, 259–272.
- 316
- 317 Haug, J. T., Haug, C., Waloszek, D. & Schweigert, G. 2011a. The importance of lithographic
318 limestones for revealing ontogenies in fossil crustaceans. *Swiss Journal of Geosciences* 104,
319 Supplement 1, S85–S98.
- 320
- 321 Haug, J. T. & Haug, C. 2016. “Intermetamorphic” developmental stages in 150 million-year-old
322 achelatan lobsters – The case of the species *tenera* Oppel, 1862. *Arthropod Structure &
323 Development* 45, 108–121.
- 324
- 325 Hyžný, M., Haug, C., & Haug, J. T. (2016). *Mesoprosopon triasinum* from the Triassic of Austria
326 revisited: The oldest eumalacostracan larva known to date and its significance for interpreting fossil
327 cycloids. *Gondwana Research*, 37, 86-97.
- 328
- 329 Jorgensen, O. M. (1925). The early stages of *Nephrops norvegicus*, from the Northumberland
330 Plankton, together with a note on the post-larval development of *Homarus vulgaris*. *Journal of the
331 Marine Biological Association of the United Kingdom (New Series)*, 13(04), 870-879.
- 332
- 333 Knight, M. D. (1970). The larval development of *Lepidopa myops* Stimpson, (Decapoda,
334 Albuneidae) reared in the laboratory, and the zoeal stages of another species of 
- 335
- 336 Leckie, R. M. (2009). Seeking a better life in the plankton. *Proceedings of the National Academy of
337 Sciences*, 106(34), 14183-14184.
- 338
- 339 Lipps, J. H. (1970). Plankton evolution. *Evolution*  2.
- 340
- 341 Luque, J. 2015 A puzzling frog crab (Crustacea: Decapoda: Brachyura) from the Early Cretaceous
342 Santana Group of Brazil: frog first or crab first? *J. Syst. Paleontol* 13, 153–166.
- 343
- 344 Martin, J. W., Olesen, J., & Høeg, J. T. (Eds.). (2014). *Atlas of crustacean larvae*. The Johns
345 Hopkins University Press, Baltimore.
- 346
- 347 Nützel, A., & Frýda, J. (2003). Paleozoic plankton revolution: evidence from early gastropod
348 ontogeny. *Geology*, 31(9), 829-831.
- 349
- 350 Palmer, A. R. (1960). Miocene copepods from the Mojave desert, California. *Journal of
351 Paleontology*,  7-452.
- 352
- 353 Pasini, G., & Garassino, A. (2009). A new phyllosoma form (Decapoda,? Palinuridae) from the Late
354 Cretaceous (Cenomanian) of Lebanon. *Atti della Società italiana di scienze naturali e del museo
355 civico di storia naturale di Milano*, 150(1), 21-28.
- 356
- 357 Perrier, V., Williams, M., & Siveter, D. J. (2015). The fossil record and palaeoenvironmental
358 significance of marine arthropod zooplankton. *Earth-Science Reviews*, 146, 146-162.
- 359
- 360 Pohle, G. & Santana W. 2014. Gebiidea and Axiidea (=Thalassinidea). In: Martin, J. W., Olesen, J.,
361 & Høeg, J. T. (eds.), *Atlas of Crustacean Larvae*, 263–271. The Johns Hopkins University Press,
362 Baltimore.
- 363
- 364 Polz, H. (1971). Eine weitere Phyllosoma-Larve aus den Solnhofener Plattenkalken. *Neues*

- 365 Jahrbuch für Geologie und Paläontologie, Monatshefte, 8, 474-488.
- 366
- 367 Polz, H.. 1972. Entwicklungsstadien bei fossilen Phyllosomen (Form A) aus den Solnhofener
- 368 Plattenkalken. Neues Jahrbuch für Geologie und Paläontologie, Monatshefte 11, 678–689.
- 369
- 370 Polz, H. 1973. Entwicklungsstadien bei fossilen Phyllosomen (Form B) aus den Solnhofener
- 371 Plattenkalken. Neues Jahrbuch für Geologie und Paläontologie, Monatshefte 5, 284–296.
- 372
- 373 Polz, H.1984. Krebslarven aus den Solnhofener Plattenkalken. Archaeopteryx 2, 30–40.
- 374
- 375 Polz, H. 1995. Ein außergewöhnliches Jugendstadium eines palinuriden Krebses aus den
- 376 Solnhofener Plattenkalken. Archaeopteryx 13, 67–74.
- 377
- 378 Polz, H.1996. Eine Form-C-Krebslarve mit erhaltenem Kopfschild (Crustacea, Decapoda,
- 379 Palinuroidea) aus den Solnhofener Plattenkalken. Archaeopteryx 14, 43–50.
- 380
- 381 Rötzer, M. A., & Haug, J. T. (2015). Larval development of the European lobster and how small
- 382 heterochronic shifts lead to a more pronounced metamorphosis. International Journal of Zoology,
- 383 art. 345172.
- 384
- 385 Rigby, S., & Milsom, C. V. (2000). Origins, evolution, and diversification of zooplankton. Annual
- 386 Review of Ecology and Systematics, 31, 293-313.
- 387
- 388 Radwańska, U. & Poirot, E. (2010). Copepod-infested Bathonian (Middle jurassic) echinoids from
- 389 northern France. Acta Geologica Polonica, 60(4), 549-555.
- 390
- 391 Radwańska, U., & Radwański, A. (2005). Myzostomid and copepod infestation of Jurassic
- 392 echinoderms: A general approach, some new occurrences, and/or re-interpretation of previous
- 393 reports. Acta Geologica Polonica, 55, 109-130.
- 394
- 395 Selden, P. A., Huys, R., Stephenson, M. H., Heward, A. P., & Taylor, P. N. (2010). Crustaceans from
- 396 bitumen clast in Carboniferous glacial diamictite extend fossil record of copepods. Nature
- 397 Communications, 1, 50.
- 398
- 399 Servais, T., Perrier, V., Danelian, T., Klug, C., Martin, R., Munnecke, A., Nowak, H., Nützel, A.,
- 400 Vendenbroucke, T.R.A., Williams, M. & Rasmussen, C. M. (2015). The onset of the ‘Ordovician
- 401 Plankton Revolution’ in the late Cambrian. Palaeogeography, Palaeoclimatology, Palaeoecology,
- 402 458, 12–28.
- 403
- 404 Signor, P. W., & Vermeij, G. J. (1994). The plankton and the benthos: origins and early history of an
- 405 evolving relationship. Paleobiology, 20(03), 297-319.
- 406
- 407 [Smith, R. S. M. (1987). The biology of larval and juvenile *Nephrops norvegicus* (L.) in the Firth of
- 408 Clyde (Doctoral dissertation, University of Glasgow).]
- 409
- 410 Stuck, K. C., & Truesdale, F. M. (1986). Larval and early postlarval development of *Lepidopa*
- 411 *benedicti* Schmitt, 1935 (Anomura: Albuneidae) reared in the laboratory. Journal of Crustacean
- 412 Biology, 6(1), 89-110.
- 413
- 414 Tanaka, G., Smith, R. J., Siveter, D. J., & Parker, A. R. (2009). Three-dimensionally preserved
- 415 decapod larval compound eyes from the Cretaceous Santana Formation of Brazil. Zoological
- 416 Science, 26(12), 846-850.

- 417
418 Tappan, H., & Loeblich, A. R. (1973). Evolution of the oceanic plankton. *Earth-Science Reviews*,
419 9(3), 207-240.
420
421 Tischlinger, H., & Arratia, G. (2013). Ultraviolet light as a tool for investigating Mesozoic fishes,
422 with a focus on the ichthyofauna of the Solnhofen archipelago. *Mesozoic Fishes*, 5, 549-560.
423
424 Wear, R. G. (1976). Studies on the larval development of *Metanephrops challenger* (Balss, 1914)
425 (Decapoda, Nephropidae). *Crustaceana*, 30(2), 113-122.
426
427
428



431 Fig. 1. Complete material of larval specimens, composite-fluorescence micrographs. All three
 432 specimens shown in the same scale to allow relative size comparison. Matrix digitally removed for
 433 clarity.
 434




Fig. 2. Specimen 1, composite-fluorescence micrographs. A. Overview of entire specimen, although incomplete showing general organisation; arrows mark small spines. B. Close-up on posterior rim of telson; arrowheads point to small hair-like structures or setae. Abbreviations: fl? = possible flagellum of antennula or antenna; hs = head shield; ms = median spine; pls = postero-lateral spine; ps3, 5 = pleon segment 3, 5; pt = pleotelson; sp = spine; ur = uropod.






Fig. 3. Specimens 2 and 3, each representing an isolated posterior part of a pleotelson (= telson) composite-fluorescence micrographs. A–D. Specimen 2. A. Overview; arrows mark small spines. B. Close-up on left lateral rim of telson; arrow points to small spine. C. Close-up on right lateral rim of telson. D. Close-up on left posterior rim of telson. E–G. Specimen 3. E. Overview; arrow marks small spines. F. Close-up on right lateral rim of telson. G. Close-up on left posterior rim of telson. Abbreviations: ms = median spine; pls = postero-lateral spine; set = hair-like structures or setae; sp = spine; ur = uropod.



462 Fig. 4. Comparison of the new fossil larva with extant forms. New fossil larva as restoration in
463 dorsal aspect; each of the others as an isolated telson in dorsal view. *Homarus gammarus* (European
464 lobster); zoea III simplified from Rötzer & Haug (2015). *Nephrops norvegicus* (scampi,
465 Kaisergranat); zoea III combined from Smith (1987) and Jorgensen (1925). *Metanephrops*
466 *challengerii*; late zoea simplified from Wear (1976). *Axius stirhynchus*; late zoea simplified from
467 Dos Santos & González-Gordillo (2005, fig. 2F). Undetermined albuneid larva; late zoea simplified
468 from Fig. 5.

469
470
471



475 Fig. 5. Extant larva of an albuneid meiuran for functional comparison; cross-polarised macro
476 images. Note the long postero-lateral spines on shield and telson and the triangular telson. A.
477 Lateral view on left side. B. Dorsal view on head shield. C. Posterior view on anterior region; dorsal
478 view on posterior pleon and pleotelson; arrows mark spines. D. Detail of the telson. Abbreviations:
479 hs = head shield; pls = postero-lateral spine; pt = pleotelson; set = hair-like structures or setae.

480
481
482