

# Is coral richness related to community resistance to and recovery from disturbance?

More diverse communities are thought to be more stable – the diversity-stability hypothesis – due to increased resistance to and recovery from disturbances. For example, high diversity can make the presence of resilient or fast growing species and key facilitations among species more likely. How natural geographic biodiversity patterns and changes in biodiversity due to human activities mediate community-level disturbance dynamics is largely unknown, especially in diverse systems. For example, few studies have explored the role of diversity in tropical marine communities, especially at large scales. We tested the diversity-stability hypothesis by asking whether coral richness is related to resistance to and recovery from disturbances including storms, predator outbreaks, and coral bleaching on tropical coral reefs. We synthesized the results of 41 field studies conducted on 82 reefs, documenting changes in coral cover due to disturbance, across a global gradient of coral richness. Our results indicate that coral reefs in more species rich regions were marginally less resistant to disturbance and did not recover more quickly. Coral community resistance was also highly dependent on pre-disturbance coral cover, probably due in part to the sensitivity of fast-growing and often dominant plating acroporid corals to disturbance. Our results suggest that coral communities in biodiverse regions such as the western Pacific may not be more resistant to natural and anthropogenic disturbances. Further analyses controlling for disturbance intensity and other drivers of coral loss and recovery could improve our understanding of the influence of diversity on community stability in coral reef ecosystems.

2 Stacy Y. Zhang<sup>1</sup>, Speare, Kelly E.<sup>1</sup>; Long, Zachary T.<sup>2</sup>; McKeever, Kimberly A.<sup>1</sup>; Gyoerkoe,  
3 Megan<sup>1</sup>; Ramus, Aaron P.<sup>1,2</sup>; Mohorn, Zach<sup>1</sup>; Akins, Kelsey L.<sup>1</sup>; Hambridge, Sarah M.<sup>1</sup>; Graham,  
4 Nicholas A. J.<sup>3</sup>; Nash, Kirsty L.<sup>3</sup>; Selig, Elizabeth, R.<sup>4</sup>, Bruno, John F.<sup>1\*</sup>

5 <sup>1</sup>Department of Biology, The University of North Carolina at Chapel Hill, Chapel Hill, NC  
6 27599-3300 USA

7 <sup>2</sup>Department of Biology and Marine Biology, University of North Carolina at Wilmington,  
8 Wilmington, NC 28403 USA

9 <sup>3</sup>Australian Research Council Centre of Excellence for Coral Reef Studies, James Cook  
10 University, Townsville, QLD 4811, Australia

11 <sup>4</sup>The Betty and Gordon Moore Center for Ecosystem Science and Economics, Conservation  
12 International, 2011 Crystal Drive, Suite 500, Arlington, VA 22202

13 \* Corresponding author: jbruno@unc.edu

## 14 Introduction

15 A large body of recent and classical ecological research has focused on how biodiversity  
16 influences the stability of communities ([May 1973](#); [McCann 2000](#); [Pimm 1984](#); Hughes and  
17 Stachowicz 2004; [Steiner et al. 2006](#); Cardinale et al. 2012). Despite this interest, there is still no  
18 consensus on whether or why diversity influences stability. This may be due in part to the  
19 multitude of ways stability has been defined ([Grimm & Wissel 1997](#); [McCann 2000](#); [Pimm](#)  
20 [1984](#)). Here, we define community stability as the ability to maintain a given state regardless of  
21 perturbation, invasion, or extinction ([McCann 2000](#)). Mechanistically, stability is made up of two  
22 components: resistance – the degree to which a community changes in response to a disturbance,  
23 and recovery or “resilience” – the rate of return to pre-disturbance conditions ([Grimm & Wissel](#)  
24 [1997](#); [Ives & Carpenter 2007](#); [May 1973](#); [McCann 2000](#)).

25 Species richness (the number of species within an area) is thought to influence community  
26 stability via several mechanisms ([Ives et al. 2005](#)). For example, species rich communities are  
27 more likely, due to chance alone, to include greater functional diversity, and redundancy of  
28 function within functional groups – the insurance hypothesis ([McNaughton 1977](#); [Tilman &](#)  
29 [Downing 1994](#)). The inclusion of species that can tolerate or benefit from disturbance could  
30 increase community resistance ([McCann 2000](#); [Tilman & Downing 1994](#); [Yachi & Loreau 1999](#)).  
31 Additionally, positive feedbacks among species via facilitation may be more common in diverse  
32 communities, which may enhance post-disturbance recovery ([Naeem 1998](#); [Naeem & Li 1997](#);  
33 [Yachi & Loreau 1999](#)).

34 Understanding the role of species richness in community stability has been a focus of a  
35 number of field studies and meta-analyses ([Duffy et al. 2001](#); [Ives & Carpenter 2007](#); [Stachowicz](#)  
36 [et al. 2007](#); [Stachowicz et al. 2008](#); Cardinale et al. 2012). Some experimental studies in  
37 terrestrial plant communities have found a positive relationship between species richness and  
38 stability ([Ives & Carpenter 2007](#); [Tilman & Downing 1994](#)). For example, more diverse plant

communities appear to be more resistant to disturbances including drought ([Tilman & Downing 1994](#)). Manipulations of marine plant and invertebrate community richness have found that diversity is positively related to resistance and recovery ([Stachowicz et al. 2007](#)). Likewise, [Boyer et al. \(2009\)](#) experimentally demonstrated that species-rich algal communities were more resistant to disturbances than species-poor communities in estuarine ecosystems of eastern North Carolina. One of the few large scale tests of the diversity-stability concept was a study of boreal forest communities in north-western Quebec ([Grandpre et al. 1997](#)), which found that greater species-richness conferred more resistance to changes in community composition and abundance following a disturbance.

We tested the hypothesis that species richness is related to community stability (resistance and recovery) in coral reef ecosystems. Coral communities are affected by a large range of natural and anthropogenic disturbances including storms, disease outbreaks, coral bleaching due to ocean warming, dynamite fishing, and predator outbreaks (e.g., *Acanthaster planci* or Crown-of-Thorns Starfish (COTS)). A recent study of coral-community recovery found that disturbance type, e.g., “physical” vs. “biological” disturbances, and other reef characteristics had little effect on recovery rate; although, recovery was influenced by post-disturbance coral cover, geographic region, and reef management ([Graham et al. 2011](#)). We expanded on this work by asking whether coral richness is related to resistance to and or recovery from disturbances at a large spatio-temporal scale using coral cover as a metric of reef state. We hypothesized that coral communities in species rich regions would exhibit greater resistance and recovery in response to disturbances. We synthesized the results of 41 field studies documenting post-disturbance changes in coral cover across a natural gradient of coral richness.

## Methods

### *Data Sources and Study Selection*

Our analyses were based on an expanded version of the coral cover change database compiled by Graham et al. (2011). We added 18 new studies (35 additional reefs), resulting in a total of 82 reefs (Text S1). Twenty-eight studies were used in the decline analyses, and 30 were included in the recovery analyses, with many studies providing both resistance and recovery data. We identified and added studies that quantitatively documented community responses following disturbances, both losses and recoveries, in percentage of benthos covered by living coral species on subtidal reefs (0.5 – 40 m depth, median survey depth = 10 m). If component values were in a graphical format, ImageJ software was used to extract values for analysis. Survey locations were categorized into four regions: Eastern Pacific, Western Pacific, Indian Ocean, and Caribbean (Fig. S1). In addition to browsing relevant coral reef and ecology journals, we also used online academic search tools (e.g., Google Scholar, ISI Web of Science) to search for published peer-reviewed sources of coral-cover change data using the search terms: “disturbance,” “coral cover,” “*Acanthaster planci*,” “bleaching,” “storm,” “cyclone,” “hurricane,” “resistance,” “resilience,” “recovery,” and “richness.”

Studies had to meet four criteria to be included in the present analysis ([Graham et al. 2011](#)). 1) A clear description of sampling methods. Most surveys were performed using a variation of the line transect method where a 10–30 m transect is placed on the reef and coral cover is estimated either *in situ* or from digital images or videos. Although manta tow data can be unbiased if extensive training is involved ([Miller & Müller 1999](#)), all manta tow data was excluded because we were unable to verify the accuracy of the majority of manta tow surveys performed. 2) Documented changes in coral cover due to one of three types of disturbance: *Acanthaster planci* outbreaks, coral bleaching, and storm (hurricanes and cyclones) impacts. Other disturbances such as dynamite fishing and ship groundings were not included due to a lack of studies and the small scale of disturbance. 3) Changes in hard coral cover were documented for resistance: pre-disturbance (initial cover) and immediately after disturbance (remnant cover); or

88 recovery: remnant cover and peak recorded cover post-disturbance (peak recovery). Where peak  
 89 recovery was assessed, post-disturbance cover was monitored for at least 3 years following  
 90 disturbance. 4) Coral cover decreased at least 10% relative to pre-disturbance cover due to  
 91 disturbance (Eqn. 1).

$$92 \quad \text{Initial cover} - \text{Post disturbance cover} \geq 0.1(\text{Initial cover}) \quad (\text{Eqn. 1})$$

93 We excluded studies in which the change in coral cover was < 10% relative to initial coral cover  
 94 because such relatively small year to year changes cannot be reliably distinguished from random  
 95 noise, e.g., stochastic variation and sampling error. Benthic survey methodologies rarely are able  
 96 to detect changes in cover < 5–10%. Graham 2011 et al. used a similar justification for excluding  
 97 such studies: “the initial drop in coral cover as a result of the disturbance exceeded 10%. Studies  
 98 recording lower initial mortality were excluded as small disturbances may have negligible impact  
 99 on hard coral cover (e.g., Edmunds 2002). Furthermore, inter-annual variation in coral cover can  
 100 be ~5% prior to a disturbance (e.g., Halford et al. 2004). A 10% relative decline reflects a sudden  
 101 change in benthic cover in response to a pulse disturbance.” However, in contrast to Graham et  
 102 al. 2011, we chose a 10% relative decline as a cutoff to account for disturbances that cause little  
 103 absolute change but may in fact have a proportionally significant impact to reefs, particularly  
 104 those with low levels of initial coral cover.

105 Predicted coral richness values in our analyses are geographic estimates based on coral  
 106 species range distributions ([Veron 2000](#); [Veron et al. 2011](#)) and have been used by previous  
 107 studies (e.g., [Roberts et al. \(2002\)](#); [Veron et al. \(2009\)](#)). Maps were originally vector data, but we  
 108 gridded them to a resolution of 50 km. The coral richness predictions that result from this  
 109 database are relatively coarse-grained, however, given the strong association between local and  
 110 regional coral richness (Karlson et al. 2004), our richness values likely correspond in a relative  
 111 sense to local richness at a site.

## Statistical Analysis

We used the R programming environment ([R Core Team 2012](#)) to perform two separate sets of regression analyses to test whether predicted coral richness was related to coral community resistance to and recovery from disturbance. Variance inflation factors (VIFs) were calculated for each pair of predictors to test for collinearity in both sets of analyses. The initial model for decline included predicted coral richness, initial (pre-disturbance) coral cover, maximum survey depth, region, and disturbance type as predictors. Initial predictors for recovery rate included predicted coral richness, post-disturbance coral cover, maximum survey depth, region, and disturbance type. We incrementally and individually removed non-significant predictors from the models using backwards stepwise elimination. Akaike Information Criteria (AIC) was used in model selection to assess which model provided the best fit. Decline (resistance) analyses were performed using square root transformed net absolute coral loss, and in the recovery analysis, we used log transformed recovery rate, calculated as the change in coral cover, generally an increase, divided by the number of years until peak coral cover was observed (Eqn 2).

$$\log\left(\frac{\text{peak recorded recovery} - \text{post-disturbance cover}}{\text{peak recovery year} - \text{post-disturbance year}}\right) \quad \text{Eqn. 2}$$

In cases where recovery was not observed and reefs continued to decline i.e. negative recovery, recovery rate was equated to zero. Transformations were based on the distribution of the raw data. All models were also performed on raw data, the results of which did not differ qualitatively from those based on transformed data. The core database (including all response and predictor variables), the R code, and model outputs were made available for editors and reviewers upon manuscript resubmission and will remain available for future readers. We did this to enable our analysis to be reproduced, to be as open as possible about our procedures, and to facilitate future analysis based on our database.

## Results

There was substantial variation among sites in predicted coral richness (24-552 species, Fig. 1), coral cover loss (Fig. 1), and recovery rate (Fig. 2). The full linear regression model testing the null hypothesis that coral species richness was not related to (sqrt transformed) post-disturbance coral cover loss (decline) found that all covariates (including richness;  $p=0.79$ ) except pre-disturbance coral cover (blcc,  $p<0.0001$ ) were unrelated to decline (Fig. S2, Table 1, Model 1, Adj.  $R^2=0.63$ , see all model results and R code in the Statistical Supplement). A simpler model (Model 2, Adj.  $R^2=0.60$ ) that included three covariates (depth, richness, and blcc) with a similar fit indicated that the richness effect was marginally significant ( $p=0.03$ ). There was no relationship (Adj.  $R^2=0.01$ ,  $p=0.50$ ) between predicted coral richness and initial cover, thus it is unlikely that collinearity between these two covariates led to the putative effects of richness on resistance. When we removed sites (two potential outliers) with extreme decline and high diversity, the relationship between relative decline and predicted species richness was still significant (in the reduced model with three covariates).

The full recovery model (Table 1, Model 4 in the Statistical Supplement, Adj.  $R^2=0.26$ ) was the best fit and indicated that none of the covariates were related to recovery rate. Predicted coral richness in particular appeared a very poor predictor of coral recovery rate ( $p=0.56$ ).

## Discussion

We did not detect an effect of region, disturbance type or depth on absolute loss in coral cover due to disturbance (Table 1, Statistical Supplement). There was a weak positive relationship between predicted species richness and coral loss (Fig. 1). However, a richness effect ( $p=0.03$ ) was only detected in the reduced model (Model 2), from which maximum depth, region, and



disturbance type were removed (Statistical Supplement). In the full model with all five covariates (Model 1), the richness effect was not significant ( $p=0.79$ ).

Concordant with several previous studies ([Graham et al. 2008](#); [Selig & Bruno 2010](#); [Selig et al. 2012](#)), reefs with greater initial cover lost the most coral following disturbance (Fig. 1, Statistical Supplement). This could be due in part to the dominance of high cover reefs in the western Pacific and Indian ocean by plating acroporid corals, that are especially sensitive to both physical and biological disturbances ([Darling et al. 2012](#)), i.e., not due to a cover effect *per se*, but instead to the relationship, both among reefs and regions, between coral cover and species composition. Acroporids are among the most sensitive coral species to most of the disturbances known to cause declines in coral cover, e.g., thermal stress, disease, predation, and physical disturbance ([Bruno et al. 2001](#); [Darling et al. 2012](#); [Willis et al. 2004](#)). Therefore reefs dominated by plating and branching acroporids – the highest cover reefs – are likely to be especially susceptible to disturbance. Although high taxonomic richness may confer greater functional diversity and redundancy, this also implies a greater number of specialists that may not be well adapted to disturbance or stressful conditions ([Dobzhansky 1950](#); [Schluter 2000](#)) and may partially explain our results. In fact, Côté and Darling (2010) argued that managing for coral cover could counter-intuitively make reefs more susceptible to disturbances including climate change.

Unlike several previous experimental studies in marine systems ([Allison 2004](#); [Reusch et al. 2005](#); [Williams 2001](#)), our mensurative study found that the rate of post-disturbance recovery was not significantly related to taxonomic richness (Table 1, Fig. 2). The large observed variance in recovery rate among sites and studies (up to 12.5 % per year) appears to be due to factors not included in our model and unrelated to estimated species richness.

Graham et al. (2011) reported that coral recovery varied somewhat among regions, with the highest observed recovery rates in the western Pacific, the most species rich region, (Roberts

et al. 2002). Despite that finding, in our analysis coral richness was a poor predictor of recovery rate (Fig. 2); although, the low richness reefs of the eastern Pacific recovered slowly, recovery of Caribbean reefs with similar richness was greater and similar to that seen in the more diverse western Pacific and Indian ocean (Fig. 2). This finding could be due in part to a publication biases towards studies that document recovery and the fact that we searched for studies documenting acute disturbances, whereas chronic perturbations are also responsible for Caribbean coral losses. Furthermore, studies conducted in the eastern Pacific and included in our analysis were mostly conducted on reefs dominated by slow-growing, massive *Porites*. Given the current state of the Caribbean (Bruno et al. 2009) there is no doubt that coral cover on most Caribbean reefs has not recovered from various natural and anthropogenic disturbances over the last several decades (Roff and Mumby 2012). Hence, our results suggest that low diversity Caribbean reefs have the potential to recover (in terms of coral cover) quite rapidly (e.g., Idjadi et al. 2006), but for a variety of reasons they rarely do (Roff and Mumby 2012).

The majority of reefs in our study were monitored for less than eight years and few were monitored long enough to recover fully (Fig. S3), given observed rates in recovery. Of the 23 reefs for which we had both initial post-disturbance and peak-year coral cover, only five (all monitored for at least 11 years after disturbance) recovered to pre-disturbance levels of cover. If recovery (coral cover increase) is indeed non-linear over time, we could have seen different results, e.g., a significant richness effect if the component studies were conducted longer. To a degree, our results could also be an artifact of the loss of Acroporid corals from Caribbean reefs in the 1980s, due primarily to disease but in some locations to storms and predation ([Aronson & Precht 2006](#)). Our earliest Caribbean recovery study began in 1987 ([Edmunds 2002](#)), after Acroporid populations had crashed regionally ([Aronson & Precht 2006](#)). Had *Acropora cervicornis* and *A. palmata* been present on Caribbean reefs when the datasets used in our analyses were collected, we might have seen more rapid recovery on low diversity reefs because

207 these corals were amongst the fastest growing corals in the Caribbean region. However, we may  
 208 also have seen even less resistance to disturbance due to their high coral cover and vulnerability  
 209 to various stressors.

210       It is possible that systematic variation in disturbance intensity could obscure a richness  
 211 effect on coral community resistance. For example, disturbance intensity may have co-varied  
 212 with coral richness or peak coral cover. Although quantification of the intensity of disturbance is  
 213 beyond the scope of this study, it would, at least in theory, be possible. Our study also addresses a  
 214 very large scale richness gradient but ignores smaller scale variation, e.g., among reefs or zones  
 215 ([Huston 1985](#)). Furthermore, we used predicted richness values from known biogeographic  
 216 ranges (Veron 2000, Veron et al. 2011), which should reflect patterns of richness at this scale, but  
 217 can make richness and region hard to tease apart and cannot account for local site level  
 218 differences due to factors such as reef zone, disturbance history, micro-climates. Diversity of fish  
 219 assemblages may also be critical for coral recovery dynamics (Bellwood et al. 2004), which was  
 220 not considered in this study. Future analyses with empirical diversity data of corals and fish,  
 221 controlling for disturbance intensity and other drivers of coral loss and recovery, could improve  
 222 our understanding of the influence of diversity on community stability in coral reef ecosystems.

# Literature Cited

- Allison G. 2004. The influence of species diversity and stress intensity on community resistance and resilience. *Ecological Monographs* 74:117-134.
- Aronson RB, and Precht WF. 2001. White-band disease and the changing face of Caribbean coral reefs. *Hydrobiologia* 460:25-38.
- Aronson RB, and Precht WF. 2006. Conservation, precaution, and Caribbean reefs. *Coral Reefs* 25:441-450.
- Bellwood, DR, TP Hughes, C. Folke, M. Nyström. 2004. Confronting the coral reef crisis. *Nature* 429: 827-833
- Boyer KE, Kertesz JS, and Bruno JF. 2009. Environmental context influences the effects of species richness on productivity but not stability of marine macroalgal communities. *Oikos* 118:1062-1072.
- Bruno JF, Siddon CE, Witman JD, Colin PL, and Toscano MA. 2001. El Niño related coral bleaching in Palau, Western Caroline Islands. *Coral Reefs* 20:127-136.
- Bruno, J. F., and E. R. Selig. 2007. Regional decline of coral cover in the Indo-Pacific: timing, extent, and subregional comparisons. *PLoS One*:e711.
- Bruno JF, Sweatman H, Precht WF, Selig ER, and Schutte VGW. 2009. Assessing evidence of phase shifts from coral to macroalgal dominance on coral reefs. *Ecology* 90:1478–1484.
- Cardinale BJ, Duffy JE, Gonzalez A, Hooper DU, Perrings C, Venail P, Narwani A, Mace GM, Tilman D et al. 2012. Biodiversity loss and its imp[act on humanity. *Nature* 486: 59-67.
- Côté IM, and Darling ES. 2010. Rethinking ecosystem resilience in the face of climate change. *PLoS Biol* 8:e1000438.
- Darling ES, Alvarez-Filip L, Oliver TA, McClanahan TR, and Côté IM. 2012. Evaluating life-history strategies of reef corals from species traits. *Ecology Letters* 15:1378-1386.
- Dobzhansky T. 1950. Evolution in the tropics. *American Scientist* 38:209-221.

- 248 Duffy JE, Macdonald KS, Rhode JM, and Parker JD. 2001. Grazer diversity, functional  
249 redundancy, and productivity in seagrass beds: an experimental test. *Ecology* 82:2417-  
250 2434.
- 251 Edmunds PJ. 2002. Long-term dynamics of coral reefs in St. John, US Virgin Islands. *Coral*  
252 *Reefs* 21:357-367.
- 253 Graham N, Nash K, and Kool J. 2011. Coral reef recovery dynamics in a changing world. *Coral*  
254 *Reefs* 30:283-294.
- 255 Graham NAJ, McClanahan TR, MacNeil MA, Wilson SK, Polunin NVC, Jennings S, Chabanet P,  
256 Clark S, Spalding MD, Letourneur Y et al. . 2008. Climate Warming, Marine Protected  
257 Areas and the Ocean-Scale Integrity of Coral Reef Ecosystems. *PLoS ONE* 3:e3039.
- 258 Grandpre D, L., Bergeron, and Y. 1997. *Diversity and stability of understorey communities*  
259 *following disturbance in the southern boreal forest*. Oxford, Royaume-Uni: Blackwell.
- 260 Grimm V, and Wissel C. 1997. Babel, or the Ecological Stability Discussions: An Inventory and  
261 Analysis of Terminology and a Guide for Avoiding Confusion. *Oecologia* 109:323-334.
- 262 Halford A, Cheal AJ, Ryan D, Williams DMcB. 2004. Resilience to large-scale disturbance in  
263 coral and fish assemblages on the Great Barrier Reef. *Ecology* 85:1892–1905.
- 264 Hughes, A. R., and J. J. Stachowicz. 2004. Genetic diversity enhances the resistance of a seagrass  
265 ecosystem to disturbance. *Proceedings of the National Academy of Sciences* 101:8998–9002.
- 266 Huston MA. 1985. Patterns of species diversity on coral reefs. *Annual Review of Ecology and*  
267 *Systematics* 16:149-177.
- 268 Huston MA. 1997. Hidden treatments in ecological experiments: re-evaluating the ecosystem  
269 function of biodiversity. *Oecologia* 110:449-460.
- 270 Idjadi, JA, SC Lee, JF Bruno, WF Precht, L Allen-Requa, and PJ Edmunds. 2006. Rapid phase-  
271 shift reversal on a Jamaican coral reef. *Coral Reefs* 25:209-211

- 272 Ives AR, Cardinale BJ, and Snyder WE. 2005. A synthesis of subdisciplines: predator-prey  
273 interactions, and biodiversity and ecosystem functioning. *Ecology Letters* 8:102-116.
- 274 Ives AR, and Carpenter SR. 2007. Stability and Diversity of Ecosystems. *Science* 317:58-62.
- 275 Karlson, RH, HV Cornell, TP Hughes. 2004. Coral communities are regionally enriched along an  
276 oceanic biodiversity gradient. *Nature* 429, 867.
- 277 May RM. 1973. *Stability and complexity in model ecosystems*. Princeton: Princeton University  
278 Press.
- 279 McCann KS. 2000. The diversity-stability debate. *Nature* 405:228-233.
- 280 McNaughton SJ. 1977. Diversity and Stability of Ecological Communities: A Comment on the  
281 Role of Empiricism in Ecology. *The American Naturalist* 111:515-525.
- 282 Miller I, and Müller R. 1999. Validity and reproducibility of benthic cover estimates made during  
283 broadscale surveys of coral reefs by manta tow. *Coral Reefs* 18:353-356.
- 284 Naeem S. 1998. Species Redundancy and Ecosystem Reliability. *Conservation Biology* 12:39-45.
- 285 Naeem S, and Li S. 1997. Biodiversity enhances ecosystem reliability. *Nature* 390:507-509.
- 286 Pimm SL. 1984. The complexity and stability of ecosystems. *Nature* 307:321-326.
- 287 R Core Team. 2012. R: A Language and Environment for Statistical Computing. Vienna, Austria:  
288 R Foundation for Statistical Computing.
- 289 Reusch TBH, Ehlers A, Hammerli A, and Worm B. 2005. Ecosystem recovery after climatic  
290 extremes enhanced by genotypic diversity. *PNAS* 102:2826-  
291 2831.
- 292 Roberts CM, McClean CJ, Veron JEN, Hawkins JP, Allen GR, McAllister DE, Mittermeier CG,  
293 Schueler FW, Spalding M, Wells F et al. . 2002. Marine biodiversity hotspots and  
294 conservation priorities for tropical reefs. *Science* 295:1280-1284.
- 295 Roff G, and Mumby PJ. 2012. Global disparity in the resilience of coral reefs. *Trends Ecol Evol*  
296 27:404-413.

- 297 Schluter D. 2000. *The ecology of adaptive radiation*: OUP Oxford.
- 298 Selig ER, and Bruno JF. 2010. A global analysis of the effectiveness of marine protected areas in  
299 preventing coral loss. *PLoS ONE* 5:e9278.
- 300 Selig ER, Casey KS, and Bruno JF. 2012. Temperature-driven coral decline: the role of marine  
301 protected areas. *Global Change Biology* 18:1561-1570.
- 302 Stachowicz JJ, Bruno JF, and Duffy JE. 2007. Understanding the effects of marine biodiversity  
303 on communities and ecosystems. *Annual Review of Ecology, Evolution, and Systematics*  
304 38:739-766.
- 305 Stachowicz JJ, Graham M, Bracken MES, and Szoboszlai AI. 2008. Diversity Enhances Cover  
306 and Stability of Seaweed Assemblages: The Role of Heterogeneity and Time. *Ecology*  
307 89:3008-3019.
- 308 Steiner CF, Long ZT, Krumins JA, and Morin PJ. 2006. Population and community resilience in  
309 multitrophic communities. *Ecology* 87:996-1007.
- 310 Tilman D, and Downing JA. 1994. Biodiversity and stability in grasslands. *Nature* 367:363-365.
- 311 Veron J. 2000. Corals of the world. *Australian Institute of Marine Science Townsville, Australia*.
- 312 Veron J, Devantier LM, Turak E, Green AL, Kininmonth S, Stafford-Smith M, and Peterson N.  
313 2009. Delineating the coral triangle. *Galaxea, Journal of Coral Reef Studies* 11:91-100.
- 314 Veron JEN. 1986. *Corals of Australia and the Indo-Pacific*. Sydney: Angus and Robertson.
- 315 Veron JT, E; DeVantier, LM; Stafford-Smith, MG; Kininmonth, S 2011. Coral Geographic.  
316 Version 1.1. ed: Aust. Inst. Mar. Sci.
- 317 Williams SL. 2001. Reduced genetic diversity in eelgrass transplantations affects both population  
318 growth and individual fitness. *Ecological Applications* 11:1472-1488.
- 319 Willis BL, Page CA, and Dinsdale EA. 2004. Coral disease on the Great Barrier Reef. In:  
320 Rosenberg E, and Loya Y, eds. *Coral Health and Disease*. Berlin: Springer-Verlag, 69-  
321 104.

322 Yachi S, and Loreau M. 1999. Biodiversity and ecosystem productivity in a fluctuating  
 323 environment: The insurance hypothesis. *Proceedings of the National Academy of Sciences*  
 324 96:1463-1468.



# Table 1 (on next page)

Table 1

Linear models performed in analyses.

# Linear model description -Decline

Raw data	AIC	R <sup>2</sup>	p-value
Richness + maxdepth + disturbance + region + blcc	351.94	0.71	4.04e-07
Richness + maxdepth + blcc	349.46	0.65	5.56e-09
Richness + blcc	414.92	0.60	2.73e-10
Richness + maxdepth + disturbance + region + blcc + maxdepth:disturbance interaction	354.75	0.71	2.85e-06

## Square root transformed

Richness + maxdepth + disturbance + region + blcc <sup>†</sup>	148.77	0.70	5.62e-07
• Intercept			0.05
• Richness			0.78
• Max depth			0.95
• Disturbance – COTS			0.69
• Disturbance – Storms			0.51
• Region – E. Pacific			0.17
• Region – Indian Ocean			0.67
• Region – W. Pacific			0.98
• BLCC			3.72e-8
Richness + maxdepth + blcc * <sup>†</sup>	147.60	0.63	1.50e-08
• Intercept			0.15
• Richness			0.03
• Max depth			0.61
• BLCC			1.02e-08
Richness + blcc	174.83	0.59	6.82e-10
Richness + maxdepth + disturbance + region + blcc + maxdepth:disturbance interaction	150.30	0.72	2.49e-06

# Linear model description - Recovery

Raw data	AIC	R <sup>2</sup>	p-value
Richness + maxdepth + disturbance + region + pdcc	264.05	0.27	0.03
Richness + maxdepth	264.11	0.12	0.04
Richness	274.63	0.062	0.07

## Log transformed

Richness + maxdepth + disturbance + region + pdcc*	104.77	0.38	0.04e-1
• Intercept			3.95e-06
• Richness			0.56
• Max depth			0.10
• Disturbance – COTS			0.35
• Disturbance – Storms			0.21
• Region – E. Pacific			0.25

• Region – Indian Ocean			0.46
• Region – W. Pacific			0.46
• PDCC			0.13
Richness + maxdepth	109.21	0.15	0.01
Richness	115.60	0.11	0.01

Table 1. Linear models performed in analyses.


\* indicates best fit model for coral cover decline and recovery based upon lowest AIC. p-values for best fit model factors listed below model description.

† indicates best fit decline models which did not differ significantly based upon AIC.

# Figure 1

Coral community richness and resistance to disturbance.

3-D scatter plot of the relationship between coral community richness, coral loss (net change in absolute percent cover in response to disturbance) and initial coral cover. See Table 1 for results of statistical analysis. Dashed lines represent the regression plane, and point coloration represents region. Refer to the Statistical Supplement for a version of this figure plotted with the response variable square root transformed.



## Figure 2

Coral community richness and recovery from disturbance.

Scatter plot of the relationship between the post-disturbance recovery of absolute coral cover and estimated coral species richness. Refer to the Statistical Supplement for a version of this figure plotted with the response variable log transformed.

