

Disturbed flow in an aquatic environment may create a sensory refuge for aggregated prey

Asa Johannesen ^{Corresp., 1, 2}, Alison M Dunn ², Lesley J Morrell ³

¹ Nesvik Marine Centre, Fiskaaling, Hvalvik, Faroe Islands

² School of Biology, University of Leeds, Leeds, United Kingdom

³ School of Environmental Sciences, University of Hull, Hull, United Kingdom

Corresponding Author: Asa Johannesen

Email address: asajoh@fiskaaling.fo

Predators use olfactory cues moved within water and air to locate prey. Because prey aggregations may produce more cue and be easier to detect, predation could limit aggregation size. However, disturbance in the flow may diminish the reliability of odour as a prey cue, impeding predator foraging success and efficiency. We explore how different cue concentrations (as a proxy for prey group size) affect risk to prey by fish predators in disturbed (more turbulent or mixed) and non-disturbed (less mixed) flowing water. We find that increasing odour cue concentration increases predation risk and disturbing the flow reduces predation risk. At high cue concentration fish were able to locate the cue source in both disturbed and non-disturbed flow, but at medium concentrations, predators only located the cue source more often than expected by chance in non-disturbed flow. This suggests that objects disturbing flow provide a sensory refuge allowing prey to form larger groups, but that group sizes may be limited by level of disturbance to the flow.

1 **Disturbed flow in an aquatic environment may create a sensory refuge for aggregated prey**

2

3 Asa Johannesen^{1,2}, Alison M. Dunn¹ & Lesley J. Morrell³

4 ¹School of Biology, University of Leeds, Leeds, United Kingdom

5 ²Nesvik Marine Centre, Fiskaaling, Hvalvik, Faroe Islands

6 ³School of Environmental Sciences, University of Hull, Hull, United Kingdom

7

8 Correspondence:

9 Nesvík Marine Centre, Fiskaaling (Aquaculture Research Station of the Faroe Islands), við Áir,
10 FO-430 Hvalvík, Faroe Islands

11

12 asajoh@fiskaaling.fo

13 +298214764

Abstract

Predators use olfactory cues moved within water and air to locate prey. Because prey aggregations may produce more cue and be easier to detect, predation could limit aggregation size. However, disturbance in the flow may diminish the reliability of odour as a prey cue, impeding predator foraging success and efficiency. We explore how different cue concentrations (as a proxy for prey group size) affect risk to prey by fish predators in disturbed (more turbulent or mixed) and non-disturbed (less mixed) flowing water. We find that increasing odour cue concentration increases predation risk and disturbing the flow reduces predation risk. At high cue concentration fish were able to locate the cue source in both disturbed and non-disturbed flow, but at medium concentrations, predators only located the cue source more often than expected by chance in non-disturbed flow. This suggests that objects disturbing flow provide a sensory refuge allowing prey to form larger groups, but that group sizes may be limited by level of disturbance to the flow.

Introduction

To avoid predation, animals use strategies from visual crypsis (Jackson *et al.*, 2005) to increased vigilance in groups (Krause and Ruxton, 2002). In cases where visual interactions between predators and prey are limited, cues such as sound (Obrist *et al.*, 1993) or detection of electric fields (Kajiura and Holland, 2002) are used instead. Olfaction is a key sense used in prey detection and location. Olfactory predators such as crustaceans (Gomez and Atema, 1996; Weissburg and Zimmer-Faust, 1993), fish (Nevitt, 1991), and molluscs (Ferner and Weissburg,

2005) can successfully track odour plumes from prey to their source. Animals use many sensing strategies, including time differences in bilateral odour detection (Gardiner and Atema, 2010), time-averaging of odour concentrations (Wilson and Weissburg, 2012), and making simultaneous comparisons of odour concentration (Page *et al.*, 2011; Vergassola *et al.*, 2007). To avoid such detection, prey may try to limit the amount of olfactory cue that they release or otherwise make it difficult for predators to detect them (Ruxton, 2009).

To reduce the risk from predators that hunt using vision, prey can group together to increase the time taken for a hunting predator to locate them. This is known as the encounter-dilution effect (Wrona and Dixon, 1991) and favours grouping as an anti-predator strategy in response to visual predators (Jackson *et al.*, 2005; Riipi *et al.*, 2001). However, if increasing group size makes prey increasingly easier for olfactory predators to find (Kunin, 1999), grouping may be counter-productive.

Larger or more numerous animals release more odour cue, eliciting a stronger reaction in the receiver (Hawkins *et al.*, 2007; Kusch *et al.*, 2004). When animals aggregate, the odour cues released interact, increasing the size and concentration of the odour plume (Villermanux and Duplat, 2003), which allows receivers to more readily track the plume to its source (Wilson and Weissburg, 2012). Grouping benefits prey avoiding olfactory predators in still water (Johannesen *et al.*, 2014). However, water movement provides a directional cue to the prey, so olfactory cues are more easily taken advantage of in flowing rather than still water (Løkkeborg, 1998).

In a review of olfactory detection distance in insects, Andersson *et al.* (2013) indicate that the increase in detection with increasing size of the source is likely to be asymptotic, although theoretical work indicates accelerating detectability may also be possible (Treisman, 1975). If the

risk of predation increases too much with group size, aggregation would be counterproductive in species that cannot otherwise defend themselves. Here, we explore this question from the perspective of three-spine sticklebacks (*Gasterosteus aculeatus*) locating odour sources of differing concentration (as a proxy for prey group size (Hill and Weissburg, 2014; Schneider *et al.*, 2014) - but see discussion) in flowing water, to test the hypothesis that increasing prey (bloodworm) cue concentration increases the risk to prey in flowing water.

Chemical cues are often detected in pulses because currents, turbulence, and other types of disturbed flow create patches of cue (Finelli and Pentcheff, 1999; Weissburg and Zimmer-Faust, 1993; Zimmer-Faust *et al.*, 1995), which may create ‘sensory refuges’ (Weissburg and Zimmer-Faust, 1993). When prey are in these refuges, predators may be less- or unable to detect them (Ferner and Weissburg, 2005), while prey may still be able to detect predators as back eddies carry odour cues ‘upstream’ (Dahl *et al.*, 1998). Prey animals occupying a sensory refuge would benefit from the reduced predation success, leading to aggregation of prey in refuge areas. If animals aggregate in sensory refuges, the sensory refuge may counteract the increased risk of detection due to larger group size. We repeat our experiment in disturbed flow to examine the additional hypothesis that disturbed flow reduces the risk to prey relative to undisturbed flow, as it creates sensory refuges (Weissburg and Zimmer-Faust, 1993). Our aim is to provide an initial exploration of the possible impact of prey aggregation and flow conditions on the detection of prey by a foraging fish.

Methods

Experimental species, transportation and housing

Two hundred three-spine sticklebacks *Gasterosteus aculeatus* (4-5cm total body length) were caught in a pond in Saltfleet, Lincolnshire in November, 2011 (53°25'59.55"N, 0°10'49.41"E) and transported in fish bags (3-5 fish per litre) to aquarium facilities in Leeds (3 hour journey). Sticklebacks can detect and locate prey using non-visual cues in still water (Johannesen *et al.*, 2012), but also occupy flowing water environments and may be able to track odour plumes to their source. Fish were housed in grey fibreglass tanks (0.5x0.5x1.0m) with gravel substrate, plastic plants, plant pots and mechanical filters. Light regime was 10:14 hours light:dark, temperature was $14 \pm 2^{\circ}\text{C}$ and fish were fed daily on defrosted frozen bloodworm. Fish were kept for six months to one year for experimentation prior to release where caught in agreement with the Home Office and Defra.

Procedure

Trials were carried out in a flow-through rectangular tank (40 cm by 53 cm, water depth 9 cm, flow velocity 3cm s^{-1}) based on a Y maze design (Ward *et al.*, 2011) (Fig 1). The stem of the maze was 33 cm in length including a 20 cm 'release zone' with a removable barrier. Each 'arm' was 20 cm long. Conditioned water was pumped from a header tank into the maze, entered the maze over a horizontal barrier in both arms of the Y, and passed through a collimator to create an even water flow ("undisturbed flow"). Flow characteristics were not measured, but pilot trials using dye indicated that the odour cue would move through the water evenly. Water left the flume through 3 mesh-covered exit holes evenly spaced across the base of the stem of the Y, and was not re-circulated. Trials were observed from behind a screen via a webcam to reduce disturbance to the fish.

Olfactory cues were created using filtered macerated bloodworm (cue concentrations; low: 5 g l⁻¹ medium: 10 g l⁻¹ and high: 20 g l⁻¹). Although spatial distribution of aggregated prey affects odour plume composition, we use prey cue concentration as a proxy for size or number of prey. This is because the interacting odour plumes from multiple prey individuals will increase cue concentration (Hawkins *et al.*, 2007; Villermanux and Duplat, 2003) and while adding the spatial factor to this experiment would be more realistic, it would also complicate the experiment. Cues were delivered to the maze using two peristaltic pumps at a rate of 10 ml min⁻¹ (source diameter 4mm, velocity 1.3cm s⁻¹). In each trial, the olfactory cue entered at one arm of the maze, and a conditioned water control (containing red food dye to copy the tint of the bloodworm cue water) entered at the other at the same rate. Cue side was allocated at random to control for side preference. After the trial, the maze was emptied and refilled with conditioned water to remove olfactory cues from the previous trial.

Sticklebacks were placed individually into the release zone and allowed to acclimatise until they resumed normal behaviour (start – stop swimming at moderate speed, five minutes minimum). Following acclimatisation, the water inlet pump was switched on and ran for two minutes (to stabilise flow) before cue pumps were turned on. The behaviour of the test fish was monitored until it had visited both sides of the stem of the Y (two minutes minimum) and the barrier was raised using a pulley system. The fish was allowed five minutes to reach the top of one arm of the Y, where its choice (cue or control) was recorded. The time taken for the fish to acclimatise (begin swimming) and the time taken to reach the top of the chosen arm were also recorded. Fish were excluded from the experiment if they did not resume normal behaviour in the release zone (N=23), did not visit both sides of the stem of the Y within 5 minutes (N = 8

123 fish) or did not make a choice ($N = 6$). Final sample sizes in undisturbed flow were: low: $N = 16$,
124 medium: $N = 16$, high: $N = 16$).

125 We subsequently investigated the effect of disturbed flow on stickleback choice in the
126 maze. Three cylinders were placed in each arm of the Y maze to create downstream disruption to
127 the flow (see Fig 1). Visualisation of the flow using food dye indicated that the cylinders caused
128 the odour plumes to split and disperse, and that the plumes appeared qualitatively different to
129 those in the experiment with no disturbance to the flow. Methods were the same as in the
130 previous experiment, but investigated only two cue concentrations: medium and high. The low
131 concentration was not used as the first experiment indicated that fish did not show a preference at
132 this concentration (see results). Eight fish were excluded from this experiment, giving final
133 sample sizes of $N = 17$ for medium cue concentration and $N = 17$ for high cue concentration in
134 disturbed flow. Each fish was used only once in the experiments, and different fish were used in
135 the two flow conditions to avoid any learning effects. The total sample size for both experiments
136 was $N = 82$.

137 *Analysis*

138 Data were analysed using R v 2.13.0 (R Core Team, 2013). Cox proportional hazards
139 survival models were used to analyse fish time to acclimatise and time to choose (survival
140 package in R; (Therneau and Lumley, 2011)). Survival models are highly flexible and useful for
141 time-to type data, especially when data do not follow a Gaussian distribution and contain
142 censored times (Therneau and Grambsch, 2000). Preference for prey cue was tested using
143 binomial exact tests against an expectation of random (0.5). A correction for false discovery rate
144 was performed (Benjamini and Hochberg, 1995) on the tests to control for multiple comparisons

and the adjusted p values are given in the results section. A binomial GLM with choice as dependent variable (cue, no cue) and cue concentration and flow type as independent variables was used to test the effect of treatments on choice.

Ethical statement

As experiments with fish fall outside of the remit of the University of Leeds Ethical Board and no licensed procedures were used, this study was not subject to ethical review.

However, laboratory experiments were carried out in accordance with University of Leeds guidelines and in agreement with Home Office licensed technical staff at the animal facility. Great care was taken to ensure optimal welfare for all fish involved in this study.

Results

Fish tested in the disturbed flow condition took less time to acclimatise than those in the non-disturbed flow condition (coxph: Chi-squared = 25.81, df = 1, $p < 0.001$), but there was no effect of cue concentration or flow type on time to choose once acclimatised (coxph: Chi-squared = 6.22, df = 5, $p = 0.29$).

In the ‘undisturbed flow’ condition, fish selected the cue arm over the control arm at medium ($N = 13/16$, $P(\text{success}) = 0.8125$, $p = 0.035$) and high ($N = 15/16$, $P(\text{success}) = 0.938$, $p = 0.003$) cue concentrations, but not at the low cue concentration ($N = 11/16$, $P(\text{success}) = 0.688$, $p = 0.26$). When flow disturbance was added, fish preferentially selected the cue arm at high (N

165 = 14/17, $P(\text{success}) = 0.824$, $p = 0.033$) but not medium ($N = 10/17$, $P(\text{success}) = 0.588$, $p =$
 166 0.629) cue concentrations (Fig 2).

167 In a test of the effects of cue concentration and flow regime, cue concentration
 168 significantly affected choice (Binomial GLM; $z = 2.235$, $N = 82$, $p = 0.025$) while the effect of
 169 water regime fell short of significance (Binomial GLM; $z = -1.814$, $N = 82$, $p = 0.070$).

170

171 Discussion

172 Our results suggest that in a Y-maze with olfactory cue presented in one arm only, fish
 173 predators can successfully choose the arm containing the cue more often than expected by
 174 chance if the concentration of the cue is high enough. At the low cue concentration, fish did not
 175 choose the cue arm more often than the non-cue arm. Interpreted in the context of our question of
 176 how increasing group size affects detection, this suggests that grouping in prey (increased cue
 177 concentration) may increase risk from olfactory predators. Adding objects to the maze to disturb
 178 the flow (i.e. create more turbulent mixing) decreased the number of successful choices,
 179 particularly at the medium cue concentration, suggesting that ‘sensory refuges’ created by
 180 disturbed flow (Weissburg and Zimmer-Faust, 1993) allow larger groups to form by countering
 181 the increased risk of detection. However, at higher cue concentrations, the benefit gained from
 182 flow disturbance may decrease, and larger groups would be more easily detected. While we did
 183 not compare multiple disturbance levels, we suggest that the level of disturbance may also
 184 influence the detection of prey cues of different concentrations.

185 We used concentration as a proxy for group size, following previous work (Schneider *et*
 186 *al.*, 2014), although the effect of group size on odour plumes is more complex. When individuals

group together, they produce a greater number of odour filaments (Monismith *et al.*, 1990; Wilson and Weissburg, 2012) that cover a wider area (Webster and Weissburg, 2001). When prey are spaced sufficiently far apart, individual odour filaments do not interact and would not result in an increase in cue concentration. In that case, the risk of detection by predators who are unable to detect any individual filament, but would detect a more concentrated one, remains low regardless of group size. However, in cases of tightly aggregated prey, filaments are likely to interact and increase the time-averaged concentration (Villermanux and Duplat, 2003) used by some predators to track plumes (Ferner and Weissburg, 2005) as well as filament concentration (Villermanux and Duplat, 2003). Therefore, tightly packed prey increase risk of detection with group size, particularly by predators who would not have detected a single individual's odour plume.

Concentration may reflect a number of other features of the prey landscape in addition to group size in tight aggregations. As cues become more diluted over distance and time, cue concentration may signal distance to prey (but see Bytheway *et al.*, 2013), or if larger individuals release more cue, the concentration could signal size of prey. Thus, predators could move towards higher concentrations because they represent better value predation opportunities, rather than because they are easier to detect. However, while the results in this experiment could reflect a perceived value of cue in the fish rather than simple detection, there is no indication of this in the latency to choose a cue arm in our data.

While turbulence or other disturbance to flow can cause odour plumes to break up (Webster and Weissburg, 2001) it can also act to mix the plumes and dilute the cue to background levels with only intermittent spikes (Webster and Weissburg, 2009) that may not be worth exploring. Either mechanism would act to make tracking the cue to the source more

difficult for the predator (Robinson *et al.*, 2011), although this may depend on the predator's sensing strategy and sensitivity (Ferner and Weissburg, 2005). Our observations with food dye suggest that the plume in disturbed flow split primarily into two meandering plumes. Assuming a fish was only exposed to one arm of the split plume, the decreased amount of cue could mean fewer or smaller prey, greater distance to prey, or the concentration might be below a detection threshold. A meandering plume will, in addition to the perceived lower reward, be more difficult to track for fish and other filament samplers, making the effort greater. While a time-averaging predator may be able to compensate for meandering plumes (Page *et al.*, 2011), a diluted and split plume signals lower reward, so inhabiting turbulent water can also benefit tightly packed prey hiding from a time-averager. However, as sticklebacks most likely do not use time-averaging sensing strategies (Nevitt, 1991; Webster and Weissburg, 2001), this has not been explicitly tested here.

Our study did not investigate the fluid mechanics and transport of olfactory cue in the different flow regimes, focusing instead on the response of the predator. Thus, we cannot speculate on the sensory mechanism, motivation, or features of water flow and cue transport that cause the different behaviours shown by our fish predators. However, the end result for the prey remains the same. If an olfactory prey cue is highly concentrated, indicating either great reward (many or large prey) or easy reward (close proximity) a predator is more likely to pursue that cue. Conversely, if the olfactory prey cue plume is somehow broken down, indicating small reward (few or small prey) or difficult reward (long distance, a plume that is difficult to track) a fish predator is less likely to pursue that cue. In the context of our question regarding aggregation, this suggests prey are able to aggregate into larger groups by taking advantage of a sensory refuge and either fooling the predator into thinking they are not worth the effort (small

reward/high effort) or decreasing the cue to avoid detection. Individuals in such aggregations would in turn benefit from greater survival chance if found due to other benefits of grouping, such as predator satiation.

The study of anti-predator aggregation has primarily focused on predators that use vision to detect their prey (Ioannou and Krause, 2008), while the effect of olfactory predators on the evolution of aggregation is less well understood. Our work suggests that group size may interact with environmental parameters, and that the evolution of grouping in response to olfactory predators may be dependent on the flow environment. However, further work is needed to fully investigate the relationship between grouping prey, detection by predators, and environmental conditions. Prey are known to aggregate in streams (Rasmussen and Downing, 1988), but aggregation decisions may depend on factors other than risk from olfactory predators, including foraging opportunities, flow speed and risk from predators relying on other sensory modalities (Ioannou and Krause, 2008; Krause and Ruxton, 2002). Experimental manipulation and characterisation of flow regimes and the response of predators and prey may help disentangle the interacting effects of group size, flow regime and aggregation in response to other resources.

Data Accessibility Section

The data presented in this paper can be found on Figshare (<http://dx.doi.org/10.6084/m9.figshare.985515>).

References

Andersson, P., Löfstedt, C. and Hambäck, P. A. (2013) How insects sense olfactory patches - the spatial scaling of olfactory information, *Oikos*, 122 (7), pp. 1009–1016. DOI:10.1111/j.1600-

- 255 0706.2012.00037.x.
- 256 Benjamini, Y. and Hochberg, Y. (1995) Benjamini Y, Hochberg Y. Controlling the false
257 discovery rate: a practical and powerful approach to multiple testing, *Journal of the Royal*
258 *Statistical Society B*, 57 (1), pp. 289–300. DOI:10.2307/2346101.
- 259 Bytheway, J. P., Carthey, A. J. R. and Banks, P. B. (2013) Risk vs. reward: how predators and
260 prey respond to aging olfactory cues, *Behavioral Ecology and Sociobiology*, 67 (5), pp. 715–725.
261 DOI:10.1007/s00265-013-1494-9.
- 262 Dahl, J., Anders Nilsson, P. and Pettersson, L. B. (1998) Against the flow: chemical detection of
263 downstream predators in running waters, *Proceedings of the Royal Society B: Biological*
264 *Sciences*, 265 (1403), pp. 1339–1344. DOI:10.1098/rspb.1998.0439.
- 265 Ferner, M. C. and Weissburg, M. J. (2005) Slow-moving predatory gastropods track prey odors
266 in fast and turbulent flow., *The Journal of Experimental Biology*, 208 (5), pp. 809–819.
267 DOI:10.1242/jeb.01438.
- 268 Finelli, C. M. and Pentcheff, N. (1999) Odor transport in turbulent flows: constraints on animal
269 navigation, *Limnology and ...*, 44 (4), pp. 1056–1071.
- 270 Gardiner, J. M. and Atema, J. (2010) The function of bilateral odor arrival time differences in
271 olfactory orientation of sharks., *Current Biology : CB*, 20 (13), pp. 1187–91.
272 DOI:10.1016/j.cub.2010.04.053.
- 273 Gomez, G. and Atema, J. (1996) Temporal resolution in olfaction: stimulus integration time of
274 lobster chemoreceptor cells, *The Journal of Experimental Biology*, 199 (Pt 8), pp. 1771–9.
- 275 Hawkins, L., Magurran, a and Armstrong, J. (2007) Innate abilities to distinguish between
276 predator species and cue concentration in Atlantic salmon, *Animal Behaviour*, 73 (6), pp. 1051–
277 1057. DOI:10.1016/j.anbehav.2006.08.011.
- 278 Hill, J. M. and Weissburg, M. J. (2014) Crabs interpret the threat of predator body size and
279 biomass via cue concentration and diet, *Animal Behaviour*, 92, pp. 117–123.
280 DOI:10.1016/j.anbehav.2014.03.025.
- 281 Ioannou, C. C. and Krause, J. (2008) Searching for prey: the effects of group size and number,
282 *Animal Behaviour*, 75 (4), pp. 1383–1388.
- 283 Jackson, A. L., Brown, S., Sherratt, T. N. and Ruxton, G. D. (2005) The effects of group size,
284 shape and composition on ease of detection of cryptic prey, *Behaviour*, 142 (6), pp. 811–826.
285 DOI:10.1163/1568539054729105.
- 286 Johannesen, A., Dunn, A. M. and Morrell, L. J. (2012) Olfactory cue use by three-spined
287 sticklebacks foraging in turbid water: prey detection or prey location?, *Animal Behaviour*, 84 (1),
288 pp. 151–158. DOI:10.1016/j.anbehav.2012.04.024.
- 289 Johannesen, A., Dunn, A. M. and Morrell, L. J. (2014) Prey aggregation is an effective olfactory
290 predator avoidance strategy, *PeerJ*, 2 (May), pp. e408. DOI:10.7717/peerj.408.
- 291 Kajiura, S. M. and Holland, K. N. (2002) Electroreception in juvenile scalloped hammerhead and
292 sandbar sharks, *The Journal of Experimental Biology*, 205 (23), pp. 3609–3621.
- 293 Krause, J. and Ruxton, G. D. (2002) *Living in Groups*. Vol. I. Oxford University Press.

- 294 Kunin, W. E. (1999) Patterns of herbivore incidence on experimental arrays and field
295 populations of ragwort, *Senecio jacobaea*, *Oikos*, 84 (3), pp. 515–525.
- 296 Kusch, R. C., Mirza, R. S. and Chivers, D. P. (2004) Making sense of predator scents:
297 investigating the sophistication of predator assessment abilities of fathead minnows, *Behavioral*
298 *Ecology and Sociobiology*, 55 (6), pp. 551–555. DOI:10.1007/s00265-003-0743-8.
- 299 Løkkeborg, S. (1998) Feeding behaviour of cod, *Gadus morhua*: activity rhythm and chemically
300 mediated food search, *Animal Behaviour*, 56 (2), pp. 371–378. DOI:10.1006/anbe.1998.0772.
- 301 Monismith, S. G., Koseff, J. R., Thompson, J. K., O’Riordan, C. A. and Nepf, H. M. (1990) A
302 study of model bivalve siphonal currents., *Limnology and Oceanography*, 35 (3), pp. 680–696.
- 303 Nevitt, G. A. (1991) Do fish sniff? A new mechanism of olfactory sampling in pleuronectid
304 flounders., *The Journal of Experimental Biology*, 157, pp. 1–18.
- 305 Obrist, M. K., Fenton, M. B., Eger, J. L. and Schlegel, P. a (1993) What ears do for bats: a
306 comparative study of pinna sound pressure transformation in chiroptera., *The Journal of*
307 *Experimental Biology*, 180, pp. 119–52.
- 308 Page, J. L., Dickman, B. D., Webster, D. R. and Weissburg, M. J. (2011) Staying the course:
309 chemical signal spatial properties and concentration mediate cross-stream motion in turbulent
310 plumes., *The Journal of Experimental Biology*, 214 (Pt 9), pp. 1513–1522.
311 DOI:10.1242/jeb.049304.
- 312 R Core Team (2013) *R: A language and environment for statistical computing*. Vienna, Austria:
313 R Foundation for Statistical Computing.
- 314 Rasmussen, J. B. and Downing, J. A. (1988) The spatial response of chironomid larvae to the
315 predatory leech *Nephelopsis obscura*, *The American Naturalist*, 131 (1), pp. 14–21.
316 DOI:10.1086/284770.
- 317 Riipi, M., Alatalo, R. V, Lindström, L. and Mappes, J. (2001) Multiple benefits of
318 gregariousness cover detectability costs in aposematic aggregations., *Nature*, 413 (6855), pp.
319 512–4. DOI:10.1038/35097061.
- 320 Robinson, E. M., Smee, D. L. and Trussell, G. C. (2011) Green crab (*Carcinus maenas*) foraging
321 efficiency reduced by fast flows., *PloS One*, 6 (6), pp. e21025.
322 DOI:10.1371/journal.pone.0021025.
- 323 Ruxton, G. D. (2009) Non-visual crypsis: a review of the empirical evidence for camouflage to
324 senses other than vision., *Philosophical Transactions of the Royal Society of London. Series B,*
325 *Biological Sciences*, 364 (1516), pp. 549–57. DOI:10.1098/rstb.2008.0228.
- 326 Schneider, J., Worischka, S., Hellmann, C., Benndorf, J. and Winkelmann, C. (2014) Flexibility
327 in feeding periodicity of a grazing mayfly in response to different concentrations of benthivorous
328 fish, *Limnologica - Ecology and Management of Inland Waters*, 45, pp. 24–32.
329 DOI:10.1016/j.limno.2013.10.002.
- 330 Therneau, T. M. and Grambsch, P. M. (2000) *Modeling Survival Data: Extending the Cox*
331 *Model*. London: Springer.
- 332 Therneau, T. M. and Lumley, T. (2011) survival: Survival analysis, including penalised

333 likelihood: R package version 2.36-5.

334 Treisman, M. (1975) Predation and the evolution of gregariousness. I. Models for concealment
335 and evasion, *Animal Behaviour*, 23, pp. 779–800. DOI:10.1016/0003-3472(75)90106-2.

336 Vergassola, M., Villermaux, E. and Shraiman, B. I. (2007) ‘Infotaxis’ as a strategy for searching
337 without gradients., *Nature*, 445 (7126), pp. 406–9. DOI:10.1038/nature05464.

338 Villermaux, E. and Duplat, J. (2003) Mixing is an aggregation process, *Comptes Rendus*
339 *Mécanique*, 331 (7), pp. 515–523. DOI:10.1016/S1631-0721(03)00110-4.

340 Ward, A. J. W., Herbert-Read, J. E. and Simpson, S. J. (2011) Diets and decisions: the potential
341 use of food protein cues in dietary, sexual and social decisions by mosquitofish, *Animal*
342 *Behaviour*, 82 (4), pp. 783–790. DOI:10.1016/j.anbehav.2011.07.010.

343 Webster, D. R. and Weissburg, M. J. (2001) Chemosensory guidance cues in a turbulent
344 chemical odor plume, *Limnology and Oceanography*, 46 (5), pp. 1034–1047.

345 Webster, D. R. and Weissburg, M. J. (2009) The Hydrodynamics of Chemical Cues Among
346 Aquatic Organisms, *Annual Review of Fluid Mechanics*, 41 (1), pp. 73–90.
347 DOI:10.1146/annurev.fluid.010908.165240.

348 Weissburg, M. J. and Zimmer-Faust, R. K. (1993) Life and death in moving fluids:
349 hydrodynamic effects on chemosensory-mediated predation, *Ecology*, 74 (5), pp. 1428.
350 DOI:10.2307/1940072.

351 Wilson, M. L. and Weissburg, M. J. (2012) Temporal and spatial sampling strategies maintain
352 tracking success of whelks to prey patches of differing distributions, *Animal Behaviour*, 84 (6),
353 pp. 1323–1330. DOI:10.1016/j.anbehav.2012.08.024.

354 Wrona, F. J. and Dixon, R. W. J. (1991) Group size and predation risk: a field analysis of
355 encounter and dilution effects, *The American Naturalist*, 137 (2), pp. 186–201.
356 DOI:10.1086/285153.

357 Zimmer-Faust, R. K., Finelli, C. M., Pentcheff, N. D. and Wethey, D. S. (1995) Odor plumes and
358 animal navigation in turbulent water flow: a field study, *Biological Bulletin*, 188 (2), pp. 111–
359 116. DOI:10.2307/1542075.

360

Figure 1(on next page)

Layout of the Y-maze (Total dimensions: length 93 cm and width 40 cm).

Water flowed over the horizontal barrier (mid-way in the 30 cm long header chamber) before entering the Y-maze through a collimator (10 cm long) to ensure even flow on both sides. The water flowed along the arms of the Y (20 cm long) before entering the stem (33 cm long), which was partitioned with a removable barrier for the release zone (20 cm long). Water flowed through the Y maze at approximately 3 cm s⁻¹ before exiting through the outlet holes (3 cm in diameter). Cue input points are marked by a black dot. Large open circles represent the cylinders (5.5 cm in diameter) added to the tank in the 'disturbed flow' treatments.




Figure 2 (on next page)

Proportion of fish choosing the prey side in a y-maze.

Stars above bars signify significant differences (binomial exact tests) from random choice of side. Dark bars are non-disturbed flow treatments and light bars are disturbed flow treatments. The horizontal dashed line indicates no preference.

Proportion of cue side choices

1.0
0.8
0.6
0.4
0.2
0.0

NS

N/A

Low

*

NS

Medium

*

*

High

Que concentration

