

Distribution and establishment of the alien Australian redclaw crayfish, *Cherax quadricarinatus*, in South Africa and Swaziland (#15514)

1

First submission

Please read the **Important notes** below, the **Review guidance** on page 2 and our **Standout reviewing tips** on page 3. When ready [submit online](#). The manuscript starts on page 4.

Important notes

Editor and deadline

Marta Sánchez / 15 Jan 2017

Files

4 Figure file(s)

3 Table file(s)

2 Raw data file(s)

Please visit the overview page to [download and review](#) the files not included in this review PDF.

Declarations

Involves a field study on animals or plants.



Please read in full before you begin

How to review

When ready [submit your review online](#). The review form is divided into 5 sections. Please consider these when composing your review:

1. BASIC REPORTING

2. EXPERIMENTAL DESIGN

3. VALIDITY OF THE FINDINGS






4. General comments

5. Confidential notes to the editor





 You can also annotate this PDF and upload it as part of your review

To finish, enter your editorial recommendation (accept, revise or reject) and submit.





BASIC REPORTING

-  Clear, unambiguous, professional English language used throughout.
-  Intro & background to show context. Literature well referenced & relevant.
-  Structure conforms to [PeerJ standards](#), discipline norm, or improved for clarity.
-  Figures are relevant, high quality, well labelled & described.
-  Raw data supplied (see [PeerJ policy](#)).

EXPERIMENTAL DESIGN

-  Original primary research within [Scope of the journal](#).
-  Research question well defined, relevant & meaningful. It is stated how the research fills an identified knowledge gap.
-  Rigorous investigation performed to a high technical & ethical standard.
-  Methods described with sufficient detail & information to replicate.

VALIDITY OF THE FINDINGS

-  Impact and novelty not assessed. Negative/inconclusive results accepted. *Meaningful* replication encouraged where rationale & benefit to literature is clearly stated.
-  Conclusions are well stated, linked to original research question & limited to supporting results.
-  Speculation is welcome, but should be identified as such.
-  Data is robust, statistically sound, & controlled.

The above is the editorial criteria summary. To view in full visit <https://peerj.com/about/editorial-criteria/>

7 Standout reviewing tips

3



The best reviewers use these techniques

Tip

Example

Support criticisms with evidence from the text or from other sources

Smith et al (J of Methodology, 2005, V3, pp 123) have shown that the analysis you use in Lines 241-250 is not the most appropriate for this situation. Please explain why you used this method.

Give specific suggestions on how to improve the manuscript

Your introduction needs more detail. I suggest that you improve the description at lines 57- 86 to provide more justification for your study (specifically, you should expand upon the knowledge gap being filled).

Comment on language and grammar issues

The English language should be improved to ensure that your international audience can clearly understand your text. I suggest that you have a native English speaking colleague review your manuscript. Some examples where the language could be improved include lines 23, 77, 121, 128 - the current phrasing makes comprehension difficult.

Organize by importance of the issues, and number your points

1. Your most important issue
2. The next most important item
3. ...
4. The least important points

Give specific suggestions on how to improve the manuscript

Line 56: Note that experimental data on sprawling animals needs to be updated. Line 66: Please consider exchanging "modern" with "cursorial".

Please provide constructive criticism, and avoid personal opinions

I thank you for providing the raw data, however your supplemental files need more descriptive metadata identifiers to be useful to future readers. Although your results are compelling, the data analysis should be improved in the following ways: AA, BB, CC

Comment on strengths (as well as weaknesses) of the manuscript

I commend the authors for their extensive data set, compiled over many years of detailed fieldwork. In addition, the manuscript is clearly written in professional, unambiguous language. If there is a weakness, it is in the statistical analysis (as I have noted above) which should be improved upon before Acceptance.

Distribution and establishment of the alien Australian redclaw crayfish, *Cherax quadricarinatus*, in South Africa and Swaziland

Ana L Nunes ^{Corresp., 1, 2, 3}, Tsungai A Zengeya ⁴, Andries C Hoffman ⁵, G John Measey ¹, Olaf LF Weyl ²

¹ Centre for Invasion Biology, Department of Botany and Zoology, University of Stellenbosch, Stellenbosch, South Africa

² Centre for Invasion Biology, South African Institute for Aquatic Biodiversity, Grahamstown, South Africa

³ Invasive Species Programme, South African National Biodiversity Institute, Kirstenbosch Research Centre, Cape Town, South Africa

⁴ South African National Biodiversity Institute, Kirstenbosch Research Centre, Cape Town, South Africa

⁵ Mpumalanga Tourism and Parks Agency, Nelspruit, South Africa

Corresponding Author: Ana L Nunes

Email address: ananunes@sun.ac.za

Background. The Australian redclaw crayfish (*Cherax quadricarinatus* von Martens), native from Australasia, has been widely translocated around the world due to aquaculture and aquarium trade. Mostly as a result of escape from aquaculture facilities, this species has established extralimital populations in Australia and alien populations in Europe, Asia, Central America and Africa. In South Africa, *C. quadricarinatus* was first reported in 2002 in the Komati River, following its escape from an aquaculture facility in Swaziland, but data on the current status of its populations is not available.

Methods. To establish a better understanding of its distribution, rate of spread and population dynamics, we surveyed a total of 46 sites in South Africa and Swaziland. Surveys were performed between September 2015 and August 2016 and involved visual observations and the use of collapsible crayfish traps.

Results. *C. quadricarinatus* is now present in the Komati, Lomati, Mbuluzi, Mlawula and Usutu rivers, and it was also detected in several off-channel irrigation impoundments. Where present, it was generally abundant, with populations having multiple size classes and containing ovigerous females. In the Komati River, it has spread over more than 112 km downstream of the initial introduction point and 33 km upstream a tributary, resulting in a minimum spread rate of 8 km.year⁻¹ downstream and 4.7 km.year⁻¹ upstream. In Swaziland, estimated downstream spread rate might go as high as 14.6 km.year⁻¹. Closer to the introduction source, individuals were generally larger and heavier, probably indicating high juvenile dispersal.

Discussion. These findings demonstrate that *C. quadricarinatus* is established in the study area in South Africa and Swaziland and that the species has spread, not only within the river of first introduction, but also between rivers. Considering the strong impacts that alien crayfish usually have on invaded ecosystems, this species can have devastating impacts on freshwater ecosystems in Southern Africa. An assessment of its potential impacts on native freshwater biota, as well as an evaluation of possible control measures are, therefore, an urgent requirement.

1 **Distribution and establishment of the alien Australian redclaw crayfish, *Cherax***
2 ***quadricarinatus*, in South Africa and Swaziland**

3 Ana L Nunes^{1,2,3}, Tsungai A Zengeya⁴, Andries C Hoffman⁵, G John Measey¹, Olaf LF Weyl²

4

5 ¹ Centre for Invasion Biology, Department of Botany and Zoology, Stellenbosch University,
6 South Africa

7 ² Centre for Invasion Biology, South African Institute for Aquatic Biodiversity, Grahamstown,
8 South Africa

9 ³ Invasive Species Programme, South African National Biodiversity Institute, Kirstenbosch
10 Research Centre, South Africa

11 ⁴ South African National Biodiversity Institute, Kirstenbosch Research Centre, South Africa

12 ⁵ Mpumalanga Tourism and Parks Agency, Nelspruit, South Africa

13


14 Corresponding Author:

15 Ana L. Nunes^{1,2,3}

16

17 E-mail address: ananunes@sun.ac.za

18 **Abstract**

19 **Background.** The Australian redclaw crayfish (*Cherax quadricarinatus*  von Martens), native
20 from Australasia, has been widely translocated around the world due to aquaculture and
21 aquarium trade. Mostly as a result of escape from aquaculture facilities, this species has
22 established extralimital populations in Australia and alien populations in Europe, Asia, Central
23 America and Africa. In South Africa, *C. quadricarinatus* was first reported in 2002 in the
24 Komati River, following its escape from an aquaculture facility in Swaziland, but data on the
25 current status of its populations is not available.

26 **Methods.** To establish a better understanding of its distribution, rate of spread and population
27 dynamics, we surveyed a total of 46 sites in South Africa and Swaziland. Surveys were
28 performed between September 2015 and August 2016 and involved visual observations and the
29 use of collapsible crayfish traps.

30 **Results.** *C. quadricarinatus* is now present in the Komati, Lomati, Mbuluzi, Mlawula and Usutu
31 rivers, and it was also detected in several off-channel irrigation impoundments. Where present, it
32 was generally abundant, with populations having multiple size classes and containing ovigerous
33 females. In the Komati River, it has spread over more than 112 km downstream of the initial
34 introduction point and 33 km upstream a tributary, resulting in a minimum spread rate of 8
35 km.year⁻¹ downstream and 4.7 km.year⁻¹ upstream. In Swaziland, estimated downstream spread
36 rate might go as high as 14.6 km.year⁻¹. Closer to the introduction source, individuals were
37 generally larger and heavier, probably indicating high juvenile dispersal.

38 **Discussion.** These findings demonstrate that *C. quadricarinatus* is established in the study area
39 in South Africa and Swaziland and that the species has spread, not only within the river of first
40 introduction, but also between rivers. Considering the strong impacts that alien crayfish usually

41 have on invaded ecosystems, this species can have devastating impacts on freshwater ecosystems
42 in Southern Africa. An assessment of its potential impacts on native freshwater biota, as well as
43 an evaluation of possible control measures are, therefore, an urgent requirement.

44 Introduction

45 Freshwater crayfish have been introduced globally, mostly for aquaculture and ornamental
46 purposes, but generally their subsequent invasions have resulted in more ecosystem losses than
47 benefits (Lodge et al., 2012). Continental Africa contains no native freshwater crayfish species,
48 but three Australasian Parastacidae species, the Australian redclaw crayfish (*Cherax*
49 *quadricarinatus*), the smooth marron (*Cherax cainii*) and the yabby (*Cherax destructor*), and a
50 single North American Cambaridae species, the red swamp crayfish (*Procambarus clarkii*), have
51 been introduced (Boyko, 2016). All four species have been introduced into South Africa, but
52 only *P. clarkii* and *C. quadricarinatus* seem to have successfully established wild populations.
53 Although *P. clarkii* has been introduced to several African countries and caused visible impacts
54 (Lowery & Mendes, 1977; Mikkola, 1996; Foster & Harper, 2006), in South Africa the species is
55 only known from a single locality and does not seem to be spreading (Nunes et al., *in press*).
56 Feral populations of *C. quadricarinatus* are more widespread in the country (du Preez & Smit,
57 2013; Coetzee et al., 2015; de Villiers, 2015) and have also been reported from Swaziland (de
58 Moor, 2002), Zimbabwe (Marufu, Phiri & Nhiwatiwa, 2014), Zambia and Mozambique
59 (Chivambo, Nerantzoulis & Mussagy, 2013; Nunes et al., 2016). Globally, *C. quadricarinatus*
60 has been translocated to non-native areas in Australia (Doupé et al., 2004; Leland, Coughran &
61 Furse, 2012) and Indonesian territories (Patoka et al., 2016), and wild populations have also been
62 introduced to Slovenia (Jaklič & Vrezec, 2011), Israel (Snovsky & Galil, 2011), Mexico
63 (Bortolini, Alvarez & Rodriguez-Almaraz, 2007; Vega-Villasante et al., 2015; Torres-Montoya
64 et al., 2016), Jamaica (Todd, 2005; Pienkowski et al., 2015), Puerto Rico (Williams et al., 2001)
65 and Singapore (Ahyong & Yeo, 2007; Belle et al., 2011).

66 *Cherax quadricarinatus* was first imported into South Africa in 1988 for research on its
67 aquaculture potential together with other *Cherax* species (Van den Berg & Schoonbee, 1991).
68 Despite considerable interest in the aquaculture of this species in the late 1990s, its import and
69 culture for commercial purposes was never permitted in South Africa. As a result, a farmer who
70 had attempted to establish an aquaculture venture in South Africa around this time managed to,
71 instead, successfully establish it in neighbouring Swaziland (de Moor, 2004). There are
72 anecdotal reports that two batches of *C. quadricarinatus* were introduced from Australia to
73 Swaziland, one for the abovementioned farm located near the Sand River Dam, close to the
74 Komati River and the other to a farm near Manzini or Big Bend, in the Usutu River catchment
75 (A Howland, 2016 - general manager of IYSIS cattle ranch, inside which the Sand River Dam is
76 situated -, pers. comm.). As is the case with many other small scale aquaculture ventures in
77 South Africa (Weyl et al., 2016), the Sand River Dam farm ended up not being economically
78 viable and was abandoned, allowing crayfish to escape into the Sand River Dam. Later, most
79 likely during the large floods that occurred in the area in 2000, it spread into the Komati River
80 via the Sand River (de Moor, 2002, 2004; A Howland, 2016, pers. comm.). From there, it spread
81 downstream and the species was first detected in the Komati River in South Africa in 2002 (de
82 Villiers, 2015). While there is no information on the outcome of the other aquaculture farm close
83 to Manzini or Big Bend (in the Usutu River catchment), in 2012 *C. quadricarinatus* was detected
84 in an outlet of Lake Nyamiti in the Ndumo Game Reserve (South Africa) (du Preez & Smit,
85 2013), which eventually connects to the Usutu River.

86 In June 2009, the species was also reported from a small wetland in a residential area close to
87 Richard's Bay, in KwaZulu-Natal Province, South Africa (R Jones, 2016 - Ezemvelo KZN
88 Wildlife -, pers. comm.), a site distant and not directly connected to the initial introduction sites.

89 Despite these initial reports of *C. quadricarinatus* in Swaziland and South Africa, no systematic
90 survey has ever been carried out to determine their distribution, spread rate and population
91 dynamics. This is of concern because crayfish invasions have generally been shown to result in
92 strong impacts on recipient ecosystems (Lodge et al., 2012) and, given the absence of native
93 crayfish on the African continent, these impacts are likely to be even stronger (de Moor, 2002;
94 Nunes et al., 2016). In this study, we assess the current distribution, rate of spread and population
95 dynamics of *C. quadricarinatus* populations in South Africa and Swaziland. In addition, for the
96 Komati River (initial main river of introduction), we further investigate if population
97 characteristics, such as abundance, biomass, sex ratio, body size and mass vary with distance to
98 the introduction source, since traits of invasive populations have been shown to vary along
99 invasion gradients (see review in Iacarella, Dick & Ricciardi, 2015).

100

101 **Materials & Methods**

102 *Field Study Permissions*

103 Permits were obtained from the Mpumalanga Tourism and Parks Agency (MPB. 5523),
104 Ezemvelo KZN Wildlife (OP 4428/2015) and Mbuluzi Game Reserve.

105

106 *Study area*

107 The study area was mainly situated in the Inkomati, Mbuluzi and Usutu River basins, all of
108 which are international river systems running through Swaziland, South Africa and
109 Mozambique. The Inkomati basin, mainly located in the Mpumalanga Province of South Africa,
110 consists of three major sub-catchments, the Komati, the Crocodile and the Sabie-Sand (MTPA,
111 2013). The Komati sub-catchment is composed of the Komati River and its tributaries, one of

112 which is the Lomati River. The Komati River rises west of Carolina in Mpumalanga, and flows
113 for 480 km in a north-easterly direction through three countries (South Africa→ Swaziland→
114 South Africa→ Mozambique). The Crocodile River is the main river in the Crocodile sub-
115 catchment, originating north of Dullstroom and flowing eastwards towards its confluence with
116 the Komati River. The Sand River Dam, where *C. quadricarinatus* was first introduced in
117 Swaziland, is located in the Inkomati catchment (Figs. 1, 2A).

118 The main river of the Mbuluzi basin is the Mbuluzi River, which originates in the Ngwenya hills
119 in northwest Swaziland, close to the border with South Africa, and flows in an easterly direction
120 through central Swaziland into Mozambique. At times, water is transferred from the Komati
121 River basin to the Mbuluzi River basin via an intricate network of approximately 40km of
122 irrigation channels (A Howland, 2016, pers. comm.; Gustafsson & Johansson, 2006). The
123 Mlawula River, located close to the border with Mozambique, is one of its tributaries, which
124 crosses several protected areas, such as the Mbuluzi Game Reserve and the Shewula Nature
125 Reserve (Fig. 2B).

126 The Usutu River basin is bordered by the Mbuluzi and Inkomati River basins to the north and the
127 Mhlathuze coastal catchment to the south. The Usutu, Pongola and Ngwavuma are its main sub-
128 catchments. The main river of the Usutu sub-catchment is the Usutu River, which rises
129 near Amsterdam, in Mpumalanga Province, and flows in a south-easterly direction through South
130 Africa and Swaziland (Beuster & Clarke, 2008). It then emerges in the province of KwaZulu-
131 Natal in South Africa where, for approximately 24 kilometres, it defines the border between this
132 country and Mozambique, along the limits of the Ndumo Game Reserve. The Ndumo Game
133 Reserve, a protected area characterised by numerous pans and wetlands, is crossed by the

134 Pongola River, which rises in Northern KwaZulu-Natal, flows eastwards until the Pongolapoort
135 Dam, from where it flows north-easterly to join the Usutu River in Mozambique (Fig. 2C).
136 Taking into account the reported sighting of *C. quadricarinatus* close to Richard's Bay, this area
137 was also surveyed, as well as two large dams in the KwaZulu-Natal Province (Albert Falls and
138 Goedertrouw Dams), where there have been anecdotal records of crayfish presence (Figs. 1, 2D).

139



140 *Sampling procedure*

141 A total of 46 sampling sites in different water bodies (main rivers, tributaries, pans, wetlands and
142 dams) were surveyed between September 2015 and August 2016 (Fig. 1). Sampling sites were
143 chosen by focusing on areas with suspected presence of *C. quadricarinatus*, according to
144 published or grey literature and to personal communications from farmers, agriculture and
145 conservation officials. Along the Komati River, which has a large number of weirs regulating its
146 flow, nine sites were sampled, six downstream and three upstream of the initial introduction
147 point (Fig. 2A). In contrast, the Lomati River is relatively less regulated and not many sites (six)
148 could be sampled on the main river or its tributaries due to difficult access. The three sampling
149 sites on the Crocodile River were located upstream of its confluence with the Komati River and
150 within the Kruger National Park (Fig. 2A). Sites on the Mbuluzi River and its tributaries were
151 concentrated close to the Mozambican border, upstream (two) and downstream (four) of the
152 potential point of introduction in this river (Fig. 2B). In the Usutu River, four points were
153 sampled in Swaziland and one in South Africa. Three sampling points were selected in the
154 Ndumo Game Reserve and two in the Pongola River, one upstream and one downstream of
155 Pongolapoort Dam (Fig. 2C). In the Richard's Bay area, two points, one where crayfish were
156 detected back in 2009 and one in a connected lake, were sampled (Fig. 2D). Finally, ten small

157 and large dams, most of which are primarily used to store water for agricultural irrigation, were
158 also sampled.

159 Overall, 34 sites were sampled in lotic habitats, spaced at least 2.5 km from each other (but
160 usually over 13 km), according to where access to the rivers was possible. Survey sites in the
161 rivers ranged between 100 and 150 m in length, depending on accessibility of the site. Twelve
162 sites were sampled in lentic habitats. Each sampling site was surveyed at least twice (each site 2-
163 4 times, except four sites where we could not return), once in the wet season (spring/summer,
164 September-March) and once in the dry season (autumn/winter, April-August), in order to
165 confirm crayfish absences and to capture differences in crayfish population dynamics over
166 seasons. The main exception were the sites in the Crocodile River inside Kruger National Park,
167 an area under strict jurisdiction of South African National Parks (SANParks) where, similarly to
168 the four sites mentioned above, we could only sample once. At each of the sampling sites, visual
169 observations of 5-10 minutes along the margins of the water body were made on arrival at the
170 location, in order to look for crayfish specimens or moults. Subsequently, around ten (range: 3-
171 15) ©Promar collapsible crayfish/crab traps (dimensions: 61 × 46 × 20 cm), baited with
172 approximately 100g of dry dog food, were set in the evening at each site, left overnight (14-16 h)
173 and checked the following morning. The number of crayfish caught in each trap, as well as their
174 cephalothorax length (to the nearest mm), mass (to the nearest g) and sex were registered.

175 Crayfish abundance was calculated based on catch per unit effort (CPUE), per sampling session.
176 Due to restrictions imposed by SANParks, traps could not be set in the Crocodile River, where
177 instead electrofishing was conducted by wading for approximately 40 minutes per site, using a
178 handheld SAMUS 725MP, with a 10 mm mesh scoop net.

179 A chi-square goodness-of-fit test was used to test whether overall  sex ratio, or per site and per
180 season, was significantly different than the expected sex ratio of 1:1. For the Komati River, we
181 also investigated a possible relationship between each site's distance from the crayfish source of
182 introduction (measured,  in km, using Google Earth, downstream from the site of initial crayfish
183 introduction and following the river's natural course) and crayfish catch per unit effort
184 (abundance and biomass), sex ratio, size and mass. This was determined using Pearson's
185 correlation coefficient or, in case the assumptions of normality or homogeneity of variances were
186 not met, the non-parametric Spearman rank correlation. The level of significance for all
187 statistical tests performed was $p < 0.05$.

188

189 **Results**

190 *Cherax quadricarinatus* was detected in 22 out of the 46 sampling sites surveyed (Figs. 1, 2,
191 Tables 1, S1). All sampling sites located on the Komati and Lomati rivers in South Africa had
192 crayfish present, but no crayfish were detected in upstream and more elevated sampling sites on
193 both rivers in Swaziland. Crayfish were also found in the Mbuluzi River, but only in sampling
194 sites downstream of the potential introduction point (interbasin transfer point between the
195 Inkomati and Mbuluzi basins) in this river (Fig. 2B). Both sites on the Mlawula River, a tributary
196 of the Mbuluzi River, also yielded crayfish. On the Usutu River, three sites close to Big Bend
197 had crayfish, but crayfish were not caught further upstream in Swaziland, or downstream in
198 Ndumo Game Reserve (Figure 2C). Crayfish were also found in six out of the 12 sampled lentic
199 habitats. However, they were not detected in the Crocodile and Pongola rivers, Ndumo Game
200 Reserve and Richard's Bay area (Fig. 2, Table S1).

201



202 A total of 577 crayfish were caught during the wet season (383 males and 194 females), with a
203 maximum of 63 individuals in a single trap (at site D01), whereas only 267 crayfish were caught
204 in the dry season (149 males and 118 females). The maximum mass that a crayfish attained was
205 250 g, for an individual caught at site K06 (Table 1). In the Komati River, average crayfish
206 abundances were quite high, ranging from 0.4 to 9.4 individuals.trap.night⁻¹ in the wet season
207 and 1.0 to 7.0 individuals.trap.night⁻¹ in the dry season. High abundances were also found in
208 dams (0.1-15.3 individuals.trap.night⁻¹), especially during the wet season. Abundances were
209 lower in the Mbuluzi (1.0-4.5 individuals.trap.night⁻¹) and Mlawula rivers (0-4.0
210 individuals.trap.night⁻¹) and much lower in the Lomati and Usutu rivers ranging, respectively,
211 from 0-0.7 individuals.trap.night⁻¹ and 0.1-0.8 individuals.trap.night⁻¹ (Table 1). Average biomass
212 was higher in the dry than in the wet season in the Komati (47.4 g.trap.night⁻¹ for dry season and
213 35.7 g.trap.night⁻¹ for wet season), Lomati (26.1 g.trap.night⁻¹ for dry season and 7.4 g.trap.night⁻¹
214 ¹ for wet season) and Mbuluzi rivers (27.8 g.trap.night⁻¹ for dry season and 22.0 g.trap.night⁻¹ for
215 wet season). On the contrary, average biomass was higher in the wet than the dry season in the
216 Mlawula River (15.5 g.trap.night⁻¹ for wet season and 5.4 g.trap.night⁻¹ for dry season), Usutu
217 River (10.9 g.trap.night⁻¹ for wet season and 5.2 g.trap.night⁻¹ for dry season) and in dams (34.8
218 g.trap.night⁻¹ for wet season and 25.6 g.trap.night⁻¹ for dry season) (Table 1).

219

220 Specimens of *C. quadricarinatus* varied widely in size, with cephalothorax lengths ranging from
221 20 to 114 mm, and individuals between 40 and 70 mm being by far the most numerous and
222 representing 73.66% of all measured crayfish. Length-frequency graphs demonstrated the
223 existence of multiple cohorts in the Komati, Mbuluzi, Mlawula and Usutu rivers, and also in

224 irrigation dams. This did not seem to be the case for the Lomati River, where only very few size
225 classes were present (Figs. 3A, B).

226 Oviparous females, or females carrying newly hatched crayfish (average size 63.8 mm, average
227 mass 58.7 g) were found in October and December 2015, at five different sampling sites, three
228 on the Komati River (K01, K02 and K03) and two in dams (D01 and D02) (Table S2). The
229 number of eggs ranged from 281 to 539 and the number of newly hatched crayfish ranged from
230 18 to 20 (many probably detached while in the traps).

231

232 In the wet season, the overall sex ratio (all sampling sites together) was significantly different
233 from the expected sex ratio of 1: 1 ($\chi^2= 58.856, p < 0.001$), with males outnumbering females,
234 while this was marginally non-significant in the dry season ($\chi^2= 3.626, p= 0.057$). Looking at
235 specific areas, in the wet season, males were significantly more numerous than females in the
236 Komati ($\chi^2= 8.022, p= 0.005$) and Mlawula rivers ($\chi^2= 3.930, p= 0.047$), as well as in dams ($\chi^2=$
237 $45.478, p < 0.001$), but not in the Mbuluzi ($\chi^2= 3.457, p= 0.063$) or the Usutu rivers ($\chi^2= 1.600,$
238 $p= 0.206$). In the dry season, sex ratios were not significantly different to the expected 1: 1
239 proportion ($p \geq 0.05$ in all cases). However, if we consider sampling sites individually, sex ratio
240 was not significantly different from the 1: 1 proportion for most of them ($p > 0.05$ for most sites),
241 except for sites K03 ($\chi^2= 13.787, p < 0.001$), D01 ($\chi^2= 45.026, p < 0.001$) and D05 ($\chi^2= 4.378, p=$
242 0.036) in the wet season and D01 in the dry season ($\chi^2= 8.257, p= 0.004$) (Table 1).

243

244 In the Komati River, crayfish were found at a maximum distance of 112 km downstream of the
245 point of introduction, indicating a minimum downstream spread rate of 8 km.year⁻¹ (using 2001
246 as the approximate year of first introduction). In the Lomati River, they were detected 93 km

247 from the source of introduction, approximately 33 km upstream from the confluence with the
248 Komati River. This indicates a total spread rate of 6.6 km.year⁻¹ or, assuming an average spread
249 rate of 8 km.year⁻¹ downstream until the confluence with the Komati River, an upstream spread
250 rate of 4.7 km.year⁻¹.

251

252 No significant correlations were found between abundance, biomass or sex ratio of *C.*
253 *quadricarinatus* during both wet and dry seasons, and distance to crayfish introduction source in
254 the Komati River (for all correlations, $p > 0.05$). However, size and mass of both females and
255 males was significantly correlated with distance to the source of crayfish introduction.
256 Interestingly, a significant positive correlation was found between these variables for females in
257 the wet season ($r = 0.344$, $N = 69$, $P = 0.004$ for size and $r_s = 0.438$, $N = 71$, $P < 0.001$ for mass),
258 while during the dry season these correlations were negative ($r = -0.686$, $N = 63$, $P < 0.001$ for size
259 and $r = -0.641$, $N = 63$, $P < 0.001$ for mass) (Fig. 4A). For males, the relationship was always
260 negative, independent of season, but only statistically significant in the dry season ($r = -0.440$, $N =$
261 66 , $P < 0.001$ for size and $r = -0.505$, $N = 66$, $P < 0.001$ for mass) (Fig. 4B).

262

263 Discussion

264 In this study we confirmed the presence of established and widespread populations of *C.*
265 *quadricarinatus* in South Africa and Swaziland. Based on the evidence that populations have
266 spread and are reproducing at multiple localities as far as 115 km from the point of introduction,
267 this species can be considered as fully invasive (category E) in these countries, according to
268 Blackburn et al. (2011). We also show how populations of this species have expanded in South
269 Africa and Swaziland since they were first detected in 2002, being now present in at least three

270 large rivers (Komati, Mbuluzi and Usutu), two tributaries (Lomati and Mlawula rivers), as well
271 as in several irrigation dams. Crayfish populations were found to be established (presence of
272 multiple cohorts and reproduction) at most sampling sites, the main exception being the Lomati
273 River, where very few individuals and size classes were found.

274 Although *Cherax quadricarinatus* were found to be capable of dispersing in an upstream
275 direction in two different tributaries (Lomati and Mlawula rivers), they were not detected in
276 upstream areas of both the Komati and Lomati rivers, which might be related with the large
277 increase in elevation in these sampling points (274-433 m a.s.l) and/or potential lower water
278 temperatures. In the Lomati River, the Driekoppies Dam might also act as a dispersal barrier.
279 Crayfish were also not detected in the Crocodile River; however, some specimens were recently
280 detected approximately 10.7 km upstream of the furthest point sampled in this study, probably as
281 the result of an escape from an adjacent invaded dam (A Hoffman, T Zengeya, 2016, pers. obs.).

282 The fact that no specimens were detected in the Ndumo Game Reserve supports the idea that
283 populations are not established in this area, which is not surprising given that only four
284 specimens were found by Du Preez and Smit (2013) back in 2012. This might be the result of an
285 extended drought period that has been occurring in the area for a long time.

286 Crayfish was not found in sites near Richard's Bay, indicating that the record from 2009 was
287 probably the result of an isolated introduction event, through release by aquarists or escape from
288 an ornamental pond. In fact, this would not be surprising, given the ease with which one can find
289 different crayfish species for sale in South Africa, either via online sources or in pet shops
290 around the country (AL Nunes, 2016, pers. obs.). The unconfirmed records of crayfish at Albert
291 Falls Dam and Goedertrouw Dam seem to be erroneous. However, it is important to note that,
292 given the extensive size of these dams, it is extremely difficult to confirm crayfish absence,

293 especially without an intensive and focused sampling, targeted specifically for these type of
294 habitats.

295 Relative abundances of *C. quadricarinatus* in the Komati River (average 3.3 indiv.trap.night⁻¹;
296 maximum 9.4 indiv.trap.night⁻¹) and in irrigation dams (average 3.7 indiv.trap.night⁻¹; maximum
297 15.3 indiv.trap.night⁻¹) were considerably higher than the ones found in other invasive
298 populations of this species in Zimbabwe (maximum of 4.0 indiv.trap.night⁻¹; Marufu, Phiri &
299 Nhiwatiwa, 2014) and Slovenia (0.09 indiv.trap.night⁻¹; Jaklič & Vrezec, 2011), reflecting a high
300 invasion potential of the species in this region. On the other hand, very low abundances were
301 found in the Lomati River (average 0.2 indiv.trap.night⁻¹), probably reflecting either a more
302 recent invasion or a less suitable habitat (Hudina et al., 2012), taking into account that this river
303 is regulated in a different way than the Komati, with few gauging weirs and a high flow velocity
304 regime. Whichever reason, crayfish populations are not yet established in the Lomati River.

305 The observed average size range of *C. quadricarinatus* collected in the various sampling sites
306 (cephalothorax length: 20-98.2 mm) was in the range of values reported for this species in other
307 invaded areas (Bortolini, Alvarez & Rodriguez-Almaraz, 2007; Jaklič & Vrezec 2011; Marufu,
308 Phiri & Nhiwatiwa, 2014).

309 The species exhibited potential to disperse both downstream the different initial invasion points
310 and upstream of two different tributaries. In the Inkomati basin, downstream and upstream
311 spread occurred at a rate of 8 and 4.7 (possibly 6.6) km.year⁻¹, respectively. However, the
312 downstream rate might be even higher, considering the high likelihood that the species has
313 already spread further downstream the Komati River into the Mozambican side (which could not
314 be sampled in this study). In the Mbuluzi River basin, which *C. quadricarinatus* most likely
315 reached via the interbasin water irrigation channels connecting it with the Inkomati basin


316 (similarly to what occurred to the spread of an alien loricariid catfish in the KwaZulu-Natal
317 province; Jones et al., 2013), the date of introduction is uncertain, but in late 2009 crayfish were
318 observed for the first time at Pequenos Libombos Dam, southern Mozambique (I Nerantzoulis,
319 2016, pers. comm.), and recorded as established in 2011 (Fig. 2B; Chivambo, Nerantzoulis &
320 Mussagy, 2013). Assuming this was the result of natural spread, and not of an exceptional
321 translocation event, this shows that in eight years, and in a downstream direction, the species
322 covered 40 km of channels between the Mbuluzi and Inkomati basins, plus 76.8 km in the
323 Mbuluzi River until the Pequenos Libombos Dam. This indicates a potential spread rate of 14.6
324 km.year⁻¹. Down and upstream dispersal have been observed for other invasive crayfish species,
325 ranging from 1.8 to 24.4 km.year⁻¹ (downstream) and 0.35 to 4 km.year⁻¹ (upstream) for
326 *Pacifastacus leniusculus* in different European countries (Bubb, Thom & Lucas, 2005; Hudina et
327 al., 2009; Weinländer & Füreder, 2009; Bernardo et al., 2011), 0.5 to 3.10 km.year⁻¹ (upstream)
328 for *P. clarkii* (Bernardo et al., 2011; Ellis et al., 2012), and 12 to 84 km.year⁻¹ (downstream) and
329 2.5 km.year⁻¹ (upstream) for *Orconectes limosus* in Eastern Europe (Hudina et al., 2009). This
330 indicates that the first estimates of dispersal rates for *C. quadricarinatus*, especially for upstream
331 movements, are high, once again suggesting a high invasion potential of the species in the study
332 area. Furthermore, irrigation dams, where crayfish populations seem to become very abundant,
333 might act as secondary sources of crayfish invasions or as stepping stones for range expansion
334 through irrigation channels or over land, facilitating subsequent establishment in new irrigations
335 dams, rivers or tributaries.

336 In the Komati River, which has been colonised for the longest time, crayfish were generally
337 larger and heavier close to the initial introduction point, with sizes decreasing as distance to the
338 invasion source increased. A similar pattern has been observed for round goby invasions in

339 Canada (Ray & Corkum, 2001; Brownscombe & Fox, 2012) and the same tendency found for
340 signal crayfish in Croatia (Hudina et al., 2012), suggesting that juveniles may disperse more
341 actively and rapidly than adults, likely due to high intraspecific competition. In the case of
342 females, this might also indicate a strategy that allocates resources to favour reproduction with
343 increased offspring closer to the source, as egg number is a function of female size (Jones, 1990).
344 However, the opposite pattern was observed for females during the wet season, with smaller
345 females found near the introduction point and larger ones further downstream. Given that sexual
346 maturity is generally reached when animals attain around 50 to 60 g (Jones, 1990),
347 corresponding to approximately 55 to 65 mm cephalothorax length in this study, this may
348 indicate that mature females are reproducing at different times of the year along the invasion
349 gradient. In sites further away from the source females are spawning in October-December (and
350 perhaps repeatedly), while reproduction might be taking place later in March-May in longer
351 established populations. In fact, in subtropical regions of Australia, *C. quadricarinatus* has a
352 natural reproductive season throughout spring and summer, with spawning occurring more than
353 once from October to March (Jones, 1990; Masser & Rouse, 1997). Alternatively, this pattern
354 might suggest that large females closer to the invasion front are more active and disperse during
355 the wet season, which might contribute to further range expansion (Brownscombe & Fox, 2012).
356 Although current legislation prohibits the importation release and movement of *C.*
357 *quadricarinatus* in South Africa (RSA, 2016), the lack of resources (manpower and financial)
358 makes it extremely challenging to enforce these regulations. Furthermore, taking into account the
359 accidental escape of *C. quadricarinatus* from an aquaculture farm in Swaziland and consequent
360 spread to South Africa and Mozambique, this study reinforces the importance of putting
361 international agreements regarding invasive species into practice, such as the SADC Protocol on

362 Fisheries, which prohibits the introduction of exotic species into aquatic ecosystems shared by
363 two states, unless the affected state parties agree to the introduction (de Moor, 2004). Clearly,
364 there is a need to strengthen and better coordinate the enforcement and effectiveness of existing
365 protocols between neighbouring countries in Africa, in what concerns introduction and spread of
366 invasive species. Taking into account that, once established, invasive crayfish populations are
367 usually impossible to eradicate, transnational cooperation should also be taken into account
368 regarding possible management actions (e.g. mechanical, physical, chemical and/or autocidal
369 methods; reviewed in Gherardi et al., 2011) to contain or hinder the spread of *C. quadricarinatus*
370 in these international river systems. These actions would need to be implemented by all countries
371 involved (South Africa, Swaziland and Mozambique), in order for the efforts of one country to
372 not be jeopardised by the other non-complying countries.

373 **Conclusion**

374 While the environmental impact of *C. quadricarinatus* in newly invaded habitats has yet to be
375 determined, local communities in South Africa have already started harvesting it (Coetzee et al.,
376 2015), increasing the risk of translocations for commercial reasons.  The possible introduction of
377 this species into new catchments in Africa is a matter of extreme concern, especially given the
378 high speed at which the species has been expanding and its potential impacts on native biota,
379 such as disease introductions, competitive interactions with native freshwater crustaceans or
380 habitat modifications (de Moor, 2002; Nunes et al., 2016). However, as no formal research has
381 been done on impacts of *C. quadricarinatus* invasive populations in any part of the world, the
382 species would be classified as ‘Data Deficient’ (current information insufficient to assess level of
383 impact) according to Blackburn’s et al. (2014) environmental impact classification for alien taxa.

384 This calls for an immediate assessment of potential impacts of this species on native freshwater
385 ecosystems in Africa.

386

387 **Acknowledgments**

388 We are greatly indebted to Len Coetzer for showing us various sampling sites in South Africa
389 and for providing us the contact details of numerous farmers and researchers in the area. We also
390 thank Vhutali Nelwamondo, Jonathan Vervaeke and Rheul Lombard for their invaluable help
391 during field work. We are grateful to the South African National Parks (SanParks), in the person
392 of Robin Petersen, for permission to sample the Crocodile River inside Kruger National Park.
393 We are grateful to Prof. CN Magagula for her essential help in contacting permitting authorities
394 in Swaziland and to Morgan Vance and All Out Africa Foundation for receiving us so well in
395 Swaziland. ALN, OLFW and GJM thank the DST-NRF Centre of Excellence for Invasion
396 Biology for their continued support.

397

398 **References**

399 Ahyong ST, Yeo DCJ. 2007. Feral populations of the Australian red-claw crayfish (*Cherax*
400 *quadricarinatus* von Martens) in water supply catchments of Singapore. *Biological Invasions*
401 9:943-946.

402 Belle CC, Wong JQH, Yeo DCJ, Tan SH, Tan HH, Clews E, Todd PA. 2011. Ornamental trade
403 as a pathway for Australian redclaw crayfish introduction and establishment. *Aquatic Biology*
404 12:69-79.

- 405 Bernardo JM, Costa AM, Bruxelles S, Teixeira A. 2011. Dispersal and coexistence of two non-
406 native crayfish species (*Pacifastacus leniusculus* and *Procambarus clarkii*) in NE Portugal
407 over a 10-year period. *Knowledge and Management of Aquatic Ecosystems* 401:28.
- 408 Beuster J, Clarke FA. 2008. Joint Maputo River Basin Water Resources Study – Moçambique,
409 Swaziland and South Africa. Tripartite Permanent Technical Committee (TPTC): Republic of
410 Moçambique, Republic of South Africa and Kingdom of Swaziland.
- 411 Blackburn TM, Essl F, Evans T, Hulme PE, Jeschke JM, Kühn I, Kumschick S, Pyšek P,
412 Rabitsch W, Ricciardi A et al. 2014. A unified classification of alien species based on the
413 magnitude of their environmental impacts. *PLOS Biology* 12:e1001850.
- 414 Blackburn TM, Pyšek P, Bacher S, Carlton JT, Duncan RP, Jarošík V, Wilson JRU, Richardson
415 DM. 2011. A proposed unified framework for biological invasions. *Trends in Ecology and*
416 *Evolution* 26:333-339.
- 417 Bortolini JL, Alvarez F, Rodriguez-Almaraz G. 2007. On the presence of the Australian redclaw
418 crayfish, *Cherax quadricarinatus*, in Mexico. *Biological Invasions* 9:615-620.
- 419 Boyko CB. 2016. Crayfish of Africa. In: Kawai T, Faulkes Z, Scholtz G, eds. *Freshwater*
420 *crayfish: A global overview*. CRC Press, Boca Raton, 583-593.
- 421 Brownscombe JW, Fox MG. 2012. Range expansion dynamics of the invasive round goby
422 (*Neogobius melanostomus*) in a river system. *Aquatic Ecology* 46:175-189.
- 423 Bubb DH, Thom TJ, Lucas MC. 2005. The within catchment invasion of the non-indigenous
424 signal crayfish *Pacifastacus leniusculus* (Dana), in upland rivers. *Bulletin Français de la*
425 *Pêche et de la Pisciculture* 376-377:665-673.

- 426 Chivambo S, Nerantzoulis I, Mussagy A. 2013. O lagostim invasor na Albufeira dos Pequenos
427 Libombos, Maputo, Moçambique: avaliação das relações tróficas. 3as Jornadas Científicas de
428 Docentes e Investigadores da Faculdade de Ciências, Maputo, Mozambique.
- 429 Coetzee HC, Nell W, Van Eeden ES, De Crom EP. 2015. Artisanal fisheries in the Ndumo area
430 of the Lower Phongolo River Floodplain, South Africa. *Koedoe* 57:1248(1-6).
- 431 de Moor I. 2002. Potential impacts of alien freshwater crayfish in South Africa. *African Journal*
432 *of Aquatic Sciences* 27:125-139.
- 433 de Moor I. 2004. Protocols for moving germplasm among countries in Africa. In: Gupta MV,
434 Bartley DM, Acosta BO, eds. *Use of genetically improved and alien species for aquaculture*
435 *and conservation of aquatic biodiversity in Africa*. WorldFish Center Conference Proceedings
436 68, Penang, Malaysia, 77-92.
- 437 de Villiers M. 2015. Freshwater crayfish found in Komati River. Lowvelder. Available at
438 <http://lowvelder.co.za/279853/crayfish-web/> (accessed 8 July 2015).
- 439 Doupe RG, Morgan DL, Gill HS, Rowland AJ. 2004. Introduction of redclaw crayfish *Cherax*
440 *quadricarinatus* (von Martens) to Lake Kununurra, Ord River, Western Australia: prospects
441 for a 'yabby' in the Kimberley. *Journal of the Royal Society of Western Australia* 87:187-191.
- 442 Du Preez L, Smit N. 2013. Double blow: Alien crayfish infected with invasive temnocephalan in
443 South African waters. *South African Journal of Science* 109:9/10(1-4).
- 444 Ellis A, Jackson MC, Jennings I, England J, Phillips R. 2012. Present distribution and future
445 spread of Louisiana red swamp crayfish *Procambarus clarkii* (Crustacea, Decapoda, Astacida,
446 Cambaridae) in Britain: Implications for conservation of native species and habitats.
447 *Knowledge and Management of Aquatic Ecosystems* 406:05.

- 448 Foster J, Harper D. 2006. Status of the alien Louisianan red swamp crayfish *Procambarus*
449 *clarkii* Girard and the native African freshwater crab *Potamonautes loveni* in rivers of the
450 Lake Naivasha catchment, Kenya. *Freshwater Crayfish* 15:189-194.
- 451 Gherardi F, Aquiloni L, Diéguez-Uribeondo, Tricarico E. 2011. Managing invasive crayfish: is
452 there a hope? *Aquatic Sciences* 73:185-200.
- 453 Gustafsson A, Johansson M. 2006. An investigation of nutrient levels along the Mbuluzi River -
454 A background for sustainable water resources management. Master of Science Thesis, Lund
455 University, Sweden.
- 456 Hudina S, Faller M, Lucić A, Klobučar G, Maguire I. 2009. Distribution and dispersal of two
457 invasive crayfish species in the Drava River basin, Croatia. *Knowledge and Management of*
458 *Aquatic Ecosystems* 394-395:09.
- 459 Hudina S, Hock K, Žganec K, Lucić A. 2012. Changes in population characteristics and structure
460 of the signal crayfish at the edge of its invasive range in a European river. *Annales de*
461 *Limnologie- International Journal of Limnology* 48:3-11.
- 462 Iacarella JC, Dick JTA, Ricciardi A. 2015. A spatio-temporal contrast of the predatory impact of
463 an invasive freshwater crustacean. *Diversity and Distributions* 21:803-812.
- 464 Jaklič M, Vrezec A. 2011. The first tropical alien crayfish species in European waters: the
465 redclaw *Cherax quadricarinatus* (Von Martens, 1868) (Decapoda, Parastacidae). *Crustaceana*
466 84(5-6):651-665.
- 467 Jones CM. 1990. The biology and aquaculture potential of *Cherax quadricarinatus*. Queensland
468 Department of Primary Industries and Fisheries. Final Report for Project No. QDPI/8860.
469 Queensland, Australia.


- 470 Jones RW, Weyl OLF, Swartz ER, Hill MP. 2013. Using a unified invasion framework to
471 characterize Africa's first loricariid catfish invasion. *Biological Invasions* 15:2139-2145.
- 472 Leland JC, Coughran J, Furse JM. 2012. Further translocation of the Redclaw, *Cherax*
473 *quadricarinatus* (Decapoda: Parastacidae), to Lake Ainsworth in northeastern New South
474 Wales, Australia. *Crustacean Research* 7:1-4.
- 475 Lodge DM, Deines A, Gherardi F, Yeo DCJ, Arcella T, Baldrige AK, Barnes MA, Chadderton
476 WL, Feder JL, Gantz CA, et al. 2012. Global introductions of crayfishes: evaluating the
477 impact of species invasions on ecosystem services. *Annual Review of Ecology, Evolution and*
478 *Systematics* 43:449-472.
- 479 Lowery RS, Mendes AJ. 1977. *Procambarus clarkii* in Lake Naivasha, Kenya, and its effects on
480 established and potential fisheries. *Aquaculture* 11:111-121.
- 481 Marufu LT, Phiri C, Nhwatiwa T. 2014. Invasive Australian crayfish *Cherax quadricarinatus* in
482 the Sanyati Basin of Lake Kariba: a preliminary survey. *African Journal of Aquatic Sciences*
483 39:233-236.
- 484 Masser MP, Rouse DB. 1997. Australian redclaw crayfish. SRAC Publication No. 244:1-8.
- 485 Mikkola H. 1996. Alien freshwater crustacean and indigenous mollusc species with aquaculture
486 potential in Eastern and Southern Africa. *South African Journal of Science* 22:90-99.
- 487 MTPA, Mpumalanga Tourism and Parks Agency. 2013. Ecstatus of the Crocodile River
488 Catchment, Inkomati River System. Report submitted to the Inkomati Catchment
489 Management Agency.
- 490 Nunes AL, Douthwaite RJ, Tyser B, Measey GJ, Weyl OLF. 2016. Invasive crayfish threaten
491 Okavango Delta. *Frontiers in Ecology and the Environment* 14:237-238.

- 492 Nunes AL, Hoffman AC, Zengeya TA, Measey GJ, Weyl OLF. *in press*. Red swamp crayfish,
493 *Procambarus clarkii*, found in South Africa 22 years after attempted eradication. *Aquatic*
494 *Conservation: Marine and Freshwater Ecosystems*.
- 495 Patoka J, Wardiatno Y, Yonvitner, Kuříková P, Petrtýl M, Kalous L. 2016. *Cherax*
496 *quadricarinatus* (von Martens) has invaded Indonesian territory west of the Wallace Line:
497 evidences from Java. *Knowledge and Management of Aquatic Ecosystems* 417:39.
- 498 Pienkowski T, Williams S, McLaren K, Wilson B, Hockley N. 2015. Alien invasions and
499 livelihoods: Economic benefits of invasive Australian red claw crayfish in Jamaica.
500 *Ecological Economics* 112:68-77.
- 501 Ray WJ, Corkum LD. 2001. Habitat and site affinity of the round goby. *Journal of Great Lakes*
502 *Research* 27:329-334.
- 503 RSA, Republic of South Africa. 2016. National Environmental Management: 395 Biodiversity
504 Act 396 (10/2004): Alien and Invasive Species List. Government Printer, Pretoria.
- 505 Snovsky G, Galil BS. 2011. The Australian redclaw crayfish *Cherax quadricarinatus* (von
506 Martens, 1868) (Crustacea: Decapoda: Parastactidae) in the Sea of Galilee, Israel. *Aquatic*
507 *Invasions* 6:S29-S31.
- 508 Todd S. 2005. The introduced red claw crayfish in Jamaica. Jamaica Clearing-House
509 Mechanism, Institute of Jamaica, Jamaica. Available at
510 <http://jamaicachm.org.jm/PDF/April2005.pdf> (accessed 13 October 2016).
- 511 Torres-Montoya EH, Salomón-Soto VM, Bucio-Pacheco M, Torres-Avendaño JI, López-Ruiz
512 M, Sánchez-González S, Castillo-Ureta H. 2016. First record of wild populations of *Cherax*
513 *quadricarinatus* (Decapoda: Parastacidae) in Sinaloa, Mexico. *Revista Mexicana de*
514 *Biodiversidad* 87:258-260.

- 515 Van den Berg RA, Schoonbee HJ. 1991. Freshwater crayfish species of *Cherax* (Decapoda:
516 Parastacidae) under investigation in the Zoology Department of the Rand Afrikaans
517 University - A preliminary report. Johannesburg, 177-185.
- 518 Vega-Villasante F, Ávalos-Aguilar JJ, Nolasco-Soria H, Vargas-Ceballos MA, Bortolini-Rosales
519 JL, Chong-Carrillo O, Ruiz-Núñez MF, Morales-Hernández JC. 2015. Wild populations of
520 the invasive Australian red claw crayfish *Cherax quadricarinatus* (Crustacea, Decapoda) near
521 the northern coast of Jalisco, Mexico: a new fishing and profitable resource. *Latin American
522 Journal of Aquatic Research* 43:781-785.
- 523 Weinländer M, Füreder L. 2009. The continuing spread of *Pacifastacus leniusculus* in Carinthia
524 (Austria). *Knowledge and Management of Aquatic Ecosystems* 394-395:17.
- 525 Weyl OLF, Daga VS, Ellender BR, Vitule JRS. 2016. A review of *Clarias gariepinus* invasions
526 in Brazil and South Africa. *Journal of Fish Biology* 89:386-402.
- 527 Williams EW Jr, Bunkley-Williams L, Lilyestrom CG, Ortiz-Corps EA. 2001. A review of
528 recent introductions of aquatic invertebrates in Puerto Rico and implications for the
529 management of nonindigenous species. *Caribbean Journal of Science* 37:246-251.

Figure 1

Study area in South Africa and Swaziland.

General overview of the study area showing the 46 sampling sites used in this study. Full circles and triangles  respectively represent river and dam sites where crayfish was found, empty circles and triangles represent river and dam sites where crayfish was not detected.




Figure 2

Detailed view of the four main study areas, with the 46 sampling sites used in this study.

(A) The Inkomati, (B) Mbuluzi and (C) Usutu river basins and (D) Richard's Bay area. The approximate point of first introduction of *C. quadricarinatus* in the Komati River and the potential point of introduction in the Mbuluzi River are indicated with red stars. Full circles and triangles respectively represent river and dam sites where crayfish was found, empty circles and triangles represent river and dam sites where crayfish was not detected. Black stars indicate sites where crayfish presence has been previously reported.




Figure 3

Length-frequency distributions of *C. quadricarinatus* in different locations of the Komati, Mbuluzi, Mlawula, Usutu and Lomati rivers and in irrigation dams.

(A) Wet season and (B) Dry season.




Figure 4

Relationship between size (cephalothorax length, in mm) and distance to crayfish introduction source for *C. quadricarinatus* in the Komati River during the wet and dry seasons.

(A) Females and (B) Males.





Table 1 (on next page)

Attributes of the 22 sites where **C. quadricarinatus** was found.

Coordinates, location, elevation (m), distance to closest crayfish introduction point (km), season, catch per unit effort (CPUE, as number of individuals and biomass), average size (cephalothorax length, mm), average mass (g) and number of males and females, for each sampling site where crayfish was found. SD stands for standard deviation, M for males, F for females, SA for South Africa and SW for Swaziland.

Site	Coordinates	Location	Elevation (m)	Distance to intro (km)	Season	CPUE (SD) (N/trap/night)	CPUE (SD) (g/trap/night)	Size M (SD)	Size F (SD)	Mass M (SD)	Mass F (SD)	M	F
K01	25°28'24.50"S 30°07'23.61"E	Komati River, SA	130	112.11	Wet	2.2 (1.39)	40.48 (26.42)	57.17 (7.28)	56 (9.59)	50.83 (19.13)	49.4 (22.49)	12	10
					Dry	3.2 (5.07)	33.01 (32.09)	53.26 (19.35)	50.97 (7.69)	34.88 (26.29)	32.93 (14.98)	17	15
K02	25°31'19.3"S 31°55'48.2"E	Komati River, SA	153	97.26	Wet	3.1 (4.36)	53.56 (53.39)	66.07 (18.21)	54.41 (9.47)	77.57 (58.17)	39.76 (21.34)	14	17
					Dry	1.43 (1.81)	39.89 (54.69)	63.35 (19.82)	46.63 (6.19)	81.33 (62.52)	32 (15.41)	6	4
K03	25°32'45.8"S 31°50'59.2"E	Komati River, SA	174	81.96	Wet	9.4 (7.73)	34.71 (16.73)	50.08 (9.04)	47.55 (7.79)	30.65 (18.63)	24.07 (12.55)	65	29
					Dry	4.43 (4.96)	66.4 (56.65)	57.80 (14.20)	49.21 (6.19)	52.67 (43.58)	28.63 (10.99)	15	16
K04	25°38'01.7"S 31°47'47.5"E	Komati River, SA	198	61.11	Wet	0.38 (0.74)	23.75 (44.01)	72.33 (16.26)	-	96 (64.09)	-	3	0
					Dry								
K05	25°43'29.4"S 31°46'49.8"E	Komati River, SA	233	44.94	Wet	2.88 (5.49)	32.42 (39.75)	56.92 (14.69)	46.55 (6.85)	53.17 (42.59)	25.09 (12.37)	12	11
					Dry	7 (11.93)	47.15 (32.96)	63.35 (11.86)	64.22 (5.11)	62.27 (31.55)	58.96 (13.97)	22	27
K06	25°51'19.4"S 31°48'27.9"E	Komati River, SA	252	21.76	Wet	1 (1.77)	29.15 (47.97)	81.83 (10.53)	46 (2.65)	138.33 (55.09)	19.5 (5.26)	6	4
					Dry	1 (1.73)	50.31 (89.19)	91.35 (8.32)	64 (0)	202.67 (45.23)	58 (0)	6	1
L01	25°36'58.6"S 31°39'48.7"E	Lomati River, SA	233	87.49	Wet	0.1 (0.32)	3 (9.49)	-	37 (0)	-	30 (0)	0	1
					Dry	0.71 (0.95)	40.14 (68.72)	76.52 (16.16)	45.56 (0)	131.5 (75.44)	18 (0)	4	1
L02	25°37'53.1"S 31°39'19.0"E	Lomati River, SA	236	89.69	Wet	0.1 (0.32)	19.2 (60.72)	95 (0)	-	192 (0)	-	1	0
					Dry	0 (0)	0 (0)	-	-	-	-	0	0
L03	25°38'55.9"S 31°40'10.7"E	Lomati River, SA	238	93	Wet	0 (0)	0 (0)	-	-	-	-	0	0
					Dry	0.2 (0.45)	38 (84.97)	-	98.24 (0)	-	190 (0)	0	1
MB01	26°08'05.6"S 31°59'48.4"E	Mbuluzi River, SW	163	23.14	Wet	4.5 (6.63)	19.78 (24.01)	62.65 (10.12)	54.4 (12.35)	53.65 (24.28)	37.3 (29.32)	17	10
					Dry	1 (1.41)	11.19 (14.48)	29.2 (7.05)	27.33 (7.02)	17.4 (12.19)	15.33 (11.68)	5	3
MB02	26°10'00.5"S 31°53'50.7"E	Mbuluzi River, SW	194	6.06	Wet	1 (1.41)	24.21 (34.31)	63.17 (18.76)	58 (5.66)	69.83 (42.83)	41 (11.31)	6	2
					Dry	1 (1.41)	44.38 (56.99)	49.2 (16.93)	49 (1.41)	120 (87.56)	81 (5.66)	5	2
ML01	26°10'34.6"S 31°59'28.8"E	Mlawula River, SW	147	47.5	Wet	1.57 (2.44)	8.99 (12.56)	49.29 (12.27)	40.75 (3.59)	30.43 (33.11)	13 (3.92)	7	4
					Dry	0 (0)	0 (0)	-	-	-	-	0	0
ML02	26°11'16.4"S 31°59'12.4"E	Mlawula River, SW	155	50	Wet	4 (3.67)	21.97 (14.24)	51.38 (8.39)	49.91 (7.18)	30.67 (16.48)	27.09 (11.47)	21	11
					Dry	0.86 (0.9)	10.71 (11.22)	29 (0)	31.2 (6.30)	13 (0)	21.8 (14.06)	1	5
US01	26°46'57.5"S 31°59'04.3"E	Usutu River, SW	79	-	Wet	0.8 (2.53)	7.84 (24.78)	71.8 (17.68)	68.33 (9.87)	85.2 (55.78)	67 (25.24)	5	3
					Dry	0.57 (1.13)	11.43 (23.61)	34.5 (9.75)	-	28.50 (23.06)	-	4	0
US02	26°51'26.8"S 31°54'29.3"E	Usutu River, SW	95	-	Wet	0.1 (0.32)	2.2 (6.96)	50 (0)	-	22 (0)	-	1	0
					Dry	0.14 (0.38)	2.43 (6.43)	31 (0)	-	17 (0)	-	1	0
USCh	26°50'51.0"S 31°54'49.8"E	Channel by Usutu, SW	125	-	Wet	0.67 (1.16)	22.67 (39.26)	67 (14.14)	-	68 (39.59)	-	2	0
					Dry	0.29 (0.76)	1.86 (4.91)	26 (0)	20 (0)	11 (0)	15 (0)	1	1
D01	25°33'08.1"S 31°54'16.0"E	Dam, SA	190	-	Wet	15.3 (19.98)	64.51 (35.17)	66.34 (10.45)	62.06 (10.78)	78.39 (37.87)	59.42 (34.24)	118	35
					Dry	1.75 (2.12)	46.87 (33.26)	65.19 (12.02)	59.5 (16.36)	73.23 (48.46)	50 (46.78)	26	9
D02	25°32'57.1"S 31°53'37.0"E	Dam, SA	186	-	Wet	3.3 (2.83)	40.89 (19.09)	59.18 (9.62)	53.86 (8.83)	53.11 (26.12)	36.67 (15.98)	18	15
					Dry	1.9 (4.09)	13.22 (23.20)	56.12 (11.36)	54.5 (8.42)	45.8 (25.45)	37.78 (16.11)	10	9
D03	25°37'14.4"S	Dam, SA	190	-	Wet	2.67 (3.68)	36.77 (49.89)	66.71 (13.33)	65.27 (14.28)	84.95 (53.81)	74.91 (49.93)	21	11

	31°51'42.3"E													
D04	25°32'41.2"S	Dam, SA	188	-	Wet	1.11 (1.27)	49.44 (52.00)	69.57 (16.27)	53.33 (15.04)	102 (52.51)	54 (6.93)	7	3	
	31°50'20.3"E				Dry	5.57 (6.45)	42.48 (21.24)	61.44 (8.49)	51.4 (11.73)	55.05 (27.89)	33.47 (21.11)	19	19	
D05	25°51'52.5"S	Dam, SA	265	-	Wet	7.4 (8.93)	16.05 (11.68)	46.44 (9.97)	42.15 (5.78)	26.35 (15.97)	17.36 (8.89)	46	28	
	31°50'00.9"E				Dry	0.9 (1.10)	19.97 (26.92)	50.97 (13.26)	53.39 (10.99)	36.8 (27.73)	36 (15.41)	5	4	
D06	25°58'43.6"S	Sand River Dam, SW	295	-	Wet	0.13 (0.35)	1.13 (3.18)	38 (0)	-	9 (0)	-	1	0	
	31°42'42.8"E				Dry	0.38 (0.74)	5.38 (10.04)	33 (0)	27 (0)	24.5 (0.71)	13 (0)	2	1	

1