

Immune responsiveness to phytohemagglutinin displays species but not sex differences in three anuran species

Zhiqiang Zhang^{Corresp., 1}, Chenchen Jin², Kangshan Qu¹, Enrique Caviedes-Vidal^{3, 4}

¹ College of Animal Science and Technology, Anhui Agricultural University, Hefei, People's Republic of China

² College of Life Science, Anhui Agricultural University, Hefei, People's Republic of China

³ Instituto Multidisciplinario de Investigaciones Biológicas de San Luis, Consejo Nacional de Investigaciones Científicas y Técnicas, and Departamento de Bioquímica y Ciencias Biológicas, Universidad Nacional de San Luis, San Luis, Argentina

⁴ Department of Forest and Wildlife Ecology, University of Wisconsin-Madison, Madison, United States

Corresponding Author: Zhiqiang Zhang

Email address: zzq-003@163.com

Phytohemagglutinin (PHA)-induced skin swelling response is widely used as a rough surrogate of integrative cell-mediated and innate immunity across multiple vertebrate taxa due to its simplification and feasibility. However, little is known whether there are sex and interspecific differences of immune responsiveness to PHA in ectotherms, especially for anurans. Therefore, we studied sex and species differences of PHA response in three anurans, Asiatic toads (*Bufo gargarizans*), Dark-spotted frogs (*Pelophylax nigromaculatus*) and Mongolian toads (*Pseudepidalea raddei*), captured in northern regions of Anhui Province (China). Footpad thickness was measured prior to (0 h) and after (6, 12, 24, 48 and 72 h) a PHA injection and normalized against saline injection in the opposite footpad. Body mass was recorded at the beginning (0 h) and end of each assay (72 h). Results showed effects of PHA assay, sex and taxa on body mass. Relative maximum swelling response (PHA_{max}) ranged from 18.58 to 29.75 %, 9.77 to 20.56 % and 21.97 to 31.78 % and its occurrence over time was apparent 10.6 to 19.72 h, 7.74 to 14.01 h and 17.39 to 23.94 h postinjection for Asiatic toads, Dark-spotted frogs and Mongolian toads, respectively. Finally, the magnitude or timing of PHA_{max} in Dark-spotted frogs was significantly thinner and faster than in Mongolian toads, and Asiatic toads had an in-between value, not different from the other two species. The magnitude of PHA_{max} was significantly positively correlated with the timing of PHA_{max} considering individuals altogether, but not when analyzed within species. Our results indicate that male and female anuran species respond similarly to PHA antigen stimulation, but the magnitude and timing of PHA_{max} is species-specific. Briefly, we provide new evidence for the suitability of PHA assay in non-model anuran species with different body sizes, and exhort the need to further investigate the nature of PHA assay at the hematological and histological levels in order to extend its application in ecoimmunological studies of amphibians.

Immune responsiveness to phytohemagglutinin displays species but not sex differences in three anuran species

Zhiqiang Zhang^{1*}, Chenchen Jin², Kangshan Qu¹ and Enrique Caviedes-Vidal^{3,4}

¹College of Animal Science and Technology, Anhui Agricultural University, Hefei, People's Republic of China

²College of Life Science, Anhui Agricultural University, Hefei, People's Republic of China

³Instituto Multidisciplinario de Investigaciones Biológicas de San Luis, Consejo Nacional de Investigaciones Científicas y Técnicas, and Departamento de Bioquímica y Ciencias Biológicas, Universidad Nacional de San Luis, San Luis, Argentina

⁴Department of Forest and Wildlife Ecology, University of Wisconsin-Madison, Madison, United States

Short title: Sex and species changes of PHA response

Correspondence: Zhiqiang Zhang, College of Animal Science and Technology, Anhui Agricultural University, 130 Changjiang West Road, Shushan District, Hefei 230036, People's Republic of China.
Email: zzq-003@163.com

ABSTRACT

Phytohemagglutinin (PHA)-induced skin swelling response is widely used as a rough surrogate of integrative cell-mediated and innate immunity across multiple vertebrate taxa due to its

simplification and feasibility. However, little is known whether there are sex and interspecific differences of immune responsiveness to PHA in ectotherms, especially for anurans. Therefore, we studied sex and species differences of PHA response in three anurans, Asiatic toads (*Bufo gargarizans*), Dark-spotted frogs (*Pelophylax nigromaculatus*) and Mongolian toads (*Pseudepidalea raddei*), captured in northern regions of Anhui Province (China). Footpad thickness was measured prior to (0 h) and after (6, 12, 24, 48 and 72 h) a PHA injection and normalized against saline injection in the opposite footpad. Body mass was recorded at the beginning (0 h) and end of each assay (72 h). Results showed effects of PHA assay, sex and taxa on body mass. Relative maximum swelling response (PHA_{max}) ranged from 18.58 to 29.75 %, 9.77 to 20.56 % and 21.97 to 31.78 % and its occurrence over time was apparent 10.6 to 19.72 h, 7.74 to 14.01 h and 17.39 to 23.94 h postinjection for Asiatic toads, Dark-spotted frogs and Mongolian toads, respectively. Finally, the magnitude or timing of PHA_{max} in Dark-spotted frogs was significantly thinner and faster than in Mongolian toads, and Asiatic toads had an in-between value, not different from the other two species. The magnitude of PHA_{max} was significantly positively correlated with the timing of PHA_{max} considering individuals altogether, but not when analyzed within species. Our results indicate that male and female anuran species respond similarly to PHA antigen stimulation, but the magnitude and timing of PHA_{max} is species-specific. Briefly, we provide new evidence for the suitability of PHA assay in non-model anuran species with different body sizes, and exhort the need to further investigate the nature of PHA assay at the hematological and histological levels in order to extend its application in ecoimmunological studies of amphibians.

INTRODUCTION

Amphibians are experiencing a global biodiversity crisis derived from diseases, habitat destruction and altered ecosystems (Chulow, Harris & Mahony, 2015). A wide range of physiological traits including innate and adaptive immunity are important to cope with the

43 increasing number of stressors and pathogen threats faced in their lifetimes (*Carey, Cohen &*
44 *Rollins-Smith, 1999*). As other physiological traits, immune defense is also energetically costly to
45 mount, and a trade-off relationship may occur within the immune system or between
46 immunological defenses and other nutrient-demanding processes such as growth, reproduction
47 and thermoregulation based on limited total energy resources (*Lochmiller & Deerenberg, 2000*).
48 Nowadays, it is still an open question what immune parameter measures are appropriate to reflect
49 an organism's defense capacity against pathogens (*Downs, Adelman & Demes, 2014*). Among
50 several immunological assays, the phytohemagglutinin (PHA) skin-swelling test is one of the
51 most widely used techniques to evaluate the magnitude of vertebrate immune response (*Sheldon*
52 *& Verhulst, 1996; Boughton, Joop & Armitage, 2011; Vinkler & Albrecht, 2011; Brock, Murdock*
53 *& Martin, 2014*). The assay consists of a subcutaneous injection of PHA that induces infiltration
54 of leukocytes and where swelling is interpreted as an integrative immune response (*Christe,*
55 *Arlettaz & Vogel, 2000; Kennedy & Nager, 2006; Martin et al., 2006; Demas et al., 2011; Brock,*
56 *Murdock & Martin, 2014*) or as an inflammation in endotherms (*Vinkler et al., 2012; Bilková,*
57 *Vinklerová & Vinkler, 2015; Bilková et al., 2016*). However, inferences using this assay should be
58 carefully evaluated because not always a greater immune response or inflammation means better
59 defenses against infections or higher survival probabilities (*Graham, Allen & Read, 2005; Vinkler*
60 *et al., 2012*). To date, the histological base of this response has been demonstrated across
61 different vertebrate taxa [e.g., birds (*Martin et al., 2006*), bats (*Turmelle et al., 2010*), amphibians
62 (*Brown, Shilton & Shine, 2011*) and crocodiles (*Finger et al., 2013*)], although no apparent
63 correlation between swelling response and leukocyte profiles was reported in the fossorial
64 mammal *Ctenomys talarum* (*Merlo, Cutrera & Zenuto, 2014*). However, more recently PHA
65 swelling has been associated to leukocyte profiles and serum bactericidal activity in endotherms
66 (*Bilková, Vinklerová & Vinkler, 2015; Zhang & Zhao, 2015*), or parasite infection in rodents
67 (*Ctenomys talarum, Merlo, Cutrera & Zenuto, 2015*). Despite all the merits of this assay, a caveat

68 must be warned because there is still some debate about its meaning. An objection is that the
69 response to PHA of an organism might not be representative of the actual response of the same
70 host to pathogens (reviewed by *Graham, Allen & Read, 2005; Owen & Clayton, 2007; Brock,*
71 *Murdock & Martin, 2014*) or that energetic costs of mounting a response to PHA may vary in
72 small mammals (*Zhang, Qiu & Wang, 2011; Merlo et al., 2014*) and birds (*Ots et al., 2001;*
73 *Martin, Scheuerlein & Wikelski, 2003; Lee, Martin & Wikelski, 2005; Nilsson, Granbom &*
74 *Råberg, 2007*).

75 PHA assay was first used in ecoimmunological studies of amphibians in the northern leopard
76 frogs, *Rana pipiens*, showing noticeable alterations of the immune response after an exposure to
77 different pesticides (*Gilbertson et al., 2003*). The sites chosen to inject PHA are diverse for post-
78 metamorphic amphibians and none is applicable to all species, for example, toe webbing (*Brown,*
79 *Shilton & Shine, 2011*), footpad (*Fites et al., 2014*), thigh (*Gervasi & Foufopoulos, 2008*) and
80 lower leg below the knee (*Clulow, Harris & Mahony, 2015*). The histological mechanism of the
81 inflammatory response induced by PHA antigen was examined in the toe webbing of cane toads,
82 *Rhinella marinus* (*Brown, Shilton & Shine, 2011*), which displayed a rapid (<12 h) infiltration of
83 neutrophils, eosinophils and macrophages at the injection site, followed by an influx of
84 lymphocytes by 24 h postinjection. In addition, marked interspecific variations of leukocytes
85 were observed in the lower leg of green and golden bell frogs, *Litoria aurea* and striped marsh
86 frogs, *Limnodynastes Peronii* (*Clulow, Harris & Mahony, 2015*). PHA assay has been also used
87 to assess the immune responsiveness under different ecologically relevant conditions, such as
88 when amphibians are forced to accelerate the metamorphosis by exposition to environment
89 desiccation (*Gervasi & Foufopoulos, 2008*), treated with protein restricted diets (*Venesky et al.,*
90 *2012*), and exposed to density stress (*Clulow, Harris & Mahony, 2015*). In different vertebrate
91 species, PHA response peaks in a wide time window, from 3 to 48 h (*de Bellocq et al., 2006;*
92 *Martin et al., 2006; Turmelle et al., 2010; Xu & Wang, 2010; Brown, Shilton & Shine, 2011;*

Finger et al., 2013; *Josserand et al.*, 2015; *Zhang, Huang & Zhao*, 2015; *Bilková et al.*, 2016), with no apparent sex differences in some small mammals (*Zhang, Qiu & Wang*, 2011; *Merlo, Cutrera & Zenuto*, 2014) and amphibians (*Brown et al.*, 2015; *Clulow, Harris & Mahony*, 2015). Most up-to-date information available on the time course and temporal dynamics of the PHA responses within and among populations or species are for birds (*Navarro et al.*, 2003) and small mammals (*de Bellocq et al.*, 2006; *Zhang, Huang & Zhao*, 2015). Conversely, data on amphibians are still scarce, especially for those anurans living in different environments (*Clulow, Harris & Mahony*, 2015). In a scenario of increasing disease and toxic threats, it is mandatory the need to provide comparable information of individual health status induced by PHA antigen on key amphibian species (*Carey, Cohen & Rollins-Smith*, 1999; *Fites et al.*, 2014; *Clulow, Harris & Mahony*, 2015).

In line with this situation, our aim was to test PHA assay response across different species with different body masses to extend its use to other non-model anuran species with different morphological and physiological adaptations to their habitats. Accordingly, we provide an evaluation of the PHA-induced skin-swelling assay among three sympatric anuran species with different habitat use in a range of body mass (30 to 90 g) of Central China (Northern Anhui province), Dark-spotted frogs (*Pelophylax nigromaculatus*), Mongolian toads (*Pseudepidalea raddei*) and Asiatic toads (*Bufo gargarizans*). In the three anuran species, both Asiatic and Mongolian toads are terrestrial with slow movements, whereas Dark-spotted frogs dwell lentic ponds or rice fields and move fast in their habitats (*Fei et al.*, 2009a, b). In addition, Asiatic toads start their reproductive activities during hibernation (*Ji et al.*, 1995; *Zou et al.*, 1996) and are easily infected by multiple parasites (*Zhao et al.*, 2014). On the other side, while Mongolian toads (*Zhou*, 1996) and Dark-spotted frogs initiate their reproduction after hibernation (*Wu*, 1965; *Cao et al.*, 1996; *Jin, Qu & Zhang*, 2014; *Jin & Zhang*, 2014), the former is not infected by multiple digenetic trematodes (*Li & Gu*, 1978), but the latter, displays a 90.38% helminth and parasite

infection rate (*Zhu et al., 2014*). We predicted that: (i) Within species, both sexes will exhibit similar swelling responses. (ii) The temporal dynamics of the response to PHA will be species-specific, as well as the magnitude of the maximum swelling at the injection point in the footpad area. In addition, we tested the effect of the PHA assay on the body mass of the individuals and its influence on the swelling response.

MATERIALS AND METHODS

Animal husbandry

All animal procedures were carried out under the guidelines of the Animal Care and Use Committee of Anhui Agricultural University (Permit number: 20120410). During April and May, 2012, we captured nine and eight males and nine and ten females Asiatic and Mongolian toads, respectively, in Mengcheng county (116.55°E 33.25°N Anhui province, China), and eight males and eight females of Dark-spotted frogs in Feixi county (117.17°E 31.70°N Anhui province, China). Immediately after capture, animals were transported to an animal indoor facility of the Anhui Agricultural University in Hefei, and housed in aquarium tanks (40 × 50 × 90 cm). They were kept under natural ambient temperature (17-21°C) and photoperiod (around 12L:12D). Fresh dechlorinated tap water was provided *ad libitum* and mealworms were provided as their food every 2-3 days during trials.

PHA-induced skin-swelling test

In all cases PHA-induced skin-swelling responses were always assayed two or three days after capture. Animals were subcutaneously injected with a dose of 50 µl of the PHA solution (PHA-P, Sigma L-8754, Sigma-Aldrich, St Louis, MO, USA, 1 mg of PHA-P dissolved in 0.3 mL of saline solution) in the footpad area pointing to the junction of the second and third digits of the right hind foot, and the opposite left footpad as an intragroup control was injected with an equal volume of sterilized saline by using a 100 µl micro injector (Shanghai Anxiang micro injector factory, China). Footpad thickness was measured with a micrometer (Tesa Shopcal, Renens,

Switzerland) to ± 0.01 mm, prior to injection (time = 0 h), and then at 6, 12, 24, 48 and 72 h postinjection. Footpad thickness was measured six times at each time point by the same operator at the point of injection and the average value was used for analysis (Smits, Bortolotti & Tella, 1999; de Bellocq et al., 2006). Body mass (± 0.001 g) was recorded at time 0 h and then, 72 h after injection in each individual of each species. In order to minimize measurement biases due to different initial footpad thickness, the expressed percentages of PHA-induced swelling responses (%) were calculated as described by Brown et al. (2015) for cane toads (*Rhinella marina*).

$$100 * \left[\frac{\text{PHA thickness at x hour} - \text{PHA thickness at 0 hour}}{\text{PHA thickness at 0 hour}} \right] - \left[\frac{\text{Saline thickness at x hour} - \text{Saline thickness at 0 hour}}{\text{Saline thickness at 0 hour}} \right]$$

Statistics

Data are presented as means ± 1 SEM. Each measurement of PHA swelling response was replicated six times. A very high repeatability of these measurements was observed using intraclass correlation coefficients ($r = 0.99$, $P < 0.001$). Therefore, hereafter the mean value of the six measurements of the swelling response for each individual was used for analyses.

Differences among group means were analyzed using analyses of the variance and covariance. Assumptions of ANOVAs and ANCOVAs were tested and when not met, the nonparametric Kruskal-Wallis H test was used. Tukey's HSD test was carried out to evaluate pairwise comparisons after ANOVAs, and Dunn's multiple comparisons test was performed after Kruskal-Wallis H. Species, sex and PHA assay effects on body mass were evaluated using a mixed-design ANOVA with an autoregressive (order 1) covariance structure (Littell, Pendergast & Natarajan, 2000) including pre and post-assay body masses, sex and species as fixed factors and subjects nested in sex and species as random factors. A mixed-design ANOVA with an autoregressive

(order 1) covariance structure was performed to test differences within individuals of the swelling response at different times (6, 12, 24, 48 and 72 h) after the PHA injection in the three species studied, using time after injection and sex as fixed factors and subjects nested in sex as a random factor. The maximum skin swelling response (PHA_{max}) after the PHA injection of an individual was defined as the highest of the five values measured along trial time (i.e., 6, 12, 24, 48 and 72 h), and the time when PHA_{max} was reached, as the time of peak. PHA_{max} was contrasted among the three species using an ANOVA, using full factorial design with taxa and sex as predictor variables.

Likelihood comparisons of the Akaike Information Criterion (AIC_c) with a second-order bias correction values produced by models using different parameters (i.e., time-after PHA injection and sex as variables and mean body mass as a covariate) for the different species studied were used as tool to select the predictors that described the best model for PHA swelling response within species (*Burnham & Anderson, 2004*). Using this procedure, variables included in the models were time-after PHA injection and sex. The same process was used to select the predictor variables, taxa and sex, to analyze PHA_{max} . Since times of peak and transformed values did not meet the normality assumption (D'Agostino's K^2 test, $P < 0.05$), a Kruskal-Wallis H test was used to test differences among taxa and sex. The correlation between PHA_{max} and timing of the peak was analyzed using the Spearman's rank correlation coefficient. In all tests, statistical significance was accepted for $\alpha < 0.05$. Statistical analyses were performed using JMP Pro 12.2.1.

RESULTS

Effects of PHA test, sex and species on body mass

A mixed-design ANOVA was conducted to test PHA assay effect on body mass in male and female individuals of the three species studied. Results showed significant body mass pre- and 72 h post- PHA injection changes within subjects (Table 1). Pairwise comparisons showed that all groups, but Asiatic toad males lost body mass after the PHA assay (Fig. 1). Between subjects'

188 effects (i.e. sex and taxa) and all interactions were statistically significant (Table 1). Sex
 189 differences of body mass before the assay were apparent for Asiatic toads, while Dark-spotted
 190 frogs and Mongolian toads did not differ. After the assay, body mass differed significantly
 191 between sexes in all species (Fig. 1). Body mass differences between taxa were also apparent,
 192 Dark-spotted frogs were lighter than Mongolian toads and Asiatic toad males, and in turn, lighter
 193 than the females of Asiatic toads (Fig. 1).

194 **Table 1.** Results of a full factorial mixed-design ANOVA with an autoregressive (order 1)
 195 covariance structure (*Littell, Pendergast & Natarajan, 2000*) run to test changes within subjects
 196 of the body mass before and after the PHA test using assay, sex and taxa as fixed factors and the
 197 subjects nested in sex and taxa as random factors.

Source	d.f.	F	P
Assay	1, 43.3	146.952	<0.0001
Sex	1, 44	7.876	0.0074
Taxa	2, 43.9	83.365	<0.0001
Sex x assay	1, 43.6	8.767	0.0049
Sex x taxa	2, 43.9	13.683	<0.0001
Taxa x time	2, 43.4	13.107	<0.0001
Sex x taxa x	2, 43.5	4.988	0.0112
assay			




Figure 1. Effects of the phytohemagglutinin (PHA) assay, sex and taxa on body mass in the three anuran species studied, Asiatic toads, Dark-spotted frogs and Mongolian toads. Bars represent mean body mass \pm 1 S.E.M. Body masses of female and male individuals of each species before the PHA injections are represented by solid gray and black bars respectively, and 72 h postinjection body masses of females and males are represented by unfilled gray and black bars, respectively. Bars or groups of bars connected by either solid (pre vs. post PHA injection), dashed-double-dot (sex) or dashed (species) lines indicate significant differences of body masses by Tukey's HSD test ($P < 0.05$).

PHA-induced skin swelling response

Skin swelling response after PHA injection showed significant differences within individuals of the three species studied (Table 2, $P_s < 0.0001$). All species increased the thickness of the footpad area following the PHA injection reaching a maximum skin swelling response (Tukey's HSD test, $P < 0.05$) and then the swelling significantly declined to a minimum at 72 h (Fig. 2). A closer

inspection of PHA_{max} and its occurrence over time (Fig. 3) showed that both parameters varied within species: the 95% CI of PHA_{max} and times of peak ranged from 18.58 to 29.75 % and 10.6 to 19.72 h, 9.766 to 20.56% and 7.738 to 14.01 h, and 21.97 to 31.78 % and 17.39 to 23.94 h for Asiatic toads, Dark-spotted frogs and Mongolian toads, respectively. Sex effect or interaction time x sex were not significant in the three species (Table 2). Between species differences in the magnitude of PHA_{max} were significant (ANOVA, taxa effect: $F_{2,49} = 4.1664$; $P < 0.022$, Fig. 3A), but not for sex ($F_{1,49} = 0.4994$; $P = 0.484$) or taxa x sex interaction ($F_{1,49} = 0.6600$, $P = 0.522$). Tukey's HSD test showed that maximum PHA-induced swelling response of Dark-spotted frogs was significantly thinner than that of Mongolian toads, and Asiatic toads PHA_{max} was not different from Mongolian and Dark-spotted frogs (Fig. 3A). Timing of PHA_{max} peaks between species was also statistically significant (Kruskal-Wallis test: $H_2 = 12.2179$, $P = 0.0022$, Fig. 3B) with score means of 23.33, 17.5 and 33.89 for Asiatic toads, Dark-spotted frogs and Mongolian toads, respectively. Dunn's multiple comparisons test showed that Dark-spotted frogs reached PHA_{max} significantly faster than Mongolian toads, and Asiatic toads had an in-between value, not different from any of the other two species (Fig. 3B). The relationship between PHA_{max} and timing of the peak between data of all individuals was significant (Spearman's rank correlation coefficient, $\rho = 0.327$, $P < 0.05$), but we fail to find a significant correlation ($P > 0.05$) within species with or without sex as a category.

Table 2. A full factorial mixed-design ANOVA with an autoregressive (order 1) covariance structure (Littell, Pendergast & Natarajan, 2000) was run to test changes of the swelling response after the PHA injection using time after PHA injection, sex and taxa as fixed factors, and the

232 subjects nested in sex and taxa as random factors. Sex and taxa were included as predictor
 233 variables and body mass excluded as a covariate based on likelihood comparisons of AICc of the
 234 different models (*Burnham & Anderson, 2004*).

Species	time after PHA injection			sex			time x sex		
	<i>d.f.</i>	<i>F</i>	<i>P</i>	<i>d.f.</i>	<i>F</i>	<i>P</i>	<i>d.f.</i>	<i>F</i>	<i>P</i>
Asiatic toads	4, 53.5	11.42	<0.0001	1, 54.4	1.07	0.306	4, 53.9	0.49	0.746
Dark-spotted frogs	4, 37.9	8.18	<0.0001	1, 14.7	1.69	0.214	4, 39	1.92	0.127
Mongolian toads	4, 46	31.80	<0.0001	1, 38.9	0.21	0.651	4, 46.5	1.41	0.246




Figure 2. Mean (\pm 1 S.E.M.) phytohemagglutinin (PHA)-induced skin swelling response, expressed as the percentage of the initial footpad area thickness corrected by the swelling that produce an injection of the same amount of saline solution in the left footpad (see Materials and methods), at 6, 12, 24, 48 and 72 h after PHA injections in Asiatic toads (A), Dark-spotted frogs (B) and Mongolian toads (C). Points not connected by the same lower case letter are significantly different (Tukey's HSD test, $P < 0.05$).




Figure 3. Mean values \pm 95% C.I. of (A) maximum phytohemagglutinin-induced skin swelling response (PHA_{max}) and (B) the times of peak of Asiatic toads, Dark-spotted frogs and Mongolian toads. Bars connected by brackets are significantly different ($P < 0.05$) by Tukey's HSD test and Dunn's multiple comparisons test in (A) and (B), respectively.

DISCUSSION

Consistent with our predictions, male and female anurans responded similarly to PHA (Table 2), suggesting that PHA assay can be used as an effective index for measuring intra-population variations in immune functions within either sex. In addition, PHA assay as an integrative response is comparable in ecoimmunological studies of amphibians with different body sizes, early or delayed primary PHA_{max} may match with their reproductive activities and parasite infections from outdoor environmental conditions.

Effects of the PHA assay, sex and taxa on body mass were marked in the three anuran species (Table 1). All groups except for Asiatic toad males lost body mass after the PHA assay (Fig. 1), suggesting that it is necessary to measure their body masses at least prior to injection (at 0 h) and

at 72 h postinjection. Compared with the toe webbing of large-bodied cane toads (*Brown, Shilton & Shine, 2011*) and the lower leg below the knee of several small-bodied tree frogs (*Chulow, Harris & Mahony, 2015*), the footpad site injected also displayed a measurable inflammation response after PHA antigen injection, which can be further applied to other anuran species with body sizes in a range of at least 30 to 90 g. As suggested from bird (*Smits, Bortolotti & Tella, 1999*) and later from amphibian (*Josserand et al., 2015*) studies on PHA measurement, a more simplified protocol without saline injection should be considered in future research in amphibians.

In the present study, no sex differences were found for immune responsiveness to PHA in the three anuran species (Table 2, Fig. 2). Similar results had been recorded in chinstrap penguins, *Pygoscelis antarctica* (*Moreno et al., 1998*), wild rodent species (*Zhang, Qiu & Wang, 2011; Merlo, Cutrera & Zenuto, 2014*), and several anuran species (*Brown et al., 2015; Chulow, Harris & Mahony, 2015*). The PHA inflammation assay uses a lectin to elicit localized inflammation that reflects an organism's capacity to mount a dynamic innate and cell-mediated immune response. In a wild rodent, immune responsiveness to primary PHA was positively correlated with the proportion of neutrophils and serum bactericidal capacity in circulating blood (*Zhang & Zhao, 2015*), and the energetic cost of mounting this response was low (*Zhang, Qiu & Wang, 2011; Merlo et al., 2014*). In amphibians, the primary PHA injection may mainly involve innate phagocytes and granulocytes, and a second PHA injection would strongly activate lymphocytes (*Brown, Shilton & Shine, 2011; Fites et al., 2014; Josserand et al., 2015*). In the present study, the activation of the innate immunity induced by a single dose of PHA antigen may represents male and female anurans responses to cope with infectious diseases and changing environmental conditions. During April and May, Asiatic toads almost terminated their reproductive activities (*Ji et al., 1995; Zou et al., 1996*), whereas Mongolian toads (*Zhou, 1996*) and Dark-spotted frogs just initiated their reproductive activities (*Wu, 1965; Cao et al., 1996; Jin, Qu & Zhang, 2014; Jin*

& Zhang, 2014). Male serum testosterone concentrations are higher during the reproductive than in the non-reproductive period in Dark-spotted frogs (Hu, 1990), or before reproduction than after reproduction in Asiatic toads (Hu, 1989). Based in life history theory, it can be hypothesized that males may show a reduced immune response and an increased susceptibility to disease than females due to the suppressive effects of androgens on the immune system (Zuk & McKean, 1996). Hence, a suppression or at least a reduction of male immune responses is a reasonable prediction if a trade-off relationship exists between immunity and reproduction (Lochmiller & Deerenberg, 2000). Moreover, male anurans might face a greater risk exposed to more infectious diseases than females because they need croak and move to find a mate during reproduction, which is energetically costly too (Hawley & Altize, 2011). Strikingly in this study, despite the different reproductive status, vocal and locomotor activity, no apparent dissimilarities were found between male and female anurans in the immune responsiveness pattern to PHA (Fig. 2). Sex differences in immune functions may be associated to the immunological parameters (Boughton, Joop & Armitage, 2011; Downs, Adelman & Demes, 2014) or vertebrate life history characters involved (Stoehr & Kokko, 2006). For example, semi-terrestrial salamanders, *Desmognathus ochrophaeus* females were reported to heal slower than males from an experimental cutaneous wound, used as a measure of an integrative immune response, and the difference in healing time was related to potential energetic constraints (Thomas & Woodley, 2015). However, it has also described that *Bufo americanus* females and larger males display similar proportions of heterophils acting as first-line defenses against bacterial infection in their leukocyte profiles (Forbes, McRue & Shutler, 2006).

In a range from fast to slow life span rodents, empirical evidences show that different components of the immune system might display similar or different changing tendencies (Martin, Hasselquist & Wikelski, 2006; Martin, Weil & Nelson, 2007; Previtali et al., 2012). As an integrative response, the temporal patterns of the PHA response might change in different

rodents, and individuals that responded faster had a lower maximum response than those with a more delayed response (*de Bellocq et al., 2006*). In twelve Lark species, environmental proxies of antigen exposure explained variation in immune investment better than proxies of pace of life (*Horrocks et al., 2015*). In anurans, unlike endotherms and regardless of body size, the primary response to PHA antigen injections is mainly attributed to innate immunity of immune system and recognized as the first-line defense to outdoor pathogens (*Brown, Shilton & Shine, 2011; Josserand et al., 2015*). The interspecific differences of the magnitude or timing of PHA_{max} were observed in the three anuran species (Table 2, Fig. 3), suggesting that a trade-off relationship between intensity and latency of the PHA response may hold among anurans. In adult anurans, PHA_{max} may be reached at 12 h (*Clulow, Harris & Mahony, 2015*), 14 h (*Josserand et al., 2015*), 24 h (*Brown, Shilton & Shine, 2011*), or 48 h (*Clulow, Harris & Mahony, 2015*) after PHA injection. For post-metamorphic amphibians, a range of ecologically meaningful environmental stressors, such as pond desiccation (*Gervasi & Foufopoulos, 2008*), different protein concentrations of diets (*Venesky et al., 2012*), and low and high population density (*Clulow, Harris & Mahony, 2015*) have great influences on the inflammation response induced by PHA antigen. Asiatic and Mongolian toads, and Dark-spotted frogs are sympatric common species, but the two formers are terrestrial and disperse slowly, and the latter inhabit lentic ponds and disperse fast by swimming or jumping (*Fei et al., 2009a, b*). It has been reported that the helminth and parasite infection rates in Dark-spotted frogs and Asiatic toads are 90.38% and 100%, respectively, and the average intensity of infection is 20.21 and 28.34, respectively in Shanghai (*Zhu et al., 2014; Zhao et al., 2014*). However, Mongolian toads were not infected by multiple digenetic trematodes from north China (*Li & Gu, 1978*). PHA-triggered inflammation was similarly impaired by *Eimeria sp.* infection alone or co-occurring with a number of gastrointestinal nematodes in the subterranean rodent, *Ctenomys talarum* (*Merlo, Cutrera & Zenuto, 2015*). In the present study, Dark-spotted frogs with an early PHA response did show a

lower peak response than Mongolian toads with a late response (Fig. 3), suggesting that parasite infections may evoke the changes of PHA response patterns. However, in *Xenopus laevis*, swelling induced by single injection of PHA was not significantly affected by the fungus disease, *Batrachochytrium dendrobatidis* supernatants, but swelling caused by a secondary injection of PHA was significantly reduced by *B. dendrobatidis* supernatants (Fites et al., 2014). It is noted that the magnitude of PHA_{max} was significantly positively correlated with the timing of PHA_{max} between data of all individuals from the three anuran species, but not for the data within species, suggesting that the magnitude and timing of PHA_{max} is species-specific.

Overall, we found that immune responsiveness to PHA did not differ between sexes for any anuran species (Table 2, Fig. 2), but PHA_{max} and its timing for toads and frogs varied among species (Fig. 3). This scenario allows PHA assay to be applied to a broad range of amphibian species under field conditions, as a simple, inexpensive and effective proxy to quantify and evaluate a facet of immune response. For future ecoimmunological studies of amphibians, PHA assay will play an important role by integrating the changes of other components of immune system facing newly emerging pathogens or environmental factors. Lastly, we caution that it is still necessary to test the validation of PHA assay in advance for a new species by using appropriate internal controls before carrying out studies under specific circumstances.

ACKNOWLEDGEMENTS

Thanks to Xinxin NIU, Zhenfei XIAN, and Shuli HUANG from Anhui Agricultural University for help in the field.

REFERENCES

Bílková B, Albrecht T, Chudíčková M, Holář V, Piálek J., Vinkler M. 2016. Application of Concanavalin A during immune responsiveness skin-swelling tests facilitates measurement interpretation in mammalian ecology. *Ecology and Evolution* **6**: 4551–4564 DOI:

- 354 10.1002/ece3.2211.
- 355 **Bílková B, Vinklerová J, Vinkler M. 2015.** The relationship between health and cell-mediated
356 immunity measured in ecology: Phytohaemagglutinin skin-swelling test mirrors blood cellular
357 composition. *Journal of Experimental Zoology Part A: Ecological Genetics and Physiology*
358 **323**: 767–777 DOI: 10.1002/jez.1990.
- 359 **Boughton RK, Joop G, Armitage SAO. 2011.** Outdoor immunology: methodological
360 considerations for ecologists. *Functional Ecology* **25**: 81–100 DOI: 10.1111/j.1365-
361 2435.2010.01817.x.
- 362 **Brock PM, Murdock CC, Martin LB. 2014.** The history of ecoimmunology and its integration
363 with disease ecology. *Integrative and Comparative Biology* **54**: 353–362 DOI:
364 10.1093/icb/icu046.
- 365 **Brown GP, Phillips BL, Dubey S, Shine R. 2015.** Invader immunology: invasion history alters
366 immune system function in cane toads (*Rhinella marina*) in tropical Australia. *Ecology Letters*
367 **18**: 57–65 DOI: 10.1111/ele.12390.
- 368 **Brown GP, Shilton CM, Shine R. 2011.** Measuring amphibian immunocompetence: validation
369 of the phytohemagglutinin skin-swelling assay in the cane toad, *Rhinella marinus*. *Methods in*
370 *Ecology and Evolution* **2**: 341–348 DOI: 10.1111/j.2041-210X.2011.00090.x.
- 371 **Burnham KP, Anderson DR. 2004.** Multimodel inference understanding AIC and BIC in model
372 selection. *Sociological Methods and Research* **33**: 261–304 DOI: 10.1177/0049124104268644.
- 373 **Cao YP, Bai M, Ma R, Chen HB, Zhang Z, Li Z, Ren SS. 2000.** Studies on the ecology and
374 morphological physiology of *Rana nigromaculata* before and after hibernation. *Sichuan*
375 *Journal of Zoology* **19**: 159–162 DOI: 10.3969/j.issn.1000-7083.2000.03.019.
- 376 **Carey C, Cohen N, Rollins-Smith L. 1999.** Amphibian declines: an immunological perspective.
377 *Developmental and Comparative Immunology* **23**: 459–472 DOI: 10.1016/S0145-
378 305X(99)00028-2.

- 379 **Christe P, Arlettaz R, Vogel P.** 2000. Variation in intensity of a parasitic mite (*Spinturnix myoti*)
380 in relation to the reproductive cycle and immunocompetence of its bat host (*Myotis myotis*).
381 *Ecology Letters* **3**: 207–212 DOI: 10.1046/j.1461-0248.2000.00142.x.
- 382 **Clulow S, Harris M, Mahony MJ.** 2015. Optimization, validation and efficacy of the
383 phytohaemagglutinin inflammation assay for use in ecoimmunological studies of amphibians.
384 *Conservation Physiology* **3**: DOI: 10.1093/conphys/cov042.
- 385 **de Bellocq JG, Krasnov BR, Khokhlova IS, Pinshow B.** 2006. Temporal dynamics of a T-cell
386 mediated immune responses in desert rodents. *Comparative Biochemistry and Physiology. A:*
387 *Molecular and Integrative Physiology* **145**: 554–559 DOI: 10.1016/j.cbpa.2006.08.045.
- 388 **Demas GE, Zysling DA, Beechler BR, Muehlenbein MP, French SS.** 2011. Beyond
389 phytohaemagglutinin: assessing vertebrate immune function across ecological contexts.
390 *Journal of Animal Ecology* **80**: 710–730. DOI: 10.1111/j.1365-2656.2011.01813.x.
- 391 **Downs CJ, Adelman JS, Demas GE.** 2014. Mechanisms and methods in ecoimmunology:
392 integrating within-organism and between-organism processes. *Integrative and Comparative*
393 *Biology* **54**: 340–352 DOI: 10.1093/icb/icu082.
- 394 **Fei L, Hu SQ, Ye CY, Tian WS, Jiang JP, Wu GF, Li J, Wang YS.** 2009a. *Anura. Amphibian*
395 *Volume 2. Fauna Sinica.* Bei Jing: Science Press. (in Chinese)
- 396 **Fei L, Hu SQ, Ye CY, Tian WS, Jiang JP, Zhong SX, Li J, Wang YS.** 2009b. *Anura Ranidae.*
397 *Amphibian Volume 3. Fauna Sinica.* Bei Jing: Science Press. (in Chinese)
- 398 **Finger JW, Amanda AL, Thomson PC, Shilton CM, Brown GP, Moran C, Miles LG, Glenn**
399 **TC, Isberg SR.** 2013. Using phytohaemagglutinin to determine immune responsiveness in
400 saltwater crocodiles (*Crocodylus porosus*). *Australia Journal of Zoology* **61**: 301–311 DOI:
401 org/10.1071/ZO13041.

- 402 **Fites JS, Reinert LK, Chappell TM, Rollins-Smith L. 2014.** Inhibition of local immune
403 responses by the frog-killing fungus *Batrachochytrium dendrobatidis*. *Infection and Immunity*
404 **82:** 4698–4706 DOI: 10.1128/IAI.02231-14.
- 405 **Forbes MR, McRuer DL, Shutler D. 2006.** White blood cell profiles of breeding American
406 toads (*Bufo americanus*) relative to sex and body size. *Comparative Clinical Pathology* **15:**
407 155–159 DOI: 10.1007/s00580-006-0623-5.
- 408 **Gilbertson MK, Haffner GD, Drouillard KG, Albert A, Dixon B. 2003.** Immunosuppression
409 in the northern leopard frog (*Rana pipiens*) induced by pesticide exposure. *Environmental*
410 *Toxicology and Chemistry* **22:** 101–110 DOI: 10.1002/etc.5620220113.
- 411 **Goto N, Kodama H, Okada K, Fujimoto Y. 1978.** Suppression of phytohemagglutinin skin
412 response in thymectomized chickens. *Poultry Science* **57:** 246–250 DOI: 10.3382/ps.0570246.
- 413 **Graham AL, Allen JE, Read AF. 2005.** Evolutionary causes and consequences of
414 immunopathology. *Annual Review of Ecology Evolution and Systematics* **36:** 373–397 DOI:
415 10.1146/annurev.ecolsys.36.102003.152622.
- 416 **Hawley DM, Altize SM. 2011.** Disease ecology meets ecological immunology: understanding
417 the links between organismal immunity and infection dynamics in natural populations.
418 *Functional Ecology* **25:** 48–60 DOI: 10.1111/j.1365-2435.2010.01753.x.
- 419 **Horrocks NPC, Hegemann A, Ostrowski S, Ndithia H, Shobrak M, Williams JB, Matson**
420 **KD, Tieleman BI. 2015.** Environmental proxies of antigen exposure explain variation in
421 immune investment better than indices of pace of life. *Oecologia* **177:** 281-290 DOI:
422 10.1007/s00442-014-3136-y.
- 423 **HU ZG. 1989.** Studies on variance of sexual hormone in blood plasma of *Bufo bufo gargarizans*
424 before and after hibernation. *Journal of Anhui Normal University*, 45-51 DOI:
425 10.14182/cnki.1001-2443.1989.03.007.

- 426 **HU ZG. 1990.** Studies on variance of sexual hormone in blood plasma of *Rana nigromaculata*
427 during reproduction. *Chinese Journal of Zoology* **25**: 15–18 DOI: 10.13859/j.cjz.1990.03.006.
- 428 **Ji Q, Li ZY, Yang Y, Zhu JA, Tang JG. 1995.** Relationship between some serum hormones and
429 glucose metabolism in *Bufo gargarizans* during pre-hibernation, hibernation and post-
430 hibernation. *Chinese Journal of Zoology* **30**: 19–22 DOI: 10.13859/j.cjz.1995.02.006.
- 431 **Jin CC, Qu KS, Zhang ZQ. 2014.** Sexual and seasonal differences of the relative fatness and
432 organ mass of *Pelophylax nigromaculata*. *Sichuan Journal of Zoology* **33**: 106–112 DOI:
433 10.3969/j.issn.1000-7083.2014.01.021.
- 434 **Jin CC, Zhang ZQ. 2014.** Sexual and seasonal differences of digestive tract mass and length in
435 *Pelophylax nigromaculata*. *Chinese Journal of Zoology* **49**: 207–214 DOI:
436 10.13859/j.cjz.2014.02.020.
- 437 **Josserand R, Troïanowski M, Grolet O, Desprat JL, Lengagne T, Mondy N. 2015.** A
438 phytohaemagglutinin challenge test to assess immune responsiveness of European tree frog
439 *Hyla arborea*. *Amphibia-Reptili* **36**: 111–118 DOI: 10.1163/15685381-00002983.
- 440 **Kennedy MW, Nager RG. 2006.** The perils and prospects of using phytohaemagglutinin in
441 evolutionary ecology. *Trends in Ecology and Evolution* **21**: 653–655
442 DOI:10.1016/j.tree.2006.09.017.
- 443 **Lee KA, Martin LB, Wikelski MC. 2005.** Responding to inflammatory challenges is less costly
444 for a successful avian invader, the house sparrow (*Passer domesticus*), than its less-invasive
445 congener. *Oecologia* **145**: 244–251 DOI: 10.1007/s00442-005-0113-5.
- 446 **Li MM, Gu CD. 1978.** Report of some digenetic trematodes from frogs and toads of China. *Acta*
447 *Zoologica Sinica* **24**: 163–169.
- 448 **Littell RC, Pendergast J, Natarajan R. 2000.** Modelling covariance structure in the analysis of
449 repeated measures data. *Statistics in Medicine* **19**: 1793–1819 DOI: 10.1002/1097-
450 0258(20000715)19:13<1793::AID-SIM482>3.0.CO;2-Q.

- 451 **Lochmiller RL, Deerenberg C. 2000.** Trade-offs in evolutionary immunology: just what is the
452 cost of immunity? *Oikos* **88**: 87–98 DOI: 10.1034/j.1600-0706.2000.880110.x.
- 453 **Martin LB, Han P, Lewittes J, Kuhlman JR, Klasing KC, Wikelski M. 2006.**
454 Phytohemagglutinin-induced skin swelling in birds: histological support for a classic
455 immunoecological technique. *Functional Ecology* **20**: 290–299 DOI: 10.1111/j.1365-
456 2435.2006.01094.x.
- 457 **Martin LB, Hasselquist D, Wikelski M. 2006.** Immune investments are linked to pace of life in
458 house sparrows. *Oecologia* **147**: 565–575 DOI: 10.1007/s00442-005-0314-y.
- 459 **Martin LB, Weil ZM, Nelson RJ. 2007.** Immune defense and reproductive pace of life in
460 *Peromyscus* mice. *Ecology* **88**: 2516–2528 DOI: 10.1890/07-0060.1.
- 461 **Merlo JL, Cutrera AP, Luna F, Zenuto RR. 2014.** PHA-induced inflammation is not
462 energetically costly in the subterranean rodent *Ctenomys talarum* (tuco-tucos). *Comparative*
463 *Biochemistry and Physiology. A: Molecular and Integrative Physiology* **175**: 90–95 DOI:
464 10.1016/j.cbpa.2014.05.021.
- 465 **Merlo JL, Cutrera AP, Zenuto RR. 2014.** Inflammation in response to phytohemagglutinin
466 injection in the Talas tuco-tuco (*Ctenomys talarum*): implications for the estimation of
467 immunocompetence in natural populations of wild rodents. *Canadian Journal of Zoology* **92**:
468 689–697 DOI: 10.1139/cjz-2013-0306.
- 469 **Merlo JL, Cutrera AP, Zenuto RR. 2015.** Parasite infection negatively affects PHA-triggered
470 inflammation in the subterranean rodent *Ctenomys talarum*. *Journal of Experimental Zoology*
471 *Part A Ecological Genetics and Physiology* **325**: 132–141 DOI:10.1002/jez.2003.
- 472 **Moreno J, de León A, Faragallo JA, Moreno E. 1998.** Breeding time, health and immune
473 response in the chinstrap penguin *Pygoscelis Antarctica*. *Oecologia* **115**: 312–319 DOI:
474 10.1007/s004420050522.

- 475 **Navarro C, Marzal A, de Lope F, Møller AP. 2003.** Dynamics of an immune response in house
476 sparrows *Passer domesticus* in relation to time of day, body condition and blood parasite
477 infection. *Oikos* **101**: 291–298 DOI: 10.1034/j.1600-0706.2003.11663.x.
- 478 **Nilsson JÅ, Granbom M, Råberg L. 2007.** Does the strength of an immune response reflect its
479 energetic cost? *Journal of Avian Biology* **38**: 488–494 DOI: 10.1111/j.0908-
480 8857.2007.03919.x.
- 481 **Ots I, Kerimov AB, Ivankina EV, Ilyina TA, Hõrak P. 2001.** Immune challenge affects basal
482 metabolic activity in wintering great tits. *Proceedings of the Royal Society B Biological*
483 *Sciences* **268**: 1175–1181 DOI: 10.1098/rspb.2001.1636.
- 484 **Owen JP, Clayton DH. 2007.** Where are the parasites in the PHA response? *Trends in Ecology*
485 *and Evolution* **22**: 228–229 DOI:10.1016/j.tree.2007.02.003.
- 486 **Previtali MA, Ostfeld RS, Keesing F, Jolles AE, Hanselmann R, Martin LB. 2012.**
487 Relationship between pace of life and immune responses in wild rodents. *Oikos* **121**: 1483–
488 1492 DOI: 10.1111/j.1600-0706.2012.020215.x.
- 489 **Sheldon BC, Verhulst S. 1996.** Ecological immunology: costly parasite defences and trade-offs
490 in evolutionary ecology. *Trends in Ecology and Evolution* **11**: 317–321 DOI: 10.1016/0169-
491 5347(96)10039-2.
- 492 **Smits JE, Bortolotti GR, Tella JL. 1999.** Simplifying the phytohemagglutinin skin-testing
493 technique in studies of avian immunocompetence. *Functional Ecology* **13**: 567–572 DOI:
494 10.1046/j.1365-2435.1999.00338.x.
- 495 **Stoehr AM, Kokko H. 2006.** Sexual dimorphism in immunocompetence: what does life-history
496 theory predict? *Behavioral Ecology*, **17**: 751–756 DOI: 10.1093/beheco/ark018.
- 497 **Thomas JR, Woodley SK. 2015.** Treatment with corticosterone delays cutaneous wound healing
498 in male and female salamanders. *General and Comparative Endocrinology* **216**: 33–38 DOI:
499 10.1016/j.ygcen.2015.04.013.

- 500 **Turmelle AS, Ellison JA, Mendonc MT, McCracken GF. 2010.** Histological assessment of
- 501 cellular immune response to the phytohemagglutinin skin test in Brazilian free-tailed bats
- 502 (*Tadarida brasiliensis*). *Journal of Comparative Physiology. B: Biochemical Systemic and*
- 503 *Environmental Physiology* **180**: 1155–1164 DOI: 10.1007/s00360-010-0486-6.
- 504 **Venesky MD, Wilcoxon TE, Rensel MA, Rollins-Smith L, Kerby JL, Parris MJ. 2012.**
- 505 Dietary protein restriction impairs growth, immunity, and disease resistance in southern
- 506 leopard frog tadpoles. *Oecologia* **169**: 23–31 DOI: 10.1007/s00442-011-2171-1.
- 507 **Vinkler M, Albrecht T. 2011.** Handling immunocompetence' in ecological studies: do we
- 508 operate with confused terms? *Journal of Avian Biology* **42**: 490–493 DOI: 10.1111/j.1600-
- 509 048X.2011.05499.x.
- 510 **Vinkler M, Schnitzer J, Munclinger P, Albrecht T. 2012.** Phytohaemagglutinin skin-swelling
- 511 test in scarlet rosefinch males: low-quality birds respond more strongly. *Animal Behaviour*
- 512 **83**: 17–23 DOI:10.1016/j.anbehav.2011.10.001.
- 513 **Wu YL. 1965.** Changes of fatness and some internal organ sizes during natural hibernation in
- 514 *Pelophylax nigromaculata*. *Chinese Journal of Zoology* **7**: 116–119 DOI:
- 515 10.13859/j.cjz.1965.03.012.
- 516 **Xu DL, Wang DH. 2010.** Fasting suppresses T cell-mediated immunity in female Mongolian
- 517 gerbils (*Meriones unguiculatus*). *Comparative Biochemistry and Physiology. A: Molecular*
- 518 *and Integrative Physiology* **155**: 25–33 DOI: 10.1016/j.cbpa.2009.09.003.
- 519 **Zhang ZQ, Huang SL, Zhao ZJ. 2015.** Response patterns of phytohemagglutinin-induced skin
- 520 swelling at different reproductive phases in female striped hamster (*Cricetulus barabensis*).
- 521 *Acta Theriologica Sinica* **35**: 74–79. (in Chinese)
- 522 **Zhang ZQ, Qiu FT, Wang DH. 2011.** Sex and seasonal differences and energetic cost of
- 523 phytohemagglutinin responses in wild Mongolian gerbils (*Meriones unguiculatus*). *Acta*
- 524 *Theriologica Sinica* **31**: 284–290. (in Chinese)

- 525 **Zhang ZQ, Zhao ZJ. 2015.** Correlations between phytohemagglutinin response and leukocyte
526 profile, and bactericidal capacity in a wild rodent. *Integrative Zoology* **10**: 302–310 DOI:
527 10.1111/1749-4877.12133.
- 528 **Zhao QP, Dong H, Han HY, Wu YL, Zhu SH, Li LJ, Wu D, Liang ST, Li S, Zhai Q, Yang**
529 **SH, Pei EL, Huang B. 2014.** A preliminary investigation on the prevalence of parasites in
530 market toads in Shanghai. *Chinese Journal of Animal Infectious Diseases* **22**: 49–53
531 DOI:10.3969/j.issn.1674-6422.2014.05.008.
- 532 **Zhou HM. 1996.** Correlation analysis on some physiologic constants in *Bufo raddei* under
533 hibernation. *Sichuan Journal of Zoology* **15**: 162–164 DOI: 10.3969/j.issn.1000-
534 7083.1996.04.007.
- 535 **Zhu SH, Dong H, Zhao QP, Wu D, Wu YL, Li LJ, Han HY, Liang ST, Li S, Zhai Q, Yang**
536 **SH, Pei EL, Huang B. 2014.** A preliminary investigation on the prevalence of helminths in
537 frogs at Shanghai farmers' markets. *Chinese Journal of Animal Infectious Diseases* **22**: 53–
538 57 DOI: 10.3969/j.issn.1674-6422.2014.06.009.
- 539 **Zou SC, Zhu WZ, Chen WD, Xie YY, Ding Y, Jia ZC. 1996.** Study on relationship among
540 several physiological constants of *Bufo gargarizans* pre- and post-hibernation. *Sichuan*
541 *Journal of Zoology* **15**: 156–158 DOI: 10.3969/j.issn.1000-7083.1996.04.005.
- 542 **Zuk M, McKean KA. 1996.** Sex differences in parasite infections: patterns and processes.
543 *International Journal of Parasitology* **26**: 1009–1023 DOI: 10.1016/S0020-7519(96)00086-0.