

# Changes in the spatial and temporal pattern of natural forest cover on Hainan Island from the 1950s to the 2010s : implications for natural forest conservation and management

Siliang Lin <sup>1</sup> , Yaozhu Jiang <sup>1</sup> , Jiekun He <sup>1</sup> , Guangzhi Ma <sup>1</sup> , Yang Xu <sup>1</sup> , Haisheng Jiang <sup>Corresp. 1</sup>

<sup>1</sup> School of Life Science, South China Normal University, Guangzhou, Guangdong, China

Corresponding Author: Haisheng Jiang  
Email address: jhs@scnu.edu.cn

The study of the past, present, and future state and dynamics of the tropical natural forest cover (NFC) might help to better understand the pattern of deforestation and fragmentation as well as the influence of social and natural processes. The obtained information will support the development of effective conservation policies and strategies. In the present study, we used historical data of the road network, topography, and climatic productivity to reconstruct NFC maps of Hainan Island, China, from the 1950s to the 2010s, using the random forest algorithm. We investigated the spatial and temporal patterns of NFC change from the 1950s to the 2010s and found that it was highly dynamic in both space and time. Our data showed that grid cells with low NFC were more vulnerable to NFC decrease, suggesting that conservation actions regarding natural forests need to focus on regions with low NFC and high ecological value. We also identified the hot-spots of NFC change, which provides insights into the dynamic changes of natural forests over time.

# **Changes in the spatial and temporal pattern of natural forest cover on Hainan Island from the 1950s to the 2010s: implications for natural forest conservation and management**

Siliang Lin <sup>a†</sup>, Yaozhu Jiang <sup>a†</sup>, Jiekun He <sup>a</sup>, Guangzhi Ma <sup>a</sup>, Yang Xu <sup>a</sup>, Haisheng Jiang <sup>a\*</sup>

<sup>a</sup> School of Life Sciences, South China Normal University, Guangzhou, Guangdong 510631, China

† These authors contributed equally to this study.

**\* Correspondence authors:** Haisheng Jiang

School of Life Science, South China Normal University, Guangzhou, Guangdong 510631, China.

Telephone: +86 159 8914 7874

E-mail: jhs@scnu.edu.cn

# **Abstract**

The study of the past, present, and future state and dynamics of the tropical natural forest cover (NFC) might help to better understand the pattern of deforestation and fragmentation as well as the influence of social and natural processes. The obtained information will support the development of effective conservation policies and strategies. In the present study, we used historical data of the road network, topography, and climatic productivity to reconstruct NFC maps of Hainan Island, China, from the 1950s to the 2010s, using the random forest algorithm. We investigated the spatial and temporal patterns of NFC change from the 1950s to the 2010s and found that it was highly dynamic in both space and time. Our data showed that grid cells with low NFC were more vulnerable to NFC decrease, suggesting that conservation actions regarding natural forests need to focus on regions with low NFC and high ecological value. We also identified the hot-spots of NFC change, which provides insights into the dynamic changes of natural forests over time.

# INTRODUCTION

Human activity is the dominant cause of contemporary environmental change worldwide (Lewis & Maslin, 2015). Land use by humans is a major component of the long-term anthropogenic global changes that have defined the “Anthropocene” as a new epoch of geologic time (Ellis et al., 2013; Dirzo et al., 2014; Lewis & Maslin, 2015). The study of land use and land cover change was initially dominated by monitoring and modelling of the ecological impacts of major land cover changes, such as deforestation and desertification, on the natural system (Lambin & Geist, 2006). However, research of land cover change has now become more integrative, focusing on both the drivers and impacts of land change, including a wider range of interacting processes. Understanding the drivers, state, trends and impacts of different land cover change in combination with the social and natural processes might help to reveal the effects of land system changes on the socio-ecological system and consequently develop effective conservation strategies.

The decrement of natural forest is an important environmental issue in the tropical environment, since these forest harbour exceptional biodiversity, providing important ecosystem services that support the livelihood of local communities (Myers et al., 2000; Sodhi et al., 2006; Page, Rieley & Banks, 2011; Arima, 2016). The study of the past, present, and future state and dynamics of the tropical natural forest cover (NFC) changes might help to better understand the manner of deforestation and fragmentation as well as the influence of social and natural processes. The obtained information will support the development of effective conservation policies and strategies, especially in areas with rapid forest decline (Romijn et al., 2015). The

first comprehensive assessment of global forest resources was reported by [Zon & Sparhawk](#) (1923), the Food and Agriculture Organization (FAO) of the United Nations has carries out regular assessments (Forest Resources Assessments; FRA) at intervals of five or ten years since 1948 ([MacDicken, 2015](#)). Although the FRA data are only partially accurate ([Grainger, 2008](#)), they shed light on NFC changes at a global scale ([MacDicken, 2015](#)). though not at a local scale, mainly because the mechanisms of NFC changes are more complicated than those reported in previous studies ([Lambin, Geist & Lepers, 2003](#)). Remote sensing by satellites has been used to tracking NFC since 1972 ([Grainger, 2008](#)). [Hansen et al. \(2013\)](#) developed a dataset of high-resolution global maps and assessed forest cover change from 2000 to 2012 using satellite data with a spatial resolution of 30 m. These results depict a globally consistent and locally relevant record of forest change. However, little is known about the long-term historical distribution of NFC prior to the 1970s ([Liu & Tian, 2010](#)). Overall, it is essential to assess the historical distribution of NFC at a specific spatial scale ([Miao et al., 2013](#)) in order to explore the underlying mechanisms ([Goldewijk, 2001](#); [Goldewijk et al., 2011](#)) and create more effective conservation policies.

Previous studies used biophysical (e.g. topography and climate) and socio-economic data (e.g. population density, road networks, migration, urban extension, and gross domestic product) to reconstruct the long-term historical distribution of NFC ([Millington, Perry & Romero-  
Calcerrada, 2007](#); [Arima et al., 2008](#); [Miao et al., 2013](#); [He, Li & Zhang, 2014](#)). However, these variables are only valid in regions with abundant historical data. Of the variables, roads have long been considered an important driver of deforestation in many tropical countries, because of

their high availability and strong correlation with NFC ([Arima et al., 2005](#); [Arima et al., 2008](#); [Gaveau et al., 2009](#); [Freitas, Hawbaker & Metzger, 2010](#); [Cai, Wu & Cheng, 2013](#); [Walker et al. 2013](#); [Li et al., 2014](#); [Arima et al., 2016](#); [Hu et al., 2016](#)). Multiple studies have focused on the effect of roads on NFC, but only a few of them have used the temporal and spatial dynamics of road networks for studying the distribution of NFC ([Perz et al., 2007](#); [Perz et al., 2008](#); [Ahmed et al., 2013](#); [Newman, McLaren & Wilson, 2014](#)).

Hainan is the largest island in the Indo-Burma biodiversity hotspot and harbours high levels of biological endemism ([Myers et al., 2000](#)). Natural forest initially covered almost the entire island, but NFC has now decreased to less than one quarter, mainly in the central mountainous area of the island, owing to the intensive exploitation and deforestation since the 1950s ([Lin et al., 2015](#)). Although the NFC changes in Hainan have been previously reported, limited information is available regarding the distribution of NFC over time, which is important to better understand the dynamics of wildlife communities ([Brook et al., 2006](#)), especially of some endangered species (e.g. Hainan Gibbon; [Xu et al., in press](#)). The deforestation of Hainan Island has led to a series of grave ecological consequences (e.g. large floods, landslides, and drought) that threatened the local communities and decreased the overall environmental quality ([Wang et al., 2014](#)). Thus, understanding the spatial and temporal dynamics of NFC is crucial to evaluate the environmental quality and restore forest ecosystems that important services to the local residents.

Here, we used historical road maps and environmental variables to simulate and document the temporal and spatial dynamics of NFC on Hainan from the 1950s to the 2010s. First, we fitted a random forest (RF) model to simulate the distribution of NFC with road-related and

environmental variables in 1975, 1995, and 2012. Next, we used the random forest model to reconstruct the NFC at a 20-years intervals from the 1950s to the 2010s. Based on our data, we aimed to: (1) explore the effect of the road network on NFC changes and (2) identify changes in the distribution of NFC from the 1950s to the 2010s.

## MATERIAL and METHODS

### Study area

This study was carried out on Hainan Island, an area of approximately 33,900 km<sup>2</sup> located in south China (Fig. 1). The island has a central mountainous region with a maximum elevation of 1,867 m (Fig. 1) and is flat in the northern and coastal regions. The climate is tropical with warm and humid weather. The average annual temperature ranges from 16°C in January to 27.5°C in July. The annual rainfall is over 1,600 mm (Francisco-Ortega et al., 2010) and unevenly distributed both between the rainy and dry season and across the island. Rainfall mainly occurs in eastern Hainan Island (Wenchang, Qionghai, Wanning, Lingshui, and Sanya Counties) due to typhoons from the Pacific Ocean. The western part of the island is characterized as dry, while the eastern part is humid (Xu et al., in press).

Prior to 1950s, the forest area on Hainan Island was largely natural, since the number of coconut and rubber plantations was limited island (Chen, 1948), whereas the non-forest area was mainly farmland. The natural forest area decreased considerably owing to logging for timber, rubber plantations, and residential expansion, with NFC reached a minimum in the 1980s. In response to this dramatic NFC loss, the Hainan Provincial Government established a natural

forest logging ban in 1994 (Standing Committee of Hainan Provincial People 's Congress, 1993; Zhang, Uusivuori & Kuuluvainen, 2000).

# **Distribution of natural forest**

In this study, arboreal forest, shrub land, and open woodland were considered as natural forest based on the standards of the National Forest Inventory of China (SFA PRC, 2004). The distribution of natural forest in 1975 was obtained from 93 topographical maps of Hainan Island and digitised using ArcInfo 9.3. (ESRI, 2008; see Appendix 1 for more details). These maps incorporated different types of land uses and outlined the distribution of arboreal forest, shrub land, and open woodland. The distribution of natural forest in 1995 and 2012 was reproduced from maps of the 5<sup>th</sup> and the 8<sup>th</sup> National Forest Resources Inventory of China (<http://www.forestry.gov.cn/gjslzyqc.html>). Three datasets of the historical natural forest distribution were primarily produced by aerial photography and validated by field observations. Field investigation data of 1997/1998 and 2012/2013 (Xu et al., in press) were used to estimate the accuracy of natural forest distribution in 1995 and 2012 (Fig. S1). To enhance data consistency, the NFC distribution in the 1970s, 1990s and 2010s was combined using the ‘Simplify polygon’ function in ArcInfo 9.3. (ESRI, 2008).

We established a 5 × 5 km grid cell system for model simulations. First, Hainan Island was divided into 5 × 5 km grid cells, and those with a land area less than 50% were excluded to minimise the bias of low NFC. A total of 1360 grid cells were used for further analysis. Next, we overlaid the polygons of natural forest in 1975, 1995 and 2012 with the grid cells, and the proportion of natural forest was considered as the NFC of grid cell. The obtained NFC datasets



were used as a response variable in model simulations.

# **Variables correlated with NFC**

We investigated three main categories of variables with a potentially high correlation with NFC: (1) topography, such as slope (SLO, °) and elevation (ELE, m), which indicate difficulty in accessibility and historical exploitation (Teixeira et al., 2009); (2) climatic potential productivity (CPP,  $\text{Kg} \cdot \text{hm}^{-2} \cdot \text{a}^{-1}$ ), which indicates the potential of forest loss due to crop farming (Pongratz et al., 2008); and (3) road network, which indicates the scale and intensity of socio-economic impact on NFC (Arima et al., 2005; Walker et al. 2013; van der Ree, Smith & Grilo, 2015; Arima et al., 2016).

Digital elevation data (approximately  $90 \times 90$  m resolution) were obtained from the Consortium for Spatial Information (<http://www.cgiar-csi.org/data/srtm-90m-digital-elevation-database-v4-1>). The slope map was created based on changes in elevation between adjacent pixels of the digital elevation data. Data on the climatic potential productivity (approximately  $1 \times 1$  km resolution) were obtained from the National Earth System Science Data Sharing Infrastructure (<http://www.geodata.cn/index.html>). The SLO, ELE, and CPP of each grid cell were calculated by averaging all cells ( $1 \times 1$  km or  $90 \times 90$  m) such that their centroid fell within each  $5 \times 5$  km grid cell.

Historical data of the road network were obtained from maps and digital databases published prior to 2013 (Table S2). Maps were digitised using ArcInfo 9.3.(ESRI, 2008) and assigned to a 20-year interval. Due to differences in the mapping scale of traffic maps, the Rural road was designated as the lowest level road, whereas Expressway, Highway, Simply-built

highway, and Cart road were designated as roads with relatively higher levels (Table S2). Based on the function and surface material of various roads, we classified these roads into two classes in the 1950s, 1970s, 1990s and 2010s: main road (Expressway, Highway, and Simply-built highway) and secondary road (Cart road and Rural road).

Four road-related variables: including the sum of road length (SRL,  $\text{km km}^{-2}$ ), the distance from the grid cell centroid to the nearest road (DNR, km), the number of nodes of road network (NON,  $\text{ea km}^{-2}$ ), and the mean node degree (MND,  $\text{ea km}^{-2}$ ; the mean number of roads connected to each node), were calculated in each grid (Table S3) to represent the density and configuration of the road network in the 1950s, 1970s, 1990s, and 2010s.

# **Reconstruction of the historical NFC from the 1950s to the 2010s**

Prior to the reconstruction of the historical NFC, we tested the predictive performance of four modelling techniques, including the generalized linear model (GLM), generalized additive model (GAM), artificial neural networks (ANN) model, and random forests (RF) model. We used the normalised mean square error (NMSE) to compare the predictive performance of the four models in simulating the NFC of each  $5 \times 5$  km grid cell (Appendix 5). NMSE is a relative measure, estimating the overall deviation between predicted and measured values and also a unitless measure in the interval [0,1]. The modelling technique with the lowest value of NMSE has the highest predictive performance. The RF model showed the lowest NMSE among the four models and consequently, was used for reconstructing the historical NFC from the 1950s to the 2010s (Table S4).

The RF model (Iverson et al., 2008) is characterized by insensitivity to multicollinearity,

steadiness across unbalanced datasets, accuracy in predicting the effects of multiple explanatory variables (Breiman, 2001; Culter et al., 2007), higher efficiency in large datasets than that of traditional machine-learning techniques, suitability in demonstrating the nonlinear effect of variables, modelling complex interactions among variables, and robustness to outliers (Li, 2013; Breiman, 2001; Culter et al., 2007). In the present study, we fitted the RF model to associate the NFC data of each grid cell with SLO, ELE, CPP, and main and secondary road networks related indicators (SRL, DNR, NON, and MND) in the 1970s, 1990s, and 2010s. Next, we used the obtained RF model to reconstruct the historical NFC of each grid cell at 20-year intervals from the 1950s to the 2010s, according to the topography, climate, and road network data in a specific period. We assumed that topography and climate did not change significantly over a relatively short time and therefore used the same SLO, ELE, and CPP in all the assessed time periods (1950s-2010s, 1950s-1970s, 1970s-1990s, and 1990s-2010s). The methodological sequence describing the input data, the modelling and validation approaches, and the generation of predictions is shown in the Fig.2.

Overall, we investigated the historical, temporal, and spatial dynamics of NFC on Hainan Island from the 1950s to the 2010s. We first applied simple linear regression to explore the relationship between each road-related indicator and the NFC change rate of each grid cell in the four assessed time periods. We used adjusted  $R^2$  to estimate the explanatory power of each road-related indicator in the simple linear regression model. The variation partition was also applied to compare the effects of main roads vs. secondary roads on NFC change. Simple linear regressions with the combined effect of the four road-related indicators of main road and secondary roads

were used to estimate the variation partition of different road types in the four assessed time periods.

All analyses were conducted in R 3.2.3 ([R Development Core Team, 2016](#)) using the ‘mgcv’ ([Wood, 2011](#)), ‘caret’ ([Kuhn, 2016](#)), ‘randomForest’ ([Liaw & Wiener, 2002](#)), ‘DMwR’ ([Torgo, 2011](#)) packages.

## Hot-spots analysis

To quantify the degree of deforestation, the Getis-Ord  $G_i^*$  statistic was computed to measure the degree of spatial clustering of a local sample and its difference from the expected value using the sum of the differences between values in the local sample and the mean, and is standardized as a Z score ([Scott & Warmerdam, 2005](#)). The Z score reveals the spatial clustering of features with either high or low values. To test the significance of the Z score the value within a specific confidence level were compared. Positive  $G_i^*$  values represent clusters that are greater than the mean (reforestation Hot-spots), whereas negative  $G_i^*$  values represent clusters that are lower than the mean (Deforestation Cold-spots; [Getis & Ord, 1996](#); [Cohen et al. 2011](#)). This statistic represents the frequency of a hotspot relative to the number of input deforested points covered the area. To identify spatial clustering patterns of NFC changes at 20-year intervals, hot-spots analysis was performed using Getis-Ord  $G_i^*$  statistics ([Mitchell, 2005](#)) with the ‘Hot Spot Analysis’ function in ArcInfo 9.3. ([ESRI, 2008](#)).

The Getis-Ord  $G_i^*$  was calculated as follows:

218

$$G_i^* = \frac{\sum_{j=1}^n w_{ij}x_j - \bar{X} \sum_{j=1}^n w_{ij}}{\sqrt{\frac{\left[ n \sum_{j=1}^n w_{ij}^2 - \left( \sum_{j=1}^n w_{ij} \right)^2 \right]}{n-1}}}$$

219

where  $x_j$  is the attribute value for feature  $j$ ,  $w_{ij}$  is the spatial weight between feature  $i$  and  $j$ ,

220

and  $n$  is equal to the total number of features. The  $G_i^*$  statistics is a Z score and thus, no further

221

calculations are required.

222

## RESULTS

223

### The temporal and spatial dynamics of NFC

224

NFC changes at 20-year intervals from the 1950s to the 2010s are shown in [Fig. 3](#). We also

225

calculated the total NFC to explore the temporal dynamics of NFC from the 1950s to the 2010s.

226

The trend of total NFC indicated a considerable decrease from 41.4% in the 1950s to 24.2% in

227

the 2010s ([Fig. 4](#);  $R^2=0.99$ ,  $p=0.004$ ).

228

In the 1950s, most grid cells with NFC >50% were identified in the central mountainous

229

regions, whereas those with NFC <20% were identified in the coastal regions ([Fig. 3](#)). The grid

230

cells with  $NFC \leq 20\%$  showed a significant increase from 433 to 839 from the 1950s to the

231

2010s ([Fig. 3](#)), whereas their distribution gradually shifted from the coastal regions to the

232

mountainous regions ([Fig. 3](#)). The grid cells with  $NFC > 50\%$  mainly distributed in the central

233

and southern mountainous region, but showed a significant decrease from 544 to 286 from the

234

1950s to the 2010s ([Fig. 3](#)). These results revealed that the natural forest decrease and also

235

fragmentized from the 1950s to the 2010s.

We used the slope of site-specific relationships between NFC and its change rate over time to identify the dynamic change patterns of NFC. The slopes were positively correlated from the 1950s to the 2010s., indicating that a lower NFC tended to correspond to a higher rate of NFC decrease (Fig. 5;  $F=122.3$ ,  $p<2.2\times 10^{-16}$ ). Assessing the pattern of NFC change rate revealed that most grid cells with low NFC change rate were distributed in the mountainous areas (Jiangfengling, Yinggeling, and Wuzhishan mountains), whereas most grid cells with high NFC change rate were distributed in the Qiongbai Platform (Fig. 5). Hot-spots analysis from the 1950s to the 2010s showed that NFC decrease mainly occurred in the platform area, whereas NFC increase occurred in the mountainous region (Fig. 6). However, NFC also increased in Wenchang County, although this area is flat and well developed. When assessed at 20-year intervals, the distribution of hot-spots with decreased NFC was observed to be shifted from the eastern regions to the western regions of Hainan Island (Fig. 6).

## **Dynamics of road expansion**

The density and configuration of the main road network markedly changed with the time from the 1950s to the 2010s. The mean SRL of main roads increased from 0.11 km km<sup>-2</sup> in the 1950s to 0.45 km km<sup>-2</sup> in the 2010s (Table 1). The standard deviation of SRL of main roads also increased, indicating that the distribution of main roads became more heterogeneous across the grid cells. The mean DNR decreased from 0.30 km in the 1950s to 0.04 km in the 2010s (Table 1). The standard deviation of DNR also decreased, indicating that the spatial distribution of main roads markedly increased. The average value of NON and MND of main roads increased from the 1950s to the 2010s (Table 1), revealing that the configuration of the road network became

more complicated across the grid cells. The standard deviation of NON of main roads increased from 0.04 ea km<sup>-2</sup> to 0.23 ea km<sup>-2</sup> from the 1950s to the 2010s, whereas the standard deviation of MND of main roads had slightly decreased from 0.05 ea km<sup>-2</sup> to 0.04 ea km<sup>-2</sup>, revealing that the configuration of the main road network became more complex and the spatial pattern of main road was uneven across the grid cells.

The SRL of secondary roads increased from 0.18 km km<sup>-2</sup> in the 1950s to 0.36 km km<sup>-2</sup> in the 1970s, but remained stable after the 1970s. The NON and MND of secondary roads also showed the same trend both clearly increase in the 1950s to 1970s, but remained stable after the 1970s (Table 1). The DNR of secondary roads had decreased from 0.11 km in the 1950s to 0.05 km in the 1970s, whereas only slightly changed after the 1970s. The standard deviation of the four road-related indicators showed that the secondary road network markedly changed in for the 1950s to 1970s, but remained stable after the 1970s (Table 1). The density and configuration of the secondary road network markedly changed from the 1950s to 1970s, whereas it changed only slightly after the 1970s (Table 1).

Both the main road and secondary road network obviously developed from the 1950s to the 2010s: however, the construction processes were faster in the former than the latter. Overall, the density and configuration of the road network was higher and more complex after nearly 60 years of construction.

### **Effects of different road expansion on NFC change**

The change of SRL of main roads explained 11.20% ( $p < 0.001$ ) of the variance in NFC change from the 1950s to the 2010s and was also the strongest indicator in the other three

assessed time periods (1950s-1970s, Adjusted  $R^2 = 14.22\%$ ,  $p < 0.001$ ; 1970s-1990s, Adjusted  $R^2 = 20.73\%$ ,  $p < 0.001$ ; and 1990s-2010s, Adjusted  $R^2 = 4.89\%$ ,  $p < 0.001$ ; [Table 2](#)). In the 1950s-2010s, the overall variation partition of main roads (Adjusted  $R^2 = 23.40\%$ ;  $p < 0.001$ ) in the simple linear regression was much higher than the secondary roads (Adjusted  $R^2 = 13.30\%$ ;  $p < 0.001$ ). The change of main roads also explained a higher degree of the variance than the change of secondary roads in the other three assessed time periods. These results indicated that the main road network had a more significant effect on deforestation than the secondary road network from the 1950s to the 2010s.

In the 1990s-2010s, the adjusted  $R^2$  of the main road network decreased from 25.2% to 5.10% ([Table 2](#)), whereas that of the secondary road network decreased from 8.99% to 0.12% ([Table 2](#)), showing that the association between the road network and deforestation weakened with time. However, the main road network still showed linkage stronger association with NFC change than the secondary road network (5.10% vs. 0.12%).

## DISCUSSION

### NFC reduction and fragmentation

Our results demonstrated that the total NFC of Hainan Island decreased significantly from the 1950s to the 2010s ([Fig. 4](#)), and the distribution of grid cells with  $NFC < 20\%$  increased in the mountainous regions and thus became more vulnerable to deforestation ([Fig. 5](#)). These patterns are consistent with those reported in other tropical regions, in which natural forest deforestation is always accompanied by fragmentation ([Pimm, 1998](#); [Laurance et al., 1998](#);



DeFries et al., 2005). Natural forest that is severely fragmented shows small patches of natural forest that are highly vulnerable to clearing (Stickler et al., 2013; Taylor, 2013). We also observed that the grid cells with NFC > 50% were located in areas with a high altitude and steep slope (Fig. 3), in which mechanisation and accessibility are limited (Freitas, Hawbaker & Metzger, 2010; Hu et al., 2016). Most nature reserves on Hainan Island were similarly distributed in the mountainous regions with high altitude and steep slope (Lin et al., 2015). Although some grid cells located in some mountainous regions showed a moderately high degree of recovery (Fig. 5), most of them with NFC > 50% was shrank and fragmented.

Natural forest fragmentation is a major issue in tropical environments, since the spatial arrangement and geometric configuration of fragments can impair ecological processes. Six general fragmented patterns have been recognised at the global scale in previous work (Geist & Lambin, 2002; Mertens & Lambin, 1997), of which five are observed in the Amazon tropical forest—rectangular, fishbone, radial, dendritic, and ‘the stem of the rose’ (Arima et al., 2005). Road network architecture plays a critical role in shaping forest fragmentation patterns in the Amazon (Arima et al., 2005; Soares-Filho et al. 2006; Walker et al. 2013). We also revealed that the road network expansion explained more than 30% of natural forest deforestation and fragmentation on Hainan Island from the 1950s to the 2010s (Table 2; Fig. 5). When the road network was overlaid by grid cells with high NFC (Appendix 6; Fig. S2), we found that the dominant fragmented pattern of Hainan Island natural forest was radial and dendritic. Unlike the Amazon, road construction combined the complex topography determined the pattern of natural forest landscape on Hainan Island (Fig. 1). Thus, forest fragmentation patterns on the Hainan

Island were highly affected by road construction and topography.

Our analysis showed that hot-spots with decreased NFC mainly occurred in the Qiongbei Platform (Fig. 5), following a dynamic spatial distribution pattern from east to west, whereas hot-spots with increase NFC occurred in Wenchang County (Fig. 5), an area that is flat and well developed with a frequent typhoon occurrence during the summer. The local residents preserved some small patches of natural forest around the villages and buildings, known as “geomantic forest”, as a protecting shield against the severe weather events.

### **Effect of road network on natural forest deforestation**

Road-related indicators are not the actual drivers of NFC decrease, but they play a role in natural forest dynamics through land-use changes and deforestation (Freitas, Hawbaker & Metzger, 2010; Newman, McLaren & Wilson, 2014). The development of roads and road networks is strongly correlated with the economic growth and is associated with the ecological disturbance and natural forest degradation (Wilkie et al., 2000; Laurance, Goosem & Laurance, 2009). In tropical forests, roads have a low direct effect on habitat loss, but a high indirect effect on the spatial patterns of deforestation (Fearnside, 2008), since they facilitate accessibility, resource extraction, and human activities (Selva et al., 2011; Hu et al., 2016). Our results were consistent with those of previous studies that also reported the significant effect of road networks on natural forest deforestation in tropical regions (Perz et al., 2007; Gaveau et al., 2009; Freitas, Hawbaker & Metzger, 2010; Cai, Wu & Cheng, 2013; Li et al., 2014; Arima et al., 2016; Hu et al., 2016). We demonstrated that road-related indicators played a substantial role in determining the deforestation of NFC in Hainan. The SRL of main roads was one of the strongest variables

associated with NFC change dynamic in the four assessed time periods([Table 2](#)). However, the power of road network in explaining the NFC reduction weakened over time, in both main and secondary roads, probably revealing the underlying socio-economic processes of road expansion itself ([Perz et al., 2007](#)).

The influence of road networks on deforestation is associated to population migration and various socio-economic events. Large populations moved to Hainan Island, since it is area rich in natural resources ([Yan, 2008](#)). After the foundation of People’s Republic of China in 1949, the economy became a key element in the national strategic plan. Both agriculture and forest farming developed rapidly, and large agricultural populations migrated to Hainan Island under the conduct of the national government. In the 1960s and 1970s, the ‘sent-down youth’ movement influenced the social activity and increased migration to Hainan. In the 1950s-1970s, roads were constructed for accessing land and timber resources and thus, the main road and secondary road network was markedly associated to natural forest deforestation ([Table 2](#)). In 1988, Hainan Island was established as a provincial agency and Economic Development Zone, leading to another population migration event, mainly of urban population. In the 1970s-1990s, roads were built for accessing land resources and linking major cities and towns; thus, the main road network had a stronger influence on deforestation than the secondary road network ([Table 2](#)). In the early 1990s, the industrial structure of Hainan Island was transformed and upgraded. The tertiary industry, especially tourism, became the core of economic development, changing the functions of road construction. Additionally, the Hainan Provincial Government established a natural forest logging ban in 1994 in response to the sharp decline in NFC ([Standing Committee](#)

of Hainan Provincial People 's Congress, 1993). Consequently, the association between the road network and deforestation weakened in 1990s-2010s (Table 2).

Deforestation was more affected by the main road network than the secondary road network in the four assessed time periods (Table 2). Population was the most important factor affecting NFC on Hainan (Zhang, Uusivuori & Kuuluvainen, 2000; Lin & Zhang, 2001). The main road network of Hainan Island includes expressways, highways, and simply-built highways built or funded by the national or provincial governments for geopolitical purposes, especially for connecting major cities. The secondary road network includes cart roads and rural roads built or funded by the city and county governments for supporting local livelihoods, connecting communities, and accessing land and other natural resources. The construction of main roads improves the connectivity of major cities, supports urbanisation (Perz et al., 2008), and thus, highly increases deforestation. The construction of secondary roads also promotes deforestation, but in a smaller scale than that of main roads (Fig. S2).

## Implications for natural forest conservation and management

Our study provided significant information for natural forest conservation and management. At a global scale, ecosystems decline and become fragmented (Saunders, Hobbs & Margules, 1991; Fischer & Lindenmayer, 2007), and thus, the small natural forest patches represent large elements of tropical natural forest (Tulloch et al., 2016). Theoretical and experimental studies have highlighted the importance of conserving large contiguous natural forest patches for maintaining biodiversity (Bender, Contreras & Fahrig, 1998; Mortelliti et al., 2014). Consequently, conservation actions have mainly focused on preserving large contiguous forest

383 patches and not relatively small fragmented forest patches (Ovaskainen, 2002; Tulloch et al.,  
 384 2016). Small fragmented forest patches contribute to short-term and long-term indigenous  
 385 species persistence; enhance the biodiversity in human-dominated fragmented landscapes  
 386 (Turner & Corlett, 1996); positively affect the diversity of forest bird and mammal species, forest  
 387 plant species, and migratory animals as well as the persistence of meta-populations (Laurance,  
 388 1994; Warkentin, Greenberg & Salgado Ortiz, 1995; Jacquemyn, Jan Butaye & Hermy, 2001);  
 389 act as refuges for plant and animal species; and help conservationists to launch a final attempt to  
 390 rescue endangered species that may serve as sources of natural forest reconstruction (Turner &  
 391 Corlett, 1996). In Hainan, small natural forest patches are formed from the fragmentation of  
 392 larger contiguous natural forest patches. Those small natural forest patches are isolated and  
 393 always have an area less than 10 ha in Hainan. Those small natural forest patches are always  
 394 located at the edge of large contiguous natural forest in the mountainous region or around the  
 395 villages in the flat region, and provide important ecosystem services for local residents. For  
 396 example, the “Geomantic forest” in Wenchang and Qionghai counties is an example of small  
 397 natural forest patches- with typical tropical forest structure, but high plant diversity (Yang & Wu,  
 398 2002), distributed in the Lingnan region that includes the modern Chinese provinces of Jiangxi,  
 399 Hunan, Guangdong, Guangxi, and Hainan. The Geomantic forest is related to the Fengshui  
 400 Theory, which includes historical and cultural features of ecological significance, and provides  
 401 important ecosystem services such as water and soil conservation as well as protection against  
 402 severe weather events (Cheng, He & Liu, 2009). Due to this forest, the NFC of Wenchang  
 403 county increased from the 1950s to the 2010s (Fig. 5). Thus, the protection of small natural

forest patches is important, and long-term data need to be used for developing effective management with specific conservation objectives. Regions with low NFC tend to be more vulnerable in natural forest loss (Fig. 5). The natural forest loss and fragmentation patterns are dynamic on Hainan (Fig. 6). Our results showed that the spatial and temporal change patterns of NFC could be accurately assessed using historical data (Newman, McLaren & Wilson, 2014). Thus, conservationists and decision-makers need to evaluate natural forest changes and develop effective management actions (Margules & Pressey, 2000). On Hainan Island, regions with low NFC, but high ecological value (Yangshan region in Haikou city; the Geomantic forest in Wenchang city; Fig. 3) located in flat areas that are well developed and have high population density, are more vulnerable than natural forest within the nature reserves. Thus, conservation frameworks and forest restoration projects are needed to preserve or recover the natural forest.

The NFC in the mountainous region increased in the past 60 years; however, regions with high NFC are vulnerable to fragmentation due to the expansion of the road network in some remote regions in the mountains (Fig. S2). The south-central mountainous region of Hainan Island is a biodiversity hotspot (Myers et al., 2000) and thus, a priority area for conservation in China (Ministry of Environmental Protection of the People 's Republic of China, 2011). Monitoring the natural forest in using remote sensing and drones could help in protecting the natural forest and predict any future changes (Paneque-Gálvez et al., 2014).

## CONCLUSIONS

This study used historical data to reconstruct NFC maps in order to better understand the

spatial and temporal change patterns. The road network was identified as an important factor of forest loss and fragmentation of tropical forests. Although the exact distribution of natural forest was not depicted, we found that the dominant fragmented pattern of Hainan Island natural forest is radial and dendritic.

Our study showed that: (1) low NFC corresponds to a high rate of NFC decrease as well as highly dynamic spatial and temporal change patterns of NFC; (2) the road network significantly affected NFC, whereas topography affected deforestation by influencing the road network construction pattern; and (3) the effect of road network on NFC was related to population migration and socio-economic events, and the main road network had a stronger relation to deforestation than the secondary road network.

Deforestation and fragmentation still occur in Hainan, and public concern remains pronounced in recent years. Thus, it is crucial to better understand implications of deforestation on biodiversity conservation, as well as the driving forces of deforestation. Conservation efforts must focus on small natural forest patches using new technologies (e.g. remote sensing and drones) that will improve monitoring and data collection.

Although our model was incomplete, since it only included topography, climate and road network data, it revealed the spatial and temporal change patterns of NFC on Hainan Island. Future modelling studies need to build an improved model, combining historical, natural, and socio-economic factors that will help to better understand the underlying mechanisms of NFC change patterns.

# **Acknowledgements**

We thank the Forestry Department of Hainan Province, the Administration Bureau of Animal and Plant Protection of Hainan Province, and the staff of Nature Reserves for the provided data and technical support. We are also grateful to the anonymous referees for their valuable comments that helped to improve our manuscript.



# References

- Ahmed SE, Souza CM, Riberio J, Ewers RM. 2013. Temporal patterns of road network development in the Brazilian Amazon. *Regional Environmental Change* 13: 927-937. DOI: 10.1007/s10113-012-0397-z.
- Arima EY, Walker RT, Perz SG, Caldas M. 2005. Loggers and forest fragmentation: behavioral models of road building in the amazon basin. *Annals of the Association of American Geographers* 95:525-541. DOI: 10.1111/j.1467-8306.2005.00473.x.
- Arima EY, Walker RT, Sales M, Souza Jr C, Perz SG. 2008. The fragmentation of space in the amazon basin: emergent road networks. *Photogrammetric Engineering and Remote Sensing* 74:699-709. DOI: 10.14358/PERS.74.6.699.
- Arima EY, Walker RT, Perz S, Jr. CS. 2016. Explaining the fragmentation in the Brazilian Amazonian forest. *Journal of Land Use Science* 11:257-277. DOI: 10.1080/1747423X.2015.1027797.
- Arima EY. 2016. A spatial probit econometric model of land change: the case of infrastructure development in western Amazonia, Peru. *PloS One* 11: e0152058. DOI: 10.1371/journal.pone.0152058.
- Bender DJ, Contreras TA, Fahrig L. 1998. Habitat loss and population decline: a meta-analysis of the patch size effect. *Ecology* 79: 517-533. DOI: 10.1890/0012-9658(1998)079[0517:HLAPDA]2.0.CO;2.
- Breiman L. 2001. Random forest. *Machine Learning* 45: 5-32. DOI: 10.1023/A:1010933404324.
- Brook BWC, Bradshaw CJA, Koh LP, Sodhi NS. 2006. Momentum drives the crash: mass

- extinction in the tropics. *Biotropica* 38: 302-305. DOI: 10.1111/j.1744-7429.2006.00141.x.
- Cai X, Wu Z, Cheng J. 2013. Using kernel density estimation to assess the spatial pattern of road density and its impact on landscape fragmentation. *International Journal Geographical Information Science* 27: 222-230. DOI: 10.1080/13658816.2012.663918.
- Chen Z. 1948. Development of Hainan Island resources. Nanjing: Zhengzhong Press. (In Chinese)
- Cheng J, He F, Liu Y. 2009. Progress of the research on Geomantic forest of Lingnan villages. *Chinese landscape architecture* 25: 93-96. (in Chinese with English abstract)
- Cohen Y, Sharon R, Sokolsky T, Zahavi T. 2011. Modified hot-spot analysis for spatio-temporal analysis: a case study of the leaf-roll virus expansion in vineyards. *Spatial 2-Spatial Data Methods for Environmental and Ecological Processes, At Foggia, Italy*. Available at: <https://aisberg.unibg.it/retrieve/handle/10446/25273/4012/40.pdf>. (Accessed 25 February 2017)
- Culter DR, Edwards TCJ, Beard KH, Cutler A, Hess KT. 2007. Random forest for classification in ecology. *Ecology* 88: 2783-2792. DOI: 10.1890/07-0539.1.
- Defries R, Hansen A, Newton AC, Hansen MC. 2005. Increasing isolation of protected areas in tropical forests over the past twenty years. *Ecological Applications* 15, 19-26. DOI: 10.1890/03-5258.
- Dirzo R, Young HS, Galetti M, Ceballos G, Isaac NJ, Collen B. 2014. Defaunation in the Anthropocene. *Science* 345: 401-406. DOI: 10.1126/science.1251817.
- Ellis EC, Kaplan JO, Fuller DQ, Vavrus S, Klein GK, Verburg, PH. 2013. Used planet: a global

- history. *Proceedings of the National Academy of Sciences of the United States of America*
- 110: 7978-7985. DOI:10.1073/pnas.1217241110.
- ESRI, 2008. ArcGIS 9.3. Environmental Systems Research Institute Inc.: Redlands, CA.
- Fearnside PM. 2008. The roles and movements of actors in the deforestation of Brazilian
- Amazonia. *Ecology and Society* 13, 23. Available at:
- <http://www.ecologyandsociety.org/vol13/iss1/art23/>.
- Fischer J, Lindenmayer DB. 2007. Landscape modification and habitat fragmentation: a
- synthesis. *Global Ecology and Biogeography* 16: 265-280. DOI: 10.1111/j.1466-
- 8238.2007.00287.x.
- Francisco-Ortega J, Wang ZS, Wang FG, Xing FW, Liu H, Xu H, Xu WX, Luo YB, Song XQ,
- Gale S, Boufford DE, Maunder M, An SQ. 2010. Seed plant endemism on Hainan Island: a
- framework for conservation actions. *The Botanical Review* 76: 346-376. DOI:
- 10.1007/s12229-010-9055-7.
- Freitas SR, Hawbaker TJ, Metzger JP. 2010. Effects of roads, topography, and land use on forest
- cover dynamics in the Brazilian Atlantic Forest. *Forest Ecology and Management* 259: 410-
417. DOI: 10.1016/j.foreco.2009.10.036.
- Gaveau DL, Wich S, Epting J, Juhn D, Kanninen M, Leader-Williams N. 2009. The future of
- forests and orangutans (*Pongo abelii*) in Sumatra: predicting impacts of oil palm plantations,
- road construction, and mechanisms for reducing carbon emissions from deforestation.
- Environmental Research Letters* 4: 034013. DOI: 10.1088/1748-9326/4/3/034013.
- Geist HJ, Lambin EF. 2002. Proximate causes and underlying driving forces of tropical

deforestation. *Bioscience* 52: 143-450. DOI:10.1641/0006-3568(2002)052[0143:PCAUDF]  
2.0.CO;2

Getis A, Ord JK. 1996. Local spatial statistics: an overview. In Longley P, Batty M, ed. *Spatial Analysis: Modeling in a GIS environment*. Cambridge: GeoInformation International, 269-294.

Goldewijk KK, Beusen A, Van Dreht G, De Vos M. 2011. The HYDE 3.1 spatially explicit database of human-induced global land-use change over the past 12,000 years. *Global Ecology and Biogeography* 20: 73-86. DOI: 10.1111/j.1466-8238.2010.00587.x.

Goldewijk KK. 2001. Estimating global land use change over the past 300 years. *Global Biogeochemical Cycles* 15: 417-433. DOI: 10.1029/1999GB001232.

Grainger A. 2008. Difficulties in tracking the long-term global trend in tropical forest area. *Proceedings of the National Academy of Sciences of the United States of America* 105: 818-23. DOI: 10.1073/pnas.0703015105.

Hansen MC, Potapov PV, Moore R, Hancher M, Turubanova SA, Tyukavina A, Thau D, Stehman SV, Goetz SJ, Loveland TR, Kommareddy A, Egorov A, Chini L, Justice CO, Townshend JRG. 2013. High-Resolution Global Maps of 21st-Century Forest Cover Change. *Science* 342: 850-853. DOI: 10.1126/science.1244693.

He F, Li S, Zhang X. 2014. Spatially explicit reconstruction of forest cover of Southwest China in the Qing Dynasty. *Geographical Research* 33: 260-269 (in Chinese with English abstract). DOI: 10.11821/dlyj201402006.

Hu X, Wu Z, Wu C, Ye L, Lan C, Tang K, Xu L, Qiu R. 2016. Effects of road network on

diversiform forest cover changes in the highest coverage region in China: An analysis of sampling strategies. *Science of The Total Environment* 565: 28-39. DOI: 10.1016/j.scitotenv.2016.04.009.

Iverson LR, Prasad AM, Matthews SN, Peters M. 2008. Estimating potential habitat for 134 eastern us tree species under six climate scenarios. *Forest Ecology and Management* 254: 390-406. DOI: 10.1016/j.foreco.2007.07.023.

Jacquemyn H, Jan Butaye J, Hermy M. 2001. Forest plant species richness in small, fragmented mixed deciduous forest patches: the role of area, time and dispersal limitation. *Journal of Biogeography* 28: 801-812. DOI: 10.1046/j.1365-2699.2001.00590.x.

Kuhn M. 2016. caret: Classification and Regression Training. R package version 6.0-73. Available at: <https://CRAN.R-project.org/package=caret>. (accessed 6 March 2017)

Lambin EF, Geist HJ. 2006. Land-Use and Land-Cover Change. Berlin: Springer.

Lambin EF, Geist HJ, Lepers E. 2003. Dynamics of land-used and land-cover change in tropical regions. *Annual Review of Environment and Resources* 28: 205-241. DOI: 10.1146/annurev.energy.28.050302.105459.

Laurance WF, Ferreira LV, Rankin-de Merona JM, Laurance SG. 1998. Rain forest fragmentation and the dynamics of Amazonian tree communities. *Ecology* 79: 2032-2040. DOI: 10.1890/0012-9658(1998)079[2032:RFFATD]2.0.CO;2.

Laurance WF. 1994. Rainforest fragmentation and the structure of small mammal communities in tropical Queensland. *Biological Conservation* 69: 23-32. DOI: 10.1016/0006-3207(94)90325-5.

555 Laurance WF, Goosem M, Laurance SGW. 2009. Impacts of roads and linear clearings on  
556 tropical forests. *Trends in Ecology and Evolution* 24: 659-669. DOI:  
557 10.1016/j.tree.2009.06.009.

558 Lewis SL, Maslin MA. 2015. Defining the Anthropocene. *Nature* 519: 171-80. DOI:  
559 10.1038/nature14258.

560 Li M, De Pinto A, Ulimwengu JM, You L, Robertson RD. 2014. Impacts of Road Expansion on  
561 Deforestation and Biological Carbon Loss in the Democratic Republic of Congo.  
562 *Environmental and Resource Economics* 60: 433-469. DOI: 10.1007/s10640-014-9775-y.

563 Li X. 2013. Using “random forest” for classification and regression. *Chinese journal of applied*  
564 *entomology* 50:1190-1197. DOI: 10.7679/j.issn.2095-1353.2013.163. (in Chinese with  
565 English abstract)

566 Liaw A, Wiener M. 2002. Classification and Regression by random Forest. *R News* 2: 18-22.  
567 Available online: <https://cran.r-project.org/doc/Rnews/> (accessed on 3 Jan. 2016)

568 Lin M, Zhang Y. 2001. Dynamic change of tropical forest in Hainan Island. *Geographical*  
569 *research* 20: 703-712. DOI: 10.11821/yj2001060008. (in Chinese with English abstract)

570 Lin S, Xu Y, He J, Ma G, Jiang H. 2015. Study on Forest Ecosystem. In: Xing Q, Shi J, Lin Z, ed.  
571 Hainan provincial biodiversity conservation strategy and action plan (2014–2030). Beijing:  
572 Sciences Press (in Chinese).

573 Liu M, Tian H. 2010. China’s land cover and land use change from 1700 to 2005: Estimations  
574 from high resolution satellite data and historical archives. *Global Biogeochemical Cycles* 24:  
575 GB3003. DOI: 10.1029/2009GB003687.

576 MacDicken KG. 2015. Global Forest Resources Assessment 2015: What, why and how? *Forest*  
577 *Ecology and Management* 352: 3-8. DOI: 10.1016/j.foreco.2015.02.006.

578 Margules CR, Pressey RL. 2000. Systematic conservation planning. *Nature* 405: 243-253. DOI:  
579 10.1038/35012251.

580 Mertens B, Lambin EF. 1997. Spatial modelling of deforestation in southern Cameroon: Spatial  
581 disaggregation of diverse deforestation processes. *Applied Geography* 17: 143-162.  
582 DOI:10.1016/S0143-6228(97)00032-5.

583 Miao L, Zhu F, He B, Ferrat M, Liu Q, Cao X, Cui X. 2013. Synthesis of China's land use in the  
584 past 300 years. *Global and Planetary Change* 100: 224-233. DOI:  
585 10.1016/j.gloplacha.2012.10.021.

586 Millington JDA, Perry GLW, Romero-Calcerrada R. 2007. Regression techniques for examining  
587 land use/cover change: a case study of a Mediterranean landscape. *Ecosystems* 10: 562-578.  
588 DOI: 10.1007/s10021-007-9020-4.

589 Ministry of Environmental Protection of the People 's Republic of China. 2013. China national  
590 biodiversity conservation strategy and action plan (2011-2030). Beijing: China  
591 Environmental Science Press. (in Chinese with English version)

592 Mitchell A. 2005. The ESRI Guide to GIS Analysis, Volume 2: Spatial Measurements and  
593 Statistics. Hongkong: Esri Press.

594 Mortelliti A, Sozio G, Driscoll DA, Bani L, Boitani L, Lindenmayer DB. 2014. Population and  
595 individual-scale responses to patch size, isolation and quality in the hazel dormouse.  
596 *Ecosphere* 5: 1-21. DOI: 10.1890/ES14-00115.1.

597 Myers N, Mittermeier RA, Mittermeier C., da Fonseca GAB, Kent J, 2000. Biodiversity hotspots  
598 for conservation priorities. *Nature* 403: 853-858. DOI: 10.1038/35002501.

599 Newman ME, McLaren KP, Wilson BS. 2014. Assessing deforestation and fragmentation in a  
600 tropical moist forest over 68 years; the impact of roads and legal protection in the Cockpit  
601 Country, Jamaica. *Forest Ecology and Management* 315: 138-152. DOI:  
602 10.1016/j.foreco.2013.12.033.

603 Ovaskainen O. 2002. Long-term persistence of species and the SLOSS problem. *Journal of*  
604 *Theoretical Biology* 218: 419-433. DOI: 10.1006/jtbi.2002.3089.

605 Page SE, Rieley JO, Banks CJ. 2011. Global and regional importance of the tropical peatland  
606 carbon pool. *Global Change Biology* 17: 798-818. DOI: 10.1111/j.1365-2486.2010.02279.x.

607 Paneque-Gálvez J, McCall MK, Napoletano BM, Wich SA, Koh LP. 2014 Small drones for  
608 community-based forest monitoring: an assessment of their feasibility and potential in  
609 tropical areas. *Forests* 5: 1481-1507. DOI: 10.3390/f5061481.

610 Perz SG, Overdevest C, Caldas MM, Walker RT, Arima EY. 2007. Unofficial road building in  
611 the brazilian amazon: dilemmas and models for road governance. *Environmental*  
612 *Conservation* 34: 112-121. DOI: 10.1017/S0376892907003827.

613 Perz SG, Walker R, Caldas M, Arima E, Souza C. 2008. Road networks and forest fragmentation  
614 in the Amazon: explanations for local differences with implications for conservation and  
615 development. *Journal of Latin American Geography* 7: 85-104. DOI: 10.1353/lag.0.0004.

616 Pimm SL.1998. Ecology-the forest fragment classic. *Nature* 393: 23-24. DOI: 10.1038/29892.

617 Pongratz J, Reick C, Raddatz T, Claussen M. 2008. A reconstruction of global agricultural areas



and land cover for the last millennium. *Global Biogeochemical Cycles* 22: GB3018. DOI: 10.1029/2007GB003153.

R Development Core Team. 2016. R: a language and environment for statistical computing. Version 3.2.3. Vienna: R Foundation for Statistical Computing. Available at: <http://www.R-project.org/>.

Romijn E, Lantican CB, Herold M, Lindquist E, Ochieng R, Wijaya A, Murdiyarso D, Verchot L. 2015. Assessing change in national forest monitoring capacities of 99 tropical countries. *Forest Ecology and Management* 352: 109-123. DOI: 10.1016/j.foreco.2015.06.003.

Saunders DA, Hobbs RJ, Margules CR. 1991. Biological consequences of ecosystem fragmentation: a review. *Conservation Biology* 5: 18-32. DOI: 10.1111/j.1523-1739.1991.tb00384.x.

Scott L, Warmerdam N. 2005. Extend Crime Analysis with ArcGIS Spatial Statistics Tools. Available at: [http://www.esri.com/news/arcuser/0405/ss\\_crimestats1of2.html](http://www.esri.com/news/arcuser/0405/ss_crimestats1of2.html). (accessed 6 March 2017)

Selva N, Kreft S, Kati V, Schluck M, Jonsson BG, Mihok B, Okarma H, Ibisch PL. 2011. Roadless and low-traffic areas as conservation targets in Europe. *Environmental Management* 48: 865-877. DOI: 10.1007/s00267-011-9751-z.

State Forestry Administration of the People's Republic of China (SFA PRC). 2004. The Technical Regulations of National Forest Inventory. Available at <http://211.167.243.162:8085/8/workdyfront/fileSingle?lm=dcgz&mfile.GUID=309CE283D9D240AB915B023020A69C0B> (accessed 5 October 2016).

- Soares-Filho BS, Nepstad DC, Curran LM, Cerqueira GC, Garcia RA, Ramos CA, Voll E, McDonald A, Lefebvre P, Schlesinger P. 2006. Modelling conservation in the amazon basin. *Nature* 440: 520-523. DOI: 10.1038/nature04389.
- Sodhi NS, Koh LP, Brook BW, Ng PK. 2006. Southeast Asian biodiversity: an impending disaster. *Trends in Ecology and Evolution* 19: 654-660. DOI: 10.1016/j.tree.2004.09.006.
- Standing Committee of Hainan Provincial People 's Congress. 1993. Statute on forest protection and management in Hainan Province. Available at: <http://www.hizw.gov.cn/data/news/2015/05/43511/>. (accessed 20 February 2017)
- Stickler CM, Nepstad DC, Azevedo AA, McGrath DG. 2013. Defending public interests in private lands: compliance, costs and potential environmental consequences of the Brazilian Forest Code in Mato Grosso. *Philosophical Transactions of the Royal Society of London Series B-Biological Science*. 368: 20120160. DOI: 10.1098/rstb.2012.0160.
- Taylor MFJ. 2013. Bushland at Risk of Renewed Clearing in Queensland. Sydney: World Wildlife Fund Australia. Available at [http://awsassets.wwf.org.au/downloads/fl012\\_bushland\\_at\\_risk\\_of\\_renewed\\_clearing\\_in\\_queensland\\_9may13.pdf](http://awsassets.wwf.org.au/downloads/fl012_bushland_at_risk_of_renewed_clearing_in_queensland_9may13.pdf) (accessed 5 October 2016).
- Teixeira AMG, Soares-Filho BS, Freitas SR, Metzger JP. 2009. Modeling landscape dynamics in an Atlantic Rainforest region: implications for conservation. *Forest Ecology and Management* 257: 1219-1230. DOI: 10.1016/j.foreco.2008.10.011.
- Torgo L. 2011. Data Mining with R: learning with case studies. Boca Raton: CRC Press. Available at: <http://www.dcc.fc.up.pt/~ltorgo/DataMiningWithR>. (accessed 6 March 2017)

- 660 Tulloch AI., Barne, MD, Ringma J, Fuller RA, Watson JEM. 2016. Understanding the  
661 importance of small patches of habitat for conservation. *Journal of Applied Ecology* 53:  
662 418-429. DOI: 10.1111/1365-2664.12547.
- 663 Turner IM, Corlett RT. 1996. The conservation value of small, isolated fragments of lowland  
664 tropical rain forest. *Trends in Ecology and Evolution* 11: 330-333. DOI: 10.1016/0169-  
665 5347(96)10046-X.
- 666 van der Ree R, Smith DJ, Grilo C. 2015. Handbook of Road Ecology. New Jersey: Wiley  
667 Blackwell.
- 668 Walker R, Arima E, Messina J, Soaresfilho B, Perz S, Vergara D, Sales M, Pereira R, Castro W.  
669 2013. Modeling spatial decisions with graph theory: logging roads and forest fragmentation  
670 in the Brazilian Amazon. *Ecological Applications* 23: 239-254. DOI: 10.1890/11-1800.1.
- 671 Wang X, Li G, Hu W, Wu X, Zhang Y. 2014. Preliminary analysis of inductive factors and the  
672 influence mechanism of Diaoluo Mountain landslide. *Journal of Natural Disasters* 1: 144–  
673 148 (in Chinese with English abstract). DOI: 10.13577/j.jnd.2014.0120.
- 674 Warkentin IG, Greenberg R, Salgado Ortiz J. 1995. Songbird use of gallery woodlands in  
675 recently cleared and older settled landscapes of the Selva Lacandona, Chiapas, Mexico.  
676 *Conservation Biology* 9: 1095-1106. DOI: 10.1046/j.1523-1739.1995.951095.x.
- 677 Wilkie D, Shaw E, Rotberg F, Morelli G, Auzel P. 2000. Roads, development, and conservation  
678 in the Congo basin. *Conservation Biology* 14: 1614-1622. DOI: 10.1111/j.1523-  
679 1739.2000.99102.x.
- 680 Wood SN. 2011. Fast stable restricted maximum likelihood and marginal likelihood estimation

681 of semiparametric generalized linear models. *Journal of the Royal Statistical Society*  
682 73:3-36. DOI: 10.1111/j.1467-9868.2010.00749.x.

683 Xu Y, Lin S, He J, Xin Y, Zhang L, Jiang H, Li Y. 2016. Tropical birds are declining in the  
684 Hainan Island of China. *Biological Conservation* (in press). DOI:  
685 10.1016/j.biocon.2016.05.029.

686 Yan J. 2008. The study on evolutionary history of Hainan Island's ecological environment.  
687 Beijing. Science Press. (in Chinese)

688 Yang X, Wu Q. 2002. Study on the characteristics of rural vegetation and the eco-environmental  
689 problems of agriculture development in Wenchang Changsa, Hainan. *Nature science*  
690 *journal of Hainan university* 20: 239-242. DOI: 1004-1729(2002)03-0239-04. (in Chinese  
691 with English abstract)


692 Zhang YQ, Uusivuori J, Kuuluvainen J. 2000. Econometric analysis of the causes of forest land-  
693 use changes in Hainan, China. *Canadian Journal of Forest Research* 30: 1913-1921. DOI:  
694 10.1139/x00-123.

695 Zon R, Sparhawk WN. 1923. *Forest Resources of the World*. New York: McGraw-Hill Book Co.

# Figure 1


Location and topography of Hainan Island, China.

JFL, Jianfengling Mountain; BWL, Bawangling Mountain; YGL, Yinggeling Mountain; WZS, Wuzhishan Mountain; DLS, Diaoluoshan Mountain.




# Figure 2

Flowchart showing the methodological sequence describing the input data, the modelling and validation approaches, and the generation of predictions.




# Figure 3

Distribution patterns and histogram statistics of natural forest cover (NFC) in the 1950s (A), the 1970s (B), the 1990s (C) and the 2010s (D) on Hainan Island.



# Figure 4

Trend of total natural forest cover (NFC) from the 1950s to the 2010s on Hainan Island.






# Figure 5


The site-specific relationships between NFC and its change rate (A), and the distribution pattern of change rate (B) from the 1950s to the 2010s.

Green represents high NFC ( $\geq 60\%$ ) in the 1950s and had NFC increased from the 1950s to the 2010s; Red represents low NFC ( $\leq 20\%$ ) in the 1950s and had NFC decreased from the 1950s to the 2010s.



# Figure 6

Hot-spots of natural forest cover (NFC) decrease and increase in four assessed periods: 1950s-2010s (A), 1950s-1970s (B), 1970s-1990s (C), and 1990s-2010s (D) on Hainan Island.



# **Table 1**(on next page)

Changes in road-related variables of main and secondary roads from the 1950s to the 2010s on Hainan Island.

Data are the means and standard deviation of all grid cells. SRL, sum of road length (km km<sup>-2</sup>); DNR, distance from the grid cell centroid to the nearest road (km); NON, number of nodes of road network (ea km<sup>-2</sup>); MND, mean number of roads connected by each node (ea km<sup>-2</sup>).

1 **Table 1**

|            |                | 1950s     | 1970s     | 1990s     | 2010s     |
|------------|----------------|-----------|-----------|-----------|-----------|
| <b>SRL</b> | Main road      | 0.11±0.17 | 0.23±0.22 | 0.27±0.24 | 0.45±0.27 |
|            | Secondary road | 0.18±0.18 | 0.36±0.23 | 0.36±0.23 | 0.36±0.23 |
| <b>DNR</b> | Main road      | 0.30±0.29 | 0.09±0.08 | 0.08±0.08 | 0.04±0.04 |
|            | Secondary road | 0.11±0.11 | 0.05±0.05 | 0.05±0.04 | 0.05±0.04 |
| <b>NON</b> | Main road      | 0.01±0.04 | 0.09±0.20 | 0.11±0.20 | 0.20±0.23 |
|            | Secondary road | 0.03±0.06 | 0.23±0.21 | 0.23±0.21 | 0.23±0.21 |
| <b>MND</b> | Main road      | 0.02±0.05 | 0.04±0.04 | 0.05±0.04 | 0.07±0.04 |
|            | Secondary road | 0.03±0.05 | 0.05±0.03 | 0.05±0.03 | 0.05±0.03 |

2

## Table 2 (on next page)

The explanatory power (Adjusted  $R^2$ , %) of the change of road-related indicators in a simple linear regression model.

SRL, sum of road length; DNR, distance from the grid cell centroid to the nearest road; NON, number of nodes of road network; MND, mean number of roads connected by each node. The  $p$  value represents the significance level of the relationship between NFC change rate and change of road-related indicators. \*, significant at  $p < 0.05$ ; \*\*, significant at  $p < 0.01$ ; \*\*\*, significant at  $p < 0.001$ .

1 **Table 2**

| Change of road-related indicators |               | Change of NFC in<br>the 1950s–2010s |                | Change of NFC<br>in the 1950s–1970s |                | Change of NFC<br>in the 1970s–1990s |                | Change of NFC<br>in the 1990s–2010s |                |
|-----------------------------------|---------------|-------------------------------------|----------------|-------------------------------------|----------------|-------------------------------------|----------------|-------------------------------------|----------------|
| Type of road                      | Indicators    | Adjusted R <sup>2</sup>             | <i>p</i> value | Adjusted R <sup>2</sup>             | <i>p</i> value | Adjusted R <sup>2</sup>             | <i>p</i> value | Adjusted R <sup>2</sup>             | <i>p</i> value |
| Main road                         | Change of SRL | 11.20                               | < 0.001***     | 14.22                               | < 0.001***     | 20.73                               | < 0.001***     | 4.89                                | < 0.001***     |
|                                   | Change of DNR | 11.07                               | < 0.001***     | 0.27                                | 0.031*         | 5.97                                | < 0.001***     | 0.45                                | 0.007***       |
|                                   | Change of NON | 4.83                                | < 0.001***     | 4.37                                | < 0.001***     | 19.03                               | < 0.001***     | 1.76                                | < 0.001***     |
|                                   | Change of MND | 5.59                                | < 0.001***     | 8.31                                | < 0.001***     | 7.35                                | < 0.001***     | 2.30                                | < 0.001***     |
|                                   | Overall       | 23.40                               | < 0.001***     | 17.67                               | < 0.001***     | 25.20                               | < 0.001***     | 5.10                                | < 0.001***     |
| Secondary road                    | Change of SRL | 0.32                                | 0.021*         | 0.14                                | 0.089          | 0.36                                | 0.015*         | < 0.01                              | 0.613          |
|                                   | Change of DNR | 7.14                                | < 0.001***     | 5.26                                | < 0.001***     | <0.01                               | 0.801          | 0.32                                | 0.021*         |
|                                   | Change of NON | 5.11                                | < 0.001***     | 2.26                                | < 0.001***     | <0.01                               | 0.932          | 0.01                                | 0.422          |
|                                   | Change of MND | 0.22                                | 0.045*         | 1.11                                | < 0.001***     | <0.01                               | 0.505          | 0.01                                | 0.605          |
|                                   | Overall       | 13.3                                | < 0.001***     | 8.99                                | < 0.001***     | 0.31                                | 0.085          | 0.12                                | 0.228          |
| Main road & Secondary road        |               | 30.7                                | < 0.001***     | 22.5                                | < 0.001***     | 26.0                                | < 0.001***     | 5.4                                 | < 0.001***     |

2