

Storm effects on intertidal invertebrates: increased beta diversity of few individuals and species (#15810)

1

First revision

Please read the **Important notes** below, the **Review guidance** on page 2 and our **Standout reviewing tips** on page 3. When ready [submit online](#). The manuscript starts on page 4.

Important notes

Editor and deadline

James Reimer / 26 Apr 2017

Files

1 Tracked changes manuscript(s)

1 Rebuttal letter(s)

6 Figure file(s)

4 Table file(s)

1 Raw data file(s)

Please visit the overview page to [download and review](#) the files not included in this review PDF.

Declarations

Involves a field study on animals or plants.



Please read in full before you begin

How to review






When ready [submit your review online](#). The review form is divided into 5 sections. Please consider these when composing your review:

- 1. BASIC REPORTING**
- 2. EXPERIMENTAL DESIGN**
- 3. VALIDITY OF THE FINDINGS**
4. General comments
5. Confidential notes to the editor

 You can also annotate this PDF and upload it as part of your review

To finish, enter your editorial recommendation (accept, revise or reject) and submit.

BASIC REPORTING

-  Clear, unambiguous, professional English language used throughout.
-  Intro & background to show context. Literature well referenced & relevant.
-  Structure conforms to [PeerJ standards](#), discipline norm, or improved for clarity.
-  Figures are relevant, high quality, well labelled & described.
-  Raw data supplied (see [PeerJ policy](#)).

EXPERIMENTAL DESIGN

-  Original primary research within [Scope of the journal](#).
-  Research question well defined, relevant & meaningful. It is stated how the research fills an identified knowledge gap.
-  Rigorous investigation performed to a high technical & ethical standard.
-  Methods described with sufficient detail & information to replicate.

VALIDITY OF THE FINDINGS

-  Impact and novelty not assessed. Negative/inconclusive results accepted. *Meaningful* replication encouraged where rationale & benefit to literature is clearly stated.
-  Data is robust, statistically sound, & controlled.
-  Conclusions are well stated, linked to original research question & limited to supporting results.
-  Speculation is welcome, but should be identified as such.

The above is the editorial criteria summary. To view in full visit <https://peerj.com/about/editorial-criteria/>

7 Standout reviewing tips

3



The best reviewers use these techniques

Tip

Example

Support criticisms with evidence from the text or from other sources

Smith et al (J of Methodology, 2005, V3, pp 123) have shown that the analysis you use in Lines 241-250 is not the most appropriate for this situation. Please explain why you used this method.

Give specific suggestions on how to improve the manuscript

Your introduction needs more detail. I suggest that you improve the description at lines 57- 86 to provide more justification for your study (specifically, you should expand upon the knowledge gap being filled).

Comment on language and grammar issues

The English language should be improved to ensure that your international audience can clearly understand your text. I suggest that you have a native English speaking colleague review your manuscript. Some examples where the language could be improved include lines 23, 77, 121, 128 - the current phrasing makes comprehension difficult.

Organize by importance of the issues, and number your points

1. Your most important issue
2. The next most important item
3. ...
4. The least important points

Give specific suggestions on how to improve the manuscript

Line 56: Note that experimental data on sprawling animals needs to be updated. Line 66: Please consider exchanging "modern" with "cursorial".

Please provide constructive criticism, and avoid personal opinions

I thank you for providing the raw data, however your supplemental files need more descriptive metadata identifiers to be useful to future readers. Although your results are compelling, the data analysis should be improved in the following ways: AA, BB, CC

Comment on strengths (as well as weaknesses) of the manuscript

I commend the authors for their extensive data set, compiled over many years of detailed fieldwork. In addition, the manuscript is clearly written in professional, unambiguous language. If there is a weakness, it is in the statistical analysis (as I have noted above) which should be improved upon before Acceptance.

Storm effects on intertidal invertebrates: increased beta diversity of few individuals and species

Guilherme N Corte ^{Corresp., 1}, **Thomas A Schlacher** ², **Helio H Checon** ¹, **Carlos A M Barboza** ³, **Eduardo Siegle** ⁴, **Ross A Coelman** ⁵, **Antonia Cecília Z Amaral** ¹

¹ Departamento de Biologia Animal, Universidade Estadual de Campinas, Campinas, São Paulo, Brasil

² School of Science and Engineering, University of the Sunshine Coast, Maroochydore DC, Queensland, Australia

³ Núcleo em Ecologia e Desenvolvimento, Universidade Federal do Rio de Janeiro, Macaé, Rio de Janeiro, Brasil

⁴ Instituto Oceanográfico, Universidade de São Paulo, São Paulo, São Paulo, Brasil

⁵ School of Life and Environmental Science, University of Sydney, Sydney, New South Wales, Australia

Corresponding Author: Guilherme N Corte

Email address: guilhermecorte@yahoo.com.br

Climate change is predicted to lead to more extreme weather events, including changes to storm frequency, intensity and location. Yet, the ecological responses to storms are incompletely understood for sandy shorelines, the globe's longest land-ocean interface. Here we document how storms of different magnitude impacted the invertebrate assemblages on a tidal flat in Brazil. We specifically tested the relationships between wave energy and spatial heterogeneity, both for habitat properties (habitat heterogeneity) and fauna (β -diversity), predicting that larger storms redistribute sediments and hence lead to spatially less variable faunal assemblages. The sediment matrix tended to become less heterogeneous across the flat after high-energy wave events, whereas β -diversity of the fauna increased after storms. This higher β -diversity was primarily driven by species losses. Significantly fewer species at a significantly lower density occurred within days to weeks after storms. Negative density and biomass responses to storm events were most prominent in crustaceans. Invertebrates assemblages appeared to recover within a short interval (weeks to months) after storms months, highlighting that most species typical of sedimentary shorelines are, to some degree, resilient to short-term changes in wave energy. Given that storm frequency and intensity are predicted to change in the coming decades, identifying properties that determine resilience and recovery of ecosystems constitute a research priority for sedimentary shorelines and beyond.

Storm effects on intertidal invertebrates: increased beta diversity of few individuals and species

Short title: Storm effects on intertidal invertebrates

Guilherme N. Corte^{1§*}, Thomas A. Schlacher^{2§}, Helio H. Checon¹, Carlos A. M. Barboza³, Eduardo Siegle⁴, Ross A. Coleman⁵, A. Cecilia Z. Amaral¹

¹Departamento de Biologia Animal, Universidade Estadual de Campinas (UNICAMP), CEP 13083-862, Campinas, SP, Brazil. hchecon@yahoo.com.br, ceamaral @unicamp.br

²School of Science and Engineering, University of the Sunshine Coast, Maroochydore DC, QLD 4558, Australia. tschlach@usc.edu.au

³Núcleo em Ecologia e Desenvolvimento, Universidade Federal do Rio de Janeiro, CEP, 27965-045Macaé, RJ. carlosambarboza@gmail.com

⁴Instituto Oceanográfico, Universidade de São Paulo (USP), CEP 05508-120 São Paulo, SP, Brazil. esiegle@usp.br

⁵Coastal & Marine Ecosystems, School of Life and Environmental Science, The University of Sydney, NSW 2006, Australia. ross.coleman@sydney.edu.au

§ both authors contributed equally to this work

*Corresponding author: guilhermecorte@yahoo.com.br

Abstract

Climate change is predicted to lead to more extreme weather events, including changes to storm frequency, intensity and location. Yet, the ecological responses to storms are incompletely understood for sandy shorelines, the globe's longest land-ocean interface. Here we document how storms of different magnitude impacted the invertebrate assemblages on a tidal flat in Brazil. We specifically tested the relationships between wave energy and spatial heterogeneity, both for habitat properties (habitat heterogeneity) and fauna (β -diversity), predicting that larger storms redistribute sediments and hence lead to spatially less variable faunal assemblages. The sediment matrix tended to become less heterogeneous across the flat after high-energy wave events, whereas β -diversity of the fauna increased after storms. This higher β -diversity was primarily driven by species losses. Significantly fewer species at a significantly lower density occurred within days to weeks after storms. Negative density and biomass responses to storm events were most prominent in crustaceans. Invertebrate assemblages appeared to recover within a short interval (weeks to months) after storms months, highlighting that most species typical of sedimentary shorelines are, to some degree, resilient to short-term changes in wave energy. Given that storm frequency and intensity are predicted to change in the coming decades, identifying properties that determine resilience and recovery of ecosystems constitute a research priority for sedimentary shorelines and beyond.

1. Introduction

Extreme weather events, including changes to storm frequency and intensity, are predicted to increase over the 21st century (IPCC 2013, Lin and Emanuel 2016, Walsh et al. 2016). These global changes to ecosystem physical and chemical conditions are having numerous and widespread biological impacts in the sea and on land (Weatherdon et al. 2016). In the global oceans, climate change is expected to substantially alter the provision of ecosystems services critical to humankind, such as coastal protection and capture fisheries (Gattuso et al. 2015), but many responses in marine ecosystems still remain incompletely understood (Hauser et al. 2016, Nagelkerken and Munday 2016).

Storms may cause massive changes to coastal environments, particularly on sedimentary shorelines (Mateo and Garcia-Rubies 2012), often causing the translocation of sediment from the beach and dunes, and the landwards movement of the coastline (Masselink et al. 2016). These large habitat changes are usually accompanied by impacts to faunal assemblages, best documented for benthic invertebrates, seagrass meadows and algal communities (Lucrezi et al. 2010, Jaramillo et al. 2012, Mateo and Garcia-Rubies 2012).

The unpredictable nature of storms generally precludes the use of a rigorous experimental design to specifically test for storm effects, meaning that nearly all published 'storm studies' are largely opportunistic (Harris et al. 2011). In addition, often only a few or no data points are available immediately before a storm, post-storm sampling can be truncated, and for large storms it is challenging or impossible to find control areas that were not affected by the event (Posey et al. 1996); arguably, this makes attribution of ecological patterns to storm effects somewhat weak. An alternative is to make *a priori* predictive hypotheses based on knowledge of the biology of species and their likely response to large disturbance events in their habitat (Harris et al. 2011).

Here, we combine oceanographic, sediment and biological data to investigate how storms can affect the sedimentary habitat of a tidal flat in Southeast Brazil and the macrobenthic assemblages inhabiting it. Specifically, we tested four complementary, predictive hypotheses:

1. Higher wave energy during storms may translocate and disperse large sediment volumes (Masselink et al. 2016). We therefore predict that habitat heterogeneity (i.e. the spatial variation in seafloor properties amongst sampling sites) would be reduced after storms.
2. Habitat heterogeneity can be a major determinant for ecological assemblages, typically promoting beta diversity (i.e. variability in species composition among sampling units for a given area)

(Anderson et al. 2006, Schlacher et al 2007, McClain and Barry 2010, Meager et al. 2011).

Therefore, we expect that storms lower beta diversity of the fauna.

3. Disturbance caused by storms has been reported to detrimentally affect populations of benthic species (Jaramillo et al. 1987, Mateo and Garcia-Rubies 2012). Accordingly, we expect lower species richness, abundance, and biomass of invertebrates after storms.
4. Given that we expect lower β -diversity (prediction 2) and reduced number of species after storms (prediction 3), we predict that changes in β -diversity may be mainly attributable to species losses rather than species replacement.

2. Material and Methods

2.1 Study area

This study was done on the intertidal flats of Araçá Bay (Brazil, 23° 49'S, 45° 24'W; Fig. 1), a sheltered and heterogeneous intertidal flat adjacent to the São Sebastião Channel, Southeast Brazil (Amaral et al. 2010). The area is relatively small (ca. 750 m wide and long) and protected from the prevailing swell by São Sebastião Island (Fig. 1). It is one of few tide-dominated environments along the southeastern coast of Brazil (Dottori et al. 2015). Hydrographic properties of Araçá Bay are subject to physical forcing by frontal systems, when current speeds can increase eightfold (Fo 1990). At the region, the highest storm waves are associated to cold fronts and reaching offshore significant wave heights of 6.4m (Pianca et al. 2010).




Fig. 1 Map showing the location of the study area (a) and the sampling sites in the intertidal area of Araçá Bay (b).

2.2 Field sampling

Field work was done during spring tides on four times, at ca. three month intervals, from September 2011 to July 2012 (25 September 2011, 5 February 2012, 7 May 2012, and 29 July 2012). Three storm events occurred during the study (22 November 2011, 06 May 2012, 18 July 2012; Fig. 2), all accompanied by torrential rain, strong winds, flooding, and building damages. We sampled on the first spring tide after the storms in May and July 2012 (one-day lag in May and 11 days in July).

During each event, field work was done early in the morning of two consecutive days, collecting fauna from 34 sites. The sampling sites were positioned to encompass habitat diversity of the tidal flat (i.e. different sediment types over a range of depths), and to achieve a reasonable dispersion and spatial coverage (Fig. 1). The position of each sampling site was recorded with a GPS (Garmin eTrex Legend, datum WGS84) and the same locations (± 1 m) were sampled during each of the four sampling events. Three faunal samples (corer: 20 cm inner diameter, 20 cm depth) and one sediment sample (corer: 3 cm inner diameter, 20 cm deep) were collected per site and event.

2.2 Biological and environmental data

Fauna cores were washed on the same day of collection through a 0.3 mm mesh sieve, and the retained fauna was fixed in 70% ethanol. Sediment granulometric analysis was performed with standard dry sieving described by Suguio (1973). Sediment statistics were calculated with SysGran software (Camargo 2006) using the classifications of Folk and Ward (1957). Organic matter content was determined by weight losses of dried samples (60°C for 24 h) after incineration (550°C for 6 h). Calcium carbonate content was determined by 10% HCl digestion.

Sediment temperature and interstitial water salinity were measured *in situ* with a digital thermometer and an analog refractometer (precision of 0.01 and 0.1 units, respectively). Wave height and period for the region were obtained for 24.5 S and 45.5 W from the global wave generation model WaveWatch III (NCEP/NOAA). Wave power (P_w) was calculated as: $P_w = \rho g^2 H^2 T / 32\pi$, where ρ is water density (1,027 kg/m³), g the acceleration due to gravity (9.81 m/s²), H the wave height (m), and T the wave period (s) (Herbich 2000). We considered wave height and power for the three days before each sampling event. This time lag was found to show the strongest correlation between wave height/power and changes in macrobenthic species in the area (Turra et al. 2016).

All work was done in accordance with permit No. 19887-1 issued by the federal environmental agency, Ministério do Meio Ambiente, Instituto Chico Mendes de Conservação da Biodiversidade.

2.3 Data analysis

We tested for differences in habitat heterogeneity and fauna β -diversity amongst times with permutational analysis of multivariate dispersion (PERMDISP, Anderson 2006). In this analysis, higher multivariate dispersion is an indication of higher variability (i.e., higher habitat heterogeneity and β -diversity) among sampling sites (Anderson 2006). For habitat heterogeneity, the test was based on Euclidean distances calculated from normalized sediment data. For fauna β -diversity, the test was based on Hellinger transformation (Legendre and De Cáceres 2013) calculated from abundance data for the full suite of species. PERMDISP analysis was done using Primer 6 software (Clarke and Gorley 2006). Ordination plots (nMDS) were computed with the vegan package in R (Oksanen 2017) to illustrate differences in habitat heterogeneity and β -diversity (i.e. dispersion of sampling sites) between sampling events.

We tested for differences in species richness, abundance and biomass of invertebrate assemblages amongst times using general linear models with 'Time' as fixed factors. Models were adjusted using the negative binomial distributions for count data (species richness and abundance) and gamma distributions for continuous data (biomass). We used Tukey post-hoc tests to examine differences among sampling times using the MASS package in R (Ripley et al. 2013). This framework was used to investigate differences in the whole assemblage and also in the main groups of intertidal macrofauna (i.e., molluscs, polychaetes and crustaceans) separately.

We used the Similarity percentage analysis (SIMPER) based on Bray-Curtis distance to investigate the contribution of each individual species to the differences in species assemblages among sampling periods. Data was $\log(x+1)$ transformed before analysis to reduce influence of abundant species. SIMPER was done in Primer 6 software (Clarke and Gorley 2006).

We used the β -diversity partitioning framework of Podani and Schmera (2011) and Carvalho et al. (2012) to investigate compositional changes of macrobenthos (i.e. β -diversity) over time. This framework calculates compositional differences among communities (β_{total}) and partitions it into β diversity attributed to species replacement (β_{repl}) and β diversity attributed to species loss or gain (β_{rich}). This analysis was done with the R package BAT (Cardoso et al. 2015)

3. Results

Seawater temperature varied seasonally, whereas salinity and organic matter content of the sediment changed relatively little over time (Table 1). The silt-, clay-, and fine sand fraction of the sediment

179 increased between Sep. 2011 and July 2012 (Table 1). Waves were higher and more powerful before
 180 samplings in May and July 2012 (Table 1, Figure 2).
 181

182 **Table 1.** Environmental parameters recorded.

	September 2011	February 2012	May 2012	July 2012
	mean (se)	mean (se)	mean (se)	mean (se)
Temperature (°C)	21.9 (0.2)	27.4 (0.2)	25.0 (0.2)	20.4 (0.1)
Salinity	32.3 (0.3)	31.7 (0.9)	30.6 (0.7)	29.9 (0.6)
Mean grain size (ϕ)	2.5 (0.7)	2.7 (0.7)	2.7 (0.5)	2.8(0.64)
Silt and clay (%)	4.2 (0.6)	4.7 (0.6)	4.8 (0.6)	5.7 (0.9)
Fine sand (%)	68.4 (3.2)	73.5 (3.2)	74.1 (3.8)	74.7 (3.1)
Coarse sand (%)	10.7 (1.6)	9.5 (1.7)	7.9 (1.3)	7.3 (1.6)
Pebbles (%)	6.2 (1.4)	3.7 (1.1)	3.2 (0.9)	3.2 (1.0)
Organic matter (%)	1.6 (0.1)	1.7 (0.2)	1.7 (0.2)	1.9 (0.2)
CaCO ₃ (%)	4.9 (0.4)	4.4 (0.4)	3.8 (0.5)	3.5(0.3)
Height of waves (m)	1.5 (0.06)	1.6 (0.04)	2.1 (0.11)	1.7 (0.04)
Power of waves (10 ⁴ W/s)	20.1 (1.7)	18.1 (7.3)	42.8 (5.3)	30.4 (3.4)

183




Fig. 2 Wave height (a) and wave power (b) during the study period (sampling events are shown by dots).

We recorded 126 species from 33,320 individuals during the study (supplementary data). Polychaetes, molluscs and crustaceans made up 94% of species (polychaetes: 67 species; molluscs: 34 species; crustaceans: 18 species). Crustaceans were the most abundant group, comprising 56.5% of all individuals, mainly because of the high number of the tanaidacean *Monokalliapseudes schubartti* (Mañé-Garzón, 1949); polychaetes made up 39.6%, and molluscs 3.7% of catches (supplementary data).

H1: Lower habitat heterogeneity after storms

Sediment properties were spatially more homogeneous after periods of higher wave power (Fig. 3a), but differences between sampling times were not significant (Fig. 4a; PERMDISP $P = 0.586$).




Fig. 3 nMDS biplot of Euclidean distance based normalized environmental data (a) and nMDS biplot of Hellinger dissimilarity matrix based on macrobenthic abundance data (b). Samples from September 2011 are located inside the polygon plotted in red, from February 2012 in yellow, from May 2012 in blue, and from July 2012 in green. Stress for these ordination are 0.10 (a) and 0.19 (b).




Fig. 4 Habitat heterogeneity (a) and β -diversity of macrobenthic assemblages (b) during four sampling events associated with significant variation in wave energy preceding each event. Measure of habitat heterogeneity and β -diversity is the distance from centroids across all sites at a time. Letters and colours denote different groups ($P < 0.05$) in permutational analysis of multivariate dispersion (PERMDISP). Error bars denote standard errors.

H2: B-diversity declines after storms due to more homogenous sediment matrix.

Macrobenthic assemblages showed a significantly higher β diversity following periods of higher wave power (Fig. 3b and 4b; PERMDISP $P = 0.001$). Species that contributed most to differences in macrobenthic assemblages among sampling periods are listed in Table 2 & 3. 'Storm effects' appear to be driven mainly by large declines in the abundance of the tanaid *Monokalliapseudes schubarti*, whereas polychaetes (i.e. *Capitella* sp.C, *Heteromastus filiformis*, *Armandia hossfeldi*) tended to increase in abundance following storm events (Tables 2 & 3).

Table 2. Similarity percentage (SIMPER) analysis showing the contribution (%) of the five most important species to differences in species assemblages among sampling periods.

	<i>Monokalliapseudes schubarti</i>	<i>Capitella</i> sp. C	<i>Heteromastus filiformis</i>	<i>Scoloplos</i> sp1	<i>Armandia hossfeldi</i>	Mean dissimilarity
Sep vs. Feb	10.98	4.96	n/a	5.34	n/a	68.1
Sep vs. May	11.14	6.55	n/a	5.46	n/a	74.9
Sep vs. Jul	9.65	6.85	5.51	5.27	6.47	69.3
Feb vs. May	16.03	8.17	5.04	5.44	n/a	75.3
Feb vs. Jul	12.75	7.9	5.9	5.37	7.24	71.6
May vs. Jul	8.29	8.97	6.63	6.22	8.04	69.9

Table 3. Temporal variation in the density (ind.m⁻²) of species that accounted for most of the assemblage-wide differences in macrobenthic assemblages among sampling events (cf. Table 2).

	September 2011	February 2012	May 2012	July 2012
	mean (se)	mean (se)	mean (se)	mean (se)
<i>Monokalliapseudes schubartii</i>	2151 (635)	3264 (571)	1776 (79)	538 (276)
<i>Capitella</i> sp.C	114 (68)	521 (257)	667 (428)	768 (336)
<i>Heteromastus filiformis</i>	23 (9)	32 (11)	47 (12)	133 (36)
<i>Scoloplos</i> sp1	112 (30)	63 (14)	35 (10)	87 (19)
<i>Armandia hossfeldi</i>	50 (25)	7 (4)	47 (19)	224 (74)

H3: Storm disturbance results in lower abundance, biomass, and species richness

Abundance, species richness and biomass were significantly lower in samples taken shortly after high-energy wave events (Fig. 5). The mean number of species per site was lowest at 9.82 species after the strongest wave event, compared with 11.82 to 14.35 species at other times (Fig. 5a). Abundance peaked at 4126 ind.m⁻² in Feb. 2012, declining to 1195 ind.m⁻² after the storm in May 2012 (Fig. 5b). Biomass declined from 6.5 to 3.5 gAFDW.m⁻² between September 2011 and May 2012 (Fig. 5c).





Fig. 5 Variation in the mean number of species per site (a), mean abundance (b) and mean biomass (c) of macrobenthic assemblages at four sampling events associated with significant variation in wave energy preceding each event. Letters and colours denote different groups in generalized linear models ($P < 0.05$). Error bars denote standard errors.

All major groups of intertidal macrofauna (i.e. molluscs, polychaetes, crustaceans) were affected by storms (Fig. 6). The mean number of species per site of all groups was significantly lower after the strongest wave event ($P < 0.05$) (Fig. 6a-c). Temporal patterns of changes in abundance and biomass did, however, differ between groups. Crustaceans showed the most pronounced density (Fig. 6f) and biomass (Fig. 6i) response, declining strongly after storms. Molluscs showed a broadly similar density

245 pattern to crustaceans, albeit being less pronounced (Figs. 6d & 6g), whereas the abundance of
246 polychaetes tended to increase following periods of higher wave energy (Figs. 6e & 6h).



24
248 **Fig. 6** Variation in the mean number of species, abundance, and biomass of molluscs (a, d & g), polychaetes (b, e &
249 h), and crustaceans (c, f & i) at four sampling events associated with significant variation in wave energy preceding
250 each event. Letters and colours denote different groups in generalized linear models ($P < 0.05$). Error bars denote
251 standard errors.

252
253
254 *H4: Species losses drive most of the change in β -diversity.*

255 Declines in species numbers accounted for most of temporal β -diversity in the macrobenthos, and its
256 contribution was higher shortly after storms (Table 4). By contrast, species replacement was less
257 important.

258

259

Table 4 – β -diversity and β -diversity partitioning among sampling periods. Higher values of β -diversity denote greater differences in the composition of species and number of individuals between two consecutive sampling times. Percentages indicate the amount of variation between periods attributable to species losses or species replacement.

	Total β diversity (β_{total})	Species replacement / substitution (β_{repl})	Species loss / gain richness differences (β_{rich})
Sep. vs Feb.	0.45	44.4%	56.4%
Feb. vs May	0.79	16.4 %	83.6 %
May vs July	0.47	11.7 %	89.3 %
mean	0.57	24.2 %	76.4 %

4. Discussion

Significant changes in macrobenthic species richness, abundance and biomass in a tropical tidal flat were associated with temporal variation in wave energy. This resulted in significant changes to fauna β -diversity over time that was mainly attributable to species losses, but, apparently, unlinked to variation in habitat heterogeneity.

Previous studies about the influence of storms on coastal soft-sediment ecosystems have shown that storms may have stronger impacts on environmental features than on the fauna (e.g. Saloman and Naughton 1984, Cochôa et al. 2006, Alves and Pezzuto 2009, Harris et al. 2011), and that offshore sediment transport is the dominant geo-morphological response of sedimentary shores to increased wave energy (Masselink et al. 2016). These studies, however, were mostly done on exposed ocean beaches, habitats with fewer species that are well adapted to high-energy conditions (Brown 1996, Schlacher et al. 2008). By contrast, our results showed that under more sheltered conditions, storm impacts were more evident in the fauna than in the environment.

The observed decrease in the number of species, individuals and biomass of macrobenthic assemblages may have been caused by accretion and redistribution of sediments, burying fauna at some site and winnowing them at others. Waves and currents may suspend fine-grained sediments (Lohrer et al. 2006), and alongshore sediment distribution within the same system or accretion of sediment in washover

deposit can occur after storms (Masselink et al. 2016). Moreover, Alcantará-Carrió et al. (2017) showed that the seaward transport of terrigenous sediment after intense rains in combination with resuspension of sediments by storm waves and transport by wind-driven currents alter the sedimentary features in the São Sebastião Channel. These hypothesized mechanisms mechanisms of fauna change are functionally supported by studies showing significant changes to the macrobenthos following sediment deposition and substantial alterations in hydrodynamic regimes (Jaramillo et al. 2012, Cummings et al. 2003, Rodil et al. 2011, Schlacher et al. 2012).

Whilst storms were followed by decreases in species richness in all major groups of the macrobenthos, richness changes were more pronounced in crustaceans which also declined strongly in abundance and biomass. This was mainly a consequence of massive (-95%) declines of the tanaid *Monokalliapseudes schubarti*. *M. schubarti* is small (ca. 5 mm) and builds tubes that rarely extend for more than 5 cm into the sediment, possibly making it more susceptible to sediment erosion (Nucci et al. 2001), In fact, morphological traits of benthic invertebrates have been suggested to modulate storms impacts (Mateo and Garcia-Rubies 2012), with small-bodied individuals and those with low mobility thought to be more susceptible to storms (Negrello Filho and Lana 2013, Urabe et al. 2013).

Fewer species of polychaetes were recorded at lower biomass after storm events. Some species, mostly small and tubiculous forms, such as the *Isolda pulchella* (Müller in Grube, 1858), were also less numerous after storms. By contrast, opportunistic polychaete species (e.g. *Capitella* sp.C, *Heteromastus filiformis* (Pearson and Rosenberg 1978)) increased in abundance after storms. We did not record significant changes in the biomass of molluscs, possibly a consequence of heavier, shelled forms of the macrobenthos being less likely to be displaced by turbulent currents associated with storms.

Storm effects appeared to be influenced by the timing of a storm in relation to the tidal regime. Masselink et al. (2016) found that storms impacts on the south-west coast of England were highest when the peak storm waves coincided with spring high tides. In our study, this was likely the case in May 2012 when the height of the storm passed during a spring tide. The observed effects of storms on the macrobenthic fauna of Araçá Bay also appeared to be stronger during a short time after storms. We found that differences in environmental and biotic characteristics were most pronounced in May 2012, when samples were taken one day after the storm had passed.

Most species typical of sedimentary shorelines are, to some degree, adapted to high-energy conditions and hence may recover relatively quickly (e.g. within days to weeks) from most storm events (Harris et al. 2011, Schlacher and Thompson 2013; Machado et al. 2016). For example, in a subtropical coastal soft-sediment ecosystem in South Brazil, Gallucci and Netto (2004) found that abundance and number of species of macrobenthic organisms declined during the passage of a cold front, but all values were back

to pre-frontal conditions 24 h later. Similarly, Machado et al. (2016) found recovery of macrobenthic assemblages inhabiting tropical ocean exposed beaches within seven weeks of a storm. It is important to emphasize, however, that recovery depends on the magnitude, spatial scale and return frequency of the disturbance events in soft-sediment environments and other marine systems (Lucrezi et al. 2010, Urabe et al. 2013, McClain and Schlacher 2015, Schlacher et al. 2015). The most powerful storms may cause ecological changes that require years to recover and may compromise the spatial and trophic structure of the ecosystems (Jaramillo et al. 1987, Mateo and Garcia-Rubies 2012).

5. Conclusion

Here we show that storms can cause significant changes to macrobenthic assemblages inhabiting a tidal flat. Decreases in species richness, abundance and biomass of invertebrate assemblages were related to increase wave power. Species losses drove changes to higher β -diversity but the fauna appeared to recover within a few weeks. Changes to habitat features were comparatively smaller. Given that storm activity, location and intensity are predicted to change over the coming decades in a warming world (Lin and Emanuel 2016, Walsh et al. 2016), ecological changes attributed to altered storm properties are likely expected. Unfortunately, the functional consequences of altered storm regimes for coastal ecosystems are fundamentally unknown, including the continued provision of ecosystem services such as coastal protection and capture fisheries. Thus, future work shall prioritise investigations of how ecological processes in coastal ecosystems respond to extreme events and which features may determine their resilience and recovery.

Acknowledgements

We thank Daniel Martin, Paulo Paiva, André Garraffoni, Maikon Di Domenico, David Schoeman, and Ronaldo Christofletti for comments and suggestions. We are also grateful to two anonymous referees that provided very constructive comments on the manuscript. Special thanks to AS Godoy, CF Silva, N Padovanni, RF Daolio, DG Filho, and R Alitto who helped with field work and species identifications.

References

- Alcántara-Carrió J, Sasaki DK, de Mahique MM, Taborda R, and de Souza LAP (2017) Sedimentary constraints on the development of a narrow deep strait (São Sebastião Channel, SE Brazil). *Geo-Marine Letters*, 1-14.
- Alves ES, and Pezzuto PR (2009) Effect of cold fronts on the benthic macrofauna of exposed sandy beaches with contrasting morphodynamics. *Brazilian Journal of Oceanography* 57:73-94.
- Amaral, A. C. Z., A. E. Migotto, A. Turra & Y. Schaeffer-Novelli, 2010. Araçá: biodiversidade, impactos e ameaças. *Biota Neotropica* 10: 219–264.
- Anderson MJ (2006) Distance-based tests for homogeneity of multivariate dispersions. *Biometrics* 62:245-253.

363 Anderson, M. J., Ellingsen, K. E., & McArdle, B. H. (2006). Multivariate dispersion as a measure of beta
364 diversity. *Ecology letters*, 9: 683-693.

365 Anderson MJ, Crist TO, Chase JM, Vellend M, Inouye BD, Freestone AL, Sanders NJ, Cornell HV, Comita LS,
366 Davies KF, Harrison SP (2011) Navigating the multiple meanings of β diversity: a roadmap for the practicing
367 ecologist. *Ecology letters* 14:19-28.

368 Brown AC (1996) Behavioural plasticity as a key factor in the survival and evolution of the macrofauna on exposed
369 sandy beaches. *Revista Chilena de Historia Natural* 69:469-474.

370 Camargo MG (2006) Sysgran: um sistema de código aberto para análises granulométricas do sedimento. *Revista*
371 *Brasileira de Geociências* 36: 371-378.

372 Cardoso P, Rigal F, and Carvalho JC (2015) BAT - Biodiversity Assessment Tools, an R package for the
373 measurement and estimation of alpha and beta taxon, phylogenetic and functional diversity. *Methods in Ecology and*
374 *Evolution* 6:232-236.

375 Carvalho JC, Cardoso P, and Gomes P (2012) Determining the relative roles of species replacement and species
376 richness differences in generating beta-diversity patterns. *Global Ecology and Biogeography* 21:760-771.

377 Clarke KR, Gorley RN (2006) PRIMER V6: user manual-tutorial. Plymouth Marine Laboratory

378 Cochôa AR, Lorenzi I, and Borzone aCAA (2006) A Influência da Passagem de uma Frente Meteorológica na
379 Distribuição da Macrofauna Bentônica Mesolitoral de uma Praia Arenosa Exposta. *Tropical Oceanography* 34:59-71.

380 Cummings V, Thrush S, Hewitt J, Norkko A, and Pickmere S (2003) Terrestrial deposits on intertidal sandflats:
381 Sediment characteristics as indicators of habitat suitability for recolonising macrofauna. *Marine Ecology Progress*
382 *Series* 253:39-54.

383 Dottori M, Siegle E, and Castro BM (2015). Hydrodynamics and water properties at the entrance of Araújo Bay,
384 Brazil. *Ocean Dynamics* 65: 1731-1741.

385 Fo C (1990) Wind driven currents in the Channel of São Sebastião: winter, 1979. *Boletim do Instituto Oceanográfico*
386 38:111-132.

387 Folk, RL and Ward WC (1957) Brazos River Bar: a study in the significance of grain-size parameters. *Journal of*
388 *Sedimentary Petrology* 27: 3-26.

389 Gallucci F, and Netto SA (2004) Effects of the passage of cold fronts over a coastal site: An ecosystem approach.
390 *Marine Ecology Progress Series* 281:79-92.

391 Gattuso JP, Magnan A, Billé R, Cheung WWL, Howes EL, Joos F, Allemand D, Bopp L, Cooley SR, Eakin CM,
392 Hoegh-Guldberg O, Kelly RP, Pörtner HO, Rogers AD, Baxter JM, Laffoley D, Osborn D, Rankovic A, Rochette J,
393 Sumaila UR, Treyer S, and Turley C (2015) Contrasting futures for ocean and society from different anthropogenic
394 CO2 emissions scenarios. *Science* 349.

395 Harris L, Nel R, Smale M, and Schoeman D (2011) Swashed away? Storm impacts on sandy beach macrofaunal
396 communities. *Estuarine Coastal and Shelf Science* 94:210-221.

397 Hauser DDW, Tobin ED, Feifel KM, Shah V, and Pietri DM (2016) Disciplinary reporting affects the interpretation of
398 climate change impacts in global oceans. *Global Change Biology* 22:25-43.

399 Herbich JB (2000) Handbook of coastal engineering. McGraw-Hill, New York

400 IPCC (2013) Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth
401 Assessment Report of the Intergovernmental Panel on Climate Change. [Stocker, T.F., D. Qin, G.-K. Plattner, M.
402 Tignor, S.K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex and P.M. Midgley (eds.)]. Cambridge University Press,
403 Cambridge, UK

404 Jaramillo E, Croker RA, and Hatfield EB (1987) Long-term structure, disturbance, and recolonization of macroinfauna
405 in a New Hampshire sand beach. *Canadian Journal of Zoology* 65:3024-3031.

406 Jaramillo E, Dugan JE, Hubbard DM, Melnick D, Manzano M, Duarte C, Campos C, and Sanchez R (2012)
407 Ecological implications of extreme events: Footprints of the 2010 earthquake along the Chilean coast. *PLoS ONE*
408 7:e35348. doi:35310.31371/journal.pone.0035348.

409 Legendre P, Cáceres M. Beta diversity as the variance of community data: dissimilarity coefficients and partitioning.
410 *Ecology Letters*. 2013 Aug 1;16(8):951-63.

411 Lin N, and Emanuel K (2016) Grey swan tropical cyclones. *Nature Climate Change* 6:106-111.

412 Lohrer A M, Hewitt JE, and Thrush SF (2006). Assessing far-field effects of terrigenous sediment loading in the
413 coastal marine environment. *Marine Ecology Progress Series* 315:13-18.

414 Lucrezi S, Schlacher TA, and Robinson W (2010) Can storms and shore armouring exert additive effects on sandy-
415 beach habitats and biota? *Marine and Freshwater Research* 61:951-962.

416 Machado PM, Costa LL, Suci MC, Tavares DC, and Zalmon IR (2016) Extreme storm wave influence on sandy
417 beach macrofauna with distinct human pressures. *Marine Pollution Bulletin*.

418 Masselink G, Scott T, Poate T, Russell P, Davidson M, and Conley D (2016) The extreme 2013/2014 winter storms:
419 Hydrodynamic forcing and coastal response along the southwest coast of England. *Earth Surface Processes and*
420 *Landforms* 41:378-391.

421 Mateo MA and Garcia-Rubies T (2012). Assessment of the ecological impact of the extreme storm of Sant Esteve's
422 Day (26 December 2008) on the littoral ecosystems of the north Mediterranean Spanish coasts. *Final Report (PIEC*
423 *200430E599)*.

424 McClain CR, Barry JP (2010) Habitat heterogeneity, disturbance, and productivity work in concert to regulate
425 biodiversity in deep submarine canyons. *Ecology* 91:964-76.

426 McClain CR, and Schlacher TA (2015) On some hypotheses of diversity of animal life at great depths on the sea
427 floor. *Marine Ecology* 36:849-872.

428 Meager J, Schlacher TA, and Green M (2011) Topographic complexity and landscape temperature patterns create a
429 dynamic habitat structure on a rocky intertidal shore. *Marine Ecology Progress Series* 428:1-12.

430 Nagelkerken I, and Munday PL (2016) Animal behaviour shapes the ecological effects of ocean acidification and
431 warming: Moving from individual to community-level responses. *Global Change Biology* 22:974-989.

432 Negrello Filho OA, and Lana PC (2013) Short-term stability of estuarine benthic assemblages: Are storms pattern-
433 defining events? *Zoologia* 30:266-272.

434 Nucci PR, Turra A., and Morgado EH (2001) Diversity and distribution of crustaceans from 13 sheltered sandy
435 beaches along Sao Sebastiao Channel, south-eastern Brazil. *Journal of the Marine Biological Association of the*
436 *United Kingdom* 81:475-484.

437 Oksanen J, Kindt R, Legendre P, O'Hara B, Stevens MH, Oksanen MJ, Suggests MA (2007) The vegan package.
438 *Community ecology package*.

439 Pianca C, Mazzini PL, Siegle E (2010) Brazilian offshore wave climate based on NWW3 reanalysis. *Brazilian Journal*
440 *of Oceanography* 58:53-70

441 Podani J, and Schmera D (2011) A new conceptual and methodological framework for exploring and explaining
442 pattern in presence - absence data. *Oikos* 120:1625-1638.

443 Posey M, Lindberg W, Alphin T, and Vose F (1996) Influence of storm disturbance on an offshore benthic community.
444 *Bulletin of Marine Science* 59:523-529.

445 Ripley B, Venables B, Bates DM, Hornik K, Gebhardt A, Firth D, Ripley MB (2013) Package 'MASS'. Cran R.

446 Rodil IF, Lohrer AM, Chiaroni LD, Hewitt JE, and Thrush SF (2011) Disturbance of sandflats by thin terrigenous
447 sediment deposits: Consequences for primary production and nutrient cycling. *Ecological Applications* 21:416-426.

448 Saloman CH, and Naughton SP (1984) Beach restoration with offshore dredged sand: effects on nearshore
449 macroinfauna. NOAA Technical Memorandum NMFS-SEFC-133

450 Schlacher TA, and Thompson L (2013) Environmental control of community organisation on ocean-exposed sandy
451 beaches. *Marine and Freshwater Research* 64:119-129.

452 Schlacher TA, Schoeman DS, Dugan JE, Lastra M, Jones A, Scapini F, and McLachlan A (2008) Sandy beach
453 ecosystems: key features, sampling issues, management challenges and climate change impacts. *Marine Ecology* 29
454 (S1):70–90. Schlacher TA, Noriega R, Jones A, and Dye T (2012) The effects of beach nourishment on benthic
455 invertebrates in eastern Australia: Impacts and variable recovery. *Science of the Total Environment* 435:411-417.

456 Schlacher TA, Weston MA, Schoeman DS, Olds AD, Huijbers CM, and Connolly RM (2015) Golden opportunities: A
457 horizon scan to expand sandy beach ecology. *Estuarine, Coastal and Shelf Science* 157:1-6.

458 Turra A, Pombo M, Petracco M, Siegle E, Fonseca M, Denadai MR (2016) Frequency, Magnitude, and Possible
459 Causes of Stranding and Mass-Mortality Events of the Beach Clam *Tivela mactroides* (Bivalvia: Veneridae). *PLoS*
460 *ONE* 11(1): e0146323. doi:10.1371/journal.pone.0146323

461 Urabe J, Suzuki T, Nishita T, and Makino W (2013) Immediate Ecological Impacts of the 2011 Tohoku Earthquake
462 Tsunami on Intertidal Flat Communities. *PLoS ONE* 8.

463 Walsh KJE, McBride JL, Klotzbach PJ, Balachandran S, Camargo SJ, Holland G, Knutson TR, Kossin JP, Lee Tc,
464 Sobel A, and Sugi M (2016) Tropical cyclones and climate change. *Wiley Interdisciplinary Reviews: Climate Change*
465 7:65-89.

466 Weatherdon LV, Magnan AK, Rogers AD, Sumaila UR, and Cheung WWL (2016) Observed and projected impacts of
467 climate change on marine fisheries, aquaculture, coastal tourism, and human health: An update. *Frontiers in Marine*
468 *Science* 3.

469

470