

1 **THE VENICE SPECIMEN OF *OURANOSAURUS NIGERIENSIS* (DINOSAURIA,**
2 **ORNITHOPODA)**

3
4 Filippo Bertozzo¹, Fabio Marco Dalla Vecchia², Matteo Fabbri³

5
6 1 Earth System Science, Vrije Universiteit Brussel, Pleinlaan 2, B-1050 Brussels/Bruxelles,
7 Belgium

8 2 Former address: Institut Català de Paleontologia Miquel Crusafont, Carrer de l'Escola Industrial
9 E-08201 Sabadell, Spain. Present address: Soprintendenza per i Beni Archeologici del Friuli
10 Venezia Giulia, Nucleo Operativo di Udine, Via Zanon, 22 I-33100 Udine, Italy, fabdalla@tin.it

11 3 Department of Geology and Geophysics, University of Yale, 210 Whitney Ave., New Haven,
12 CT06611, USA

13
14 Corresponding Author: Filippo Bertozzo¹

15 Email: filippo.bertozzo@gmail.com
16

Formatted: Italian (Italy)

ABSTRACT

Ouranosaurus nigeriensis is an iconic African dinosaur recorded by a few specimens, including a couple of nearly complete skeletons from the Lower Cretaceous Gadoufaoua locality of the Ténéré desert in Niger. Only the holotype was completely described in the unique paper dedicated to this important dinosaur, although a few bones of the paratype were also included in the description. A mounted skeleton of *Ouranosaurus nigeriensis* is exposed at the Natural History Museum of Venice, Italy since 1975. It was never explicitly established whether it is the paratype and second nearly complete skeleton reported in literature or a third, unreported specimen. We disentangle herein its complex history, (thanks also to an unpublished field map of the paratype) and describe it. ~~It~~ [This mounted specimen](#) includes the paratype material (found in 1970 and collected in 1972), with the exclusion of the left femur and the right coracoid (which were replaced with plaster copies) and possibly the manual phalanges. Some elements (e.g., the right femur, the right tibia, two dorsal vertebrae and some pelvic bones) were plausibly added from other individual/s. The vertebral column of the paratype is in a fair state of articulation and a better reference for the vertebral count of the taxon ~~respect to~~ [than](#) the holotype. Some differences are observed between the latter and the Venice specimen. For example, the first dorsals in the Venice specimen are unlike those reported or hypothesized in the holotype, suggesting that the dorsal count could be 15 and the trunk would be consequently shorter; there are five to six ~~more additional~~ proximal caudals in the Venice specimen ~~than in the holotype~~; the metacarpus is ~~slightly~~ [visibly](#) different in the two specimens. The linear size of the Venice specimen is about 90% ~~the linear size~~ [that](#) of the holotype. The osteohistological analysis (the first one for this taxon) of some long bones, a rib and a dorsal neural spine reveals that the specimen is a sub-adult ~~individual~~; this is ~~also~~ supported ~~also~~ by somatic evidence of immaturity. The ~~size and shape of its~~ dorsal 'sail', formed by the elongated neural spines of the dorsal, sacral and proximal caudal vertebrae, is unique ~~to this taxon as for size and shape~~ among ornithomimids; a display role is the most probable function for this bizarre structure.

INTRODUCTION

Ouranosaurus nigeriensis and *Spinosaurus aegyptiacus* are [among](#) the most iconic African dinosaurs because of their outstanding hypertrophic neural spines. *O. nigeriensis* comes from the upper part of the El Rhaz Formation at the Gadoufaoua locality of the Sahara Desert, located 145 km east of Agadez, Niger (Taquet, 1976). The El Rhaz Formation of Niger has yielded a rich dinosaur association including theropods (*Suchomimus*, *Cristatosaurus*, *Kryptops* and *Eocarcharia*), sauropods (*Nigersaurus*) and the ornithomimids *Ouranosaurus* and *Lurdusaurus* (Le Loeuff et al., 2012). ~~Its age was considered to be Aptian by~~ Taquet (1976) [suggested an Aptian age for this formation, and Aptian–Albian by](#), whereas Sereno et al. (1999) [subsequently proposed](#) Sereno et al. (1999); [an Aptian–Albian age, and](#) ~~but~~ Le Loeuff et al. (2012) [have recently proposed](#) a Barremian age. The detailed anatomical description of *Ouranosaurus nigeriensis* was published by Taquet (1976). Only a few specimens are formally referred to *O. nigeriensis* in that paper: the holotype GDF 300, the paratype GDF 381 and two isolated bones (GDF 301 and 302; Taquet, 1976, p. 58), although the discovery of several in situ specimens is mentioned in that paper (Taquet, 1976, p. 14–15). No other scientific works have been dedicated to this taxon since then, although a few papers dealt with it (Rasmussen, 1998; Dean-Carpentier, 2008; Taquet, 2012). Despite ~~to this and~~ the difficult access to the original holotype material, ~~it is always~~ [Ouranosaurus is frequently included in the cladistic phylogenetic analyses](#) of iguanodontian dinosaurs (e.g., Sereno, 1986; Norman, 2004, 2015; McDonald et al., 2010a, b, 2012). Since 1975, a nearly complete and mounted skeleton of *O. nigeriensis* ~~has been~~ [is](#) exhibited at the Museo di Storia Naturale (Natural History Museum) of Venice, Italy. Apparently, the Venice

specimen is not mentioned in Taquet (1976) as well as in any other scientific papers dealing with *Ouranosaurus*. Therefore, it had to be considered as still undescribed.

The aim of our work ~~was is~~: 1) ~~to~~ disentangle~~ing~~ the complicated history of the *Ouranosaurus* specimens with particular focus on the Venice specimen; 2) to describe the latter and compare its skeletal elements with those published by Taquet (1976) in order to ~~notice~~ highlight differences and get information useful for point 1; 3) to perform the first osteohistological analysis on *O. nigeriensis* sampling some significative bones and get information on the ontogenetic stage of the Venice specimen; 4) to establish whether the bones of the nearly complete mounted skeleton can ~~be~~ reliably ~~be~~ referred to a single individual or to more individuals.

Institutional abbreviation: CEA, Commissariat à l'Énergie Atomique et aux énergies alternatives, France; CSRL, Centro Studi e Ricerche Ligabue, Venice, Italy; GDF, Muséum National du Niger, Africa; IRSNB, Institut Royal des Sciences Naturelles de Belgique; MNHM, Muséum National d'Histoire Naturelle, Paris, France; MSNVE, Museo di Storia Naturale di Venezia (Natural History Museum of Venice), Venice, Italy; TMP, Royal Tyrrell Museum of Palaeontology, Drumheller, Canada.

MATERIALS AND METHODS

The object of this paper is the Venice specimen of *O. nigeriensis* (MSNVE 3714; Fig. 1), ~~which is catalogued with the number MSNVE 3714~~. It is apparently a nearly complete skeleton, ~~which is exhibited at the MSNVE~~ mounted in a bipedal posture. ~~It was the object of the bachelor thesis of one of us (FB) under the supervision of FMDV at the University of Bologna in years 2012-2013 (Bertozzo, 2012)~~. The specimen was donated to the MSNVE by the Italian entrepreneur and philanthropist Giancarlo Ligabue (founder of the Centro Studi e Ricerche Ligabue, Venice) in 1975. ~~then, Since that date, it has been~~ exhibited ~~to in~~ the public in the Museum. According to the available information, the specimen underwent two distinct restoration phases. The first preparation of the bones used to assemble the mount was done by French preparators at the Muséum National d'Histoire Naturelle of Paris, France, before 1975 when the skeleton was mounted in Venice. A make-up and remount of the specimen was performed by an ~~Italian private firm~~ in ~~the years~~ 1999-2000. No reports or any kind of available documentation exist about the restoration ~~done on~~ of the bones before 1975 and in 1999-2000. A list of the original material is also unavailable.

The history of the *Ouranosaurus* specimens ~~and, including~~ MSNVE 3714 ~~in particular was was~~ traced back based on ~~the~~ papers dealing with the paleontological expeditions to Gadoufaoua (Ligabue et al., 1972; Ligabue & Rossi-Osmida, 1975; Taquet, 1970; 1976; 1998; Boccardi & Boccazzi, 1978), the information supplied by the Centro Studi e Ricerche Ligabue of Venice, and the personal communication from Philippe Taquet. Ronan Allain, MNHN, kindly made available to us a copy of the map of the in situ specimen of *O. nigeriensis* found in 1970 along the landing strip (indicated as GDF 381 in Taquet, 1976), which was drawn by Philip Taquet and is deposited at MNHN.

The specimen is mounted on a metal ~~support frame~~. In order to photograph and describe them, all the bones were removed from the ~~support frame~~, with the exception of the sacrum, which is fixed to it. In order to have a complete photographic documentation of the specimen, we took pictures of every single bone in its cranial (anterior), caudal (posterior), dorsal, ventral, lateral, and medial views. We used a camera Canon EOS 600D, lens Tamron 17-5 mm F2.8, focal 50 mm and sensitivity 100 ISO. The photographs are stored in the archive of the MSNVE, which is accessible to researches by contacting the responsible for Research and Scientific Divulcation of the Museum. We used a caliper 200 mm long with measurement error of 0.01 mm and a metric string 100 cm long (measurement error of 0.1 cm) to measure the bones. A table with all the measurements is ~~reported included~~ in the Supplementary Information. In order to ~~detect~~ identify the reconstructed

Formatted: French (France)

Commented [PG1]: You should name it as it is still existing: cfr Flavio Bacchia

parts ~~in each element~~, we took pictures under UV-light using a Wood Lamp (SKU 51029, emitting ultraviolet light at 4 W).

We consider as proximal caudal vertebrae those with pleurapophyses (often improperly reported as caudal ribs or transverse processes in the literature); middle caudals lack pleurapophyses but have hemapophyses; distal caudals lack pleurapophyses and hemapophyses. The cervical-dorsal transition in the vertebral column was identified following Norman (1986). Centrum height was measured in the caudal (posterior) articular facet; neural spine height was measured as the straight line from the mid-point of the spine in correspondence of the dorsal margin of the postzygapophysis to the apex of the spine. The orientation of the skeletal elements is that hypothesized for the living animal, unless otherwise specified.

~~Different phylogenetic hypothesis on the iguanodontian ornithopods have been published in the last decade. We choose that by~~ McDonald et al.'s (2012) ~~phylogenetic context was chosen~~ as a reference ~~for the present study~~, when needed.

Bone surface texture, degree of fusion of the elements and obliteration of the sutures in skulls and vertebrae are the most common approaches to assess the ontogenetic stage of fossil tetrapods ~~individuals~~ (e.g., Brochu, 1996; Werning, 2012). However, histological analysis ~~still remains~~ remains the most reliable methodology ~~to for~~ establishing it and ~~to obtain an estimation of for estimating~~ the absolute age of ~~the an~~ individual (e.g. Chinsamy 2005; Erickson et al. 2004; Erickson 2005). The left humerus, the right femur, the right tibia, the neural spine of the dorsal vertebra 14, and the right dorsal rib 15 were selected for the osteohistological analysis. Core samples were taken from the long bones following the method described ~~in by~~ Stein & Sander (2009) and using an electric drill press Timbertech Kebo01 and a cylindrical diamond drill bit (16 mm in diameter, 80 mm in height and with a 2 mm-thick wall). Samples were taken from the diaphysis of the long bones. Only areas lacking evident superficial erosion and surface cracks were selected. The shaft of the rib was transversally cut proximally. This area was selected because it is considered to preserve the most complete growth record (Erickson 2005). The neural spine was cross-sectioned at three different levels: at the base, in the middle, and in the apical part. Samples were then mounted on glass slides, polished up to a thickness of ~70 microns and finally analyzed with Leica DMLP and Nikon Optiphot2-pol microscopes. The type of the microstructure, the density and type of vascular canals, the amount of remodeling, the number of Lines of Arrested Growth (LAGs) and the presence or absence of an External Fundamental System (EFS) are the proxies used in this study to evaluate the ontogenetic stage of the sampled skeletal elements. The definition of the type of arrangement of the vascular canals was based on the orientation of the main axis. LAGs were identified and counted when the arrest in bone deposition was visible at different magnifications and when the interruption was continuous along the slide. When two or more LAGs were tightly spaced in the inner cortex, these were considered as annuli and counted as a single year.



Figure 1: MSNVE 3714, *Ouranosaurus nigeriensis*. The mounted specimen as exhibited today at the Museo di Storia Naturale of Venice. As for scale, the right femur is 920 mm long.

RESULTS

Historical background of the Venice specimen

The historical background of the Venice specimen is quite complicated. Between 1965 and 1972, five French paleontological expeditions searched for dinosaurs in the Gadoufaoua area of Sahara Desert in Niger (Taquet, 1976). The first one took place in January-February 1965; eight iguanodontian specimens were identified in the site named "niveau des Innocents" and located east of the Emechedoui wells. Two further iguanodontian skeletons, acronymized GDF 300 and GDF 381, were found 7 km south-east of Elrhaz in the Camp des Deux Arbres locality (geographic coordinates 16° 42'/lat., 9° 20'/long). During the second expedition (February 25th-April 7th, 1966), GDF 300 (a nearly complete but disarticulated and scattered skeleton) and GDF 381 ("un squelette aux deux tiers complet [a skeleton two thirds complete]", p. 54) were collected. The following year, those specimens were carried to Paris for preparation and study. GDF 300 ~~would become later~~ became the holotype of *Ouranosaurus nigeriensis* (Taquet, 1976, p. 57). According to Taquet (1976, p. 14), the other specimen (GDF 381) is not described in Taquet (1976), so it cannot be the paratype, although the latter is reported as MNHN-GDF 381 at p. 58. In fact, the skeleton GDF 381 that was found in 1965 100 m from GDF 300 and collected in 1966 is ~~indicated-referred to~~ as ~~the~~ "*Iguanodontidé trapu* [ponderous Iguanodontid]" ~~at p. 54 by Taquet (1976)~~ and ~~would become later subsequently became~~ the holotype of *Lurdusaurus arenatus* (see Taquet and Russell 1999), ~~as~~ ~~This was~~ confirmed by P. Taquet (pers. comm. to FMDV, 2012). However, Taquet and Russell (1999) contributed to increase the confusion, ~~because they labelled indicating~~ the holotype of *Lurdusaurus arenatus* ~~(which is described as a nearly complete skeleton found in 1965; p. 86) with the acronym as~~ MNHN GDF 1700 and referring an isolated right coracoid of *L. arenatus* (probably found in 1970, see Taquet, 1976, p. 54) as GDF 381. ~~According to Taquet and Russell (1999), the material of *Lurdusaurus arenatus* is at the MNHN.~~

Commented [PG2]: This sentence is particularly cryptic ! I really do not understand what you mean

Commented [PG3]: Of course !

184 The third expedition in 1969 found some dinosaur material at the In Gall locality (actually outside
 185 the Gadoufaoua area), but no *Ouranosaurus* is reported from there.
 186 During the fourth expedition (January 5th-March 23rd, 1970), a skeleton of *Ouranosaurus*
 187 *nigeriensis* without ~~the its~~ skull, but in better state of articulation than GDF 300, was discovered 4
 188 km south of the "niveau des Innocents" at the margin of the landing strip built by the CEA, (p. 58);
 189 apparently, it also received the field number GDF 381 (see Taquet, 1976, pl. IX, fig. 2). Other
 190 vertebrate fossils were collected there, including postcranial elements of the giant crocodile
 191 *Sarcosuchus imperator*. ~~Notice It should benoticed~~ that the paratype of *O. nigeriensis* ("un squelette
 192 presque complet auquel manque le crane" ["a nearly complete skeleton lacking the skull]) is said to
 193 come from "4 km Sud du niveau des Innocents, Bordure Est terrain d'aviation, lat. 16°26', long. 9°8'
 194 " ["4 km south of the niveau des Innocents, eastern margin of the airfield, 16°26'/lat., 9°8'/long.]
 195 (Taquet, 1976, p. 58) and ~~has received~~ the number GDF 381-MNHN (Taquet, 1976, p. 58). So, the
 196 specimen discovered during the fourth French expedition is the paratype. Much confusion is caused
 197 ~~by the fact that because~~ Taquet (1976, p. 15) does not mention this discovery ~~when he presenteding~~
 198 the results of the 1970 expedition.
 199 During the fifth expedition (January 5th - February 25th, 1972), the *Ouranosaurus* found in 1970
 200 (GDF 381) was excavated and brought to Paris (Taquet, 1976, p. 15 and 60). Apparently, this is the
 201 third and last ornithopod skeleton excavated and brought to France by French expeditions, together
 202 with GDF 300 and the GDF 381 found in 1965 and collected in 1966.
 203 In 1971, Giancarlo Ligabue and Cino Boccazzi knew about the Gadoufaoua locality during a travel
 204 across the Sahara desert (Ligabue et al., 1972). Ligabue led a first Italian expedition (February 3rd-
 205 22nd, 1972; at the same time as the fifth French expedition), actually a prospection in order to
 206 establish the basis for a future expedition (Ligabue et al., 1972; Boccardi & Boccazzi, 1978). This
 207 occurred the following year (November 4th - December 11th, 1973) and was an Italian-French
 208 expedition led by Giancarlo Ligabue and Philippe Taquet. A field report and a list of the excavated
 209 material was published in Ligabue & Rossi-Osmida (1975); the list included "1 [sic] *Ouranosaurus*
 210 *nigeriensis*" (p. 80). ~~According to Per~~ Rossi-Osmida (2005), all ~~the~~ fossils collected during the
 211 Italian expeditions were brought to the MNHM of Paris where they were prepared, restored and
 212 casted.
 213 In the formal description of the new species *Ouranosaurus nigeriensis*, Taquet (1976, p. 58)
 214 indicated GDF 300 as the holotype, " GDF 381- MNHN " as the paratype, and GDF 301 and GDF
 215 302, ~~respectively~~ a large coracoid and a femur, ~~respectively~~, as referred material. Despite being
 216 reported as a practically complete skeleton missing just the skull (p. 58), only the elements of the
 217 paratype that are not preserved in the holotype were described by Taquet (1976); a description or a
 218 list of the bones preserved in the paratype was never published. The holotype was brought back to
 219 Niger after ~~the its~~ study (Taquet, 1976) and ~~it results to be~~ exhibited at the National Museum of
 220 Niger in Niamey (Taquet, 1976, pl. IX, fig. 1); the MNHN ~~has only kept~~ a plaster copy (Currie &
 221 Padian, 1997, p. 369; A. McDonald, pers. comm. to FB, 2011). No further reference to the
 222 paratype is made in the literature, which according to the acronym used by Taquet (1976) should be
 223 at the MNHN. ~~It is Wworthy~~ of note; ~~that Taquet (1976) made~~ no mention ~~to of~~ the 1973 Italian-
 224 French expedition and ~~of the Ouranosaurus specimen said to have been collected during this field~~
 225 ~~campaign in that occasion is made in Taquet (1976).~~
 226 As ~~anticipated above already mentioned~~, Giancarlo Ligabue donated a nearly complete skeleton of
 227 *O. nigeriensis* (the skull and lower jaw were missing and replaced by copies) to the city of Venice
 228 after preparation at MNHN. It was exhibited to the public in 1975 in a room of the MSNVE along
 229 with other vertebrate specimens (including a complete skull of the crocodyliform *Sarcosuchus*
 230 *imperator*) supposed to have been collected during the 1973 expedition. ~~Since that date, it has been~~
 231 ~~exhibited to the public in the Museum.~~ At the end of the ~~1990's~~ of the last century, the skeleton was
 232 restored and remounted.

The paratype GDF 381 - MNHN apparently disappeared, as well as the referred specimens GDF 301 and 302. According to A. McDonald (pers. comm., to FB, 2011), and Currie & Padian (1997, p. 369), the MNHM has only a cast of the holotype. According to Currie & Padian (1997, p. 369), the only original specimen of *O. nigeriensis* other than the holotype is the Venice specimen, indirectly confirming that it is the paratype.

In order to clarify ~~once for ever~~ this issue, we asked P. Taquet whether the specimen exhibited at the MSNVE is the paratype of *O. nigeriensis* (GDF 381 - MNHN in Taquet, 1976) discovered in 1970 and collected in 1972 (as suggested by the above reconstruction, but never expressed explicitly) or another one. He confirmed that it is the paratype and that the missing bones were casted from the holotype (P. Taquet, 2012, pers. comm. to FMDV and FB). He also told us that he mapped the paratype bones in the field and ~~that~~ the map is ~~still kept~~ at the MNHN.

The map of the Venice specimen "pro parte"

R. Allain sent us a copy of ~~this~~ field map ~~that P. Taquet told us to refer to the excavation of the paratype in 1972~~. It is labeled "Ouranosaurus nig - Camp aviation - 1970 - (specimen musée Venice pro parte)" ["Ouranosaurus nig[er]iensis] - Airfield - 1970 - (specimen Venice Museum pro parte)"]. ~~Unfortunately, the map that we received is divided into eight sheets distributed in two pdf files, with incomplete information about how to assemble them. Each bone in the sheets is identified by a number, probably to identify the elements to recombine the skeleton once in the laboratory.~~

One main sheet is the scan of part of the original map made of brownish cardboard with the drawing of an articulated dorsal and sacral segment of a vertebral column adjacent to a slightly displaced proximal part of the caudal segment and some limb and girdles elements. It is evidently the map of the specimen pictured in Taquet, 1976, pl. IX, fig. 2; 1998, fig. 12) and identified as GDF 381, i.e. the paratype of *O. nigeriensis*.

A second sheet, also a scan of the original brownish cardboard, nearly totally overlaps to this one.

A third sheet is the total drawing of the caudal vertebral column segment, overlapping the main sheet; however, it is a poor-quality scan of the photocopy of the original map (the latter is no more available).

A fourth sheet is the map of the cervical segment of the vertebral column and some girdle and limb elements; it does not overlap the main sheet, but comparison with the images in Taquet (1976, pl. IX, fig. 2; 1998, fig. 12) allows to reliably connect the two sheets. We assembled these sheets (Fig. 2); we consider the resulting map as that of the main cluster of bones (that we name cluster 1) of the map of the Venice specimen "pro parte". It includes the bones numbered from 1 to 137, belonging to a semi-articulated skeleton exposing its right side. Numbers 96 (possibly a bone near the ulna) and 97 (fragment of ilium) and relative elements are not in the figure, but they are mentioned in an handwritten note on the map. Also a distal caudal vertebra was not numbered and is not drawn in the map, but it is mentioned in an handwritten note.

Two further sheets are scans of the photocopies of parts of the original map; they represent partly disarticulated elements from the pelvic region with part of the tail and some limb elements of an ornithopod skeleton (Fig. 2). A mark along the margins of the sheets allows joining them, but there is no indication on their spatial relationships with the other sheets. The sheets contain many handwritten notes (in French, of course), but the poor quality of the photocopy/scan does not allow us to understand many of them. Possibly, they give info about the spatial relationships of the sheets, but the fact that also people at the MNHN are unable to assemble the whole map (R. Allain, 2016, pers. comm. to FMDV) suggests that this is not the case. We consider this as the cluster 2 of the map of the Venice specimen "pro parte". As the photos in Taquet (1976, pl. IX, fig. 2) and Taquet (1998, fig. 12) show that other bones occurs "dorsally" respect to the main cluster, we tentatively located there the second cluster, but distance between the two clusters and the orientation of the second could be inaccurate. This cluster includes bones numbered 200, 202-209, 212, 214-231, 233-

Commented [PG4]: Bis repetita placent !

256 and 258-286. Numbers 210-211, 213 and 257 are in another sheet (see below), while the remaining are apparently missing, supposing a progressive numeration of the specimens. Also numbers from 138 to 200 are not present in the sheets that we receive from the MNHN, suggesting that one or more sheets are missing or that the two clusters are actually not related. The seventh sheet is also a poor-quality scan of the photocopy of part of the original map and represents only four bones (numbers 210-11, 213 and 257), including a complete ilium. Also in this case, its spatial relationships with the clusters 1 and 2 are unknown. However, the numeration indicates that this sheet is related to those of the cluster 2 and the location of the numbers in this cluster suggests that its position is possibly that in Fig. 2. Finally, one sheet reports only the hand writing "Ouranosaurus nig - Camp aviation - 1970 - (specimen musée Venice pro parte)".

We assembled the sheets, excluded the last one, remarking the original drawing line to obtain a line with a consistent and appreciable width (Fig. 2). The poor quality of the scans of the photocopies sometimes prevented the identification of portions of the original lines. We do not know whether they were drawn or not; they simply cannot be seen or they were originally missing in the map. In those cases, we avoided interpretations and left the lines interrupted. We also typewrote the original field identification of the bones by P. Taquet reported in the map, translated into English and sometimes abbreviated; when different from our identification or dubious, we report them in dark gray colour instead of black. Of course, field identification of the skeletal elements could be wrong and the mistake recognized only once the bone is freed from the enclosing rock. Some of the handwritten notes translated into English are also reported typewritten in dark gray. Finally, the sheets of the two clusters and the single sheet with the four bones do not seem to be at the same scale (lengths of some skeletal elements are reported, anyway).

This map confirms once again that the paratype is MSNVE 3714 (although "pro parte") and shows that the generic and specific name was created at least four years before its publication. The presence of a total of three pubes and possibly three ilia, scapulae and ulnae, as well as a duplication of segments of the caudal vertebral column indicates that at least two individuals are represented in the assembled map and cluster 1 and 2 belong to distinct skeletons.

The correspondence of the bones reported in the map and those occurring in the mounted skeleton is checked below; the implications of the word "pro parte" are discussed in the section Discussion.

Commented [PG5]: Sorry, but this part of the manuscript, like it is written, is not really interesting ! If the photocopies provided by R. Allain are of poor quality, please consult the original documents in Paris (1hour and a half from Brussels by Thalys).





Figure 2: *Ouranosaurus nigeriensis*, the field map of the remains partly used to mount MSNVE 3714. The map was drawn by P. Taquet and refers to the specimen found in 1970 ("*Ouranosaurus nigeriensis*" - Airfield - 1970 - (specimen Venice Museum pro parte)). The map is made of some assembled sheets (see text for the explanation); the three resulting sheet are not to the same scale, so scale bar is not reported. It was redrawn following the original lines as much as it was possible. Some original handwritten notes have been translated into English and typewritten in dark gray, as well as the lines originally added by Taquet to cancel the wrongly drawn elements. Dark gray names of the bones are the original identifications when their correctness is dubious or not testable; black abbreviations are ours and are partly our bone identification and partly unambiguous original identifications. "Near ulna" and "fragment of ulna" is an handwritten note that refers to collected elements numbered 96 and 97, which were not drawn in the map. Abbreviations: ca, calcaneum; co, coracoid; fe, femur; fi, fibula; h, humerus; il, ilium; mc, metacarpal; mt, metatarsal; ph, manual phalanx; pph, pedal phalanx; pu, pubis; ra, radius; sv, sacral vertebra; sc, scapula; st, sternal plate; ti, tibia; u, ulna . When the elements are reported as left in the original map, they are in brackets.

SYSTEMATIC PALEONTOLOGY

331
332 Dinosauria Owen, 1842
333 Ornithischia Seeley, 1887
334 Ornithopoda Marsh, 1881
335 Iguanodontia Dollo, 1888
336 Ankylopollexia Sereno, 1986
337 Styracosterna Sereno, 1986
338 Hadrosauriformes Sereno, 1997 *sensu* McDonald 2010
339 Hadrosauroida Cope 1869 *sensu* Sereno, 1986
340
341 *Ouranosaurus nigeriensis* Taquet, 1976
342
343 Note: the name *Ouranosaurus nigeriensis* was firstly published by Taquet in Ligabue & Rossi-
344 Osmda (1975, p. 41), without a formal description.
345
346 **Holotype:** GDF 300, a nearly complete skeleton, lacking the left maxilla; the right lacrimal; the
347 right quadratojugal; the stapes; the articulars; the dorsal vertebrae 1 and ?14; the centrum of caudal
348 vertebra 1 and caudals 25-26 and 30-31; most of the distal elements of the tail and some distal
349 chevrons; both femora (only the distal condylar end of one of them was found); the left tibia; the
350 left astragalus and calcaneum; the left metatarsals; and eight pedal phalanges. The skeletal elements
351 in situ were scattered on a 15 m² surface. The specimen is exhibited at the Musée National du
352 Niger, Niamey.
353
354 **Paratype:** GDF 381- MNHN (MSNVE 3714, "pro parte", see below), partial skeleton without
355 skull, but [preserving-with](#) the vertebral column in fairly good conditions of articulation and
356 probably missing only the atlas and the distal segment of the tail.
357
358 **Referred material:** GDF 301, large coracoid, and GDF 302, femur (present location unknown;
359 GDF 302 possibly added to MSNVE 3714, see below).
360
361 **Horizon and Locality:** level GAD 5, upper part of the Elrhaz Formation, Tégama Series, Aptian,
362 Aptian-Albian, or possibly Barremian, Early Cretaceous. All specimens are from the Gadoufaoua
363 area of Niger. The holotype comes from the Camp des Deux Arbres locality, 7 km south east of
364 Elrhaz, 16°42' lat. 9°20' long; the paratype was found 4 km south of the Niveau des Innocents
365 locality, along the eastern border of the airfield, 16°26' lat, 09°08' long. The exact locality for GDF
366 301 and GDF 302 is unknown, but probably it is the same as the holotype.
367
368 **Emended diagnosis:** Basal hadrosauroid dinosaur with the following autapomorphies: thickened,
369 paired domes on nasals, so that nasals extend further dorsally than frontals; dorsal 'sail' made of
370 extremely tall neural spines of the dorsal, sacral and proximal caudal vertebrae (up to 7 times the
371 height of the centrum in the proximal mid-dorsal vertebrae), with a sinusoidal outline (lower peak in
372 the sacral segment); tallest neural spines of dorsal vertebrae flare apically (i.e., the cranial and
373 caudal margins of the spine are not parallel but they diverge regularly basoapically).
374
375 **Phylogenetic relationships of *Ouranosaurus nigeriensis***
376 There is no agreement on the phylogenetic relationships of *Ouranosaurus nigeriensis* (e.g., Sereno,
377 1986; Norman, 2004, 2015; McDonald et al., 2010a, b, 2012). On non-cladistic bases, Taquet (1976)
378 considered *Ouranosaurus* as a derived "iguanodontid" that is closely related to *Probactrosaurus*
379 *gobiensis*, although he recognized some derived features shared with the hadrosaurids. In his

380 phylogeny of Ornithischia, Sereno (1986) found *Ouranosaurus* to be a member of the
 381 Hadrosauroidea and sister taxon of the Hadrosauridae. According to Norman (2004), *Ouranosaurus*
 382 is more derived than *Iguanodon bernissartensis* and *Mantellisaurus atherfieldensis*, and basal to
 383 non-hadrosaurid iguanodontians (*P. gobiensis*, *Eolambia caroljonesa*, *Protophadros byrdi* and
 384 *Altirhinus kurzanovi*). In McDonald et al.'s (2012) analysis, *Ouranosaurus* falls within the
 385 Hadrosauroidea between a politomy of *Mantellisaurus atherfieldensis* and *Hypselospinus fittoni*
 386 (less derived) and *Altirhinus kurzanovi* (more derived) (Fig. 3A). Thus, it is more derived than
 387 *Iguanodon bernissartensis*. In the latest published phylogeny by Norman (2015), it occurs in a more
 388 basal position (Fig. 3B) as the sister taxon of the clade 'Iguanodontoids' + Hadrosauriformes. The
 389 different phylogenetic affinities of the taxon found by the different authors are probably due to the
 390 presence of a mix of primitive (e.g., presence of palpebral bone, large and coarsely denticulated
 391 dentary teeth with two primary ridges, conical ungual phalanx on manual digit I, and double fibular
 392 process of the tibia) and derived (e.g., anterior end of the premaxilla expanded transversely; oral
 393 margin of the premaxilla reflected dorsally; long rostral diastema in the dentary; and greatly
 394 dorsoventrally expanded distal portion of the prepubic plate) characters.

Commented [PG6]: I am not sure that it is necessary to keep this paragraph... you do not provide new information.



396
 397
 398 **Figure 3: Relationships of *Ouranosaurus nigeriensis*.** Cladograms from McDonald et al. (2012;
 399 Adam consensus tree) (A) and Norman (2015) (B), redrawn. The basal part of the tree is not
 400 reported in both cases. Legend: 1 = Hadrosauriformes; 2 = Hadrosauroidea; 3 = Styracosterna; 4 =
 401 "Iguanodontoids".

404 **Description of MSNVE 3714 and comparison with the holotype**
 405 MSNVE 3714 is a partial skeleton lacking the whole skull and mandible; the ossified elements of
 406 the hyoid apparatus; the atlas; all the cervical ribs; right dorsal ribs 1, 2, 4 - 10, 13 and 15; left
 407 dorsal ribs 6, 11, 12 and 14; caudal vertebrae 27 to 31 and those posterior to caudal vertebra 38
 408 (vertebrae 27-31 and 39-43 are plaster copies); the right coracoid; the right carpus; the left
 409 metacarpals; the digit I (thumb spike), phalanges II-1 and 2, III-2, IV-1 and V-2 to 4 (ungual) of the
 410 left manus; phalanx II-3 (ungual), III-2 and 3 (ungual), IV-2, and V-3 and 4 (ungual) of right
 411 manus; the left femur; and the whole right pes (Fig. 4). All these elements, excluded those with the
 412 exception of the hyoid apparatus, the atlas and the cervical ribs, have been reconstructed. Most of

the other bones have also been partly reconstructed and restored (Fig. 4). Elements of the right side are more weathered than those of the left side, because the skeleton lied mostly on the left side (as it is shown by the map). In this section, the skeletal elements MSNVE 3714 are described and compared with those reported in the map as well as with those preserved in the holotype.



Figure 4: MSNVE 3714, *Ouranosaurus nigeriensis*, original and reconstructed parts in the right (A) and left (B) views. The reconstructed parts are in red.

AXIAL SKELETON

The axial skeleton of MSNVE 3714 is composed of 77 vertebrae, but 10 caudals are totally reconstructed with plaster, so only 66 are actually preserved (Fig. 5A-D). They are all more or less restored, ~~in particular~~ particularly the base of the neural arch, the zygapophyses and the transverse processes of the dorsals and caudals ~~are mostly reconstructed~~. Curiously, ~~the holotype also preserves remains of~~ 66 vertebrae ~~are also preserved in the holotype~~, but eight elements occurring within those preserved are missing according to Taquet (1976), so the original vertebral column would be at least 74 (but see below). ~~As Because~~ the axial skeleton of the paratype is in a better state of anatomical articulation respect to the holotype (see Fig. 2), it is a better reference for the vertebral count ~~of in~~ *Ouranosaurus nigeriensis*.

Commented [PG7]: I really do not understand what you mean ! Please rephrase.




Figure 5: MSNVE 3714 and holotype, vertebrae. MSNVE 3714, the cervical series (A); the dorsal series (B); the sacrum (C); and the caudal series (D). Holotype, the cervical series (E); the dorsal series (F); the sacrum (G); and the caudal series (H). Numbers are progressive within each series. White vertebrae are those of the mount that are totally reconstructed. E-G are redrawn from Taquet (1976). Scale bar equals 50 cm.

Cervical vertebrae. The axis and the following ten presacral vertebrae are preserved, while the atlas is missing (Fig. 5A). The presacral vertebra 11 has the parapophysis that appears to be cut by the neurocentral suture (Fig. 6A), so it is a cervical and not the first dorsal, according to the definition of first dorsal vertebra by Norman (1986). The presacral vertebra 12 in the mounted skeleton has a relatively tall neural spine and a parapophysis that seems to be at the very base of the neural arch (Fig. 6B), so it is the first dorsal vertebra. Therefore, it can be confidently estimated that MSNVE 3714 had 11 cervical vertebrae. The holotype preserves 11. Eleven cervical vertebrae are

Commented [PG8]: Please add GDF 300 in all the the figures

also preserved in the holotype (presacrals 1 to 11; Fig. 5E), but the neck was totally disarticulated in situ and presacral vertebrae 12 to 14 are not preserved according to Taquet (1976). The map of the paratype shows nine articulated cervicals (probably cervicals 2-10), a further centrum that is slightly separated from the adjacent centra (but without the possibility of the presence of another vertebra missing in between) and one that is articulated with a [string series](#) of 13 dorsal vertebrae with very tall neural spines. ~~The first centrum is probably the cervical 11, while the second is the first dorsal.~~ *Iguanodon bernissartensis*, *Equijubus normani* and *Jinzhousaurus yiangi* also have 11 cervicals, *Mantellisaurus atherfieldensis* 10 or 11 (Norman, 2004; Wang et al., 2010; McEDonald et al., 2014). So, a cervical count of 11 is supported for *Ouranosaurus*.




Figure 6: MSNVE 3714, cervical-dorsal transition. The last (11) cervical (A); the first dorsal (B). Both are in left lateral view. Reconstructed parts are in dark gray color. Abbreviations: dia, diapophysis; ncs, neurocentral suture; ns, neural spine; par, parapophysis; poz, postzygapophysis; prz, prezygapophysis. Scale bar equals 10 cm.

Formatted: French (France)

In the axis (Fig. 7A1-4), the odontoid process, the prezygapophyses, the cranial portion of the neural arch above the pedicels and parts of the latter, and parts of the centrum were missing and have been reconstructed (prezygapophyses were not reconstructed and lack). The diapophyses are small and knob-like like those of the holotype, but they are mostly reconstructed. The neural spine is low and sub-triangular in lateral outline, with a rounded dorsal margin. A broad circular depression occurs in the middle of both right and left sides of the spine. The neural spine in the axis of the holotype has a different M-like lateral outline (i.e., the dorsal margin is concave in the middle) and seems to lack the lateral depressions (Taquet, 1976, fig. 37B). Cervicals 3-11 (Figs. 5A, 7B and 7C) do not appear to differ significantly from those of the holotype and those of advanced iguanodontians in general (e.g., Norman, 1980; 1986). However, most of their processes are reconstructed. For example, the long diapophyses and parapophyses are totally reconstructed in cervicals 3 and 4; the long postzygapophyses are both reconstructed in cervical 3, while only the right one is original in cervical 4. In the cervical 5, the long processes bearing the prezygapophyses and diapophyses are made of resin. Small fragments of the neural spine are preserved in the cervicals 8-10. So, the centra are practically all what we have of them for comparison. They are deeply opisthocelous and ~~bea~~ have a median, longitudinal and blunt keel ventrally.




Figure 7: MSNVE 3714, cervical vertebrae. Axis in cranial (A1), right lateral (A2), left lateral (A3), and caudal view (A4); cervical vertebra 3 in cranial (B1), right lateral (B2), left lateral (B3), and caudal (B4) views; cervical vertebra 11 in cranial (C1), right lateral (C2), left lateral (C3), and caudal (C4) views. Abbreviations: dia, diapophysis; od, odontoid process; ns, neural spine; par, parapophysis; poz, postzygapophysis; prz, prezygapophysis. Scale bar equals 10 cm.

Dorsal vertebrae. MSNVE 3714 has 17 dorsals (Fig. 5B). As said above, a string series of 14 dorsals in relative anatomical connection is identifiable in the map of the in situ paratype (Fig. 2). A further centrum seems to occur caudal to the last vertebra of the string series and displaced ventrally. Thus, MSNVE 3714 has at least two dorsal vertebrae more than the mapped paratype. There are two explanations for this: 1) the two additional vertebrae could have been preserved there, but they were not recognized before preparation or at least one of them could have been covered by a broad element present in that area (plausibly an ilium); 2) two vertebrae from another specimen were added to the paratype material because the holotype was supposed to have 17 dorsals (Fig. 5F). The first explanation seems to be the less probable because that is also the position of the sacrum, which is only partly exposed, as it is composed of six vertebrae but only three are drawn in the map (one is cancelled).

Commented [PG9]: You said 13 before . Please be consistent

Commented [PG10]: Again, I do not understand...

502 In the dorsals of MSNVE 3714, the peduncles of the neural arches, parapophyses, transverse
 503 processes and relative diapophyses are all reconstructed (presumably, taking as reference for
 504 proportions and morphology those of the holotype). Also the neural spines, which are the most
 505 important feature of those skeletal elements, are heavily restored (Fig. 8).
 506 The centra of ~~the~~ dorsals 1 and 2 are opisthocelous; from the dorsal vertebra 3 ~~onwards~~, the
 507 centrum becomes slightly amphicoelous to amphiplatyan. ~~The C~~centra range ~~from~~ 87 (vertebra 12)
 508 to 112 mm (vertebra 17) in length and are slightly longer than high (elongation is ~~more better~~
 509 marked in ~~the~~ dorsal 17; Fig. 9C3). The centrum of dorsal 1 has a ventral longitudinal keel and sub-
 510 circular articular surfaces, like those of the cervicals (Fig. 9A); ~~the centrum and~~ is smaller than that
 511 of the last cervical (Fig. 6). The relatively small parapophyses are located just above the
 512 neurocentral suture (Fig. 6B). In ventral view, all other dorsal centra are spool-shaped with a keeled
 513 ventral margin (Fig. 9A3 and B3); only centrum 17 ~~seems to apparently~~ lacks ~~a the~~ keel (Fig. 9C3).
 514 ~~AsBecause~~ the sacral vertebrae have a faint ventral keel or lack it, dorsal 17 could actually be a
 515 dorsosacral. The articular facets of the centra 2 to 16 are higher than wide, while it is the reverse in
 516 centrum 17.
 517 The morphology of the tall neural spines shows a certain variability (Fig. 8). The spine of dorsal 1 is
 518 straight, inclined caudally (60°) and it slightly tapers apically in its basal part, while the apical half
 519 has parallel caudocranial margins and does not flare apically (Figs. 6B and 9A1-2); it is only 1.41
 520 times the height of its centrum. The spine of dorsal 2 is incomplete apically. The preserved part is
 521 2.7 times the height of the centrum; as reconstructed, it is about four times the height of the
 522 centrum. It is narrow craniocaudally, slightly sloping caudally (about 80°) and slightly recurved; as
 523 the cranial and caudal margins are parallel, ~~probably~~ it was ~~probably~~ not expanded apically (so, the
 524 reconstruction is faithful). The spine of dorsal 3 is taller and craniocaudally longer than that of
 525 dorsal 2, but it is also incomplete apically. It is straight with nearly parallel craniocaudal margins
 526 and it slightly slopes caudally (about 83°). The apex and part of the apical ~~tract~~ of the caudal margin
 527 of the spine are reconstructed, but ~~probably~~ the spine was ~~probably~~ not ~~sensibly~~ expanded
 528 craniocaudally. Dorsal 4 has a neural spine that is taller and craniocaudally longer than that of
 529 dorsal 3; it is incomplete apically too. It is straight, with its minimum craniocaudal length just
 530 below the mid-shaft; ~~its~~ cranial and caudal margins diverge above the point of minimum
 531 craniocaudal length, so the apex was probably slightly expanded. Unlike the preceding vertebra, the
 532 spine slopes cranially (about 85°). Dorsal 5 has a spine that is nearly complete apically and is
 533 apparently taller than that of the preceding vertebra. It is straight and only slightly sloping cranially.
 534 Its minimum craniocaudal length occurs in the lower third, but it is unclear whether this is a real
 535 feature or an artifact of preparation. The cranial and caudal margins diverge above that point, so the
 536 spine flares sensibly toward its apex. The latter is ~~not squared~~. The spine of dorsal 6 is straight ~~and~~
 537 vertical. It is incomplete apically, but it is anyway at least as tall as the preceding one. The cranial
 538 and caudal margins diverge above the lower third, so the spine flares sensibly toward the apex.
 539 Unlike that of the preceding vertebra, the spine of dorsal 7 is recurved cranially. Like spine 6, it
 540 flares apically. As the apical portion is partly reconstructed, its squared outline is just hypothetical.
 541 The cranial curvature cannot be a real feature, because it would prevent the zygapophyseal and
 542 central articulation with the preceding vertebra, unless the spines of the two vertebrae were
 543 overlapping laterally. The spine of dorsal 8 is unlike those of the preceding and following vertebrae.
 544 It appears to be craniocaudally narrower and is slightly sloping caudally. It also flares above its
 545 basal third. The apex is reconstructed, so its squared outline is just hypothetical and its total height
 546 is unknown. The spine of dorsal 9 is slightly curved basally, but the rest is straight vertical (Fig.
 547 9B1-2). Flaring starts in the basal part of the spine. If ~~the~~ reconstruction is ~~faithful correct~~ ~~(a-part-of~~
 548 the apex is ~~partly~~ preserved but a basal portion is reconstructed), the spine is the tallest (it is seven
 549 times the height of its centrum). The apex is partly reconstructed, so its squared outline is just
 550 hypothetical. The curvature of the basal part of the spine is more marked in vertebra 10 than in the

Commented [PG11]: ????

Commented [PG12]: Is rounded ???

preceding [one vertebra](#). This spine also flares starting from the basal portion; its apical third is mostly reconstructed, so nothing can be said about its real outline. The whole spine of vertebra 11 seems to be slightly recurved, but its basal part is reconstructed, so this feature could be an artifact. The preserved portion of the apical part shows that this spine was lower than spine 9. Spine 12 seems to be arched basally and straight from mid-shaft on. Its apical portion is mostly reconstructed, so its real height and the shape of its apex are unknown. The following neural spines 13-17 are all arched (less in the spine 15, which is poorly preserved) and flares apically like the preceding ones, although proportionally less than in the mid-dorsals (Fig. 9C1-2). Their craniocaudal length decreases slightly moving caudally. Their height decreases markedly; spine 14 is just slightly lower than spine 11, but the decrease is marked in the following vertebrae 15-17. The basal part of the spine in vertebrae 5, 7, 8, and 16-17 shows a cranial bump that is made by the cranially expanded prespinal lamina (Figs. 8 and 9C1) and is observed also in the distal dorsals of the holotype (Fig. 5F). All spines are narrow transversely and they do not thicken apically.




Figure 8: MSNVE 3714, dorsal vertebrae in lateral view. Above, the left side; below, the right side. The parts of the neural spine that have been reconstructed or just covered by resin are highlighted in white. Reconstructed parts of the centrum, transverse processes, zygapophyses and pedicels of the neural arch are not highlighted. Numbers are progressive. Scale bar equals 10 cm.

[In the text](#), Taquet (1976, p. 109) [saidys](#) that 13 dorsals are preserved in the holotype and that the cervical series is separated from the first preserved dorsal by a gap that could be filled by other dorsals or just be caused by displacement. He [opteds](#) for the first hypothesis, saying that the first four dorsals are probably missing. However, [he figures](#) only three dorsal vertebrae [as are](#) missing (the presacral vertebrae 12 to 14) in [his](#) figure 38 (here Fig. 5F). Furthermore, Taquet (1976, p. 109)

says stated that the dorsals following the gap are not displaced (i.e., they are in anatomical connection), but figure 9 of that paper shows this is true only for a segment of just nine mid-dorsal vertebrae. So, it is unclear how he established that four dorsals are missing and that the total count of the dorsals is 17 (see Taquet, 1976; figs 38 and 40). The much better articulated vertebral column of the paratype GDF 381 (MSNVE 3714 pro parte) shows that Taquet (1976) is wrong in his reconstruction of the holotype dorsal vertebral ~~series~~ ~~ing~~. Only one of the supposedly missing proximal dorsals of the holotype (see Fig. 5F) is present in MSNVE 3714; it corresponds to the first dorsal. The second dorsal of MSNVE 3714 corresponds to ~~the~~ vertebra 4 of the holotype (see Fig. 5F). Dorsal vertebra 5 of the holotype has a cranially sloping neural spine that would cause the crossing with the neural spine of the preceding vertebra when the two vertebrae are in anatomical articulation; furthermore, the spine tapers apically. There is no such a vertebra in MSNVE 3714: dorsal 3 is morphologically like vertebra 6 of the holotype. Dorsal 4 of MSNVE 3714 corresponds to dorsal 7 of the holotype in the relative height and slight cranial slope of the neural spine, but the latter has a paddle-like neural spine. In order to have the same number (17) of dorsal vertebrae as the holotype, MSNVE 3714 must consequently have more mid-posterior elements, as it can be appreciated by comparison of Figures 5B and F. In both cases, the first mid-dorsals tend to have straight vertical neural spines that are craniocaudally expanded apically, while the last mid-dorsals and distal dorsals have arched spines whose craniocaudally expansion decreases moving caudally. However, the number of those elements is different in the two specimens. It is evident that the spines of dorsals 7 and 8 in MSNVE 3714 are different from those of the contiguous vertebrae (Fig. 8); this suggests that those two vertebrae were added to maintain the count of 17 vertebrae reported for the holotype, which is also suggested by the vertebral count in the field map of the paratype. Probably the number of 17 dorsals was originally established because iguanodontians like *Mantellisaurus atherfieldensis* (at that time referred to *Iguanodon*) and *Iguanodon bernissartensis* have 17 or more dorsals (see Taquet, 1976, p. 111). However, the comparison between the holotype, the field map of the paratype and MSNVE 3714 suggests that *Ouranosaurus nigeriensis* had a shorter torso (possibly with 14 dorsals and one dorsosacral) and that the tallest neural spine is that of dorsal vertebra 7.

Some inconsistencies regarding the dorsal vertebrae are found in Taquet (1976). The dorsals 10-12 are reported to have the highest neural spines (p. 112), but the tallest is actually the spine of ~~the~~ dorsal 9 (like MSNVE 3714; see Figs. 5F and 8) and height decreases gradually in the following vertebrae, according to ~~the~~ figures 38 and 40 and measurements reported at pages 178-179. Vertebra 9 is the sixth preserved vertebra in the holotype and could be dorsal 7, if only the first dorsal is missing of the preceding dorsals. The highest neural spine is said to be 3.9 times the height of its centrum (p. 112), but it is actually nearly seven times according to figure 38 (7.11 according to measurements at p. 178). The centrum of the dorsal with the highest spine is reported to be 160 mm high, but it is less than 90 mm high in dorsal 8 according to the scale bar in figure 41 and that of dorsal 9 is 90 mm according to measurements at p. 178.



Figure 9: MSNVE 3714, dorsal vertebrae. Dorsal vertebra 1 in right lateral (A1), caudal (A2), and ventral view (A3); dorsal vertebra 9 in left lateral (B1), cranial (B2), and ventral view (B3) views; dorsal vertebra 17 in left lateral (C1), caudal (C2) and ventral (C3) views. Abbreviations: bpl, 'bump' of the prespinal lamina; kl, keel; par, parapophysis. Scale bars equal 10 cm.

Dorsal ribs. Only four out of 17 right dorsal ribs are not totally reconstructed and 12 out of 17 on the left side. Their state of preservation is anyway poor and large portions of the shafts are reconstructed in most of them. Practically, they are mostly shaft segments glued together. The *tuberculum* is just a small knob placed dorsolaterally to the *capitulum* forming with it an angle of about 90°. The cross-section of the proximal half of the rib is T-shaped. The map of the in situ paratype shows remains of at least 13-14 dorsal ribs.

Sacral vertebrae. Like the sacrum of the holotype, that of MSNVE 3714 (Figs. 5C, and 10) is composed of six fused vertebrae. The transverse processes of the vertebrae and the sacral ribs are completely reconstructed. A shallow longitudinal keel ~~occurs in~~ extends along the ventral surface of the sacral centra 1-2, becomes very faint in centra 3-4 and lacks in centra 5-6; centrum 6 has a nearly flat ventral side. The neural spines are straight, vertical, and only slightly craniocaudally

635 longer ~~apically at their apex than at their basal part~~. The spines of sacral vertebrae 1-3 are of similar
636 height; they increase in height from sacral 4 up to 6, which bears the tallest spine. Therefore, the
637 last two sacral spines form the beginning of the caudal hump of the 'sail'. ~~The S~~spines are regularly
638 separated ~~regularly~~, ~~except luded~~ the last one; ~~the distance between spines 5 and 6 is twice -whose~~
639 ~~gap from the preceding spine is twice~~ the distance between spines 4 and ~~spine~~ 5.
640 In the map of the in situ paratype (Fig. 2), only three neural spines can be seen in the sacral region
641 ~~of the sacrum~~; the drawing of a distal fourth one was cancelled as it were erroneously drawn.
642 Possibly, the other three sacral were ~~embedded in~~~~covered by~~ rock.
643 Apparently, the trend in neural spine height is the reverse in the holotype: height decreases from
644 sacral 1 to sacral 5 (the spine of sacral 6 is not preserved). Thus, there is an abrupt depression ~~with~~
645 marginal steps in the 'sail' of the holotype in correspondence of the sacrum (Fig. 5F-H). This
646 condition is probably unnatural and that ~~one~~ in MSNVE 3714, showing a smoother passage from
647 the dorsal to the sacral and from the sacral to the caudal spines (Figs 1 and 5B-D), ~~is perhaps~~
648 appears more reliable.
649

Commented [PG13]: I do not really understand.



Figure 10: MSNVE 3714, the sacrum. Left lateral view. Scale bars equal 10 cm.


Caudal vertebrae. The tail is composed of 43 caudal vertebrae, but five vertebrae (caudals 27 to 31) and the terminal [string series](#) of five vertebrae (caudals 39-43) are made of plaster (Fig. 5D). Thus, [there are 33](#) preserved vertebrae ~~are 33~~. The total caudal count was surely higher (see the caudal count of several dinosaur taxa in Hone, 2012), possibly as high as in the tail of *Iguanodon bernissartensis*, which is composed of 46 caudal vertebrae (Norman, 1980; but it is incomplete) or even much higher (the count is over 75 in TMP 98.58.01, an indeterminate hadrosaurid; FMDV and MF, pers. obs.). There are 20 proximal and 12 (17 including the five that are totally reconstructed) middle caudal vertebrae. The last preserved caudal (caudal 38) seems to be a distal element (see below).

663 As for the dorsals, 33 caudals are preserved also in the holotype, but two vertebrae between ~~the~~
 664 caudals 24 and 27 are missing as well as two between ~~the~~ caudals 29 and 32 (Taquet, 1976, p. 118;
 665 Fig. 5H). However, this reconstruction is hypothetical because the tail was partly disarticulated
 666 (Taquet 1976, fig. 9). The holotype has 14-15 proximal caudals (15 according to the text, 14
 667 according to fig. 44) and more than 12 (16 considering the hypothetical ones) mid-caudals. The last
 668 nine caudals appear to be distal ones (Taquet 1976, fig. 44); in facts, ~~the haemapophyseal~~ facets ~~for~~
 669 ~~the chevron occur~~ can be observed up to the posterior portion of caudal 31 according to the text
 670 (p.119), but actually caudal 31 is not preserved in fig. 44!
 671 According to the field map, the first 24 caudals of the paratype are articulated ~~and~~ with their relative
 672 chevrons; after a gap containing just an element identified as a phalanx in the handwritten note (but
 673 apparently it is ~~another further~~ centrum because of its size and position) and ~~in-line~~ aligned with the
 674 first segment, another segment of six vertebrae ~~occurs can be observed~~. ~~Another further~~ distal
 675 centrum is slightly displaced and one labeled as "without number" is not drawn. So, the vertebral
 676 count fits with that of MSNVE 3714. The five vertebrae that are totally reconstructed in MSNVE
 677 3714 correspond with those missing in the gap. Therefore, the count obtained from the
 678 paratype/MSNVE 3714 seems to be more reliable than that from the holotype.
 679 *Iguanodon bernissartensis* has 14 proximal, about 22-24 middle and at least 8-10 distal caudal
 680 vertebrae (Norman, 1980). *Mantellisaurus atherfieldensis* (IRSNB 1551; Norman, 1986) has 15
 681 proximal and at least 17 middle caudals (the tail is incomplete distally). So, *Ouranosaurus* has four
 682 or five more proximal caudals than *Iguanodon bernissartensis* and *Mantellisaurus atherfieldensis*
 683 and possibly a ~~more~~ caudally ~~prolonged longer~~ *M. caudifemoralis* (Persons & Currie 2011).
 684 MSNVE 3714 seems to have a comparatively low mid-caudal count. However, the last and only
 685 distal caudal element is probably the one labeled "+1 without n(umber)" in the map, which is not
 686 drawn, it was not articulated to the others vertebrae and probably was collected far away from the
 687 others. Thus, the actual mid-caudal count of the paratype/MSNVE 3714 "pro parte" could be
 688 higher. The mid-caudal counts of the Venice specimen, *Iguanodon bernissartensis* and
 689 *Mantellisaurus atherfieldensis* suggest that also the count of the mid-caudals in the holotype is
 690 wrong and probably higher than that reported in Taquet (1976).
 691 In MSNVE 3714, centra are slightly amphicoelous in the proximal and first middle caudals to
 692 become amphiplatyan caudally. The caudal surface of proximal and middle caudals is more squared
 693 than the rounded cranial one because of the presence of the facets for the hemapophysis (Figs.
 694 11A1, B1 and C2). The latter appear in the third caudal, so the first chevron occurs between the
 695 caudal 3 and 4. Centra are constricted in the middle and hourglass-like; they are shorter than tall up
 696 to vertebra 17 and shorter than broad up to vertebra 19 (Tab. 1). The centrum of the caudals 1 and 2
 697 has a longitudinal ventral keel, which is faintly developed in ~~the~~ caudal vertebra 3; the following
 698 centra seem also to be convex ventrally, but they are deformed, poorly preserved or reconstructed.
 699 From vertebra 11 up to vertebra 33, the ventral side of the centrum has a broad, median and
 700 longitudinal sulcus. This change in convexity seems to occur when the two ~~articular~~
 701 ~~haemapophyseal~~ facets ~~for the haemapophysis~~ are placed only caudally in the vertebra; before
 702 vertebra 11 the chevrons articulate on facets placed both on the cranial and the caudal ventral
 703 extremities of the centrum. This is related also to the relative increase in elongation of the centrum.
 704 As ~~anticipated mentioned~~ above, the last preserved vertebra (38) seems to lack the articular facets
 705 for the hemapophysis. This suggests that it is a distal caudal. A handwritten note in the field map of
 706 the paratype informs that the most distal vertebra of the sample is not figured in the map (Fig. 2);
 707 probably it was scattered ~~respect to the others~~ and possibly it was the only collected distal element,
 708 which was later attached to the last preserved mid-dorsal. Another possibility is that vertebra 38 is
 709 just one of the last mid-caudals and the articular facets for the ~~ha~~emapophysis were weathered
 710 away.

711 The lateral surface of the centrum near its articular facets is rough, with longitudinal grooves in
 712 some proximal and in all mid-caudal vertebrae (caudals 11-25; Fig. 12A), suggesting the presence
 713 of a cap of cartilage. A neurocentral suture is observed in proximal and middle caudal vertebrae
 714 (Fig. 12B-D) up to the caudal 25.
 715 The neural arches of vertebrae 32-38 are totally reconstructed.
 716 The pleurapophysis of the proximal caudals is flattened dorsoventrally and scarcely projecting
 717 laterally (Fig. 13). It occurs at the base of the neural arch on a ventral expansion of the pedicel
 718 overlapping the centrum laterally. It decreases in size along the series becoming knob-like; it
 719 disappears totally in the caudal vertebra 21 (Fig. 13C), but in caudals 19 and 20, it is just a small
 720 bump (Fig. 13B). It is rarely completely preserved, being often made partly or totally of resin.
 721 The prezygapophyses are reconstructed in all caudals except the right ones in vertebrae 17 and 21.
 722 The articular surfaces are oval and face dorsomedially; those of the postzygapophyses are also oval
 723 and face lateroventrally.
 724 Neural spines decrease gradually in height moving caudally (Fig. 5D). However, apical portions are
 725 reconstructed in spines 1, 3, 5 and 7, as well as segments of the shaft in many others. Spines are
 726 mostly spatulate-like in lateral view, with a slight craniocaudal apical expansion (Figs. 5D and 11).
 727 They are inclined caudally to a different degrees. For example, the spine of caudal vertebra 2 is only
 728 slightly sloping (77.2°), while those of vertebrae 7, 9 and 16 slope 52.7°, 58.4° and 48.6°,
 729 respectively (Fig. 5D). The proximal spines are generally straight (Fig. 11A), but spines 6, 10
 730 (Fig. 11B2-3) -12, 14-15 and those posterior to spine 17 (Fig. 11C1-2) are arched with concavity
 731 facing cranially (Fig. 5D). Non-harmonic variability in sloping may be a restoration bias, because
 732 the proximal portion of some neural spine was broken into several pieces that have been glued
 733 together and missing portions have been reconstructed. Also the basal arching of spines 3 and 4
 734 (which is not observed in preceding and following spines; Fig. 5D) could be a consequence of
 735 restoration.
 736 Neural spine inclination and morphology in the caudals of the holotype are more regular than in
 737 MSNVE 3714 (Taquet 1976, figs 40 and 43-44; Fig. 5H); also, the spines of vertebrae 1-4 are
 738 slightly arched backward (Fig. 5H), unlike those of the Venice specimen. Proximal caudals 2-7
 739 present a cranially projecting bump of the basal part of the prespinal lamina that occurs only in
 740 caudals 1-3 of the holotype.
 741 **Hæmapophyses** (Fig. 11E). There are 26 hæmapophyses (chevrons), but seven are totally
 742 reconstructed. Hæmapophyses 1-18, 20, and 23 are original, although they all present reconstructed
 743 parts; chevrons 19, 21-22 and 24 to the last one are totally artificial.
 744 The first hæmapophysis is located between caudals 3 and 4, like the holotype, while the last occurs
 745 between vertebrae 28 and 29, but it is artificial like the two vertebrae. Hæmapophyses are
 746 elongated and forked proximally into two pedicels bearing the articular facets. There are two
 747 articular facets per pedicel in the chevrons of the first proximal caudal vertebrae, as the pedicel
 748 articulates on two centra. The dorsoventral length of the hæmapophyses tends to decrease caudally.
 749 However, chevron 14 is shorter than chevron 15, both unbroken distally; possibly chevron 14 is in
 750 the wrong position and should be placed in a more distal position. The spines are craniocaudally
 751 narrow and laterally compressed. They appear to be spatulate-like, although many distal portions
 752 are damaged. The spine of the hæmapophysis 1 is straight, while those of the following elements
 753 up to hæmapophysis 5 are arched; only the pedicels of chevron 6 are preserveds-only-its-pedicels;
 754 chevrons 7 and 8 are only weakly arched; chevron 9 seems to be straight, but the proximal segment
 755 of the shaft is reconstructed; chevron 10 is nearly straight; only the distal end is slightly recurved
 756 backward in chevron 11; chevron 12 is straight; chevrons 13-17 are slightly recurved; the small last
 757 chevrons are poorly preserved and partly reconstructed.
 758 In the field map of the paratype, only the pedicels of the first two chevrons are drawn, between
 759 caudals 3 and 4 and 4 and 5, respectively. The following chevrons appear to be entire (with the


760 exception of chevron 5) up to chevron 17 that is followed by five vertebrae with incomplete or very
 761 small chevrons; no chevrons are associated with the last nine vertebrae.
 762 Morphology and sloping of the neural spines and chevrons in the tail of the Venice specimen are
 763 less regular and harmonic than in the holotype and, in general, in tetrapod tail skeletons. This is
 764 probably a cause of the breakage of the long and thin apophyses and ~~consu~~ subsequent restoration.
 765 However, the absence of ~~most~~the largest part of the chevrons 1 and 2 in the paratype map (if not
 766 due to their partial concealment into the rock) would suggest that some chevrons were replaced with
 767 material from an individual distinct from the paratype, possibly from the cluster 2 occurring near
 768 the paratype cluster (Fig. 2).
 769 In both basal and derived iguanodontians, the first chevrons (from just chevron 1 up to chevron 6)
 770 taper distally (e.g., *Tenontosaurus tilletti*, Forster 1990, fig. 5A; *Iguanodon bernissartensis*,
 771 Norman 1980, fig. 47; *Mantellisaurus atherfieldensis*, Norman 1986, fig. 39; *Xuwulong yueluni*,
 772 You et al 2011, fig. 2; *Tethyshadros insularis*, Dalla Vecchia 2009, fig. 1; *Kritosaurus*
 773 *incurvimanus*, Parks, 1920, pl. 1; *Brachylophosaurus canadensis*, Prieto-Marquez, 2001, fig. 52;
 774 *Corythosaurus casuarius*, Brown, 1916, fig. 2). ~~and~~ are more inclined caudally than the following
 775 chevrons and touch each other (*Iguanodon bernissartensis*, Norman 1980, fig. 47; *Xuwulong*
 776 *yueluni*, You et al 2011, fig. 2; *Tethyshadros insularis*, Dalla Vecchia 2009, fig.1; *Kritosaurus*
 777 *incurvimanus*, Parks, 1920, pl. 1; *Brachylophosaurus canadensis*, Prieto Marquez, 2001, fig. 52).
 778 This is not the case of both the holotype and MSNVE 3714 (see Figs 1 and 5H), suggesting that the
 779 tails of those skeletons were recomposed and mounted in ~~the a~~ wrong way.
 780 Unlike other iguanodontians, no ossified tendons are preserved with MSNVE 3714. According to
 781 Taquet (1976, p. 113), they were represented by a few fragmentary remains in the holotype and
 782 possibly by their traces on the neural spines of the distal dorsals. Thus, the characteristic lattice
 783 occurring laterally on the dorsal to proximal caudal vertebrae of the iguanodontians was at best
 784 scarcely developed in this high-spined taxon (contra Organ 2006b).

Commented [PG14]: Please do not forget the taphonomic processes. Preservation of the tendons requires exceptional fossilisation conditions ! I saw many hadrosaur tails without preserved tendons.



786
 787

788 **Figure 11: MSNVE 3714, caudal vertebrae.** Vertebra 6 in caudal (A1) and right lateral (A2)
789 views; vertebra 10 in cranial (B1), right lateral (B2), left lateral (B3) and ventral (B4) views;
790 vertebra 21 in left lateral (C1), caudal (C2) and ventral (C3) views; vertebra 35 in left lateral (D1)
791 and caudal (D2) views; hemapophysis 8 in left lateral (E1) and cranial (E2) views. Abbreviations:
792 af, articular facets of the hemapophysis; afh, articular facet for the hemapophysis; bpl, bump of the
793 prespinal lamina; ns, neural spine; pla, pleurapophysis; poz, postzygapophysis; prz,
794 prezygapophysis; vld, ventral longitudinal depression of the centrum. Scale bar equals 10 cm.
795



796
797
798 **Figure 12: Evidences of osteological immaturity in the caudal vertebrae of MSNVE 3714.**
799 Rough surface in vertebral centrum 24 (A); neurocentral suture in vertebra 8 (B); vertebra 10 (C);
800 and vertebra 21 (D). Vertebrae are figured in left lateral view. Arrows point to the grooved surface
801 in A and to the neurocentral suture in B-D. Scale bar equals 5 cm in A and 10 cm in B-D.
802




Figure 13: Proximal to mid-caudal transition in MSNVE 3714. Vertebra 18 (A); vertebra 20 (B); vertebra 21 (C). They are shown in dorsal view. Abbreviations: ns, neural spine; pla, pleurapophysis; prz, prezygapophysis. Scale bar equals 5 cm.

APPENDICULAR SKELETON

Coracoid (Fig. 14A). The right coracoid is a plaster copy, while the left one is original, and smaller than the right one (it is about 8% shorter), its size being only 125 in length and 130 mm in height. The left coracoid is a squared element with a convex, but partly damaged, dorsal margin and a concave ventral margin; the cranial margin is shallowly convex, whereas the middle part of the caudal margin between the humeral glenoid facet and the scapular sutural facet is concave (however, this zone is partly reconstructed; Fig. 14A3). The medial surface is shallowly concave, whereas the lateral surface is slightly convex. The caudodorsal part of the element is thickened, and bears supporting the rough and nearly triangular sutural surface for the scapula. The coracoid contribution to the humeral glenoid occurs isset in the caudoventral corner of the bone. The coracoid foramen opens near the caudal margin. The sternal process is damaged and mostly missing.

Taquet (1976) described the right coracoid of the paratype (GDF 381) instead of that from the holotype, because of the bad preservation of the latter. The right coracoid of the Venice specimen, is a plaster replica of the one described and figured by Taquet (1976, p.124, fig. 48). Only one

element is identified as a coracoid by the handwritten notes in the field map of the paratype, but it is not specified whether it is a left or a right one. The smaller size of the left one respect the right one suggests that it is not the coracoid GDF 301 of the referred material, which is reported by Taquet (1976, p. 58) as "coracoïde de grandes dimensions" (large-sized coracoid).

Scapula (Fig. 14B). Both scapulae are preserved, but the blade of the right one is mostly reconstructed. The left scapula is 640 mm long (3% longer than the scapula from the holotype). It is an elongate bone with a thickened and dorsoventrally-expanded proximal portion and a strap-like blade. The scapular contribution to the humeral glenoid is an oval and deep depression, ~~which is~~ bordered caudally by a triangular, prominent and ventrocaudally-pointing scapular labrum. The sutural surface for the coracoid is located cranially in the middle of the expanded proximal portion. The prominent acromion process has a rectangular outline (it is longer than high) in lateral and medial views. The scapular blade has a concave ventrocaudal margin and a convex dorsocranial margin; ~~its proximal part~~ is slightly arched medially ~~in its proximal part. The latter is and~~ relatively thick, whereas the distal portion of the blade is quite thin. The neck is poorly defined. The ventrocaudal and dorsocranial margins are nearly parallel along 2/3 of the blade length and diverges ~~in along~~ the last third, so the blade is narrow, expanding into ~~distal spatulae-symmetrical spatula-like shape only distally~~. The deltoid ridge crosses diagonally the blade ~~in of~~ the right scapula; it is barely visible in the left scapula, ~~where and~~ its distal portion ~~does not seem to~~ apparently does not cross diagonally the blade (Fig. 14B2). This suggests that the relative segment of the broken blade (Fig. 14B3) was joined to the others in the wrong way and belongs to a right scapula. The deltoid fossa is ~~evident visible~~ ventral to the proximal segment of the deltoid ridge. Two elements (both expanded proximal parts) are identified as scapulae ~~by in~~ the handwritten notes in the field map of the paratype (cluster 1); one complete scapula ~~seems to occur~~ is also apparently present in the cluster 2.

Sternals bones (Fig.14C). ~~Both Ssternals elements~~ are ~~both~~ preserved in MSNVE 3714, although they are partly reconstructed (Fig. 14C3). They are hatchet-shaped with an expanded and broad proximomedial portion (the sternal 'paddle') and a rod-like caudolateral process (the sternal 'handle'). ~~Their M~~maximum length is 330 mm (6% longer than the holotype sternals). The medial margin of the 'paddle' in both elements is reconstructed, being smooth instead of rough as it is usually observed in iguanodontians-should be; many missing fragments, fractures and holes were replaced or filled by resin. The 'paddle' has a triangular caudoventral process. The 'handle' is slightly expanded and thickened to the tip. Two elements are identified as sternal bones ~~by in~~ the handwritten notes ~~in accompanying~~ the field map of the paratype.




Figure 14: MSNVE 3714, pectoral girdle elements. Left coracoid in caudal (A1) and dorsomedial (A2) views; drawings of the left coracoid in caudal and dorsomedial views with the reconstructed parts evidenced in dark gray color. Left scapula in cranial (B1) and lateral (B2) views; drawing of the left scapula in lateral view with the reconstructed parts evidenced in dark gray color. Left sternal plate in dorsomedial (C1) and ventrolateral (C2) views; drawings of the left sternal plate in dorsomedial and ventrolateral views with the reconstructed parts evidenced in dark gray color. The segment of the scapular blade that seems to belong to another individual is marked in yellow. Abbreviations: ap, acromion process; cof, coracoid foramen; clp, caudolateral process ('handle'); crmp, craniomedial plate; cvp, caudoventral process; cos, coracoid sutural surface; df, deltoid fossa; dr, deltoid ridge; gl, glenoid; scl, scapular labrum; scs, scapular sutural surface; sp, broken sternal process. Scale bar equals 10 cm.

Humerus (Fig. 15A). Both humeri are original in MSNVE 3714. This element (510 mm long; 92% the length of the humerus from the holotype) is relatively slender and slightly sigmoid in cranial and caudal views. The caput humeri (humeral head) is a bulge located in the middle of the proximal end of the element and skewed to the caudal surface (Fig. 15A1); it extends along the caudal surface of the proximal portion of the humerus as a distally tapering buttress. Lateral to the caput is a distinct and sub-spherical greater (outer) tuberosity; medial to the caput is a scarcely defined medial (inner) tuberosity. The deltopectoral crest is much longer than wide (i.e., it is scarcely prominent) and symmetric (the apical point is in the middle of the crest and there is no steep distal margin). Its blunt apex is in the proximal half of the humerus. The distal end of the bone bears the radial and ulnar condyles, which are twisted laterally. The first is rounded, pointing more cranioventrally than that of the holotype. The radial condyle is larger than the ulnar one. The condyles are separated by an intercondylar groove.

Two elements (both expanded proximal parts) are identified as scapulae ~~by in~~ the handwritten notes ~~in accompanying the~~ field map of the paratype (cluster 1); one complete scapula ~~is seems to occur~~ also ~~apparently present~~ in ~~the~~ cluster 2.

Two elements are identified as humeri by in the handwritten notes in of the field map of the paratype and are in their anatomical position (Fig. 2).

Ulna and radius (Fig. 15B). Both radius-ulna pairs are preserved; the elements of each pair are stuck together by glue. The ulna is 415 mm long (88% the length of the holotype ulna) and its shaft is straight in all views. The Its proximal end is expanded craniocaudally and mediolaterally, and is bears formed by three processes. The well-developed-robust olecranon is the more robust. It is larger and more well-formed better developed than that of the ulna from the holotype (see Taquet, 1976, fig. 51a, c). Two nearly triangular flanges bordering the wide depression for the radius ~~are present~~ are present distal to the olecranon; the lateral flange is shorter and points craniolaterally, whereas the medial one is longer and points craniomedially. The convex distal articular surface is bean-shaped.

The radius is 350 mm long (84% the length of the radius from the holotype) and its shaft is also straight in all views. Its proximal end is mediolaterally expanded and ovoid in proximal view; its distal end is craniocaudally expanded and is wider than the proximal end. The distal surface is nearly triangular, with a convex lateral margin and a straight medial margin. The proximal and distal portions of the radius lie on cranial depressions of the ulna.

Two elements are identified as ulnae by in the handwritten notes in of the field map of the paratype; one (probably the left one) is completely exposed, the other apparently fragmentary or just partially cropping out from the rock. Only one radius (probably the left one, which is parallel to the relative ulna) is identified as-by in those notes.




Figure 15: MSNVE 3714, forelimb, long bones. Left humerus in caudal (A1), cranial (A2), proximal (A3) and distal (A4) views; left radius-ulna in caudal (B1), cranial (B2), medial (B3), lateral (B4), proximal (B5) and distal (B6) views. Abbreviations: ch, caput humeri; dpc, deltopectoral crest; gtb, greater tuberosity; itb, inner tuberosity; lf, lateral flange; mf, medial flange; ol, olecranon; ra, radius; rc, radial condyle; u, ulna; uc, ulnar condyle. Scale bar equals 10 cm.

Carpals (Fig. 16A). Carpals are not preserved in the right forelimb and were not replaced by artificial copies, whereas original elements ~~occur~~ are present in the left one. There is a proximal row of three carpals, which are stuck by glue to their relative metacarpals (which, however, are artificial); there are no distal carpals. One carpal apparently articulates with phalanx I-1 and metacarpal II, therefore it is in the position of the radiale (which is usually fused with the metacarpal I in basal ankylopollexians; references); the carpal in the middle articulates with metacarpal III and possibly partly with metacarpal IV and can be identified as the intermedium; the ~~other third~~ carpal articulates with the metacarpal IV and should be the ulnar. The 'radiale' has a blocky shape, with a sub-quadrangular outline, an irregular proximal surface and a slightly concave distal surface. The 'intermedium' is the largest ~~of the three carpal~~ and has a quadrangular outline in proximal and lateromedial views (it is dorsoventrally higher than lateromedially wide). A groove divides the medial side of the element into two parts in a way that they appear as two separate elements in that view. The 'ulnare' is a quadrangular blocky element that is proximodistally longer than ventrodorsally high in lateromedial view.

The main cluster of the field map (Fig. 2) shows two rounded elements ~~of the main cluster~~ that are labeled as "carpals" in a handwritten note.

The wrist of the holotype is completely different, being formed by two large and proximodistally short proximal elements. The wrist elements of MSNVE 3714 more closely resemble ~~more~~ those of *Iguanodon bernissartensis* (see Norman 1980, fig. 60) and *Mantellisaurus atherfieldensis* (see Norman 1986, fig. 50) in their relatively small size and block-like aspect, but the largest element in both taxa is the radiale, ~~which is~~ fused with metacarpal I. We suspect that those mounted in MSNVE 3714 are not actually carpals or at least they do not preserve their original shape.

Metacarpals (Fig. 16). Metacarpals from the right manus are artificial, whereas those of the left one are original with minor restoration (Fig. 16A2 and B2). Metacarpals III and IV are the longest, metacarpal III being only slightly longer than metacarpal IV (113 and 108 mm, respectively, in both our and Taquet's measurements). The shortest is metacarpal V (68 mm long; 74 mm according to Taquet's measurements).

Scattered manual ~~remains~~ elements, including two or three metacarpals, can be recognized in the region of the forelimb bones in the map of cluster 1 (Fig. 2); because of their position, they plausibly belong to the left forelimb, which is fairly well-articulated.

The orientation of the manus in the following description is the standard one for digitigrade quadrupedal amniotes, although *Ouranosaurus* probably kept the palms facing somewhat medially. Metacarpal II (85 mm long in both our and Taquet's measurements) is straight, relatively slender (it is constricted in the middle), ~~and~~ with expanded extremities. Its proximal end is expanded craniopalmarly and mediolaterally. Its proximal surface has a squared outline and is flat. A longitudinal ridge ~~runs~~ extends along the medial margin from the proximomedial corner to mid-shaft. The distal end is pointed at its lateropalmar margin; the distal surface has a sub-oval outline and is concave. As underlined by Taquet (1976, p. 131), ~~the~~ metacarpal II from the holotype is more slender and mediolaterally flattened than this metacarpal. Metacarpal III is straight and expanded at both extremities. ~~Its~~The proximal end is expanded mediolaterally; ~~its~~the proximal surface has a sub-rectangular outline and is gently convex. ~~Its~~The proximal end is more pointed laterally ~~than~~ medially. The craniomedial corner of the proximal surface is craniomedially pointed. The distal end is expanded mediolaterally; a shallow palmodistal groove divides it into two rounded condyles.

Commented [PG15]: Where? In 1976 monograph, in handwritten notes?

According to Taquet (1976, p. 131), this metacarpal is unlike that of the holotype, but probably it is a typo: it is written 3 instead of 5. Actually, the metacarpal III of the holotype is only longer, according to figures 56 and 57a. Metacarpal IV is straight, with proximal and distal ends that are moderately expanded mediolaterally. The proximal surface has an elliptical outline (it is craniopalmarly compressed) and is convex. The lateral surface of the shaft presents a ridge that extends from the proximal to the distal end. The proximal portion of metacarpal IV of the holotype differs in being mediolaterally flattened (Taquet, 1976, fig. 56). Metacarpal V is stout and hourglass-shaped in craniopalmar view, with the distal extremity that is wider than the proximal one. The proximal surface has a rounded outline and is shallowly concave in the middle. The palmar surface is slightly concave, whereas the cranial one is convex. The distal end is expanded palmarly, whereas the proximal one is expanded cranially. A small knob occurs near the distal extremity of the metacarpal at the palmomedial corner. The distal surface has a sub-rectangular outline, with a shallowly concave central part. As underlined by Taquet (1976, p. 131), the metacarpal V of the holotype is completely different, being much slender, nearly as long as metacarpal IV and mediolaterally flattened.

In conclusion, the palm of MSNVE 3714 is sensibly different from that of the holotype, as already noticed by Taquet (1976).

Manual phalanges (Fig. 16). The right manus of MSNVE 3714 preserves seven original phalanges, whereas the left one has six. The field map (Fig. 2) shows two elements of the cluster 1 that are labeled as "phalanx" in handwritten notes, both on the side of the forelimbs. One is an ungual phalanx and occurs 2.5 metres away from the cluster; the other is another, larger and arrow-shaped ungual phalanx. Also a group of four smaller phalanges occurs in the ?left hand region. Phalanx II-1 (right hand) is hourglass-shaped in palmar and dorsal views, with proximal and distal extremities that are mediolaterally expanded. Its distal end is more dorsopalmarly expanded than the proximal end. The proximal surface has a sub-circular outline and is flat; the distal surface has a sub-triangular outline and is also flat. Its dorsal surface is covered by a film of resin. Phalanx II-2 (right hand) is a small and mediolaterally short element, with a slightly concave palmar surface. Its dorsal surface is covered by a film of resin. Its proximal surface has a bean-like outline and is slightly convex; its distal surface has a sub-elliptical outline and is flat. Phalanx II-3 (the ungual; left hand) is hoof-like and elongated; its proximal surface has an elliptical outline and a depression at its center. Phalanx III-1 (preserved in both hands) is a stout and hourglass-shaped element in palmar and dorsal views. Its proximal surface is flat with a sub-elliptical outline. The outline of the distal surface is bean-like with a concave central part. The distal termination bears two rounded condyles that are separated by a broad and shallow depression. The right phalanx III-1 of the paratype figured in Taquet (1976, fig. 57b) is mediolaterally narrower than that in the mounted skeleton (about 28 and 34 mm, respectively); probably Taquet (1976) figures the left phalanx III-1 of the paratype (which is 28 mm wide) instead of the right. Phalanx III-2 is missing in both hands. The ungual phalanx of digit III (phalanx III-3; left hand) is hoof-like and long (it is the longest ungual phalanx). It shows two longitudinal grooves that are parallel to its lateral and medial edges; they start in the upper third of the bone and reach its distal tip. They anchored the keratin sheath of the ungual. The proximal surface has a sub-elliptical outline and a concave center. Phalanx IV-1 (right hand) is hourglass-shaped in palmar and dorsal views, with the distal extremity that is more mediolaterally expanded than the proximal one. The proximal surface has a sub-elliptical outline and is flat. The distal one has a sub-elliptical outline and a slightly convex surface. The sub-triangular distal condyles are separated by a shallow depression that is less broad than that of phalanx III-1. The proximal surface has an elliptical outline and is flat. The distal surface has also an elliptical outline and is slightly convex. This phalanx is more slender than the phalanx IV-1 of the paratype figured by Taquet (1976, fig. 57b). Phalanx IV-2 (left hand) is hourglass-shaped and relatively stout in palmar and dorsal views. As it is partly reconstructed and/or covered by resin,

1011 other morphological details are not reliable. The ungual of digit IV (IV-3; preserved in both hands)
 1012 is hoof-like and elongated. The proximal articular surface has an elliptical outline and a concave
 1013 center. Phalanx V-1 (preserved in both hands) is the stoutest manual phalanx and is hourglass-
 1014 shaped in palmar and dorsal views. The proximal surface has a bean-like outline and a shallowly
 1015 concave center. The proximal extremity is more dorsopalmarily expanded than the distal one. The
 1016 palmar surface of the phalanx is slightly concave and presents some thin longitudinal grooves
 1017 (which are more evident in the right element than in the left one), possibly for muscular insertion or
 1018 just evidence of a cartilage covering (like in the lateral side of the caudal vertebrae near their
 1019 articular faces). The distal surface has also a bean-like outline and is slightly convex. Phalanx V-2
 1020 (right hand) is also hourglass-shaped in palmar and dorsal views. Its proximal surface has a sub-
 1021 elliptical outline and a slightly concave center; its distal surface has a bean-like outline and a
 1022 slightly concave center.
 1023 The description of the paratype phalanges by Taquet (1976) is messy. He does not describe or list
 1024 all the phalanges preserved in that specimen. He just says that seven phalanges from the paratype
 1025 left hand allowed him to complete his description of the manus (p. 132), but then he refers those
 1026 phalanges to the right hand in the caption of figure 57. In figure 57c, Taquet (1976) shows the right
 1027 ungual phalanx V of the paratype, but that phalanx is not preserved in MSNVE 3714. In *Iguanodon*
 1028 and *Mantellisaurus*, this element is strongly reduced (Norman, 1980, 1986), thus Taquet's
 1029 assignment is possibly wrong. That phalanx is similar to the left ungual phalanx IV of MSNVE
 1030 3714, but with a reversed curvature. The Venice specimen preserves two right phalanges that
 1031 Taquet (1976) does not utilize in the description of *Ouranosaurus* and does not figure: IV-3 and V-
 1032 2. Phalanges II-1, III-1, IV-1 in the right manus of MSNVE 3714 are morphologically unlike the
 1033 corresponding paratype phalanges figured by Taquet (1976). It is unclear whether this is related to
 1034 the quality of the drawing (which is however unlikely) or they are actually different phalanges.
 1035 Possibly, the phalanges described by Taquet (1976) were replaced in MSNVE 3714 with other
 1036 phalanges and the original were sent back with the holotype to Niamey or kept in the MNHN
 1037 collection as it is the case of other bones from the paratype that were described in 1976 (i.e., the
 1038 right coracoid and the left femur).
 1039




Figure 16: MSNVE 3714, forelimb, *mani* in dorsal (cranial) view. Left (A1) and right manus (B1). A2 and B2 are the drawings of the left and right manus, respectively, with the totally reconstructed parts evidenced in dark gray color and those just covered by a film of resin coloured in pale gray; also minor reconstructed portions are in pale gray. Abbreviations: int, intermedium; mc II-V, metacarpals II-V; mk, medial distal knob on the metacarpal V; ph II-V, phalanges of manual digits II-V (the last phalanx of each digit is the ungual); rad, radiale; uln, ulnare. Elements without abbreviation are reconstructed. Scale bar equals 10 cm.

Ilium (Fig. 17A). Both ilia are preserved, but the left is more complete than the right one. The left ilium is 770 mm long (91% the length of the holotype ilium). It is lateromedially compressed and sigmoid in dorsal view. The dorsal margin of the iliac blade between the preacetabular and postacetabular processes is straight in lateral and medial views and thickened. The preacetabular process is long and curved downward; its rostral tip is slightly upturned pointing cranioventrally, like that of the ilium of *Iguanodon bernissartensis* (Norman 1980, figs. 63-64), and tapers rostrally. The preacetabular process is bent laterally in dorsal and ventral views. A prominent ridge crosses diagonally the proximal half of the process in its medial surface, bordering dorsally a broad shelf (medial shelf). The body of the ilium is sub-rectangular and bears facets for the articulation with the sacrum medially. In the left element, the articular ridge is relatively thick and is arched with ventral concavity, apparently not extending caudal to the level of the caudal end of the acetabulum; it is shallower, thinner, straight and extending more caudally in the right ilium. As at least one ilium was detached from the sacrum (as shown in the field map; Fig. 2, see below), probably the pelvis and sacrum were not coossified. The iliac contribution to the acetabulum is a shallow and caudocranially short notch along the ventral margin of the iliac body. The pubic peduncle is broken and mostly missing in both ilia; the ischial peduncle is scarcely projecting, but its articular surface is quite long. The ischial peduncle is separated from the postacetabular process by a notch. The knob-like supracetabular process (antitrochanter) is located just ventral the dorsal beginning of the postacetabular process, skewed caudally with respect to the acetabulum. The postacetabular process

is large, deep and arrowhead-shaped in lateral and medial views. Ventrally, it bears a distinct *brevis* shelf.

Only one element can be referred to an ilium in the cluster 1 of the field map (Fig. 2). It was labeled as "ilium" in a handwritten note that was cancelled as it were wrong. Notwithstanding, that element can plausibly be an ilium because of its outline and position. It seems to lack the long preacetabular process like the right ilium of MSNVE 3714. The left ilium could still be totally inside the rock; this remains speculative in absence of a list of the bones obtained after preparation of the collected blocks in the MNHN laboratories. The word "ilium" is handwritten in a zone of the cluster 2 close to the pubes and at least one ischium, but the outline of the element is not recognizable; another isolated ilium (not labeled as such) is clearly present in this cluster and preserves the long preacetabular process as the left ilium of MSNVE 3714. It could have been used to complete the individual of the cluster 1.

The ilia of MSNVE 3714 are quite similar in lateral and medial views to that of the holotype figured by Taquet (1976). However, the supracetabular process does not correspond with the dorsal hump at the beginning of the postacetabular process, but it occurs in a more cranial position.

Pubis (Fig. 17B). In both pubes, the distal portions of the prepubic blade and the posterior pubic ramus are reconstructed. As reconstructed, they measure 790-810 mm long from the rostral tip of the blade to the tip of the posterior ramus. The neck of the prepubic process is stout, taller than long. The plate-like prepubic blade is much expanded dorsoventrally; it is asymmetrically more expanded dorsally than ventrally. The iliac peduncle is massive and broad in lateromedial view; the ischial peduncle is much smaller and short. Those peduncles form the cranioventral margin of the acetabulum. The preserved proximal portion of the posterior pubic ramus is rod-like and slightly tapers distally; its basal portion is placed medially with respect to the ischial peduncle. The ischial peduncle and the posterior pubic ramus border a notch that should be partially closed caudally by a small process ascending from the pubic ramus to form the obturator foramen, but this is not the case and an obturator foramen is not evident.

Only one element in the cluster 1 of the field map is labeled as "pubis" by a handwritten note (Fig. 2); it is close to the forelimb and apparently is more complete than those of MSNVE 3714. Two apparently complete and parallel ilia (labeled as such by a handwritten note) occur in the cluster 2 close to at least one ischium and a ilium. The considerations made for the ilia are valid also for the pubes.

Differences with the holotype are in the expansion of the prepubic blade, but that part is reconstructed in MSNVE 3714.

Ischium (Fig. 17C). Both ischia are preserved, but some portions are reconstructed, in particular the pubic and obturator processes. They measure 880 mm (105% the length of the ischia from the holotype). The iliac peduncle is large, fan-shaped in lateral and medial views and is directed craniodorsally. Its distal articular surface is oval in outline. The shaft is straight; it tapers up to mid-length then it enlarges gradually up to the distal extremity. The latter has a mediolaterally compressed, dorsoventral expanded and ventrally-pointing 'boot'.

Only one element in the cluster 1 of the field map is labeled as "ischium" by a handwritten note (Fig. 2); it is close and parallel to the left fibula and has not the appearance of an ischium. One ischium (labeled as such by a handwritten note) occur in the cluster 2 close to the paired pubes and in its anatomical position respect one of them. The proximal part of another ischium seems to occur in a slightly displaced position, but it is not labeled as such by an handwritten note. The considerations made for the ilia and pubes are valid also for the ischia.

The iliac peduncle of MSNVE 3714 is more robust than that of the ilium from GDF 300 (Taquet, 1976, fig. 60). Furthermore, the shaft is sinuous in the holotype.




Figure 17: MSNVE 3714, pelvic girdle elements. Left ilium in medial (A1); lateral (A2); dorsal (A4); and ventral (5) views. Left pubis in lateral (B1) and medial (B2) views. Left ischium in medial (C1) and lateral (C2; upside-down) views. A3, B3 and C3 are the drawings of the elements of figures A1-B1, B1-B2 and C1-C2, respectively, with the reconstructed parts evidenced in dark gray color. Abbreviations: ac, acetabulum; bs, *brevis* shelf; ilp, iliac peduncle of ilium and pubis; isf, distal 'foot' of the ischium; isp, ischial peduncle of ilium and pubis; ms, medial shelf; no, notch; obf, obturator foramen; obp, obturator process; pop, postacetabular process; por, posterior pubic ramus (pubis s.s.); ppb, prepubic process; ppn, neck of the prepubic process; prp, preacetabular process; pup, pubic peduncle of ilium and ischium; sap, supracetabular process; sf, facet for the

1128 articulation with sacrum. Scale bar in A1-B1, B1-B2 and C1-C2 equals 10 cm.

1129

1130 **Femur** (Fig. 18A). The right femur is original, while the left one is a plaster replica. Taquet (1976,
1131 pl. 24, figs. 1 and 3) used the left femur of the paratype to describe the femur of *Ouranosaurus*
1132 because those of the holotype are partially and poorly preserved. According to the field map, the left
1133 femur was the only one preserved in the cluster 1; none occurs in the cluster 2. As the left femur of
1134 MSNVE 3714 is just a copy of that figured by Taquet (1976), the original bone was probably sent
1135 back to Niger with the holotype or remained in Paris, like the right coracoid and possibly the
1136 manual phalanges, although there is no written record about this.
1137 The right femur is 920 mm-long, being 8% longer than the left one and much longer than the tibia
1138 (129.5%). Its morphology is that common to femora of all advanced iguanodontians. Its shaft is
1139 straight in lateral and medial views except for the distal third, which gently curves caudally; it is
1140 slender and straight in cranial and caudal views. The caput femoris (head of the femur) is rounded
1141 in medial view, nearly triangular in cranial and caudal views, medially directed and set off from the
1142 rest of the femur by a distinct neck. The greater trochanter points dorsocaudally in lateral view
1143 extending dorsally slightly above the caput, and it is mediolaterally compressed. It forms the whole
1144 lateral half of the proximal part of the femur. A shallow depression separates it from the caput. Like
1145 in other advanced iguanodontians, the lesser (cranial) trochanter is a mediolaterally flattened, long
1146 and tongue-shaped process that is located laterocranially respect to the greater trochanter and
1147 separated by a cleft. In the right femur, the lesser trochanter is placed slightly more distally than in
1148 the left one. The fourth trochanter is located midway along the mediocaudal margin of the femoral
1149 shaft and points caudally. It is not pendent as that in the left femur of the paratype figured by Taquet
1150 (1976, fig. 62C): its caudal margin is slightly sigmoid (Fig.18A2) like that of *Hypselosaurus* cf.
1151 *fittoni* in Norman (2015, fig. 18). The distal articular end of the femur is craniocaudally expanded
1152 and divided into two condyles separated by the cranial intercondylar (for extensor tendons) and the
1153 caudal intercondylar (for flexor tendons) grooves (Fig. 18A7). The condyles are more expanded
1154 caudally than cranially. The medial condyle is larger than the lateral one and projects distally
1155 beyond it. Its cranial extremity is bent laterally partly encircling medially the cranial intercondylar
1156 groove. In distal view, this condyle has a roughly rectangular outline. The medial condyle of the left
1157 femur (figured by Taquet 1976, p. 142) has a sub-triangular outline, tapers cranially and its cranial
1158 extremity is not bent laterally. The lateral condyle expands caudally with a mediolaterally flattened
1159 condylid that reaches the level of the caudal end of the medial condyle. The intercondylar grooves
1160 are deep; the caudal is slightly deeper and narrower than the cranial one.

1161 **Tibia** (Fig. 18B). Both tibiae are entirely preserved in MSNVE 3714. In the field map, the cluster 1
1162 preserves only one tibia and fibula, plausibly the left one. A tibia and at least part of a fibula occur
1163 in the cluster 2; this might be the right one of MSNVE 3714. The left tibia is 710 mm long, which is
1164 90% the length of the holotype tibia. The tibial morphology is quite consistent within
1165 iguanodontians. It is a straight bone that is craniocaudally expanded at its proximal end and
1166 mediolaterally expanded at its distal end. A prominent and mediolaterally compressed crest (the
1167 cnemial crest) projects cranially from the proximal expansion and curves laterally. The cranial
1168 portion of the proximal expansion bears two condyles, a caudomedial condyle (larger) and a lateral
1169 one, which are separated each other by a deep and narrow groove. The shaft is straight and ovoid in
1170 cross-section, but with a twist of the main axis of the section from craniocaudal proximally to
1171 transverse (lateromedial) distally. The distal end is divided into two malleoli, the lateral one
1172 extending more distally than the medial one and articulating with the caudoproximal articular facet
1173 of the calcaneum. The medial malleolus is broader than the lateral one in craniocaudal view, it is
1174 slightly bent cranially in lateral view and articulates with the astragalus.

1175 **Fibula** (Fig. 18B). Both fibulae are preserved and are glued to the relative tibiae and cannot be
1176 removed. The left fibula is glued in a slightly wrong position, as it should rest in a shallow lateral

1177 depression of the proximal part of the tibia, bordered by the cnemial crest (Norman 1986, fig. 58F;
1178 see also Godefroit et al 1998, p. 44) not on the tibial condyles. The fibula is a straight and relatively
1179 slender bone that is slightly shorter than the tibia. Their extremities are expanded craniocaudally,
1180 the proximal more than the distal one. The proximal expanded portion is lateromedially flattened
1181 and slightly crescentic in proximal view, with a convex lateral margin and a concave medial margin
1182 (Fig. 18B5). The proximal end shows a cranially pointing wing in lateral and medial views. The
1183 distal end is club-shaped, leans against the lateral malleolus of the tibia and articulates with the
1184 calcaneum. In caudal view, a straight longitudinal ridge runs from near the proximal end to the
1185 middle of the shaft. The distal surface is not visible because it is articulated with the calcaneum.
1186



Figure 18: MSNVE 3714, hind limb: femur, tibia, fibula and tarsals. Right femur in medial (A1), lateral (A3), cranial (A4), caudal (A5), proximal (A6) and distal (A7) views; A2 is a particular of the fourth trochanter. The left tibia and fibula in medial (B1), lateral (B2), cranial (B3), caudal (B4), proximal (B5) and distal (B6) views. B5 is the mirrored proximal view of the right tibia-fibula, because the proximal part of the left tibia-fibula is poorly preserved and badly mounted. Abbreviations: as, astragalus; ca, calcaneum; caig, caudal intercondylar groove; cf, caput femoris

(femoral head); cmc, proximal caudomedial condyle of tibia; cnc, cnemial crest; cod, condylid; crig, cranial intercondylar groove; ftr, fourth trochanter; gtr, greater trochanter; hn, neck of the femoral head; lc, proximal lateral condyle of tibia; lcd, distal lateral condyle of femur; lm, lateral malleolus; ltr, lesser trochanter; mcd, distal medial condyle of femur; mm, medial malleolus. Scale bar equals 10 cm.

Pes (general). The left pes (Fig. 19) presents original elements, whereas the right one is totally reconstructed. All pedal elements are glued together in the mounted skeleton, so description is limited to the exposed features.

In the field map (Fig. 2), cluster1 shows elements only from the left pes, which are close to the distal end of the paired tibia and fibula. They include bones indicated as astragalus and calcaneum by handwritten notes, and some phalanges; although only one large metatarsal seems to be present in the associated pedal bones, two much smaller elements are identified as metatarsals by handwritten notes. Such a big difference in size must be considered as an 'artistic license'.

Tarsals (Fig. 18B). Astragalus and calcaneum are glued to tibia and fibula, therefore they cannot be observed entirely. The astragalus is quadrangular, but broader medially than laterally, in distal view and cup-shaped in medial view; it is sensibly larger than the calcaneum. Its proximal surface is concave and receives the medial malleolus of tibia. The proximal margin sends two ascending processes: one occurs in the craniolateral corner and the other in the caudomedial corner; both are moderately developed.

The calcaneum is kidney-like in distal view, longer than wide; it is sub-triangular in lateral view, with a convex distal margin. Its proximal articular surface is divided into two deep facets, a cranial one for the fibula and a caudal one for the lateral malleolus of the tibia. It is exactly like the calcaneum of the holotype (Taquet 1976, fig. 67).


Metatarsals (Fig. 19). The morphology of the metatarsus is that common to metatarsi of all advanced large iguanodontians. Metatarsal III is the longest and most robust of the three. The proximal half of metatarsal II (185 mm long) is mediolaterally flattened; the distal half is slightly bent laterally and the distal end is mediolaterally expanded (the articular surface faces ventromedially). Both the proximal and distal ends are craniocaudally expanded. Most of the lateral face is occupied by the broad and slightly concave articular facet for metatarsal III; this surface is bordered cranioventrally by a crest (the "external blade" of Taquet, 1976, fig. 68B). The proximal articular surface is elliptical, longer than wide, with a convex medial margin and a slightly concave lateral margin. The dorsal craniolateral corner sends a craniolaterally-directed small flange that partially overlaps the corresponding corner of the proximal end of metatarsal III.

Metatarsal III (215 mm long) has a straight and robust shaft that gently curves medially near its proximal end. Its proximal portion is expanded craniocaudally and flattened lateromedially where it bears the convex articular surfaces for the metatarsals IV and II, laterally and medially respectively. The distal end is lateromedially expanded and is divided into lateral and medial condyles by a shallow and wide craniocaudal furrow. Metatarsal IV is slightly longer than II (it is 190 mm long). It is arched laterally and its distal articular surface faces ventrolaterally. Medially, its proximal half presents a deeply concave articular surface for metatarsal III. The cranial half of this surface is on a broad craniomedial flange of the metatarsal that overlaps the metatarsal III. A large knob occurs on the shaft just below the articular surface for metatarsal III. The distal end is transversely expanded and bears two condyles that are separated by a wide and shallow furrow.

Phalanges (Fig. 19). Digits II-IV have three, four and five phalanges, respectively, but II-3 and IV-2 to 4 are totally reconstructed. All terminal phalanges are unguals.

Phalanx II-1 is stout, gently arched laterally and hourglass-shaped in dorsoplantar view. Its distal end is more expanded lateromedially than the proximal end. The proximal surface has a quadrangular outline and is flat; the distal end is divided into two condyles that are separated by a

1244 wide and shallow furrow. Phalanx II-2 is much smaller than phalanx II-1 and has a quadrangular
 1245 outline in dorsoplantar view. Its proximal surface has a sub-triangular outline; the distal end bears
 1246 two scarcely developed articular condyles. Phalanx III-1 is large, stout and as long as wide. Its
 1247 proximal surface has a sub-elliptical outline and is slightly concave. Its distal end bears two scarcely
 1248 developed articular condyles. Phalanx III-2 is disc-like, much proximodistally shorter than wide
 1249 lateromedially. Both proximal and distal surfaces have a sub-triangular outline; the proximal one is
 1250 slightly convex, whereas the distal is concave. Phalanx III-3 has a shape similar to that of phalanx
 1251 III-2, but it is transversely narrower. The long phalanx III-4 (the ungual phalanx, 70 mm long) is
 1252 spade-like, dorsoplantarly flattened and slightly arched plantarily; the medial expansion of its distal
 1253 end is more developed than the lateral one. Phalanx IV-1 is hourglass-shaped in dorsoplantar view
 1254 and resembles phalanx II-1. Its proximal end is more dorsoplantarly expanded than the distal end.
 1255 Its proximal surface has a sub-rectangular outline. The distal end is divided into two condyles that
 1256 are separated by a wide and shallow furrow. The ungual phalanx IV-5 is a tiny element with a
 1257 squared outline in dorsoplantar view and is gently arched plantarily.
 1258 According to Taquet (1976, p. 153), seven pedal phalanges are preserved in the paratype; eight are
 1259 actually present in the Venice specimen, including unguals III and IV (which should not be present
 1260 according to Taquet, 1976). Lengths reported by Taquet (1976) correspond with those measured in
 1261 MSNVE 3714 for the phalanges II-1, II-2, III-1 and III-2. The description of the paratype pedal
 1262 phalanges by Taquet (1976) contains further mistakes. All those phalanges are considered to be
 1263 from the right pes in the caption of figure 71; in the text, phalanges II-1, III-1 are reported as left,
 1264 III-2 as possibly right, while the provenance of phalanges II-2, IV-2 and III-3 is not established.
 1265 Furthermore, phalanx IV-2 in referred to the holotype in the text, while it reported to belong to the
 1266 paratype in the caption of the figure. The only ungual phalanx is reported as a right IV-5 in the
 1267 figure, while the text says only that it is from the right foot; that phalanx does not correspond with
 1268 the ungual of digit IV of MSNVE 3714 and resembles more the ungual placed at the end of digit III.



1270

1271 **Figure 19: MSNVE 3714, hind limb: pes.** Left pes in dorsocranial view. To the right, drawing of
1272 the left pes with the totally reconstructed parts evidenced in dark gray colour and minor
1273 reconstructed portions in pale gray. Abbreviations: mtII-IV, metatarsals II-IV; pphII-IV, phalanges
1274 of pedal digits II-IV (the last phalanx is always the ungual one). Elements without abbreviation are
1275 reconstructed. Scale bar equals 10 cm.

1276
1277 **Osteohistology of MSNVE 3714**

1278 The elements that were sampled for osteohistological analyses are the humerus, the femur, the tibia,
1279 the neural spine of a distal dorsal vertebra, and a distal dorsal rib (see Materials and Methods
1280 chapter for details).

1281
1282 **HUMERUS**

1283 The left humerus was sampled on its caudal side at mid-shaft. The cylindrical core sample was 23
1284 mm thick and its diameter was 14 mm (Fig. 20A). The medullary cavity shows relatively few
1285 trabeculae, which are generally not connected to each other. Trabecular density is lower in
1286 comparison to that present in the femur (see below). The edge of the medullary cavity is neat (Fig.
1287 20B). Erosional cavities are present in the inner compacta; they show a minor density than in the
1288 femur (see below). Those cavities have an elliptical or rounded outline and occur in the inner one
1289 fifth of the compacta; their density decreases moving toward the outer surface of the cortex. The
1290 microstructure of the cortex is fibrolamellar with a matrix composed of woven bone (Fig. 20C-D).
1291 Compacted coarse cancellous bone (CCCB; *sensu* Hübner, 2012) occurs in the inner cortex
1292 surrounding the erosional cavities (Fig. 20B). Vascularization (*sensu lato*; see Chinsamy, 2005) is
1293 mainly composed of well-developed (i.e., large and with many lamellae) primary osteons that
1294 become more and more organized (i.e., regularly arranged in the space) towards the outer cortex
1295 (Fig. 20C-D). Vascularization has a laminar circumferential arrangement (Fig. 20C-D). The
1296 distance between the single vascular canals is relatively high (Fig. 20C-D). Zonation is present.
1297 Four to six LAGs are recognized in the compacta (Fig. 20A); the spacing between successive LAGs
1298 decreases moving towards the outer surface of the bone. Secondary osteons and EFS are absent
1299 (Fig. 20A-D).

1300





Figure 20: MSNVE 3714, left humerus, thin section. Panoramic view under lambda filter (the outer surface of the bone is at the top of the figure) (A); detail of the progressive transition between the compacta and the medullary cavity, characterized by erosional cavities (B); detail of the fibrolamellar bone and longitudinal vascularization forming the primary bone (C); detail of the microstructure of the outermost cortex showing the absence of an EFS (D); no secondary osteons are observed in the inner cortex. Green arrows point to the LAGs. Abbreviations: lb, lamellar bone; mc, medullary cavity; ps, periosteal surface; vc, vascular canals; wb, woven bone. Scale bars equal 10 mm in A and 1 mm in B-D.

FEMUR

The right femur was sampled on the craniolateral side at mid-diaphysis. The cylindrical core sample was 21 mm thick and its diameter was 14 mm (Fig. 21A). The medullary cavity is characterized by isolated trabeculae. The passage between the medullary cavity and the compact cortex is gradual, because of the presence of many resorption cavities in the inner cortex (Fig. 21B). Resorption cavities tend to decrease in density moving towards the external surface of the cortex, and their outline changes from irregular to rounded or elliptical. The compacta is composed of two different types of bone: the primary and the CCCB. The primary bone is composed of fibrolamellar bone with woven bone forming the matrix (Fig. 21C-D). CCCB bone is present in the inner most compact bone wall, especially in the areas surrounding the resorption cavities (Fig. 21B). Primary osteons are abundant through all the compact bone. Vascularization is irregularly organized and a clear orientation is not evident. Primary vascular canals are still open, although infilling of lamellar bone is present. Secondary osteons (Haversian systems) cannot be identified in the thin section. No LAGs or annuli can be observed and there is no EFS (Fig. 21A-D).




1327
1328

1329 **Figure 21: MSNVE 3714, right femur, thin section.** Panoramic view under lambda filter (the
1330 outer surface of the bone is at the top of the figure) (A); the gradual transition between the compacta
1331 and the medullary cavity made of CCB and erosional cavities (B); detail of the plexiform
1332 vascularization and fibrolamellar bone forming the primary bone and the absence of zonation and
1333 LAGs within the compacta (C); detail of the microstructure of the outermost cortex showing the
1334 absence of an EFS (D). Abbreviations: ec, erosional cavities; lb, lamellar bone; ps, periosteal
1335 surface; vc, vascular canals; wb, woven bone. Scale bars equal 10 mm in A and 1 mm in B-D.

1336
1337 **TIBIA**

1338 The right tibia was sampled craniolaterally in the diaphysis, slightly below mid-shaft. The
1339 cylindrical core sample was 26 mm thick and its diameter was 14 mm (Fig. 22A). Within the
1340 medullary cavity, a typical spongiosa is absent: trabeculae are rarefied (Fig. 22A-B). The boundary
1341 between the medullary cavity and the cortex is abrupt and uneven and there is a thin endosteal
1342 lamella. The primary bone microstructure is fibrolamellar with woven bone constituting the matrix
1343 (Fig. 22C-D). In the inner cortex, vascularization is irregular in its orientation, density and
1344 organization. Primary osteons are well developed and generally with a laminar circumferential
1345 orientation. Locally, reticular arrangement of the primary vascular canals is observed. Organization


1346 of primary vascular canals increases towards the outer surface of the bone. Infilling of lamellar bone
 1347 is present in those canals, which, however, are not completely filled. Secondary osteons are
 1348 abundant in the innermost cortex, extending over one fifth to one sixth of the compact bone wall
 1349 thickness (Fig. 22A). Unlike all other sampled long bones, erosional cavities are absent. Six LAGs
 1350 occur in the compact bone wall. There is no EFS (Fig. 22D).
 1351



1352 **Figure 22: MSNVE 3714, right tibia, thin section.** Panoramic view under lambda filter (the outer
 1353 surface of the bone is at the top of the figure) (A); gradual transition between the compacta and the
 1354 medullary cavity made of CCB and erosional cavities (B); detail of the deeper cortex, showing
 1355 zonation of the primary bone, irregular vascularization and fibrolamellar bone (C); detail of the
 1356 outermost cortex showing the absence of an EFS (D); note the remodeling in the inner compacta).
 1357 Abbreviations: lb, lamellar bone; mc, medullary cavity; ps, periosteal surface; so, secondary
 1358 osteons; vc, vascular canals; wb, woven bone. Green arrows point to the LAGs. Scale bars equal 10
 1359 mm in A and 1 mm in B-D.
 1360

1361
 1362 NEURAL SPINE
 1363 The neural spine of dorsal vertebra 14 was cross-sectioned in the basal third, in the middle, and in
 1364 the apical third (Figs. 23A, 24A and 25A). The cross-section is oval proximally becoming
 1365 rectangular in the central and distal segments. The medullary cavity is filled with spongiosa (Fig.
 1366 23B, 24B and 25B). The boundary between the medullary cavity and the cortex is not abrupt, as
 1367 erosional cavities occur between the trabecular structure of the medullary cavity and the compact
 1368 cortex (Figs. 23C, 24C and 25C). The compacta becomes thinner and thinner moving from the
 1369 proximal part of the spine towards its apical part. The microstructure is fibrolamellar and tends to
 1370 become more organized moving towards the outer surface of the bone. Evidence of the presence of
 1371 Sharpey's fibers is observed on the lateral surfaces of the spine in the proximal section. No
 1372 Sharpey's fibers are found in the other two thin sections. Primary vascularization is prevalently
 1373 longitudinal and becomes more organized and rarified moving through the outer cortex towards the
 1374 outer surface. Haversian systems are generally present in the inner half of the compacta (Figs. 23B-
 1375 C, 24 B-C and 25B-C). Six, four, and three LAGs are identified in the proximal, median and distal
 1376 sections, respectively. We consider six or seven LAGs the most reliable count to establish the age of
 1377 the individual, because the base of the neural spine is expected to preserve the most complete
 1378 growth record. The spacing between the zones decreases moving towards the outer surface of the
 1379 bone. An EFS is absent (Figs. 23D, 24D and 25D).

1380



1381
 1382 **Figure 23: MSNVE 3714, neural spine of dorsal vertebra 14, transverse thin section of the**
 1383 **basal third.** Panoramic view of the cranial half of the section (A); Haversian systems in the inner
 1384 most cortex (B); transition between the compacta and the medullary cortex with erosional cavities
 1385 and remodeling (C); detail of the outer cortex showing absence of an EFS and outermost LAGs (D).
 1386 Abbreviations: lb, lamellar bone; ps, periosteal surface; so, secondary osteons; tb, trabeculae; vc,
 1387 vascular canals; wb, woven bone. Green arrows point to the LAGs. Scale bars equal 10 mm in A, 1
 1388 mm in B and C, and 500 microns in C.




Figure 24: MSNVE 3714, neural spine of dorsal vertebra 14, thin section of the median part. Panoramic view of the cranial half of the section (A); Haversian systems in the inner most cortex and endosteal bone (B); gradual transition between the compacta and the medullary cortex with erosional cavities and marked remodeling of the primary bone (C); detail of the outer cortex showing absence of an EFS and zonation of the primary bone (D). Abbreviations: eb, endosteal bone; ec, erosional cavities; ps, periosteal surface; so, secondary osteons; tb, trabeculae; vc, vascular canals; wb, woven bone. Green arrows point to the LAGS. Scale bars equal 10 mm in A and 1 mm in B-D.




Figure 25: MSNVE 3714, neural spine of dorsal vertebra 14, transversal thin section of the apical third. Panoramic view of the cranial half of the section; notice how the compact cortex becomes thinner trending through the top of the neural spine (A); detail of the Haversian systems in the inner most cortex and endosteal bone (B); gradual transition between the compacta and the medullary cortex (C; notice the erosional cavities and deep remodeling of the primary bone); detail of the outer cortex with outermost LAGs but without an EFS (D). Abbreviations: eb, endosteal bone; ec, erosional cavities; lb, lamellar bone; ps, periosteal surface; so, secondary osteons; wb, woven bone. Green arrows point to the LAGs. Scale bars equal 10 mm in A and 1 mm in B-D.

DORSAL RIB

The transversal cross section of the proximal part of the dorsal rib 15 has an oval outline. The cortex of the lateral side is heavily eroded; its maximum thickness is 17 mm (Fig. 26A). The medullary cavity is filled with spongiosa. The boundary between the spongiosa and the compacta is gradual, the erosional cavities in the inner cortex becoming smaller and fewer moving toward the outer surface of the cortex (Fig. 26B). The microstructure is fibrolamellar with a matrix of woven bone. Primary osteons are well developed and abundant in the outer cortex and have a longitudinal orientation. Vascular canals are partially infilled with lamellar bone, so they are still opened. Primary vascularization is more organized and less dense in the outer cortex. Secondary osteons are present in the innermost cortex (Fig. 26C). Six or seven LAGs can be identified (Fig. 26A); they tend to be more and more closely spaced moving towards the outer surface of the cortex. An EFS is absent (Fig. 26D).




Figure 26: MSNVE 3714, dorsal rib, transverse thin section. Cranio-lateral intersect of the dorsal rib (A); transition between the outer cortex and the medullary cavity (B; note erosional cavities and deep remodeling of the primary bone); Haversian systems in the inner cortex (C); detail of the outer most cortex showing zonation and the absence of an EFS (D). Abbreviations: ec, erosional cavities; ps, periosteal surface; so, secondary osteons; vc, vascular canals; wb, woven bone. Green arrows point to the LAGS. Scale bars equal 10 mm in A, 1 mm in B and D, and 500 microns in C.

DISCUSSION

Is MSNVE 3714 a composite?

The Venice specimen is undoubtedly the paratype GDF 381- MNHN figured in Taquet (1976, pl. 9, fig. 2) and figured in the field map that we received from the MNHN and is labeled "Ouranosaurus nig[er]iensis] - Airfield - 1970 - (specimen Venice Museum pro parte)". However, the latter depicts two clusters of bones belonging to two different individuals (Fig. 2). Cluster 1 is most of a quite well articulated skeleton without skull and lower jaw and apparently lacking also the distal caudals, at least half of the pelvic elements (only an ilium, a pubis and an ischium are present at best), one

hind limb, elements of the hands and possibly one coracoid. Cluster 1 is labeled "GDF 381 Today in Venice" in the map. That note was clearly added after the mapping and is a further prove that cluster 1 is the paratype and represents the skeleton exhibited in Venice. Cluster two is a partial skeleton with the pelvic elements, the proximal part of the caudal segment of the vertebral column, part of a hind limb, and possibly the sacrum and a scapula.

The words "Pro-parte" used in the caption could mean that only part of the mapped bones was used in the mount of the Venice skeleton and the other elements remained in Paris or were sent back to Niamey with the holotype.

Some of the paratype bones were replaced in MSNVE 3714 with plaster casts, namely the left femur and the right coracoid. Also the original manual phalanges of GDF 381 have apparently been replaced with other phalanges in MSNVE 3714. All those bones were used by Taquet in his 1976 description of *Ouranosaurus*. We hypothesize that they were sent back to Niamey with the holotype in order to keep all the described material in the same place. It is unclear whether this was the fate of the pedal phalanges too, because of their confusing description in Taquet (1976). Of course, this remains speculative in absence of a written documentation attesting it.

Some elements of MSNVE 3714 are not present in the field map of GDF 381: the left coracoid, at least one ilium (probably the left), one pubis, one ischium, the right femur and the right tibia. Furthermore, the distal portions of the first two chevrons lack in the map but are present in the mounted elements and there are two dorsal vertebrae more in the mount than in the mapped skeleton.

The left coracoid could be one of the unidentified or partly emergent from the rock elements of the quite well-articulated shoulder girdle. Other elements were possibly still covered by rock when the map was drawn (e.g., the left ilium and the distal portion of the chevrons). However, it is not plausible that the right femur and tibia were still covered by rock because the skeleton exposed its right side and the exposed hind limb was the left one. We hypothesize that the well-preserved pelvic elements from the cluster 2 were used to integrate the pelvic region of GDF 381 in the mount. Also the right tibia could come from cluster 2, while the right femur could be GDF 302 of the referred material. The two dorsals in excess could come from another specimen or from part of cluster 2 that results to be poorly-traced in the field map and were added to the original vertebral column to maintain the dorsal count of 17 supposed for the holotype. Of course, this remains speculative in absence of a written documentation attesting it.

Therefore, MSNVE 3714 is most probably a composite (as most of the dinosaur mounts in the museums), although it is mostly made of the paratype of *Ouranosaurus nigeriensis*.

Also the osteohistological analysis suggests that the right femur is from an individual that is distinct from the paratype. According to a skeletocronological study on *Hypacrosaurus* (Horner, Ricqlés & Padian, 1999), the best growth record is recorded in the femur, followed by the tibia, dorsal ribs and neural spines. The humerus, the tibia and the dorsal rib growth records in MSNVE 3714 are comparable to each other suggesting that they could belong to the same individual (although the tibia is probably from a different one). The neural spine shows one or two LAGs more than the humerus, the tibia and the dorsal rib. However, this could be due to the lower amount of remodeling and lower density of Haversian systems found in the thin sections of the neural spine. So, also the neural spine could belong to the same individual ~~of as the~~ humerus, the tibia and the dorsal rib.

Following Horner, Ricqlés & Padian (1999), the femur should present a higher number of LAGs, higher density of Haversian systems and a more advanced remodeling respect to humerus, tibia, dorsal rib and dorsal neural spine, if at the same growth stage, but it does not. The lack of outer LAGs and an EFS could be explained with superficial abrasion, but weathering cannot account for their total absence. Of course, we do not think that the cortex of a 920 mm long femur could be produced in a single year. Possibly, LAGs are absent because we sampled a part of the bone were they are missing. However, the lack of Haversian systems and the higher density of vascularisation

1492 still suggest that the femur belong to a more immature individual than that from which the humerus,
1493 the tibia, the neural spine and the dorsal rib come from.

1495 **Differences with the holotype**

1496 Resuming what exposed in the description of MSNVE 3714, this and the holotype differ in some
1497 respects. Furthermore, comparison suggests that some mistakes are probably present in the
1498 description of the holotype and in the mount of the Venice specimen. MSNVE 3714 and holotype
1499 differs in:

- 1500 1) The outline of the axis neural spine;
- 1501 2) The morphology of the first dorsal vertebrae, suggesting that of the reconstruction of the
1502 proximal tract of the dorsal segment of the vertebral column of the holotype by Taquet (1976) is
1503 wrong. Probably, the dorsal count is 15 (14 dorsals and one dorsosacral) instead of 17 in both
1504 specimens and the trunk is consequently shorter;
- 1505 3) The relative height of the spines of the sacral vertebrae;
- 1506 4) Five or six more proximal caudals in MSNVE 3714 than in the holotype (four to five more than
1507 in *Iguanodon* and *Mantellisaurus*);
- 1508 5) Some minor features of the caudal neural spines;
- 1509 6) The regularity of the sloping in the neural spines and chevrons in the tail (minor in MSNVE
1510 3714);
- 1511 7) A more developed ulnar olecranon in MSNVE 3714;
- 1512 8) The structure of the carpus (that of MSNVE 3714 is probably incomplete or wrongly
1513 assembled/reconstructed);
- 1514 9) The shape of the metacarpals;
- 1515 10) The position of the supracetabular process of ilium respect to the dorsal hump at the beginning
1516 of the postacetabular process (it occurs in a more cranial position in the holotype);
- 1517 11) The robustness of the iliac peduncle of the ischium (more robust in MSNVE 3714) and the
1518 curvature of the shaft (straight in MSNVE 3714).

1519 Most of them are minor differences probably due to intraspecific variability and possibly
1520 ontogeny (however, we do not know the ontogenetic stage of the holotype). Others are caused by
1521 mistakes in the preparation or assemblage of the skeletal elements in both specimens. Puzzling is
1522 the morphological difference between the elements of the metacarpus. It could be speculatively
1523 explained with the intraspecific variability.

1524 Finally, the description of the paratype manual phalanges by Taquet (1976) does not correspond
1525 with what observed in the Venice specimen. Possibly, the phalanges described by Taquet (1976)
1526 were replaced in MSNVE 3714 with other phalanges and the original were sent back with the
1527 holotype to Niamey or kept in the MNHN collection.

1529 **Ontogenetic stage of MSNVE 3714**

1530 Describing *Ouranosaurus*, Norman (2015, p. 62) writes that "this animal attained a length of 6–7 m
1531 when mature." Actually, nobody attempted to establish the ontogenetic stage of the individuals of
1532 *Ouranosaurus nigeriensis*. Taquet (1976) does not discuss the ontogenetic stage of GDF 300; he
1533 just reports that the elements of the axis of the holotype and the bones of the neurocranium are
1534 fused and their boundaries are difficult to recognize. Possibly, these features were considered as an
1535 evidence of maturity. However, no information about the obliteration of the neurocentral sutures in
1536 the other vertebrae are reported and no osteohistological investigation was attempted. Furthermore,
1537 the universal validity of obliteration of cranial sutures as evidence of osteological maturity has been
1538 argued by Bailleul et al (2016).

1539 The holotype was estimated to be seven metres long (Taquet 1976, p. 175). The length of the
1540 mounted MSNVE 3714 is about 6.5 metres from the tip of the snout to the last preserved caudal

Commented [PG16]: You could check it! Paris is not the end of the world!

vertebra. The total length was probably higher because the total count of the caudal vertebrae in large ornithomimids usually ranges between 50 and at least 75 (Hone 2012; FMDV and MF, pers., obs.). The humerus, ulna and tibia of the Venice specimen are 92, 88 and 90% the length of those of the holotype, respectively. This is in agreement with the total length. The linear size of the first is therefore about 10% smaller than that of the holotype. Relative proportions of the two individuals can be seen in Figure 27. Also the size of the other/s individual/s used to complete MSNVE 3714 is approximately the same.

Establishing the ontogenetic stage of MSNVE 3714 was undertaken by the macroscopic observation of the evidence of osteological immaturity of the skeletal elements and by their osteohistological features. The neurocentral sutures in the mid-tail caudals (Fig. 12B-D) and in at least the last cervical vertebra (Fig. 6A) are not obliterated. The rough surface texture of the centrum near its articular facets in mid-caudal vertebrae (Fig. 12A) also indicates a cartilage covering and incomplete ossification. This is suggestive of immaturity (Bennett, 1993; Brochu, 1996; Irmis, 2007). The osteological immaturity of the paratype is supported also by the unfusion of ilia and sacrum, which is the case of the holotype too.

The increasing organization of vascular canals toward the outer surface, the presence of Haversian systems, the decreasing spacing between LAG's and the absence of an EFS observed in the thin sections of tibia, neural spine and dorsal rib suggest that these skeletal elements belonged to a sub-adult individual (sensu Horner et al., 2000). In the humerus, the increasing organization of vascular canals toward the outer surface, the decreasing spacing between LAG's and the absence of EFS, but the absence also of Haversian systems, is also compatible with a sub-adult growth stage sensu Horner et al. (2000).

The conclusions are that a 7 metres-long individual of *Ouranosaurus nigeriensis* (including a minimum of 20 missing distal caudals) was not fully grown, although probably close to adulthood. In crocodiles, the obliteration of the neurocentral suture during ontogeny starts in the tail and ends in the neck (posterior-anterior sequence of neurocentral closure; Brochu, 1996). This is the case of most ornithomimids, although not all (e.g., Zheng et al., 2012), while that pattern was not followed in several saurischian dinosaurs (Irmis, 2007). Actually, there is considerable variation of both the sequence and timing of neurocentral suture closure within archosaurs (Irmis, 2007). The condition in *Ouranosaurus* suggests that it did not follow a simple posterior-anterior or anterior-posterior sequence of neurocentral closure and that obliteration occurred relatively late in the ontogeny, when the individual was close to adulthood.




Figure 27: Size of the holotype and MSNVE 3714. Holotype (A) and MSNVE 3714 (B). Redrawn and modified from Taquet (1976). Scale bar equals 50 cm.

1579 **Osteohistological comparison with other ornithopods**

1580 The osteohistology of many ornithopod taxa has been studied, including *Hypsilophodon* (Reid,
1581 1984; Chinsamy, et al. 1998); the “Proctor Lake ornithopod” (Winkler, 1994); *Orodromeus* (Horner
1582 et al., 2001; Horner et al., 2009); a “hypsilophodontid” from Dinosaur Cove/Flat Rocks, Victoria,
1583 Australia (Chinsamy et al., 1998; Woodward et al., 2011); *Rhabdodon* (Nopcsa, 1933; Reid, 1984,
1584 1990; Ősi et al., 2012); *Mochlodon* (Ősi et al., 2012); *Zalmoxes* (Benton et al. 2010; Ősi et al.,
1585 2012); *Dryosaurus* (Horner et al., 2001; Horner et al., 2009); *Dysalotosaurus* (Chinsamy, 1995;
1586 Hübner, 2012); *Valdosaurus* (Reid, 1984); *Camptosaurus* (Horner et al., 2009); *Tenontosaurus*
1587 (Werning, 2012); *Iguanodon* (de Ricqlès et al., 2012); '*Telmatosaurus*' (Benton et al., 2010);
1588 *Edmontosaurus* (Reid, 1985); *Maiasaura* (Barreto et al., 1993; Barreto, 1997; Horner et al., 2000,
1589 2001; Woodward et al., 2015); and *Hypacrosaurus* (Horner et al., 1999; Cooper et al., 2008).
1590 *Orodromeus* shows longitudinal arrangement of the vascular canals, far less dense and complex
1591 than that present in more derived ornithopods like *Maiasaura* and *Hypacrosaurus* (Werning, 2012).
1592 Remodeling is scarce at the adult ontogenetic stage (Horner et al., 2009). LAGs occur in juvenile
1593 individuals; the highest LAGs number is two in sections with EFS (Horner et al., 2009; Werning,
1594 2012). *Orodromeus* is therefore considered as characterized by slow growth (Horner et al. 2009;
1595 Werning, 2012). Rhabdodontids (*Zalmoxes* and *Rhabdodon*) and *Tenontosaurus* show higher
1596 growth rates than *Orodromeus*, as suggested by the woven bone forming the microstructure, more
1597 complex orientation of the vascular canals (radial or circumferential) and the absence of LAGs
1598 during the young ontogenetic stage (Werning, 2012). Remodeling is generally present only during
1599 the late growth (late sub-adult and adult ontogenetic stages) (Werning, 2012). Moreover,
1600 vascularization generally tends to decrease through ontogeny, showing a progressive passage
1601 between the initial rapid growth characterized by woven bone to the later slow growth indicated by
1602 lamellar bone leading to an EFS (Werning, 2012). More derived Iguanodontians (e.g. *Dryosaurus*,
1603 *Dysalotosaurus*, *Valdosaurus*, *Camptosaurus*, *Iguanodon*, *Telmatosaurus*, *Edmontosaurus*,
1604 *Maiasaura* and *Hypacrosaurus*) show the presence of fast deposited woven bone and vascular
1605 canals with a more complex and dense pattern (reticular or circumferential canals) (Reid, 1984;
1606 Horner et al., 1999; Horner et al., 2000, 2009; Benton et al., 2010; Hübner, 2012; Werning, 2012).
1607 Remodeling starts earlier during ontogeny in comparison to the other taxa reported above and
1608 widely spaced zones are still present during the sub-adult and adult ontogenetic stages (Reid, 1984;
1609 Horner et al., 1999, 2000, 2009; Benton et al., 2010; Hübner, 2012; Werning, 2012). Moreover,
1610 LAGs count observed in derived iguanodontians is generally lower than that generally found in the
1611 more basal taxa reported above, indicating that the somatic maturity was reached earlier (Reid,
1612 1984; Horner et al., 1999, 2000, 2009; Benton et al., 2010; Hübner, 2012; Werning, 2012). The
1613 disorganized tissue type is still present during the sub-adult and adult ontogenetic stage and the
1614 passage to the EFS is abrupt (Reid, 1984; Horner et al., 1999, 2000, 2009; Benton et al., 2010;
1615 Hübner, 2012; Werning, 2012).
1616 As expected, *Ouranosaurus* shares similar microstructural patterns with derived iguanodontians.
1617 The woven bone and the high vascular density with an alternated reticular and circumferential
1618 arrangement present in the bone microstructure suggest a fast growth. Remodeling is already
1619 present in the sub-adult ontogenetic stage. A fast growth is also supported by the widely spaced
1620 LAGs with the presence of the same bone structure and type of vascularization within the zones.
1621 The faster growth is phylogenetically coincident with the taxonomical diversification of the derived
1622 iguanodontians and their increase in body size (Werning, 2012). It is still unclear whether the higher
1623 growth rates are a consequence or a cause of the increase in body size in the clade (Werning, 2012).
1624 However, the large body size of *Tenontosaurus*, coupled with slow growth rates in comparison to
1625 those of dryomorphs, suggest that faster growth is a consequence of the body size and not the
1626 opposite, because *Ouranosaurus* has an estimated length comparable to that of *Tenontosaurus*, but

shows faster growth. As an alternative, *Tenontosaurus* may represent the maximum size an ornithomimid could grow with the basal slow growth rates (Werning, 2012).

The function of the back 'sail' of *Ouranosaurus*

Among the Dinosauria, hyperelongation of the neural spines reaches its maximum in the dorsal vertebrae of the theropod *Spinosaurus aegyptiacus* (see Stromer, 1915; Ibrahim et al., 2014), which also lived in northern Africa during the Cretaceous, although 15-20 million years later than *Ouranosaurus*. Its neural spines start to elongate from the first dorsal, reaching the maximum height in the last dorsal; spine height decreases from the first sacral on. They formed a sort of a 'sail' on the back of the animal. The basal segment of those dorsal neural spines is greatly expanded craniocaudally (Ibrahim et al., 2014). The middle segment is narrow and the apical one is expanded with the craniocaudal margins diverging apically, but to a lesser extent than in *Ouranosaurus* (Ibrahim et al., 2014). The spine height is up to ten times the centrum height. The apical portion of the spine has sharp cranial and caudal edges; it is marked by thin vertical striae, and is spaced away from adjacent spines (Ibrahim et al., 2014). According to Ibrahim et al. (2014), the 'sail' had probably a social function, given the low density of vascularization of the bone pointing against a thermoregulatory function. Recently, new skeletons of the giant ornithomimid *Deinocheirus mirificus* have shown that this, weird-looking theropod has tall neural spines in the middle dorsals (Lee et al., 2014). The neural spines of the proximal dorsals are relatively low, but spine height increases progressively up to the last dorsal, which has a neural spine that is 8.5 times taller than its centrum height. The base of the neural spines is not craniocaudally expanded as in *Spinosaurus*. All six sacral neural spines are tall as well, and, except for the first sacral, the apical parts of the spines are fused into a midline plate of bone with a straight dorsal margin in lateral view (Lee et al., 2014). No hypotheses have been proposed about the function of the 'sail' in this dinosaur.

In *Ouranosaurus nigeriensis*, the elongation of the neural spines is not restricted to the trunk as in *Spinosaurus* and *Deinocheirus*, but extends to the proximal caudal region. Occasionally, the base of the neural spine (actually, the prespinal lamina) is slightly expanded in the mid-posterior dorsals like in *Spinosaurus*, but this expansion occurs only cranially. The apical portions of the neural spines seem to have vertical striations, like in *Spinosaurus* and *Deinocheirus*. The external cortex is relatively thinner than in *Spinosaurus* (Ibrahim et al., 2014), with a thicker spongiosa. Sharpey's fibers (which are related to the attachment of muscles and ligaments) occur only in the proximal (basal) part of the neural spine in MSNVE 3714. Based on comparison with the musculature of crocodiles and birds (Tsuihiji, 2005; Organ, 2006a), this suggests that muscles of the *M. transversospinalis* group attached just above the base of the neural spines, connecting them at their cranial and caudal edges. The absence of Sharpey's fibers in the thin sections from the middle and apical portions of the spine, plus the absence of muscle insertion marks on the bone surface may indicate that muscles of the *M. transversospinalis* group were not attached to the entire surface of the spines. In extant archosaurs, some muscles of the *M. transversospinalis* group insert on the dorsal margin of the spines; insertion is supposed to occur at the base of the spines in the low-spined synapsid *Sphenacodon* (see Huttenlocker et al., 2010). In *O. nigeriensis*, the apparent basal attachment of those muscles may be related to the hyperelongation of the neural spine. However, caution is due because the absences of Sharpey's fibers and superficial insertion marks could be caused by taphonomic factors and preparation and restoration as well. Ossified tendons have not been found along the vertebral column of the paratype and are reported as scarcely present in the holotype (Taquet 1976). Ossified tendons along the epiaxial skeleton are an ornithischian general feature, are usually abundant in iguanodontians and organized in a rhomboidal lattice structure (see, for example, Forster, 1990; Organ, 2006a, 2006b; Norman, 2011; Wang et al., 2010). They are associated with the subunits of *M. transversospinalis* (Organ, 2006a). Their scarce development in

Commented [PG17]: Again, be careful about taphonomy!

1675 *Ouranosaurus* also supports a peculiar development of the muscles of the *M. transversospinalis*
 1676 group in this dinosaur.
 1677 In iguanodontian ornithopods the dorsal neural spines are usually much shorter than those of the
 1678 theropods *Spinosaurus* and *Deinocheirus*. In *Iguanodon bernissartensis*, the taller neural spines are
 1679 2.43 times their centrum height in mid-dorsals (Prieto-Marquez, 2008). Neural spines are
 1680 proportionally taller in a few other ornithopod taxa, but never as in *Ouranosaurus*: the ratio
 1681 spine/centrum height is > 4.3 in *Morelladon beltrani* (see Gasulla et al., 2015), 4.5 in GPIT 1802/1-
 1682 7 (*Iguanodontia* indet.; Pereda-Suberbiola et al., 2011) and 4.18 in *Barbsoldia sicinskii* and
 1683 *Hypacrosaurus altispinus* (see Prieto-Marquez, 2008).
 1684 In *Ouranosaurus*, the 'sail' reaches its maximum height in the mid-proximal dorsals, then decreases
 1685 up to the sacrum to slightly increase again in the last sacral and first caudals, decreasing gradually
 1686 in the rest of the tail (a sinusoidal outline; Figs 1 and 5). In *Barbsoldia sicinskii* holotype, which
 1687 preserves the dorsal, the sacral and the proximal caudal vertebrae, the outline of the dorsal portion
 1688 of the 'sail' is nearly semicircular and only slightly asymmetrical; the sacral and caudal spines
 1689 reduce their height gradually moving caudally. Due to the fragmentary condition of the specimens,
 1690 the shape of the 'sail' is unknown in the other taxa mentioned above. The vertebral column of
 1691 *Hypacrosaurus* is still undescribed. The cast of a composite juvenile individual of *H. altispinus*
 1692 exhibited at the Royal Tyrrell Museum of Drumheller (Canada) shows a 'sail' where the mid-distal
 1693 dorsal, the sacral and the first caudal spines have the same height (the curvature of the 'sail' is
 1694 actually that of the vertebral column). If that assemblage is reliable, its 'sail' is unlike that of
 1695 *Ouranosaurus*. At the present state of knowledge, the combination of size and shape of the 'sail' of
 1696 *O. nigeriensis*, is therefore unique.
 1697 The only work dealing also with the function of iguanodontian 'sails' is that by Bailey (1997), who
 1698 supported a thermoregulation role.
 1699 The tall spines of *Ouranosaurus* were plausibly a support for a structure like a membrane or a hump
 1700 (Bailey, 1997). Of course, the definition of such a structure is hampered by the lack of preservation
 1701 of the soft tissues that were covering the spines. However, the presence of a keratinous covering
 1702 directly on the bone can be excluded, because of the absence of Sharpey's fibers in the middle and
 1703 apical portions of the neural spine (Huttenlocker et al., 2010). Vascularization is not particularly
 1704 dense in neural spines; this does not support a relationship between elongation and increase of
 1705 blood input through the bone for thermoregulatory (contra Bailey, 1997) or display purposes. A
 1706 thermoregulatory role of the 'sail' to keep high and constant the body temperature as in ectotherm
 1707 tetrapods would be unnecessary, if the relatively high growth rates observed in *Ouranosaurus* are
 1708 related with homeothermy. A social (display) role of the structure like that hypothesized for
 1709 *Spinosaurus* is possible but, obviously, it is speculative and cannot be tested.
 1710
 1711 CONCLUSIONS
 1712 MSNVE 3714, the Venice specimen of *Ouranosaurus nigeriensis* is the paratype of the species
 1713 GDF 381- MNHN, found in 1970 and collected in 1972 by a French team, although it lacks some of
 1714 the original bones (i.e., the left femur and the right coracoid), which were replaced by plaster
 1715 copies. The field map of the bones found in 1970 shows the presence of at least two individuals, the
 1716 most complete of which was the paratype. Probably, some skeletal elements from the second
 1717 individual and possibly other sources (i.e., femur GDF 302) were added to the paratype material to
 1718 complete the mounted skeleton for exhibit purposes. Portions of most of the original skeletal
 1719 elements have been reconstructed or covered by resin during the restoration process.
 1720 The Venice specimen shows some minor differences with the holotype that probably reflect
 1721 intraspecific variability. Its carpus appears to be badly reconstructed and the original manual
 1722 phalanges possibly replaced by other elements. Other differences are caused by mistakes in the
 1723 reconstruction of the vertebral column. Probably *O. nigeriensis* had 14 dorsal and one dorsosacral

vertebrae, 20 proximal caudal vertebrae, more than the supposed 17 mid-caudal vertebrae and a total caudal count much higher than that represented by the preserved caudal vertebrae. Puzzling are the morphological differences in the metacarpus. Based on histological analysis, (the first performed on *Ouranosaurus*, a fast growth rate is assumed for this taxon. The samples show features suggesting a sub-adult ontogenetic stage for the paratype and the other/s individual/s used to assemble MSNVE 3714. Immaturity is suggested also by unfusion and superficial texture of some skeletal elements. The study of MSNVE 3714 suggests also some thoughts about the way scientifically important specimens are managed in the process leading from their discovery to their storage or exhibit in a museum. Loss of scientific information should be avoided as much as it is possible. Often preparation for exhibition purposes reduces the scientific value of the specimen, ~~hiding~~ osteological features. Even worse, the scientific value of the specimen is often dramatically affected by the total absence or availability of documentation about the material prior and during preparation (including detailed information on the field provenance of each skeletal element, photographs, detailed field notes, detailed description of the preparation and restoration works and materials used in them; Dalla Vecchia et al., 2015). That process must be always under scientific control and documented. All necessary information should be attached to the specimen as the pedigree is attached to a purebred dog.

Commented [PG18]: In my opinion, you should delete this paragraph. All palaeontologists are well aware of this problem for a long time, this is really not a brand new discovery! Or, in other words, this is useless preachifying.

Supporting Information

Tab.1 - Measurements of the skeletal elements of MSNVE 3714, *Ouranosaurus nigeriensis*.

ACKNOWLEDGEMENTS

We would like to thanks dr. Mauro Bon, responsible for Research and Scientific Divulgateion of the Museo di Storia Naturale of Venice, for allowing to study the specimen under his care; Barbara Favaretto and the staff of the Museo di Storia Naturale of Venice for assistance during the study the specimen. This study was partially supported by the MSNVE. Dr. Mauro Bon provided access and historical information on MSNVE 3714. Prof. Philippe Taquet (MNHN) provided historical information on the paratype and the French expeditions in Gadoufaoua in 1970s; Dr. Ronan Allain (MNHN) sent us a copy of the field map of the paratype and allowed for its use. The Fondazione Giancarlo Ligabue (former Centro Studi e Ricerche Ligabue) gave us additional information about the Italian expeditions in Gadoufaoua during the 1970s. Dr. Karen Poole (Stony Brook University, New York) and Andrew McDonald (St. Louis Science Center, St. Louis) shared information on the codings of *Ouranosaurus* in their phylogenetic analyses of Ornithopoda. Donald Henderson (TMP), and Albert Prieto-Marquez (The Field Museum, Chicago) are kindly acknowledged for sharing pictures and information about comparative materials.

REFERENCES

- Bailey J B. 1997. Neural spine elongation in dinosaurs: Sailbacks or buffalo-backs? Journal of Paleontology 71(6):1124-1146.
- Bailleul A M, Scannella J B, Horner J R, Evans D C. 2016. Fusion patterns in the skulls of modern archosaurs reveal that sutures are ambiguous maturity indicators for the Dinosauria. PloS One 11(2):e0147687.
- Barreto C. 1997. Dinosaur growth plates and dinosaur bone growth. In: Wolberg DL, Stump E, Rosenberg G, editors. DINOfest international: Proceedings of a symposium held at Arizona State University. Philadelphia: The Academy of Natural Sciences. pp. 95-100.
- Barreto C, Albrecht RM, Bjorling DE, Horner JR, Wilsman NJ. 1993. Evidence of the growth plate and the growth of long bones in juvenile dinosaurs. Science 262(5142):2020-2022.

1773 - Bennett SC. 1993. The ontogeny of *Pteranodon* and other pterosaurs. *Paleobiology*, 19(01): 92-
1774 106.

1775 - Benton MJ, Csiki Z, Grigorescu D, Redelstorff R, Sander PM, Stein K, Weishampel DB. 2010.
1776 Dinosaurs and the island rule: The dwarfed dinosaurs from Hateg Island. *Palaeogeography*,
1777 *Palaeoclimatology*, *Palaeoecology* 293(3):438-454.

1778 - Bertozzo F. 2012. *Ouranosaurus nigeriensis* (Ornithischia, Ornithopoda) esposto al Museo di
1779 Storia Naturale di Venezia: storia del reperto e revisione anatomica. Bachelor Thesis. University of
1780 Bologna.

1781 - Boccardi V, Bottazzi C. 1978. Il cimitero dei dinosauri: il racconto di una straordinaria scoperta
1782 nel deserto del Ténéré. SugarCo Edizioni s.r.l., Milano.

1783 - Brochu CA. 1996. Closure of neurocentral sutures during crocodilian ontogeny: implications for
1784 maturity assessment in fossil archosaurs. *Journal of Vertebrate Paleontology* 16:49-62.

1785 - Brown B. 1916. *Corythosaurus casuarius*: skeleton, musculature and epidermis. Order of the
1786 Trustees, American Museum of Natural History.

1787 - Chinsamy A. 1995. Ontogenetic changes in the bone histology of the Late Jurassic ornithopod
1788 *Dryosaurus lettowvorbecki*. *Journal of Vertebrate Paleontology*, 15(1):96-104.

1789 - Chinsamy A. 2005. The microstructure of dinosaur bone: deciphering biology with fine-scale
1790 techniques. John Hopkins University Press.

1791 - Chinsamy A, Rich T, Vickers-Rich P. 1998. Polar dinosaur bone histology. *Journal of Vertebrate*
1792 *Paleontology* 18(2):385-390.

1793 - Cooper LN, Lee AH, Taper ML, Horner JR. 2008. Relative growth rates of predator and prey
1794 dinosaurs reflect effects of predation. *Proceedings of the Royal Society of London B* 275:2609-
1795 2615.

1796 - Currie PJ, Padian K. 1997. *Encyclopedia of dinosaurs*. San Diego: Academic Press.

1797 - Dalla Vecchia FM. 2009. *Tethyshadros insularis*, a new hadrosauroid dinosaur (Ornithischia) from
1798 the Upper Cretaceous of Italy. *Journal of Vertebrate Paleontology*, 29(4):1100-1116.

1799 - Dalla Vecchia FM, Bosch R, Fortuny J, Galobart À. 2015. The pterodactyloid pterosaur from the
1800 Lower Cretaceous of Brazil at the CosmoCaixa Science Museum (Barcelona, Spain). *Historical*
1801 *Biology* 27(6):729-748.

1802 - Dean-Carpentier M. 2008. New Information on the teeth and jaw bones of the rare hadrosauriform
1803 *Ouranosaurus nigeriensis*. *Journal of Vertebrate Paleontology* 28(3)A.

1804 - Erickson GM. 2005. Assessing dinosaur growth patterns: a microscopic revolution. *Trends in*
1805 *Ecology & Evolution* 20(12):677-684.

1806 - Erickson GM, Makovicky PJ, Currie PJ, Norell MA, Yerby SA, Brochu CA. 2004. Gigantism and
1807 comparative life-history parameters of tyrannosaurid dinosaurs. *Nature* 430 (7001):772-775.

1808 - Forster CA. 1990. The postcranial skeleton of the ornithopod dinosaur *Tenontosaurus tilletti*.
1809 *Journal of Vertebrate Paleontology* 10:273-294.

1810 - Gasulla JM, Escaso F, Narváez I, Ortega F, Sanz JL. 2015. A New Sail-Backed Styracosternan
1811 (Dinosauria: Ornithopoda) from the Early Cretaceous of Morella, Spain. *PloS One*,
1812 10(12):e0144167.

1813 - Godefroit P, Dong ZM, Bultynck P, Li H, Feng L. 1998. New *Bactrosaurus* (Dinosauria:
1814 Hadrosauroidae) material from Iren Dabasu (Inner Mongolia, PR China). *Bulletin de l'Institut*
1815 *Royal des Sciences Naturelles de Belgique, Sciences de la Terra*, 68(Supplement):3-70.

1816 - Hone DW. 2012. Variation in the tail length of non-avian dinosaurs. *Journal of Vertebrate*
1817 *Paleontology* 32:1082-1089.

1818 - Horner JR, de Ricqlès A, Padian K. 1999. Variation in dinosaur skeletochronology indicators:
1819 implications for age assessment and physiology. *Paleobiology* 25:295-304.

Formatted: French (France)

1820 - Horner JR, de Ricqlès A, Padian K. 2000. Long bone histology of the hadrosaurid dinosaur
 1821 *Maiaasaura peeblesorum*: growth dynamics and physiology based on an ontogenetic series of
 1822 skeletal elements. *Journal of Vertebrate Paleontology* 20(1):115-129.
 1823 - Horner JR, Padian K, de Ricqlès A. 2001. Comparative osteohistology of some embryonic and
 1824 perinatal archosaurs: developmental and behavioral implications for dinosaurs. *Paleobiology*
 1825 27(01):39-58.
 1826 - Horner JR, de Ricqlès A, Padian K, Scheetz RD. 2009. Comparative long bone histology and
 1827 growth of the “hypsilophodontid” dinosaurs *Orodromeus makelai*, *Dryosaurus altus*, and
 1828 *Tenontosaurus tilletii* (Ornithischia: Euornithopoda). *Journal of Vertebrate Paleontology*
 1829 29(3):734-747.
 1830 - Hübner TR. 2012. Bone histology in *Dysalotosaurus lettowvorbecki* (Ornithischia:
 1831 Iguanodontia)—variation, growth, and implications. *PLoS One* 7:e29958.
 1832 - Huttenlocker AK, Rega E, Sumida SS. 2010. Comparative anatomy and osteohistology of
 1833 hyperelongate neural spines in the spheonacodontids *Sphenacodon* and *Dimetrodon* (Amniota:
 1834 Synapsida). *Journal of Morphology* 271:1407-1421.
 1835 - Ibrahim N, Sereno PC, Dal Sasso C, Maganuco S, Fabbri M, Martill DM, Zouhri S, Myhrvold N,
 1836 Iurino DA. 2014. Semiaquatic adaptations in a giant predatory dinosaur. *Science* 345:1613-1616.
 1837 - Irmis RB. 2007. Axial skeleton ontogeny in the Parasuchia (Archosauria: Pseudosuchia) and its
 1838 implications for ontogenetic determination in archosaurs. *Journal of Vertebrate Paleontology*
 1839 27:350-361.
 1840 - Lee Y-N, Barsbold R, Currie PJ, Kobayashi Y, Lee H-J, Godefroit P, Escuillié F, Chinzorig T.
 1841 2014. Resolving the long-standing enigmas of a giant ornithomimosaur *Deinocheirus mirificus*.
 1842 *Nature* 515(7526):257-260.
 1843 - Ligabue G, Pinna G, Azzaroli A, Taquet P. 1972. I dinosauri del Ténéré. Italia: Longanesi & C.
 1844 - Ligabue M, Rossi-Osmida G. 1975. Gadoufaoua: Uadi dei dinosauri. Relazione della spedizione
 1845 scientifica “Ligabue” - Sahara 1973. Italia: Ed. Armona.
 1846 - Le Loeuff J, Läng E, Cavin L, Buffetaut E. 2012. Between Tendaguru and Bahariya: on the age of
 1847 the Early Cretaceous dinosaur sites from the Continental Intercalaire and other African formations.
 1848 *Journal of Stratigraphy* 36:486-502.
 1849 - McDonald AT, Barrett PM, Chapman SD. 2010a. A new basal iguanodont (Dinosauria:
 1850 Ornithischia) from the Wealden (Lower Cretaceous) of England. *Zootaxa*, 2569:1-43.
 1851 - McDonald AT, Wolfe DG, Kirkland JJ. 2010b. A new basal hadrosauroid (Dinosauria:
 1852 Ornithopoda) from the Turonian of New Mexico, *Journal of Vertebrate Paleontology* 30(3):799-
 1853 812.
 1854 - McDonald AT, Espilez E, Mampel L, Kirkland JJ, Alcalá L. 2012. An unusual new basal
 1855 iguanodont (Dinosauria: Ornithopoda) from the Lower Cretaceous of Teruel, Spain. *Zootaxa*
 1856 3595:61-76
 1857 - McDonald AT, Maidment SCR, Barrett PM, You H-l, Dodson P. 2014. Osteology of the Basal
 1858 Hadrosauroid *Equijubus normani* (Dinosauria, Ornithopoda) from the Early Cretaceous of China. In
 1859 Eberth DA & Evans DC, editors, *Hadrosaurs*, pp. 44-72, Indiana University Press, Bloomington,
 1860 Indiana.
 1861 - Nopcsa F. 1933. 11. On the Histology of-the Ribs in Immature and Half-grown Trachodont
 1862 Dinosaurs. *Proceedings of the Zoological Society of London* 103:221-226.
 1863 - Norman DB. 1980. On the ornithischian dinosaur *Iguanodon bernissartensis* from the Lower
 1864 Cretaceous of Bernissart (Belgium). *Mémoire de l'Institut Royal des Sciences Naturelles de*
 1865 *Belgique* 178:1-105.
 1866 - Norman DB. 1986. On the anatomy of *Iguanodon atherfieldensis* (Ornithischia: Ornithopoda).
 1867 *Bulletin-Institut royal des sciences naturelles de Belgique. Sciences de la terre* 56:281-372.

Formatted: French (France)

Formatted: French (France)

1868 - Norman DB. 2004. Basal Iguanodontia; pp. 413-437 in Weishampel DB, Dodson P, Osmólska H,
1869 editors, The Dinosauria, 2nd edition, University California Press, Berkeley, CA, USA, pp. 413-437.
1870 - Norman DB. 2011. On the osteology of the lower Wealden (Valanginian) ornithopod *Barilium*
1871 *dawsoni* (Iguanodontia: Styracosterna). Special Papers in Palaeontology 86:165-194.
1872 - Norman DB. 2015. On the history, osteology, and systematic position of the Wealden (Hastings
1873 Group) dinosaur *Hypselospinus fittoni* (Iguanodontia: Styracosterna). Zoological Journal of the
1874 Linnean Society 173:92-189.
1875 - Organ CL. 2006a. Thoracic epaxial muscles in living archosaurs and ornithopod dinosaurs. The
1876 Anatomical Record (Part A) 288:782-793.
1877 - Organ CL. 2006b. Biomechanics of ossified tendons in ornithopod dinosaurs. Paleobiology
1878 32(4):652-665.
1879 - Ósi A, Prondvai E, Butler R, Weishampel DB. 2012. Phylogeny, Histology and Inferred Body
1880 Size Evolution in a New Rhabdodontid Dinosaur from the Late Cretaceous of Hungary. PLoS One
1881 7(9):e44318. doi:10.1371/journal.pone.0044318
1882 - Parks WA. 1920. The osteology of the trachodont dinosaur *Kritosaurus incurvimanus*: University
1883 of Toronto. Studies in Geology(11): 1-74.
1884 - Pereda-Suberbiola X, Ruiz-Omenaca JJ, Fernández-Baldor FT, Maisch MW, Huerta P, Contreras,
1885 R, Izquierdo LA, Huerta MD, Montero VU, Welle J. 2011. A tall-spined ornithopod dinosaur from
1886 the Early Cretaceous of Salas de los Infantes (Burgos, Spain). Comptes Rendus Palevol, 10(7):551-
1887 558.
1888 - Persons WS IV, Currie PJ. 2011. The Tail of *Tyrannosaurus*: Reassessing the Size and
1889 Locomotive Importance of the *M. caudofemoralis* in Non-Avian Theropods. The Anatomical
1890 Record 294:119-131.
1891 - Prieto-Marquez A. 2001. Osteology and variation of *Brachylophosaurus canadensis* (Dinosauria,
1892 Hadrosauridae) from the Upper Cretaceous Judith River formation of Montana. Doctoral
1893 dissertation, Montana State University-Bozeman, College of Letters & Science.
1894 - Prieto-Marquez A. 2008. Phylogeny and historical biogeography of hadrosaurid dinosaurs. PhD
1895 Thesis. Florida State University.
1896 - Rasmussen ME. 1998. Notes on the morphology and the orientation of the forelimb of
1897 *Ouranosaurus nigeriensis*. Oryctos 1:127-130.
1898 - Reid REH. 1984. The histology of dinosaurian bone and its possible bearing on dinosaurian
1899 physiology. In Symposia of the Zoological Society of London 52:629-663. Cambridge University
1900 Press.
1901 - Reid REH. 1985. On supposed Haversian bone from the hadrosaur *Anatosaurus*, and the nature of
1902 compact bone in dinosaurs. Journal of Paleontology 59(1):140-148.
1903 - de Ricqlès A, Godefroit P, Yans J. 2012. Histological Assessment of Vertebrate Remains in the
1904 2003 Bernissart Drill. In Bernissart Dinosaurs and Early Cretaceous Terrestrial Ecosystems,
1905 Godefroit P, editor, Indiana Indiana University Press, 137-146.
1906 - Rossi-Osmida G. 2005. Per piste dimenticate. Scritti di Giancarlo Ligabue. Il Punto Edizioni,
1907 Trebaseleghe.
1908 - Sereno PC. 1986. Phylogeny of the bird-hipped dinosaurs (Order Ornithischia). National
1909 Geographic Research, 2(2):234-256.
1910 - Sereno PC, Beck AL, Dutheil DB, Larsson HCE, Lyon GH, Moussa B, Sadleir RW, Sidor CA,
1911 Varricchio DD, Wilson GP, Wilson JA. 1999. Cretaceous Sauropods from the Sahara and the
1912 uneven rate of skeletal evolution among dinosaurs. Science 286:1342-1347.
1913 - Stein K, Sander PM. 2009. Histological core drilling: a less destructive method for studying bone
1914 histology. In Brown MA, Kane JF, Parker WG, editors,. Methods in Fossil Preparation, Proceedings
1915 of the First Annual Fossil Preparation and Collections Symposium, 69-80.
1916 - Stromer E. 1915. Ergebnisse der Forschungsreisen Prof. E. Stromers in den Wüsten Ägyptens. II

Formatted: French (France)

1917 Wirbeltierreste der Baharije-Stufe (unterstes Cenoman). 3. Das original des Theropoden
 1918 *Spinosaurus aegyptiacus* n. gen., n. sp. Abhandlungen der Königlich Bayerischen Akademie der
 1919 Wissenschaften Mathematisch- physikalische Klasse 28:1-31.
 1920 - Taquet P. 1970. Sur le gisement de Dinosauriens et de Crocodiliens de Gadoufaoua (République
 1921 du Niger). Comptes Rendus de l'Académie des Sciences à Paris, Série D: 271:38-40.
 1922 - Taquet P. 1976. Géologie et Paléontologie du Gisement de Gadoufaoua (Aptien du Niger).
 1923 Editions du Centre National de la Recherche Scientifique, Paris, 191 pp.
 1924 - Taquet P. 1998. Dinosaur Impressions: Postcards from a Paleontologist. Translated by Padian K.
 1925 Cambridge University Press, Cambridge, 244 pp.
 1926 - Taquet P. 2012. The African Cousins of the European Iguanodontids. In: Godefroit P, editor,
 1927 Bernissart Dinosaurs and Early Cretaceous Terrestrial Ecosystems, 283-292, Indiana University
 1928 Press, Bloomington.
 1929 - Taquet P, Russell DA. 1999. A massively-constructed iguanodont from Gadoufaoua, Lower
 1930 Cretaceous of Niger. Annales du Paléontologie 85:85-96.
 1931 - Tsuihiji T. 2005. Homologies of the transversospinalis muscles in the anterior presacral region of
 1932 Sauria (crown Diapsida). Journal of Morphology, 263(2):151-178.
 1933 - Wang X, Pan R, Butler RJ, Barrett PM. 2010. The postcranial skeleton of the iguanodontian
 1934 ornithomimid *Jinzhousaurus yangi* from the Lower Cretaceous Yixian Formation of western Liaoning,
 1935 China. Earth and Environmental Science Transactions of the Royal Society of Edinburgh 101:135–
 1936 159.
 1937 - Werning S. 2012. The ontogenetic osteohistology of *Tenontosaurus tilletti*. PLoS One
 1938 7(3):e33539.
 1939 - Winkler DA. 1994. Aspects of growth in the Early Cretaceous Proctor Lake ornithomimid. Journal of
 1940 Vertebrate Paleontology 14:53A.
 1941 - Woodward HN, Rich TH, Chinsamy A, Vickers-Rich P. 2011. Growth dynamics of Australia's
 1942 polar dinosaurs. PLoS One 6(8):e23339.
 1943 - Woodward HN, Fowler EAF, Farlow JO, Horner JR. 2015. *Maiasaura*, a model organism for
 1944 extinct vertebrate population biology: a large sample statistical assessment of growth dynamics and
 1945 survivorship. Paleobiology, 41(04):503-527.
 1946 - You H, Li D, Liu W. 2011. A new hadrosauriform dinosaur from the Early Cretaceous of Gansu
 1947 Province, China. Acta Geologica Sinica (English Edition) 85:51–57.
 1948 - Zheng W, Jin X, Shibata M, Azuma Y, Yu F. 2012. A new ornithomimid dinosaur from the
 1949 Cretaceous Liangtoutang Formation of Tiantai, Zhejiang Province, China. Cretaceous Research
 1950 34:208–219.