

Waterbirds in the Baltic Sea are changing wintering patterns because of climate warming: A study of selected species in the Odra River Estuary - a key European wintering site (#15149)

1

Second revision

Please read the **Important notes** below, the **Review guidance** on page 2 and our **Standout reviewing tips** on page 3. When ready [submit online](#). The manuscript starts on page 4.

Important notes

Editor

James Roper

Files

1 Tracked changes manuscript(s)

1 Rebuttal letter(s)

2 Figure file(s)

4 Table file(s)

3 Other file(s)

Please visit the overview page to [download and review](#) the files not included in this review PDF.

Declarations

Involves vertebrate animals.



Please read in full before you begin

How to review

When ready [submit your review online](#). The review form is divided into 5 sections. Please consider these when composing your review:

- 1. BASIC REPORTING**
- 2. EXPERIMENTAL DESIGN**
- 3. VALIDITY OF THE FINDINGS**
4. General comments
5. Confidential notes to the editor

 You can also annotate this PDF and upload it as part of your review

To finish, enter your editorial recommendation (accept, revise or reject) and submit.

BASIC REPORTING

-  Clear, unambiguous, professional English language used throughout.
-  Intro & background to show context. Literature well referenced & relevant.
-  Structure conforms to [PeerJ standards](#), discipline norm, or improved for clarity.
-  Figures are relevant, high quality, well labelled & described.
-  Raw data supplied (see [PeerJ policy](#)).

EXPERIMENTAL DESIGN

-  Original primary research within [Scope of the journal](#).
-  Research question well defined, relevant & meaningful. It is stated how the research fills an identified knowledge gap.
-  Rigorous investigation performed to a high technical & ethical standard.
-  Methods described with sufficient detail & information to replicate.

VALIDITY OF THE FINDINGS

-  Impact and novelty not assessed. Negative/inconclusive results accepted. *Meaningful* replication encouraged where rationale & benefit to literature is clearly stated.
-  Data is robust, statistically sound, & controlled.
-  Conclusions are well stated, linked to original research question & limited to supporting results.
-  Speculation is welcome, but should be identified as such.

The above is the editorial criteria summary. To view in full visit <https://peerj.com/about/editorial-criteria/>

7 Standout reviewing tips

3



The best reviewers use these techniques

Tip

Example

Support criticisms with evidence from the text or from other sources

Smith et al (J of Methodology, 2005, V3, pp 123) have shown that the analysis you use in Lines 241-250 is not the most appropriate for this situation. Please explain why you used this method.

Give specific suggestions on how to improve the manuscript

Your introduction needs more detail. I suggest that you improve the description at lines 57- 86 to provide more justification for your study (specifically, you should expand upon the knowledge gap being filled).

Comment on language and grammar issues

The English language should be improved to ensure that your international audience can clearly understand your text. I suggest that you have a native English speaking colleague review your manuscript. Some examples where the language could be improved include lines 23, 77, 121, 128 - the current phrasing makes comprehension difficult.

Organize by importance of the issues, and number your points

1. Your most important issue
2. The next most important item
3. ...
4. The least important points

Give specific suggestions on how to improve the manuscript

Line 56: Note that experimental data on sprawling animals needs to be updated. Line 66: Please consider exchanging "modern" with "cursorial".

Please provide constructive criticism, and avoid personal opinions

I thank you for providing the raw data, however your supplemental files need more descriptive metadata identifiers to be useful to future readers. Although your results are compelling, the data analysis should be improved in the following ways: AA, BB, CC

Comment on strengths (as well as weaknesses) of the manuscript

I commend the authors for their extensive data set, compiled over many years of detailed fieldwork. In addition, the manuscript is clearly written in professional, unambiguous language. If there is a weakness, it is in the statistical analysis (as I have noted above) which should be improved upon before Acceptance.

Waterbirds in the Baltic Sea are changing wintering patterns because of climate warming. A study of selected species in the Odra River Estuary - a key European wintering site.

Dominik Marchowski ^{Corresp., 1}, Łukasz Jankowiak ¹, Dariusz Wysocki ¹, Łukasz Ławicki ², Józef Girjatowicz ³

¹ Department of Vertebrate Zoology and Anthropology, Institute for Research on Biodiversity, Faculty of Biology, University of Szczecin, Szczecin, Poland

² West- Pomeranian Nature Society, Szczecin, Polska

³ Hydrography and Water Management Unit, Faculty of Earth Science, University of Szczecin, Szczecin, Poland

Corresponding Author: Dominik Marchowski
Email address: marchowskid@gmail.com

Some species of birds react to climate change, by wintering in places closer to their breeding areas. We investigated the responses of two groups of waterbirds to factors associated with climate change. The Odra River Estuary (SW Baltic Sea) is of key importance for wintering waterfowl. The most abundant birds here comprise two ecological groups: benthic feeders and fish feeders. We showed that numbers of all benthivores, but not piscivores, were negatively correlated with the presence of ice. We predicted that, with ongoing global warming, this area would increase in importance for benthic feeders and decrease for fish feeders. The maximum range of ice cover in the Baltic Sea has a weak and negative effect on both groups of birds. Five of the seven target species are benthivores (Greater Scaup *Aythya marila*, Tufted Duck *A. fuligula*, Common Pochard *A. ferina*, Common Goldeneye *Bucephala clangula* and Eurasian Coot *Fulica atra*), and two are piscivores (Smew *Mergellus albellus* and Goosander *Mergus merganser*). Local changes at the level of particular species vary for different reasons. The local decline of Common Pochard may be a reflection of the species' global decline. Climate change may be responsible for some of the local changes in the study area; thereby increasing the importance of the area for Greater Scaup and Tufted Duck while reducing it for Smew.

Waterbirds in the Baltic Sea are changing wintering patterns because of climate warming. A study of selected species in the Odra River Estuary – a key European wintering site.

Dominik Marchowski^{1,*}, Łukasz Jankowiak¹, Dariusz Wysocki¹, Łukasz Ławicki², Józef Girjatowicz³

¹Department of Vertebrate Zoology and Anthropology, Institute for Research on Biodiversity, Faculty of Biology, University of Szczecin, Szczecin, Poland

²West-Pomeranian Nature Society, Szczecin, Poland

³Hydrography and Water Management Unit, Faculty of Earth Science, University of Szczecin, Szczecin, Poland

*Corresponding author: Dominik Marchowski marchowskid@gmail.com

19

20 Abstract

21 Some species of birds react to climate change, by wintering in places closer to their
 22 breeding areas. We investigated the responses of two groups of waterbirds to factors associated
 23 with climate change. The Odra River Estuary (SW Baltic Sea) is of key importance for wintering
 24 waterfowl. The most abundant birds here comprise two ecological groups: benthic feeders and
 25 fish feeders. We showed that numbers of all benthivores, but not piscivores, were negatively
 26 correlated with the presence of ice. We predicted that, with ongoing global warming, this area
 27 would increase in importance for benthic feeders and decrease for fish feeders. The maximum
 28 range of ice cover in the Baltic Sea has a weak and negative effect on both groups of birds. Five
 29 of the seven target species are benthivores (Greater Scaup *Aythya marila*, Tufted Duck *A.*
 30 *fuligula*, Common Pochard *A. ferina*, Common Goldeneye *Bucephala clangula* and Eurasian
 31 Coot *Fulica atra*), and two are piscivores (Smew *Mergellus albellus* and Goosander *Mergus*
 32 *mergamser*). Local changes at the level of particular species vary for different reasons. The local
 33 decline of Common Pochard may be a reflection of the species' global decline. Climate change
 34 may be responsible for some of the local changes in the study area; thereby increasing the
 35 importance of the area for Greater Scaup and Tufted Duck while reducing it for Smew.

36

37 Introduction

38 Migration distance has declined in several species of aquatic (and other) birds as a result
 39 of climate change (Musil et al. 2011; Lehikoinen et al. 2013; Meller 2016). The distances that

birds migrate from their breeding areas in northern and eastern Europe to their central European wintering areas are shorter during mild winters (Lehikoinen et al. 2013; Pavón-Jordan et al. 2015); conversely birds may change their wintering sites to warmer regions during colder periods because they may perceive local manifestations of ~~large scale atmospheric features~~ (Newton 2008). Reducing migration distance can provide several benefits associated with earlier arrival at the breeding grounds and greater survival (Coppack and Both 2002; Jankowiak et al. 2015a; 2015b). Food resources of wintering sites may also influence migration decisions (Cresswell 2014; Aharon-Rotman et al. 2016). Although winter site fidelity is ~~very high~~ among waterfowl (Newton 2008), this can change in response to weather, habitat and competition (Cresswell 2014). ~~The changing of winter sites should thus be seen as~~ trade-off between the costs of finding a new site and the benefits it offers (Aharon-Rotman et al. 2016). At sub-zero temperatures, shallow waters freeze over; ~~birds therefore~~ expend more time and energy searching for food in deeper waters, ~~with obvious consequences for their energy balance.~~ Shallow waters of offshore lagoons create ideal conditions for three functional groups of waterbirds: piscivores, herbivores and benthivores. Our study area, the Odra River Estuary (ORE), accommodates large numbers of waterbirds because of food resources (Marchowski et al. 2015; Marchowski et al. 2016). Two groups of waterbirds – ~~fish feeders and benthic feeders~~ – are among those most commonly wintering here. Because of its position on the 0°C isotherm in winter (van Erden and de Leeuw 2010), the study area is subject to significant variations in bird habitat conditions. Yet even relatively small variations in temperature, causing ice cover to form or disappear, can lead to the displacement of waterbirds. Changes in abundance and community structure in the ORE may reflect the impact of climate change. Analysis of the dates of the appearance of ice-related phenomena in the Szczecin Lagoon and of their frequency over time reveals a distinct pattern

endorsing recently observed trends in climate warming (Girjatowicz 2011). In this paper we test whether abundance of some species in the ORE is changing in response to climate warming. We predict that benthic feeding birds will be more sensitive and fish feeding birds less sensitive to the factor related to climate, namely ice cover. If our prediction is correct, elevated temperatures and the correspondingly shorter period of ice cover should provide relatively better conditions for benthic feeders, which should increase in abundance. Feeding areas where sedentary mussels are abundant tend to be in shallow waters (Marchowski et al. 2015), so surface ice cover reduces food availability. Fish, on the other hand, remain available even if the ice cover is considerable, since unfrozen, deeper, areas, located further from the shore, may still be rich in fish. During cold winters the water in the ORE never freezes over entirely: even during periods of sustained below-zero temperatures, patches of water remain free of ice, where large aggregations of fish feeders (Smew and Goosander) have been observed (Kaliciuk et al. 2003; Czeraszewicz et al. 2004; Marchowski and Ławicki 2011; Guentzel et al. 2012; Marchowski and Ławicki 2012; Marchowski et al. 2013).

Study area

The study area lies in the south-western Baltic Sea and forms the Polish part of the Odra River Estuary system, which includes the Great Lagoon (the Polish part of the Szczecin Lagoon), Świna Backward Delta, Kamień Lagoon, Dziwna Strait and Lake Dąbie with a total area of 522.58 km² (Fig. 1). The whole area has been designated as four interconnected Important Bird Areas (IBA) and also a Natura 2000 area (Wilk et al. 2010). The average and maximum depths of the estuary are 3.8 and 8.5 m, respectively; the dredged shipping lane passing through the

85 estuary from the Baltic Sea to the port of Szczecin is 10.5 m deep (Radziejewska and
 86 Schernewski 2008). ~~The~~ waters of the Szczecin Lagoon, Kamień Lagoon and Lake Dąbie are
 87 brackish. ~~The~~ salinity in the central part of the estuary varies from 0.3 psu to 4.5 psu (mean = 1.4
 88 psu) and declines with increasing distance from the sea (Radziejewska and Schernewski 2008).
 89 ~~Periodic backflows of water from the Pomeranian Bay (salinity 7 psu) take place through the~~
 90 ~~Świna Strait and, to a lesser extent, through the Dziwna and Peene Straits (the latter situated in~~
 91 ~~the German part of the ORE). The~~ average winter temperature is 0.3° C (Weatherbase 2016). The
 92 ORE is subject to ~~strong anthropogenic pressure manifested by~~ high levels of eutrophication
 93 (Radziejewska and Schernewski 2008). ~~The~~ communities of benthic organisms are typical of
 94 freshwater bodies, and the fauna includes large populations of zebra mussels *Dreissena*
 95 *polymorpha*, which were introduced in the mid-19th century. By the 1960s, the biomass of zebra
 96 mussels in the Szczecin (Great) Lagoon was estimated at 110 000 metric tons (Wiktor 1969,
 97 Wolnomiejski and Woźniczka 2008). ~~At present, this appears to be fairly stable; in the early~~
 98 ~~2000s the estimated biomass was 94 280 metric tons (Marchowski et al. 2015).~~ The distribution
 99 of the zebra mussel is extremely uneven (see the map in Marchowski et al. 2015). The average
 100 density of the zebra mussel in the ORE is 0.18 kg /m², but the vast majority ~~of these resources~~
 101 occupies around 10% of the ~~entire water body bed~~, where the mean density is 2.05 kg/m²
 102 (Stańczykowska et al. 2010). ~~The fish consist~~ mainly ~~of~~ freshwater species such as roach *Rutilus*
 103 *rutilus*, bream *Abramis brama*, pike *Esox lucius*, perch *Perca fluviatilis* and ruff *Gymnocephalus*
 104 *cernua*; there are also anadromous fish including smelt *Osmerus eperlanus* and occasionally
 105 herring *Clupea harengus* among others (Wolnomiejski and Witek 2013).

106

107 **Methods**

108 **Bird censusing**

109 Our study covers two functional groups of waterbirds: benthivores – Greater Scaup
 110 (*Aythya marila* – hereafter Scaup), Tufted Duck (*A. fuligula*), Common Pochard (*A. ferina* –
 111 hereafter Pochard), Common Goldeneye (*Bucephala clangula* – hereafter Goldeneye) and
 112 Eurasian Coot (*Fulica atra* – hereafter Coot); piscivores – Smew *Mergellus albellus* and
 113 Goosander *Mergus merganser*. ~~The study site is known to regularly host large numbers of the~~
 114 ~~biogeographic populations of the above species~~ (Kaliciuk et al. 2003; Czeraszewicz et al. 2004;
 115 Wilk et al. 2010; Marchowski and Ławicki 2011; Guentzel et al. 2012; Marchowski and Ławicki
 116 2012; Marchowski et al. 2013) ~~and meets the criteria for classification as a key site for them~~
 117 (Wilk et al. 2010). ~~Here, by biogeographic population we mean that part of the global population~~
 118 ~~associated with a specific flyway region~~. These subpopulations are: Pochard – north-east Europe
 119 / north-west Europe; Tufted Duck – north-west Europe (wintering); Scaup – northern Europe /
 120 western Europe; Goldeneye – north-west and central Europe; Smew – north-west and central
 121 Europe (wintering); Goosander – north-west and central Europe (wintering); Coot – north-west
 122 Europe (wintering) (Wetlands International 2016).

123 Censuses were conducted using standard methods for non-breeding season waterbird
 124 counts (Komdeur et al. 1992; Wetlands International 2010). Birds were counted during 17
 125 seasons (1991/1992 to 1993/1994 and 2001/2002 to 2015/2016) during the migration and
 126 wintering periods between November and April. From 1991/1992 to 1993/1994 three censuses
 127 were carried out per season in November, January, and March or April; there was one midwinter
 128 count in January in 2001/2002. Altogether we analysed the results of 44 counts. Most counts
 129 were done on foot. Each observer was equipped with 10x binoculars and tripod-mounted spotting
 130 scopes. Observers walked along the same routes, and the same counting method was used during

every census every year. Additionally, fourteen aerial counts were made at an average speed of about 100 km/h and an altitude of about 80 m above the water (see supplementary materials – S1 Table for the method of data collection: aerial or ground). In the early 1990s counts were aerial, whereas in 2009-2015 parallel aerial and ground counts were carried out (to compare methods). In ice-free conditions the species covered in this study can be assigned to a group with just a small error between methods (<6%), one species – Coot had a moderate error (16%), the ground method estimated greater numbers than the aerial one. During periods with more than 70% ice cover, abundance from the air was greater than abundance from the ground (Dominik Marchowski pers. com.). The count method was treated as a random effect in the model. The detailed methodology and results of the counts are given elsewhere (Meissner and Kozakiewicz 1992; Meissner et al 1994; Kaliciuk et al. 2003; Czeraszewicz et al. 2004; Marchowski and Ławicki 2011; Guentzel et al. 2012; Marchowski and Ławicki 2012; Marchowski et al. 2013). Where large numbers of unidentified *Aythya* species were counted – 26 000 ducks in November 2009, 13 000 in November 2010, 6 000 in January 2012, 3 300 in March 2012 and 13 500 in November 2015 – they were estimated to be in the ratio of 1:0.8 (scaup:tufted) based on observations in other studies. ~~This research involved observations of birds from a distance, which do not disturb the birds. In Poland, such studies do not need special permission or approval.~~

Statistical analysis

The dependent variable was the percentage occurrence of a given species in relation to the total estimated population size in a given year. This approach was taken because the population sizes of the species covered by our study follow different trends. For example, the population of Scaup is decreasing, that of Smew is increasing and that of Goosander is more or

154 less stable (Nagy et al. 2014; Wetland International 2016). Thus, if we showed the trend of
 155 absolute numbers in our area, the resulting error would be the larger, the greater the changes in
 156 the size of the entire population. Therefore, we indicate the numbers of a species by means of a
 157 coefficient calculated as the percentage of the biogeographic population present in the study area
 158 during a particular count. We obtained ~~the data relating to the biogeographic populations~~ from
 159 1992 to 2012 from Nagy et al. (2014); for the period 2013-2016 we used the flat trend calculated
 160 by Nagy et al. (2014) (Table 1). Initially, we placed the different species in ecological groups.
 161 The benthivores (denoted by B) included Scaup, Tufted Duck, Pochard, Goldeneye and Coot,
 162 and the piscivores (P) contained Smew and Goosander. We used the minimum temperatures
 163 averaged over the 15 days leading up to the count day. The climate data were obtained from the
 164 Szczecin weather station (53.395 N, 14.6225 E, <http://tutiempo.net>). Another climate covariate
 165 was ice cover in the study area; data relating to this were published by the Polish Institute of
 166 Meteorology and Water Management. These data are from the observation point at Miroszewo
 167 on the shore of the Szczecin Lagoon (53.734 N, 14.331 E, <http://www.imgw.pl/>). We compared
 168 the number of days with 100% ice cover in the period from 0 to 15 days prior to the bird counts.
 169 The ice cover of 100% refers specifically to the Miroszewo observation point. This estimate is a
 170 good approximation for the region. In practice, however, the ORE is never completely covered
 171 by ice (Girjatowicz 1991; 2005; see the Discussion for an explanation) and birds are still present
 172 in such conditions. We also utilized the maximum ice extent in the Baltic Sea (max ice) (data
 173 obtained from the website of The European Environment Agency (EEA 2017)). Apart from
 174 climatic variables, we also wanted to test the changes in species occurrence during the survey
 175 years, so we used season as covariate. Prior to the final analysis, we checked ~~the~~
 176 multicollinearity ~~correlation~~ between the above variables using the variation inflation factor

(VIF). ~~If VIFs were high, the relevant variables were excluded from the analysis. The~~ VIFs of all variables were in acceptable limits, minimal temperatures (VIF = 2.1), max ice (VIF = 1.03), ice cover (VIF = 2.07) and season (VIF = 1.04). However, we found a moderate linear significant relationship between ~~minimal~~ temperature and ice cover ($r = 0.52$, $p < 0.001$) and after exclusion of minimal temperature VIF showed no multicollinearity issue between variables – ice cover (VIF = 1.04), max ice (VIF = 1.03), season (VIF = 1.03) – and these were used in the subsequent analyses. We used a general linear mixed model (GLMM) to test the hypothesis relating to the different patterns of occurrence of benthivores and piscivores in the ORE. The percentage of the entire biogeographic population present in the study area, estimated by species, was used as a target variable using the normal distribution response distribution and identity link function. Mixed models permitted repetition across survey months, methods (aerial and ground counts) and species (random intercept). Thus, to test our hypotheses we included the following interactions: feeding group*season, feeding group*ice cover and feeding group*max ice. Selection of the best model structure for the dependent variable was based on the Akaike information criterion (AIC) (Zuur et al. 2009). All possible models were carried out (they are listed in Table S1 in Supplementary material). As the final models we assumed those in which $\Delta AIC < 2$ (Burnham & Anderson 2002) and in our case it was only a general model with all the tested variables. To demonstrate interactions at the level of particular species we produced another GLMM model (with month and method as random factors) and assessed the following interactions: species*season, species*ice cover and species*max ice. The parameters of this model are listed in Table S3 in the Supplementary material. The predicted values of this model for each species are shown on Figure 2 and predicted values were back-transformed. We used

IBM SPSS Statistic version 20 software for the statistical analysis. $P < 0.05$ was considered statistically significant.

Results

Benthic feeding species in the study area were more sensitive to lower temperatures and left sooner when colder weather increased ice cover, whereas numbers of fish feeding species did not change, regardless of the extent of ice cover (Tables 2, Fig. 2B). ~~According to our most optimal model (Table S2), we found that~~ interactions between feeding group and season, feeding group and ice cover, and feeding group and maximum ice extent on the Baltic sea were all significant. However, the strongest effects were interactions with ice cover, then interaction with season. The effect of maximum ice extent was very small (Table 2). Our results show changes in population indices in the ORE over the last 25 years. These indices increased in the case of benthic feeding species but decreased for fish feeders (Table 2, Fig 2A). Ice cover across the whole Baltic Sea had the same, though weak, impact on both functional groups of birds. Numbers of birds in the ORE declined with expanding ice cover in the Baltic (Table 2, Fig 2C).

Where particular species are concerned, the situation is more complex. The population indices of Scaup and Tufted Duck in the ORE exhibited an increasing trend, despite the general decline in their entire northern and western European populations; numbers of both species in the ORE were adversely affected by ice cover in that region but not by ice cover in the whole Baltic. Relative numbers of Pochard in the ORE have declined, but so has the whole northern European population; ice cover in the study area was detrimental to abundance there, but ice cover in the whole Baltic had no effect. For Goldeneye, the index for the ORE population was unchanged, despite the increase in the European population; abundance was negatively impacted by ice

cover in the study area, but not by ice cover in the entire Baltic. Relative numbers of Coot in the ORE remained unchanged, despite the slight increase in the European population; abundance was negatively impacted by ice cover in both the study area and in the entire Baltic. The ORE population index for Smew decreased, despite the increase in its biogeographic population; abundance in the ORE was unaffected by ice cover either in the study area or in the Baltic as a whole. Finally, relative numbers of Goosander in the ORE remained unchanged, like those of the whole population wintering in north-western and central Europe; abundance in the ORE was unaffected by ice cover either in the study area or in the Baltic as a whole. The details relating to all these species are listed in Tables 1 and 2, Fig. 2 and Table S3. Table 3 summarizes the changes in the importance of the ORE for wintering populations of diving waterbirds in the last 25 years.

Discussion

The phenomenon of freezing in our study area has decreased over time (Girjatowicz 2011), so that target birds species should tend to feed more recently more often than in the past. However, two functional groups of waterbirds – benthivores and piscivores – react differently to ice cover, a factor that is directly connected to climate change; this has consequences for the wintering patterns of these species. Benthic feeding birds (Scaup, Tufted Duck, Pochard, Goldeneye and Coot) tend to be more sensitive to ice cover in the study area than fish feeders (Smew and Goosander). Benthivorous birds feed in the ORE mainly on mussels of the genus *Dreissena* (Marchowski et al. 2015, 2016); the highest quality of this food resource is found

primarily in water 1-2 m deep (Wolnomiejski and Witek 2013). Shallow water freezes over faster, displacing birds to deeper unfrozen areas where food is accessible only with difficulty. In addition, when ice cover is present, the abundance of food in unfrozen areas declines owing to its greater exploitation, because the birds congregate on a limited area. In the case of piscivorous birds we predicted that increasing ice cover would not affect their numbers: our results substantiate that prediction. The ORE is never completely covered by ice: the shipping lane between Świnoujście and Szczecin is kept free of ice (Girjatowicz 1991; 2005), and there are always other areas free of ice, especially at the mouths of the small rivers flowing into the estuary. These ice-free areas may still abound in fish and provide food for fish feeders. In general, we have demonstrated the growing importance of the study area for all the benthivores. With respect to particular species, the two most numerous ones have increased in numbers, whereas another three do not follow the general trend. In addition, we have shown that the study area is decreasing in importance to piscivores and that Smew is decreasing in numbers.

An interesting result is the negative effect of maximum ice cover in the entire Baltic Sea on the numbers of all species in our study (Fig. 2). This is unexpected, since our study area is in the warmer south-western Baltic, where one would anticipate an increase in the number of waterbirds in such circumstances (Alerstam 1990). The explanation for this relationship is not easy and certainly goes far beyond the scope of this work, but it may inspire further research. However, we can speculate on possible scenarios. Maps showing the maximum range of ice cover in the Baltic Sea show clearly that when the northern Baltic, i.e. the Gulf of Bothnia and the Gulf of Finland, is completely frozen over, the entire Pomeranian Bay (SW Baltic) (see the map – Fig.1) together with the ORE is also covered with ice (Finnish Meteorological Institute 2017). These areas freeze over quickly because of their shallowness and low salinity, the latter

being due to the considerable influence of fresh water from the Odra river basin. Consequently, during harsh winters, birds from northern Baltic move to the south and west, but they by-pass our study area as it is covered by ice. Under such circumstances there may sometimes be better conditions for waterbirds in areas farther north, e.g. the southern coast of Sweden, where there is no ice cover (Finnish Meteorological Institute 2017). Worth noting here, however, is that such cold weather causing the entire Pomeranian Bay and Odra River Estuary to freeze over is rare and becoming rarer (EEA 2017). Nevertheless, if we consider the impact of ice cover of the whole Baltic within species, we can see differences between them and the non-significant impact of this phenomenon on e.g. Smew and Goosander, which corresponds with the local results (Fig. 2).

The global temperature has risen about 1°C over the last 130 years, and Northern Hemisphere temperatures of the last 30 years have been the highest in over 800 years (Stocker et al. 2013). The extent and duration of ice cover in the Baltic have decreased on average by 50% over the last 36 years (Schröder 2015). There is evidence that the range and occurrence of migratory birds have changed in response to climate change and that some species have shortened their migratory movements by wintering closer to their breeding areas (Musil et al. 2011; Lehtikoinen et al. 2013; Pavon-Jordan et al. 2015; Meller 2016). Assuming continued climate warming, the negative correlation between numbers of benthic feeding birds and the number of days with ice cover indicates that the ORE is becoming more important for this group of birds. Climate change seems to be the primary reason for increases (in the study area) in numbers of Scaup and Tufted Duck and decreases in numbers of Smew; this corresponds with the findings of Lehtikoinen et al. (2013) in the case of Tufted Duck and of Pavón-Jordan et al. (2015) in the case of Smew. Our results are important for conservation planning. Declines in the

populations of species such as Scaup and Tufted Duck, even though the importance of our study area to these species is increasing, but at the same time there is an increase in exposure to locally emerging threats. The biggest threats to these species in the area include fishery bycatches (Žydelis et al. 2009; Bellebaum et al. 2012). The ecology of diving ducks makes this type of threat responsible for the extra mortality of all species covered by this study. Comparison of a species' estimated total population numbers (Nagy et al. 2014) with numbers for the ORE is interesting, since local trends and European trends do not always concur. The different responses of particular species to the factors investigated are also worth examining. We grouped the species by trends in the study area and discuss these for each species below.

Species with increasing population index in the study area

Between the late 1980s and 2012, the population of Scaup wintering in northern and western Europe declined at an annual rate of 3.57%/year (Nagy et al. 2014). Around 41% of the Scaup from this population spent the winter in the Baltic Sea region (Skov et al. 2011), and this, in turn, declined by 60% from 1991 to 2010 (Aunins et al. 2013). At the same time we found that the importance of the ORE for this species was increasing. Scaup numbers increased by 300% in the Szczecin Lagoon (the biggest part of Odra River Estuary – see the map – Fig. 1) and the eastern coastal areas of Germany, as opposed to declines further west along the German coast, where some areas (Wismar Bay and Travelförde) had fewer birds than 15 years earlier (Skov et al. 2011). A similar trend was found in Sweden, where the number of wintering Scaup increased between 1971 and 2015 (Nilsson and Haas 2016). But farther west, in the Netherlands, Hornman et al. (2012) recorded decreases at the most important wintering sites since 1980/1981. All of these studies confirm that Scaup is shifting its wintering range northwards and eastwards, closer

to its breeding areas: this is the reason for the heightened importance to this species of the ORE, even as its overall population wintering in northern and western Europe is declining.

Tufted Duck populations wintering in north-western Europe have recently been decreasing by 0.98%/year (Nagy et al. 2014). Lehtikoinen et al. (2013) showed that the population estimated for the North-West Europe flyway remained relatively stable between 1987 and 2009, a situation confirmed by Wetlands International (2016). In the Baltic Sea region, too, there were no significant changes in numbers between 1991 and 2010 (Aunins et al 2013). We have found that our study area has increased in importance for this species, although not to the same extent as for Scaup. By comparison, Nilsson and Haas (2016) showed Swedish populations to have increased between 1971 and 2015, and Lehtikoinen et al. (2013) reported a rapid increase in the last three decades for Finland. Tufted Ducks in the ORE behave in the same way as Scaup in that they form mixed flocks consuming the same type of food (Marchowski et al. 2016). At a larger scale, Tufted Ducks have a different migration and wintering strategy: Scaup concentrate in a few hot spots, moving jump-wise between them, whereas the distribution of Tufted Ducks is more diffuse (van Erden and de Leeuw 2010; Skov et al. 2011; Carboneras and Kirwan 2016a; Carboneras and Kirwan 2016b). This could cause Tufted Ducks to disperse to smaller water bodies outside our study area, e.g. the numerous lakes in the Pomeranian Lake District in northern Poland (~34 000 km²), whereas Scaup remain almost exclusively in the ORE (e.g. Marchowski and Ławicki 2011; Marchowski et al. 2013). The results of the Wintering Waterbird Monitoring programme also show the greater prevalence in Poland of Tufted Duck (29.5%) than Scaup (7.8%) (Neubauer et al. 2015). Scaup is known to concentrate in big flocks during migration and wintering, and the whole biogeographic population may be concentrated in a few hot-spots such the ORE (Marchowski et al. 2015): this is important in the context of species

conservation planning. We have shown an increase in the importance of ORE for Scaup, but at the same time there is an increase in exposure to locally emerging threats such as bycatches in fishing nets (Bellebaum et al. 2012). Taking into account the above pattern of Scaup behaviour and our results, there is a justified fear that locally operating threats in the ORE may affect the entire biogeographic population of the species. This is one of the most important messages of our work.

Species with decreasing population index in the study area

Pochard populations from north-east / north-west Europe have declined rapidly at an annual rate of 3.35%/year (Nagy et al. 2014). Pochard numbers in the Baltic Sea region also declined by 70% between 1991 and 2010 (Aunins et al. 2013). In 1995 there were an estimated 300 000 Pochard in the north-east/north-west European population (Delany et al. 1999). With a constant decline of 3.35%/year, the total population should now be less than 150 000 (Nagy et al. 2014). Numbers of Pochard were expected to be higher in the ORE because of the reduced ice cover. However, we found a significant reduction in the importance of the estuary to this species (Table 3), corresponding with its global decline (Aunins et al. 2013; Nagy et al. 2014; Wetlands International 2016). Pochard behaves more like Tufted Duck than Scaup over winter in being more dispersed and occurring on smaller bodies of water (e.g. Marchowski and Ławicki 2011; Marchowski et al. 2013; Neubauer et al. 2015). This implies that individuals may also be wintering outside the study area, e.g. on the numerous water bodies of the Pomeranian Lake District, like Tufted Duck. This local decline, however, seems to be driven by the species' global decline, despite the emergence of better conditions for wintering that might favour population growth.

Smew populations wintering in northern, western and central Europe increased at 1.97%/year between the late 1980s and 2012 (Nagy et al. 2014); in the Baltic Sea region numbers increased by 30% between 1991 and 2010 (Aunins et al. 2013). Although Smew cannot be classified as a piscivore in the same way as Goosander (and Red-breasted Merganser), it does feed on very small fish and on small invertebrates (Carboneras & Kirwan 2016 c). Though more dependent on shallow water than Goosander, Smew generally forages on mobile types of food. So even if shallow waters freeze over, it may remain on site and search for food in deeper water, which is what we have observed. We found that nowadays, the ORE is of less importance to Smew (Table 3). This statement is underpinned by the northward and eastward shift in wintering area boundaries due to climate warming, as already demonstrated by Pavon-Jordan et al. (2015). Confirmation of this process is provided by the significant increase in numbers of Smew in 1971-2015 in places to the north of our study area, in Sweden (Nilsson and Haas 2016).

Species with no changes in the population index in the study area

Coot populations wintering in north-west Europe increased by 0.19%/year between the late 1980s and 2012 (Nagy et al. 2014), but in the Baltic region there was a 60% decline between 1991 and 2010 (Aunins et al. 2013). We have found no changes in Coot numbers in the ORE over the last 25 years (Table 3). Likewise, no changes in numbers were recorded between 1975 and 2010 at wintering sites in warmer areas to the south-west (the Netherlands) (Hornman et al. 2012). Long-term figures for Sweden (1971-2015), while not revealing any distinct increase, do show that Coot populations fluctuated, rising during mild periods and falling during cold periods (Nilsson and Haas 2016). The expected increase in numbers due to improvements in habitat quality did not happen. Factors such as pressure from American mink *Neovison vison*, which are responsible for the decline of Coot in many places (e.g. Ferreras and Macdonald 1999), may

have held back potential increases. Moreover, compared to the bottom-diving ducks, Coot is more sensitive to cold weather: a study by Fredrickson (1969) demonstrated high mortality after periods of severe weather (also reflected in the results of Swedish breeding bird surveys – Leif Nilsson pers. com.) but that the population recovered during mild winters. This factor may also be the reason for the different reactions of Coot and diving ducks to the cold.

Goldeneye populations wintering in north-west and central Europe increased at 0.26%/year between the late 1980s and 2012 (Nagy et al. 2014) and increased in the Baltic Sea region by 50% between 1991 and 2010 (Aunins et al. 2013). This corresponds to the data provided by Lehtikoinen et al (2013), which show an increase in numbers in the northern Baltic wintering area (Finland and N Sweden), but a decline in the southern part of its wintering range (Switzerland, France). In our work we found the relative number of Goldeneye in the ORE to be stable in the period 1992-2016 (Table 3). This again tallies with the findings of Lehtikoinen et al. (2013) that duck abundances are independent of temperature in the central part of the flyway. This is probably why the shift in wintering range is not perceptible in our study area but is more pronounced at other, e.g. Swedish wintering sites, where numbers have increased (Nilsson and Haas 2016) but not in the Netherlands, where they have declined (Hornman et al. 2012).

Goosander populations wintering in north-west and central Europe have been stable since the early 1990s (Nagy et al. 2014); moreover, numbers in the Baltic Sea between 1991 and 2010 did not change significantly (Aunins et al. 2013). We also found non-significant changes in the ORE, so it must be regarded as stable (Table 3). As in the case of Goldeneye, the explanation is that in the central part of the flyway, species abundances are independent of temperature. In other areas, observations indicate a shift farther to the north and east in the wintering range as a result of climate warming (Hornman et al. 2012; Lehtikoinen et al. 2013; Nilsson and Haas 2016).

Conclusion

Our study has confirmed the part played by climate change in shaping the distribution of waterbirds. Apart from climate changes, however, feeding ecology, fishery, interspecific competition and human-related disturbance may be also important and should be taken into consideration (Quan et al. 2002, Žydelis et al. 2009, Clavero et al., 2011; Eglington & Pearce-Higgins, 2012). We show that the protected areas covered by our study will be more important for some species (Scaup and Tufted Duck) but less so for others (Smew). Taking into account the large abundance of the target species regularly present in the ORE, conservation measures applied here will have a large impact on whole populations; this applies primarily to Scaup. Shifts in species distributions should be accounted for in future management plans for Special Protection Areas of the European Natura 2000 network. We believe that our results add new insight to the problem of wintering waterbird protection and can help to shape conservation policy in the southern Baltic.

Acknowledgements

We thank all the people who took part in the fieldwork – mainly members of the West-Pomeranian Nature Society – but especially the most active among them during the entire study period: Michał Barcz, Ryszard Czeraszewicz, Sebastian Guentzel, Michał Jasiński, Zbigniew Kajzer, Jacek Kaliciuk, Krzysztof Kordowski, Aneta Kozłowska, Wojciech Mrugowski, Arkadiusz Oleksiak, Bartosz Raclawski, Tomasz Rek, Artur Staszewski, Marcin Sołowiej, Piotr Siuda, Paweł Stańczak and Maciej Przybysz. We are grateful to Leif Nilsson and the two anonymous reviewers for their valuable comments on the first version of the manuscript, and to Peter Senn, who kindly improved our English.

References

- Aharon-Rotman Y., Buchanan K.L., Clark N.J., Klaassen M., Buttemer W.A. 2016. Why fly the extra mile? Using stress biomarkers to assess wintering habitat quality in migratory shorebirds. *Oecologia* 182: 385-395.
- Alerstam T. 1990. Bird migration. Cambridge University Press, Cambridge.
- Aunins A., Nilsson L., Hario M., Garthe S., Dagys M., Petersen I.K., Skov H., Lehikoinen A., Roos M.M., Ranft S., Stipniece A., Luigujoe L., Kuresoo A., Meissner W., Korpinen S. 2013. Abundance of waterbirds in the wintering season. HELCOM Core Indicator Report. Online. Available: <http://www.helcom.fi/>.
- Bellebaum, J., Schirmeister, B., Sonntag, N., Garthe, S. 2012. Decreasing but still high: bycatch of seabirds in gillnet fisheries along the German Baltic coast. *Aquat. Conservat. Mar. Freshwat. Ecosyst.* 23, 210–221.
- BirdLife International 2004. Birds in Europe, population estimate, trends and conservation status. Cambridge UK: BirdLife Conservation Series No.12.
- BirdLife International 2015 a. European Red List of Birds. Luxembourg: Office for Official Publications of the European Communities.
- BirdLife International. 2015 b. *Aythya ferina*. The IUCN Red List of Threatened Species 2015:e.T22680358A82571892. <http://dx.doi.org/10.2305/IUCN.UK.2015-4.RLTS.T22680358A82571892.en>. Downloaded on 23 October 2016.
- Blums P., Mednis A., Bauga I., Nichols J.D., Hines J.E. 1996. Age-specific survival and philopatry in three species of European ducks: a long-term study. *Condor* 98: 61-74.
- Burnham, K.P. & Anderson, D.R. 2002. Model Selection and Multimodel Inference: A Practical Information-Theoretic Approach. Springer, New York/
- Carboneras C. & Kirwan G.M. 2016 a. Greater Scaup (*Aythya marila*). In: del Hoyo, J., Elliott, A., Sargatal, J., Christie, D.A. & de Juana, E. (eds.). *Handbook of the Birds of the World Alive*. Lynx Edicions, Barcelona. (retrieved from <http://www.hbw.com/node/52912> on 23 October 2016).
- Carboneras, C. & Kirwan, G.M. 2016 b. Tufted Duck (*Aythya fuligula*). In: del Hoyo, J., Elliott, A., Sargatal, J., Christie, D.A. & de Juana, E. (eds.). *Handbook of the Birds of the World Alive*. Lynx Edicions, Barcelona. (retrieved from <http://www.hbw.com/node/52910> on 23 October 2016).
- Carboneras, C. & Kirwan, G.M. 2016 c. Smew (*Mergellus albellus*). In: del Hoyo, J., Elliott, A., Sargatal, J., Christie, D.A. & de Juana, E. (eds.). *Handbook of the Birds of the World Alive*. Lynx Edicions, Barcelona. (retrieved from <http://www.hbw.com/node/52927> on 21 October 2016).
- Carboneras, C. & Kirwan, G.M. 2016 d. Goosander (*Mergus merganser*). In: del Hoyo, J., Elliott, A., Sargatal, J., Christie, D.A. & de Juana, E. (eds.). *Handbook of the Birds of the World Alive*. Lynx Edicions, Barcelona. (retrieved from <http://www.hbw.com/node/52931> on 21 October 2016).
- Carboneras, C., Christie, D.A. & Kirwan, G.M. 2016. Common Goldeneye (*Bucephala clangula*). In: del Hoyo, J., Elliott, A., Sargatal, J., Christie, D.A. & de Juana, E. (eds.). *Handbook of the Birds of the World Alive*. Lynx Edicions, Barcelona. (retrieved from <http://www.hbw.com/node/52925> on 23 October 2016).

- Clavero, M., Villero, D. & Brotons, L. 2011 Climate change or land use dynamics: do we know what climate change indicators indicate? PLoS ONE, 6, e18581.
- Coppack, T., Both, C., 2002. Predicting life-cycle adaptation of migratory birds to global climate change. Ardea 90: 369–378.
- Cramp, S. & Simmons, K.E.L. eds. 1977. The Birds of the Western Palearctic. Vol. 1. Oxford University Press, Oxford.
- Cresswell, W. 2014. Migratory connectivity of Palaearctic – African migratory birds and their responses to environmental change: the serial residency hypothesis. Ibis 156: 493–510. doi:10.1111/ibi.12168
- Czeraszewicz R., Haferland H.J., Oleksiak A. 2004. The results of waterbirds count in the Western Pomerania in the season 2003/2004. In: Czeraszewicz R., Oleksiak A. Waterbirds in Western Pomerania. The results of counts in 2003/2004, ecology and conservation, ZTO–PZŁ, Szczecin. 2004. pp. 5–16. (in Polish with German summary).
- Guentzel S., Ławicki Ł., Marchowski D., Kajzer Z., Barcz M., Raclawski B., Sołowiej M., Staszewski A. 2012. The Szczecin Lagoon PLB320009. In: Ławicki Ł., Guentzel S. editors. Important Bird Areas in Poland – the inventory of non-breeding species in 2011/2012 season. ECO-EXPERT, Szczecin. (In Polish).
- De Leeuw J.J. 1999. Food intake rates and habitat segregation of Tufted Duck *Aythya fuligula* and Scaup *Aythya marila* exploring Zebra Mussels *Dreissena polymorpha*. Ardea 87: 15–31.
- Delany S.N., Reyes C., Hubert E., Phil S., Rees E., Haanstra L., van Strien A. 1999. Results from the International Waterbird Census in the Western Palearctic and Southwest Asia 1995 and 1996. Wetland International Publ. Wageningen, The Netherlands.
- Dillingham P.W., Fletcher D. 2008. Estimating the ability of birds to sustain additional human-caused mortalities using a simple detection rule and allometric relationship. Biological Conservation 141: 1783–1792.
- EEA - The European Environment Agency data obtained from: <http://www.eea.europa.eu/data-and-maps/figures/maximum-extents-of-ice-cover> 10.02.2017
- Eglington, S.M. & Pearce-Higgins, J.W. 2012. Disentangling the relative importance of changes in climate and land-use intensity in driving recent bird population trends. PLoS ONE, 7, e30407
- Ferreras P., Macdonald D. W. 1999. The impact of American mink *Mustela vison* on water birds in the upper Thames. J Appl Ecol 36: 701–708.
- Finnish Meteorological Institute data obtained from: <http://en.ilmatietaenlaitos.fi/> 10.02.2017
- Girjatowicz J.P. 1990. Ice cover atlas for Polish Baltic Coastal waters. Akademia Rolnicza, Szczecin (in Polish).
- Girjatowicz J.P. 2005. Ice conditions in Szczecin Lagoon and the Pomeranian Bay. In: Borówka R.K., Musielak S. (eds), Środowisko przyrodnicze wybrzeży Zatoki Pomorskiej i Zalewu Szczecińskiego. Uniwersytet Szczeciński, Szczecin, pp. 126–138 (in Polish).
- Girjatowicz J.P. 2011. Ice Conditions on the Southern Baltic Sea Coast. J. Cold Reg. Eng. 25:1–15.
- Hornman M., van Roomen M., Hustings F., Koffijberg K., van Winden E., Soldaat L. 2012. Population trends in wintering and migrating waterbirds in The Netherlands in 1975–2010 (in Dutch with English summary). Limosa 85: 97–116

- 512 Jankowiak, Ł., Antczak, M., Kwieciński, Z., Szymański, P., Tobolka, M., Tryjanowski, P.,
513 2015a. Diurnal raptor community wintering in an extensively used farmland. *Ornis Fennica*
514 92: 76-86.
- 515 Jankowiak, Ł., Polakowski, M., Kułakowski, T., Świętochowski, P., Tumił, T., Broniszewska,
516 M., Takács, V., 2015b. Habitat and weather requirements of diurnal raptors wintering in
517 river valleys. *Biologia (Bratisl)*. 70, 1136–1142.
- 518 Newton, I., 2008. The migration ecology of birds. Academic Press, London, United Kingdom.
- 519 Kaliciuk J., Oleksiak A., Czeraszewicz R. 2003. The results of counts of waterbirds in the
520 Western Pomerania in the season 2002/2003. In: Czeraszewicz R., Oleksiak A.
521 Waterbirds in Western Pomerania. The results of counts in 2002/2003, ecology and
522 conservation. ZTO–PZŁ, Szczecin. 2003. pp. 14–25. (in Polish with German summary).
- 523 Kear, J. ed. 2005. Ducks, geese and swans. Oxford University Press, Oxford.
- 524 Lehtikoinen A., Jaatinen K., Vahatalo A.V., Preben C., Crowe O., Deceuninck B., Hearn R., Holt
525 C.A., Hornman M., Keller V., Nilsson L., Langendoen T., Tomankova I., Wahl J., Fox
526 A.D. 2013. Rapid climate driven shifts in wintering distributions of three common
527 waterbird species. *Global Change Biology*: 19, 2071–2081, doi: 10.1111/gcb.12200
- 528 Ludwiczowski, I., Barker, R. & Bräger, S. 2002. Nesting area fidelity and survival of female
529 Common Goldeneyes *Bucephala clangula*: are they density-dependent? *Ibis* 144: 452–
530 460.
- 531 Ławicki Ł., Czeraszewicz R., Guentzel S., Jasiński M., Kajzer Z., Kaliciuk J., Oleksiak A. 2008.
532 Wintering of Waterbirds in the Western Pomerania in the years 2002-2008. *Not. Orn.* 49:
533 235–244. (in Polish with English summary).
- 534 Komdeur J., Bertelsen J., Cracknell G. 1992. Manual for aeroplane and ship surveys of
535 waterfowl and seabirds. IWRB Special Publication No.19. IWRB, Slimbridge, U.K.
- 536 Larsen K.1992. New figures of seaduck winter population in the Western Palearctic. IWRB
537 Seaduck Bulletin No.1 Jan.1992.
- 538 Marchowski D., Ławicki Ł. 2011. Numbers of Waterfowl in Western Pomerania in the
539 2009/2010 season. *Ptaki Pomorza* 2: 159–166 (in Polish with English summary).
- 540 Marchowski D., Ławicki Ł. 2012. Numbers of Waterfowl in Western Pomerania in the season
541 2010/2011. *Ptaki Pomorza* 3: 129–134 (in Polish with English summary).
- 542 Marchowski D., Ławicki Ł., Guentzel S. 2013. Numbers of Waterfowl in Western Pomerania in
543 the season 2010/2011. *Ptaki Pomorza* 4: 149–169. (in Polish with English summary).
- 544 Marchowski D., Neubauer G., Ławicki Ł., Woźniczka A., Wysocki D., Guentzel S., Jarzowski
545 M. 2015. The importance of non-native prey, the Zebra Mussel *Dreissena polymorpha*, for
546 the declining Greater Scaup *Aythya marila*: a case study at a key European staging and
547 wintering site. *PLoS ONE* 10(12): e0145496.
- 548 Marchowski D., Jankowiak Ł., Wysocki D. 2016. Newly demonstrated foraging method of
549 Herring Gulls and Mew Gulls with benthivorous diving ducks during the nonbreeding
550 period. *Auk* 133: 31–40.
- 551 Meller K. 2016. The impacts of temperature on the long-term variation in migration and breeding
552 performance of birds. PhD dissertation. The Helsinki Lab of Ornithology The Finnish
553 Museum of Natural History Luomus Zoology Unit University of Helsinki Finland.
554 Available at:
555 https://helda.helsinki.fi/bitstream/handle/10138/160724/The_impa.pdf?sequence=1

- Musil P., Musilová Z., Fuchs R., Poláková S. 2011. Long-term changes in numbers and distribution of wintering waterbirds in the Czech Republic, 1966– 2008. *Bird Study* 58: 450–460.
- Nagy S., Flink S., Langendoen T. 2014. Waterbird trends 1988-2012. Results of trend analyses of data from the International Waterbird Census in the African-Eurasian Flyway. Wetlands International. Ede, the Netherlands.
- Neubauer G., Meissner W., Chylarecki P., Chodkiewicz T., Sikora A., Pietrasz K., Cenian Z., Betleja J., Gaszewski K., Kajtoch Ł., Lenkiewicz W., Ławicki Ł., Rohde Z., Rubacha S., Smyk B., Wieloch M., Wylegała P. Zieliński P. 2015. Monitoring of Birds in Poland in years 2013-2015. *Newsletter of Nature Monitoring* 13: 1-92.
- Nilsson L., Haas F. 2016. Distribution and numbers of wintering waterbirds in Sweden in 2015 and changes during the last fifty years. *Ornis Svecica* 26:3–54.
- Pavón-Jordan D., Fox A., D., Clausen P., Dagys M., Deceuninck B., Devos K., Hearn R. D., Holt C. A., Hornman M., Keller V., Langendoen T., Ławicki Ł., Lorentsen S., H., Luiguj L., Meissner W., Musil P., Nilsson L., Paquet J. Y., Stipniece A., Stroud D. A., Wahl J., Zenatello M., Lehtikainen A. 2015. Climate-driven changes in winter abundance of a migratory waterbird in relation to EU protected areas. *Diversity Distrib.* 21: 571-582.
- Pearce J.M., Reed J.A., Flint P.L. 2005. Geographic variation in survival and migratory tendency among North American Common Merganser. *Journal of Field Ornithology* 76: 109-118.
- Perdeck A. C. 1998. Poisson regression as a flexible alternative in the analysis of ring-recovery data. *Euring Newsletter* 2: 37-42.
- Radziejewska T., Schernewski G. 2008. The Szczecin (Oder) Lagoon. In: Schiewer U, editor. *Ecology of Baltic Coastal Waters*. Berlin Heidelberg: Springer, pp. 116–117.
- Runge M. C., Sauer J.R., Avery M.L., Blackwell B.F., Koneff M.D. 2009. Assessing allowable take of migratory birds. *Journal of Wildlife Management*. 73: 556-565.
- Skov H., Heinänen S., Žydelis R., Bellebaum J., Bzoma S., Dagys M., Durinck J., Garthe S., Grishanov G., Hario M., Kieckbusch J.J., Kube J., Kuresoo A., Larsson K., Luigujoe L., Meissner W., Nehls H. W., Nilsson L., Petersen I.K., Roos M. M., Pihl S., Sonntag N., Stock A., Stipniece A., Wahl J. 2011. Waterbird Populations and Pressures in the Baltic Sea. Nordic Council of Ministers, Copenhagen.
- Schröder J. 2015. Does ice coverage in the Baltic Sea affect numbers of diving ducks wintering in the Netherlands? *Limosa* 88: 22-30.
- Stańczykowska A., Lewandowski K., Czarnoleski M. 2010. Distribution and densities of *Dreissena polymorpha* in Poland – past and present. In: Van der Velde G, Rajagopal S, Bij de Vaate A, editors. *The Zebra Mussel in Europe*. Margraf Publishers, Weikersheim: pp. 119-132.
- Stempniewicz L., Meissner W. 1999. Assessment of the zoobenthos biomass consumed yearly by diving ducks wintering in the Gulf of Gdańsk (southern Baltic Sea). *Ornis Svecica* 9: 143–154.
- Stocker T.F., Qin D., Plattner G.K., Tignor M., Allen S.K., Boschung J., Navels A., Xia Y., Bex V. Midgley P.M. eds. 2013. *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge University Press.
- Taylor, B. & Kirwan, G.M. 2016. Common Coot (*Fulica atra*). In: del Hoyo, J., Elliott, A., Sargatal, J., Christie, D.A. & de Juana, E. (eds.). *Handbook of the Birds of the World Alive*.

- 601 Lynx Edicions, Barcelona. (retrieved from <http://www.hbw.com/node/53695> on 21
- 602 October 2016).
- 603 van Erden M. R., de Leeuw J.J. 2010. How *Dreissena* sets the winter scene for water birds:
- 604 dynamic interaction between diving ducks and zebra mussels. In: van der Velde G.,
- 605 Rajagopal S., bij de Vaate A. The Zebra Mussel in Europe. 2010. Backhuys Publ., Leiden,
- 606 The Netherlands.
- 607 Wacker A. 2010. Careless youth? Food in the early life-stages of zebra mussels. In: Van der
- 608 Velde G., Rajagopal S., bij de Vaate A. The Zebra Mussel in Europe. 2010. Backhuys
- 609 Publishers, Leiden, The Netherlands.
- 610 Weatherbase. 2016.: Historical Weather for Szczecin, Poland. (access: 29.01.2016), available at:
- 611 <http://www.weatherbase.com/>.
- 612 Wetlands International. Guidance on waterbird monitoring methodology: Field Protocol for
- 613 waterbird counting. Wetlands International, Wageningen. 2010. Available from:
- 614 <http://www.wetlands.org>.
- 615 Wetlands International 2016. "*Waterbird Population Estimates*". Retrieved
- 616 from wpe.wetlands.org on Tuesday 11 Oct 2016
- 617 Wiktor J. 1969. Biology of *Dreissena polymorpha* (Pall.) and its importance on Szczecin
- 618 Lagoon. Stud. Mater. Mor. Inst. Ryb., Gdynia. (in Polish).
- 619 Wilk T., Jujka M., Krogulec J., Chylarecki P. (eds.) 2010. Important Bird Areas of international
- 620 importance in Poland. OTOP, Marki.
- 621 Wolnomiejski N., Witek Z. 2013. The Szczecin Lagoon Ecosystem: The Biotic Community of
- 622 the Great Lagoon and its Food Web Model. Versita Ltd. London.
- 623 Wolnomiejski N., Woźniczka A. 2008. A drastic reduction in abundance of *Dreissena*
- 624 *polymorpha* Pall. in the Skoszeńska Cove (Szczecin Lagoon, River Odra estuary): effects
- 625 in the population and habitat. Ecological Questions 9: 103–111.
- 626 Žydelis, R., Bellebaum, J., Österblom, H., Vetemaa, M., Schirmeister, B., Stipniece, A., Dagys
- 627 M., van Eerden M., Garthe S. 2009. Bycatch in gillnet fisheries – An overlooked threat to
- 628 waterbird populations. Biol. Conserv. 142, 1269–1281.
- 629 Zuur, A., Ieno, E.N., Walker, N., Saveliev, A.A., Smith, G.M. 2009. Mixed Effects Models and
- 630 Extensions in Ecology with R. Springer-Verlag, New York.
- 631 Quan, R.C., Wen, X.J., Yang, X.J., 2002. Effects of human activities on migratory waterbirds at
- 632 Lashihai Lake. Biological Conservation 108, 273–279.

Table 1. Biogeographic populations and annual trends (after Nagy et al. 2014) for seven species of waterbirds using the Odra River Estuary.

(1) Target species. (2) Functional group: B – benthivores, P – piscivores. (3) Estimated number of individuals from biogeographic population in 1992, the numbers are presented in thousands. (4) Estimated number of individuals from biogeographic population in 2012, the numbers are presented in thousands. (5) Population trend % per annum - long term assessment. (6) Significances of changes.

Species (1)	Functional group (2)	Number of individuals (1992) (3)	Number of individuals (2012) (4)	Population trend % p.a. (5)	Significance of changes (6)
Greater Scaup	B	300	150	-3.57	Large decline
Common Pochard	B	280	150	-3.35	Large decline
Tufted Duck	B	1,100	820	-0.98	Large decline
Goosander	P	130	100	-0.09	Stable
Eurasian Coot	B	990	950	+0.19	Moderate increase
Common Goldeneye	B	210	240	+0.26	Moderate increase
Smew	P	13	24	+1.97	Large increase

Table 2. Results of general linear mixed models for seven species showing the influence of ice cover, maximum ice extent [km²] in the Baltic Sea (max ice) and season on the percentage of occurrence of benthivores (denoted by B, Scaup, Tufted Duck, Pochard, Goldeneye, Coot) and piscivores (denoted by P, Smew, Goosander) in the Odra River Estuary. Species, method and month were treated as random effects in relation to their regional breeding populations.

Model Term	Coefficient	Std. Error	t	P
Intercept	26.553	11.619		
Ice cover	0.014	0.006	2.375	0.018
Season	-0.013	0.006	-2.204	0.028
Max ice	-0.001	0.000	-2.824	0.005
Feed[B]	-38.751	11.959	-3.240	0.001
Season*Feed[B]	0.019	0.006	3.212	0.001
Ice cover*Feed[B]	-0.044	0.007	-6.623	<0.001
Max ice*Feed[B]	0.001	<0.001	2.071	0.039

Species (r)	0.074	0.048
Method (r)	0.015	0.020
Month (r)	0.001	0.002

705

706

707

708

709

710

Table 3. Population index trends in the Odra River Estuary (ORE) for the biogeographic population (b.p.) of diving waterbirds showing the percentage of the biogeographic population in 1992; the percentage of the biogeographic population in 2016; the mean percentage of the biogeographic population in the period 1992 – 2016 $\pm$ standard error; and the trend in the period 1992 – 2016.

Species	%b.p.1992	%b.p.2016	Mean1992– 2016 $\pm$ SE	Trend in ORE
Greater Scaup	5.68	12.60	14.17 $\pm$ 2.84	↑
Tufted Duck	2.87	4.79	2.61 $\pm$ 0.25	↑
Common Goldeneye	4.48	0.63	1.21 $\pm$ 0.14	→
Eurasian Coot	0.86	0.68	0.61 $\pm$ 0.07	→
Goosander	12.59	1.80	6.85 $\pm$ 1.01	→
Smew	7.04	2.76	7.01 $\pm$ 1.27	↓
Common Pochard	1.84	0.20	0.62 $\pm$ 0.09	↓

Figure 1. The Odra River Estuary, north-western Poland.




Figure 2 A-C. Predicted results of the general linear mixed model showing the influence of season, ice cover and maximum ice extent [km²] in the Baltic Sea (max ice) on the percentage of the population of the target species in the Odra River Estuary. The predicted values were obtained from the model where we added species as a fixed variable. The model's parameters are listed in Table S1 in the Supplementary material.

