

Changes in zooplankton community, and seston and zooplankton fatty acid profiles at the freshwater/saltwater interface of the Chowan River, NC

Deborah Lichti ^{Corresp., 1}, Jacques Rinchard ², David G. Kimmel ^{1, 3}

¹ Department of Biology, East Carolina University, Greenville, North Carolina, United States

² Department of Environmental Science and Biology, State University of New York College at Brockport, Brockport, New York, United States

³ Alaska Fisheries Science Center, National Oceanic and Atmospheric Administration, Seattle, Washington, United States

Corresponding Author: Deborah Lichti
Email address: lichtid12@students.ecu.edu

The variability in zooplankton fatty acid composition may be an indicator of larval fish habitat quality as fatty acids are linked to fish larval growth and survival. We sampled an anadromous fish nursery, the Chowan River, during spring of 2013 in order to determine how the seston fatty acid composition varied in comparison with the zooplankton community composition and fatty acid composition during the period of anadromous larval fish residency. The seston fatty acid profiles showed no distinct pattern in relation to sampling time or location. The mesozooplankton community composition varied spatially and the fatty acid profiles were typical of freshwater species in April. The Chowan River experienced a saltwater intrusion event during May, which resulted in brackish water species dominating the zooplankton community and the fatty acid profile showed an increase in polyunsaturated fatty acids (PUFA), in particular eicosapentaenoic acid (EPA) and docosahexaenoic acid (DHA). The saltwater intrusion event was followed by an influx of freshwater due to high precipitation levels in June. The zooplankton community composition once again became dominated by freshwater species and the fatty acid profiles shifted to reflect this change; however, EPA levels remained high, particularly in the lower river. We found correlations between the seston, microzooplankton and mesozooplankton fatty acid compositions. Salinity was the main factor correlated to the observed pattern in species composition, and fatty acid changes in the mesozooplankton. These data suggest that anadromous fish nursery habitat likely experiences considerable spatial variability in fatty acid profiles of zooplankton prey and that are correlated to seston community composition and hydrodynamic changes. Our results also suggest that sufficient prey density as well as a diverse fatty acid composition is present in the Chowan River to support larval fish production.

Changes in zooplankton community, and seston and zooplankton fatty acid profiles at the freshwater/saltwater interface of the Chowan River, NC

Deborah A. Lichti^{1*}, Jacques Rinchar², and David G. Kimmel^{1,3}

¹Department of Biology, East Carolina University, N108 Howell Science Complex, Greenville, NC, 27858, USA

²Department of Environmental Science and Biology, College of Brockport- SUNY 350 New Campus Drive, Brockport, NY 14420, USA

³*Present Address:* NOAA Alaska Fisheries Science Center, 7600 Sand Point Way NE, Seattle, WA 98115, USA

Abstract

The variability in zooplankton fatty acid composition may be an indicator of larval fish habitat quality as fatty acids are linked to fish larval growth and survival. We sampled an anadromous fish nursery, the Chowan River, during spring of 2013 in order to determine how the seston fatty acid composition varied in comparison with the zooplankton community composition and fatty acid composition during the period of anadromous larval fish residency. The seston fatty acid profiles showed no distinct pattern in relation to sampling time or location. The mesozooplankton community composition varied spatially and the fatty acid profiles were typical of freshwater species in April. The Chowan River experienced a saltwater intrusion event during May, which resulted in brackish water species dominating the zooplankton community and the fatty acid profile showed an increase in polyunsaturated fatty acids (PUFA), in particular eicosapentaenoic acid (EPA) and docosahexaenoic acid (DHA). The saltwater intrusion event was followed by an influx of freshwater due to high precipitation levels in June. The zooplankton community composition once again became dominated by freshwater species and the fatty acid profiles shifted to reflect this change; however, EPA levels remained high, particularly in the lower river. We found correlations between the seston, microzooplankton and mesozooplankton fatty acid compositions. Salinity was the main factor correlated to the observed pattern in species composition, and fatty acid changes in the mesozooplankton. These data suggest that anadromous fish nursery habitat likely experiences considerable spatial variability in fatty acid profiles of zooplankton prey and that are correlated to seston community composition and hydrodynamic changes. Our results also suggest that sufficient prey density as well as a diverse fatty acid composition is present in the Chowan River to support larval fish production.

53

54 **Introduction**

55 Estuaries are considered important nursery habitat for many ecologically and
 56 commercially important fish and invertebrates (Beck et al. 2001, Boesch and Turner 1984,
 57 Sheaves et al. 2015, Sheaves 2016). Estuaries function as fish nurseries because they are highly
 58 productive, support large planktonic populations across multiple size ranges, and fish within
 59 estuaries generally have higher growth rates compared to other habitats (Beck et al. 2001).
 60 Hence, many fish have evolved life-history strategies whereby larvae and juvenile stages have
 61 residency periods in estuaries (McHugh 1967; Boehlert and Mundy 1988; Beck et al. 2001; Able
 62 2005; Walsh et al. 2005). Beck et al. (2001) defined nursery habitat as an area that contributes to
 63 higher production of individuals that move to juvenile habitat compared to other areas. Higher
 64 growth rates of larval fish are possible because of zooplankton prey that are present during their
 65 critical transition from yolk sac to free-living, feeding larvae (Hjort 1914; Mullen et al. 1986;
 66 Rulifson et al. 1993; Cooper et al. 1998; Martino and Houde 2010; Binon 2011). However,
 67 spatial and temporal overlap between predators and prey does not completely explain how fish
 68 nurseries function mechanistically. The quality of prey can play a major role in determining the
 69 effectiveness of a nursery for early stages of fish.

70 The quality (chemical composition) of zooplankton prey can influence fish growth,
 71 development, and survival (Fraser et al. 1989; Webster and Lovell 1990; Copeman et al. 2002;
 72 Rossi et al. 2006; Malzahn et al 2007, Paulsen et al. 2014). Fatty acids are chemically diverse,
 73 often incorporated into organisms unmodified, and different organisms have distinct profiles
 74 (Dalsgaard et al. 2003). Fatty acids are one class of compounds found in lipids that are
 75 particularly important, impacting neural and vision development in fish (Gulati et al. 1997;
 76 Müller-Navarra et al. 2000; Kainz et al. 2004; Masclaux et al. 2012). Fatty acids may act as both

dietary tracers in the food web and indicators of overall food quality (Iverson et al. 2004). The majority of organisms need specific dietary fatty acids for somatic development and fitness (Masclaux et al. 2012). These fatty acids, 18:3 ω -3, α -linolenic acid (ALA), and 18:2 ω -6, linoleic acid (LA), are labeled essential fatty acids because they cannot be directly synthesized by heterotrophic organisms and must come from the diet (Arts et al. 2009). Polyunsaturated fatty acids (i.e., 20:5 ω -3, eicosapentaenoic acid (EPA), 22:6 ω -3, docosahexaenoic acid (DHA), and 20:4 ω -6, arachidonic acid (ARA)) are required for all organisms and play a role in health and cell function (Dalsgaard et al. 2003). Thus, an organisms' fatty acid signature may indicate dietary consumption and nutritional quality of its prey (Goncalves et al. 2012).

Fatty acids are present in estuaries as a result of *de novo* synthesis by phytoplankton and the delivery of detrital material of plant origin (Dalsgaard et al. 2003). The free-floating portion of organic and inorganic particles is termed seston (Postel et al. 2000). The organic portion of seston is important because it forms the origin point for the propagation of fatty acids through the pelagic food web. Zooplankton assimilate fatty acids from the seston through direct consumption of phytoplankton cells, detritus and/or consumption of microzooplankton that graze phytoplankton or detritus (Wacker and von Elert 2001; Kainz, Arts, and Mazumder 2004; Vargas et al. 2006). To date, studies of the relationship between seston and zooplankton fatty acid composition have shown variable patterns. Persson and Verde (2006) demonstrated in laboratory studies that broad zooplankton taxonomic groups (cladoceran vs. copepods) have different fatty acid profiles independent of the food source. The seston has been correlated to the fatty acid profile of zooplankton *in situ* as well (Goulden and Place 1990; Brett et al 2006; Taipale et al. 2009; Gladyshev et al. 2010; Ravet, Brett, and Arhoditsis 2010). The mismatch of fatty acids in seston to zooplankton has also been shown in many studies (Desvillettes et al. 1997; Persson and

Vrede 2006; Rossi et al. 2006; Smyntek et al. 2008). Most marine zooplankton species cannot convert precursor fatty acids and only obtain longer-chained fatty acids from their diet, compared to freshwater zooplankton species that can convert precursor fatty acids (Rossi et al. 2006; Persson and Verde 2006). However, it is clear that the fatty acid composition of the seston changes as a result of local conditions, e.g., the salinity and temperature, nutrient concentration, and the degree of autotrophy or heterotrophy in the system (Farkas and Herodek 1964; Desvillettes et al. 1997; Wacker and von Elert 2001; Goncalves et al. 2012). Therefore, combined knowledge of the changing nature of seston fatty acid composition, zooplankton community composition changes, and fatty acid profiles forms a useful base for assessing the quality of fish nursery habitat.

Zooplankton community composition in estuaries has been intensely studied and abiotic factors are thought to structure zooplankton communities (Ambler et al. 1985; Orsi 1986; Cervetto et al. 1999; Mouny and Dauvin 2002; Kimmel and Roman 2004; Lawrence et al. 2004; Islam et al. 2005). Zooplankton community composition in temperate estuaries is dominated by crustaceans in general, and copepods and cladocerans in particular (Tackx et al. 2004, Marques et al. 2006; Winder and Jassby 2011; Chambord et al. 2016). Cladocerans are characterized by high levels of EPA and ARA and this is thought to be related to a life history strategy focused on high rates of somatic growth (Persson & Vrede 2006). In contrast, copepods have higher relative DHA levels because this fatty acid is critical for nervous system development (Arts et al. 2009). Copepods feature more developed nervous systems compared to cladocerans and this is a function of active hunting of prey, mate location, and predator avoidance (Dalsgaard et al. 2003). Carnivorous crustacean zooplankton have shown to be richer in PUFAs and this is thought to be

related to their food source (rotifers and smaller bodied cladoceran/copepods compared to phytoplankton) (Arts et al. 2009).

Here we explore the species composition and variability in fatty acid composition of the lower food web at the freshwater/saltwater interface of an estuarine fish nursery, the Chowan River, North Carolina, USA. The Chowan River is considered a critical habitat for larval and juvenile blueback herring (*Alosa aestivalis*), alewife (*A. pseudoharengus*), collectively known as river herring (NCDMF 2007). The river herring are of interest because they have been severely overfished and a moratorium on harvest is in place at various locations along the eastern United States, including North Carolina (ASMFC 2012). The Chowan River also serves as a nursery habitat for American shad (*A. sapidissima*) and striped bass (*Morone saxatilis*); however, the status of the habitat for the latter species is unknown (Greene et al. 2009).

The overall goal of our study was to determine if species and fatty acid composition of the lower food web could be used to indicate habitat quality of an estuarine fish nursery. In order to achieve this goal, we examined the spatial and temporal variability in species composition of microzooplankton and mesozooplankton as well as the fatty acid composition of the seston, microzooplankton, and mesozooplankton during the period of larval fish residency in the Chowan River. Our specific objectives were to determine: 1) if differences in the species composition and fatty acid composition were present in the system; 2) if so, were there patterns in species composition and fatty acid composition in time and space; 3) if particular species were related to the species composition patterns and if particular fatty acids were related to the fatty acid composition patterns; 4) if changes in species and fatty acid composition were related to changes in salinity dynamics; 5) if patterns in fatty acid composition correlated across trophic levels. We hypothesized that species composition would be related to salinity and that fatty acid

profiles of micro- and mesozooplankton would relate to species composition and would reflect that of the seston. If supported, this would suggest that the quality of the larval fish forage, based on fatty acids, could be used to assess fish nursery quality.

Materials and Methods

Study Site

The Chowan River is one of the largest tributaries that drains into the Albemarle Sound (Fig. 1a and 1b) and is the 12th largest river basin in North Carolina (NCDENR 2006). It is mainly a freshwater estuary that experiences intermittent salinity intrusion, mainly in the winter months (Leech et al. 2009). The Chowan River was classified as “nutrient sensitive waters” in 1979 (NCDENR 2006) and has routinely experienced algal blooms and low dissolved oxygen levels ($< 3.0 \text{ mg L}^{-1}$). The entire river is classified as a Strategic Habitat Area for larval and juvenile river herring (NCDMF 2007). Sampling took place south of Holiday Island on a 34 km transect of Chowan River (Fig. 1c). Seven locations (4 km apart) were sampled between Holiday Island and the river mouth (Fig 1c). Sampling occurred on 10-11 April, 31 May, and 25 June 2013, dates that span the residency for larvae of alewife, blueback herring, and striped bass. For our study, we divided the river into three sections: upper, middle, and lower. The main differences among these sections were (1) the distance from the Albemarle Sound and (2) the potential influence of salinity.

Sample collection

Water column properties

Vertical profiles of temperature ($^{\circ}\text{C}$) and salinity were measured with a conductivity, temperature, and depth sensor (CTD, Yellow Springs Instruments, Castaway). Water samples were collected at a depth of 3 meters with a Niskin water sampler.

168 *Zooplankton*

169 Water depths ranged from 5.27 to 7.56 m during zooplankton sampling. Two horizontal
 170 net tows were made with 0.5 m diameter nets of two different mesh sizes (60 and 200 μm). Two
 171 mesh sizes were used in order to generate an adequate representation of the zooplankton for the
 172 size range $> 60 \mu\text{m}$. The zooplankton samples between 60 and 200 μm are designated
 173 microzooplankton and the $> 200 \mu\text{m}$ zooplankton samples are designated mesozooplankton
 174 throughout the remainder of the paper. The zooplankton net was towed obliquely through the
 175 water for 1 minute (species composition) and 2 minutes (fatty acid composition) at an average
 176 boat speed of 0.75 m s^{-1} . The volume filtered was calculated using the volume of a cylinder (
 177 $V = \pi r^2 L$), where r was the radius of the plankton net (0.25 m) and L was determined using the
 178 boat speed (m s^{-1}) and the tow time (s). Each identification and count sample, depending on mesh
 179 size, was filtered through a 60 or 200 μm filter, and zooplankton for composition were preserved
 180 in a 120 mL glass jar with 10 ml of 10% buffered formaldehyde, sucrose, and filtered water. The
 181 addition of sucrose to the formalin helps to reduce ballooning of cladoceran bodies and inflation
 182 of their carapace (Haney and Hall 1973). The 60 μm sample had a half tablet of Alka Seltzer
 183 added to keep rotifers from pulling in critical body parts (legs and arms) to ease identification
 184 (Chick et al. 2010). The zooplankton samples for fatty acid analysis were collected at seven sites
 185 on the Chowan River for April and June, and three sites in May. Due to limitations related to
 186 sampling preparation in May, a subset of the field sites was sampled for fatty acid analyses. The
 187 zooplankton samples for fatty acids were placed in a 1000 mL plastic container on ice, and
 188 processed in the laboratory.

189 *Laboratory Processing*

190 *Zooplankton Identification*

191 Samples were filtered through a sieve (60 or 200 μm) to remove the sugar formalin
 192 solution, and then added to a beaker with a known volume of water. A total of three subsamples
 193 (2 mL per subsample for microzooplankton and 5 mL per subsample for mesozooplankton) were
 194 analyzed for community composition using a Hensen-Stempel pipette. Organisms were
 195 identified using a dissecting microscope and enumerated using a Ward counting wheel. The
 196 zooplankton were identified to genus except for the freshwater copepods that were identified to
 197 order. Copepod nauplii were grouped together because identification can be difficult at this stage
 198 (Johnson and Allen 2012). If a species in a subsample comprised greater than 500 individuals,
 199 then that species was not counted for the other two subsamples. Species abundances (A) were
 200 determined using the equation: $A = A_s(V_t V_s^{-1})$ where A_s is the number of individuals in the
 201 subsample, V_t is the total volume of water in the beaker, and V_s is the volume of the subsample.

202 *Lipid and fatty acid samples*

203 The water samples (300 mL) were concentrated on a 0.7 μm WhatmanTM GF/F filter (47
 204 mm diameter) and stored at -80°C until ready to process, which constituted the seston material.
 205 The zooplankton samples were filtered through 60 and 200 μm sieves stacked to collect species
 206 based on size. Each sample was visually analyzed to determine the dominant species with a
 207 dissecting microscope, and detritus and phytoplankton were removed. The samples were
 208 concentrated on a GF/F filter (47 mm diameter) by mesh size (60, 200 μm), and stored at -80°C
 209 until ready to process.

210 Total lipids were extracted with chloroform-methanol (2:1, v/v) containing 0.01%
 211 butylated hydroxytoluene as an antioxidant (Folch et al. 1957). The organic solvent was

evaporated under a stream of nitrogen and lipid concentration determined gravimetrically. Transmethylation of fatty acids was done according to the method described by Metcalfe and Schmitz (1969). A known amount of nonadecanoate acid (19:0) dissolved in hexane at a concentration of 8 mg ml⁻¹ (Nu Check Prep Inc.) was added as internal standard. The fatty acid methyl esters (FAME) were separated by gas chromatography (Agilent 7890A Gas Chromatograph, Agilent Technologies, Inc.) using a 7693 mass spectrometer detector (Agilent Technologies, Inc.), a capillary column (Omegawax™ 250 fused silica capillary column, 30 mm x 0.25 mm and 0.25 mm film thickness, Supleco®), and a 7890A autoinjector (Agilent Technologies, Inc.). Helium was used as the carrier gas at a flow of 1.3 ml min⁻¹ and the injection volume was 2 mL. Initial temperature of the oven was 175°C for 26 min, which was increased to 205°C by increments of 2°C min⁻¹, then held at 205°C for 24 min. The source and analyzer for the mass spectrometer was set at 230°C. The individual fatty acid methyl esters were identified by comparing the retention times of authentic standard mixtures (FAME mix 37 components, Supleco) and quantified by comparing their peak areas with that of the internal standard (Czesny and Dabrowski 1998). The results of individual fatty acid composition are expressed in percentage of total identified FAME.

Statistical Analysis

We performed a series of multivariate analyses to address our specific objectives. We used PERMANOVA a part of the PRIMER 6 statistical software package (Clarke and Gorley 2006), to test for overall differences between the microzooplankton and mesozooplankton community composition, and fatty acid profiles of seston, microzooplankton, and mesozooplankton. PERMANOVA is a non-parametric technique related to ANOVA, but uses permutations and fewer assumptions compared to the traditional ANOVA approach (Anderson

2001). As such, it is particularly well suited to multivariate data sets that violate the traditional assumptions of ANOVA and also have low sample sizes, as was our case (Anderson 2001).

If PERMANOVA detected differences, we then generated separate, Bray-Curtis similarity matrices for microzooplankton species composition (60 μm mesh), mesozooplankton species composition (200 μm mesh), seston fatty acid composition, microzooplankton fatty acid composition, and mesozooplankton fatty acid composition. A separate cluster analysis was performed using PRIMER 6 in order to reveal patterns over time and space for each similarity matrix. Each individual sample was associated with a location in the river (upper, middle, lower) and month (April, May, June) and these labels were used for visualization of samples in the cluster dendrogram. Next, a similarity percentage analysis (SIMPER) test was used to compare similarities within groups and determine the species or fatty acids that contributed to each grouping from the cluster analysis (Clarke and Gorley 2006). The SIMPER test was set at 70% cumulative contribution.

We then wanted to determine if salinity and temperature were related to the observed patterns and we used redundancy analysis for this purpose (Legendre and Legendre 1998). The redundancy analysis was carried out in the R environment (R v3.2.3, R Core Development Team 2015) using rda function (Oksanen et al. 2013) in the vegan package (Oksanen et al. 2017). Finally, we used a Mantel matrix comparison to correlate fatty acid profiles between the three groups (seston, microzooplankton, and mesozooplankton). The mantel.rtest function in the ade4 package (Oksanen et al. 2017) in the R environment (R v3.2.3, R Core Development Team 2015) was used.

Results

Salinity and Temperature

Salinity was near zero (0.02 to 0.04) throughout the river in April. During May, salinities in the upper river remained low (0.07), but the water column became stratified in the middle and lower river, with salinities ranging from 0.28-1.66. The river returned to freshwater (0.04 to 0.08) again in June due to a tropical storm that brought heavy rains for a two-week period. North Carolina experienced the second wettest June since 1895 with rainfall that ranged from 15.2 to 19.05 cm in the study area (Hiatt 2013). Water temperatures increased during the study period April ($15.7 \pm 1.1^{\circ}\text{C}$), May ($24.0 \pm 1.0^{\circ}\text{C}$) and June ($26.2 \pm 0.3^{\circ}\text{C}$).

Zooplankton Community Composition

There were significant differences between the overall microzooplankton and mesozooplankton community composition (PERMANOVA, $p = 0.001$). Microzooplankton could be separated into two distinct groups by cluster analysis at 65% similarity (Figure 2a). Group 1 consisted of the vast majority of the samples collected across April, May, and June throughout the river (Figure 2a). Group 2 consisted of two samples collected in the middle and lower river in June (Figure 2a). Both groups were dominated by rotifers and copepod nauplii, but group 2 had a higher contribution of copepod nauplii (Figure 2b and Table 1).

Three groups of mesozooplankton were differentiated at 50% similarity using cluster analysis (Figure 3a). Group 3 consisted of samples from the May collection only, Group 4 consisted of a mixture of April and June samples in primarily the upper and middle river, and Group 5 consisted of one April upper site, May upper river section and June samples throughout the river (Figure 3a). Group 3 mesozooplankton percent composition was dominated by *Acartia* spp. with a minor contribution by Bosminidae, Group 4 consisted primarily of equivalent percentages of Cyclopoida and Bosminidae, and Group 5 mesozooplankton percent community

composition was characterized by higher percentages of *Leptodora* spp. and Calanoida (Figure 3b and Table 1).

Fatty Acid Composition

A total of 24 specific fatty acids were found in all samples (Tables A1-A3). Fatty acids were first separated into broad categories: saturated fatty acids (SFA), monounsaturated fatty acids (MUFA), and polyunsaturated fatty acids (PUFA) (Figure 4a). Seston had a higher percent of SFA, and lower percent of MUFA and PUFA compared to micro- and mesozooplankton (Figure 4a). Mesozooplankton and microzooplankton had a similar percent composition of SFA, MUFA, and PUFA (Figure 4a). There were eight dominant fatty acids found in all the samples, but the percent composition varied (Figure 4b). The most common SFA was palmitic acid (16:0), the most common MUFAs were palmitoleic acid (16:1 ω -7) and oleic acid (18:1 ω -9), and the most common PUFAs were ALA, 18:4 ω -3, EPA, and DHA (Figure 4b; Tables A1-A3). A comparison of MUFAs and PUFAs among the seston and zooplankton showed that seston had the lowest overall percentages of MUFAs and PUFAs (PERMANOVA, $p = 0.001$, Figure 4b). There was a difference between the microzooplankton fatty acid profile and the mesozooplankton fatty acid profile (PERMANOVA, $p = 0.025$). The microzooplankton fatty acid profile was characterized by a higher percentage of 18:1 ω -9 compared to the other MUFAs and PUFAs. In contrast, the mesozooplankton had the highest percent composition attributed to two PUFAs, EPA and DHA (Figure 4b).

Seston

Three groups were designated at 60% similarity using cluster analysis for the seston fatty acid composition (Figure 5a). The groups showed no distinct pattern in terms of sampling time or location (Figure 5a). Seston fatty acid composition of Group A was characterized by 18:1 ω -9,

16:1 ω -7, ALA, and EPA, Group B by 16:1 ω -7, 18:1 ω -9, 18:2 ω -6 and EPA, and Group C had similar percentage composition of MUFAs and PUFAs, except 18:2 ω -6 (Figure 5b and Table 2).

Microzooplankton

Three groups were designated at 70% similarity using cluster analysis for the microzooplankton fatty acid composition (Figure 6a). The groups segregated temporally, with Group D and E consisting of April samples only, and Group F consisted of May and June samples only (Figure 6a). The Group D fatty acids were dominated by 18:1 ω -9 and to a lesser extent, 18:2 ω -6, Group E showed similar percent composition among the fatty acids, with higher percentages of 18:1 ω -9 and EPA, and Group F also had similar percent composition of fatty acids; however, the PUFAs EPA and DHA had significant contribution (Figure 6b and Table 2).

Mesozooplankton

Four groups were designated by cluster analysis at 77% similarity for the mesozooplankton fatty acid composition (Figure 7a). Groups G and I consisted of April samples only whereas Group J consisted of May and June samples only (Figure 7a). Group H showed spatial separation, consisting of primarily upper river locations across all of the months (Figure 7a). The Group G and H fatty acids were dominated by 18:1 ω -9, Group 8 by 16:1 ω -7, 18:1 ω -9, and EPA, Group I had similar percentages of fatty acids with ALA, DHA and EPA having higher percentages, and Group J had DHA and EPA as dominant components of the fatty acids (Figure 7b and Table 2).

Redundancy analysis and Mantel matrix correlations

Salinity was the main factor found to be correlated to observed patterns in the species composition of both microzooplankton and mesozooplankton (Redundancy, $p = 0.004$). However, salinity was not correlated to changes in the seston (Redundancy, $p = 0.490$) or

microzooplankton (Redundancy, $p = 0.27$) fatty acid profiles. Salinity was associated with the changes in the mesozooplankton fatty acid profiles (Redundancy, $p = 0.034$). Seston fatty acid profiles were correlated to the microzooplankton fatty acid profiles ($p = 0.013$) based on the Mantel matrix comparison, but not correlated to the mesozooplankton fatty acid profiles ($p = 0.340$). The microzooplankton fatty acid profiles were correlated to the mesozooplankton fatty acid profiles (Mantel, $p = 0.059$).

Discussion

We found temporal and spatial differences in the species and fatty acid composition of the lower food web that were mainly related to a salinity intrusion that occurred during the study period during May. Prior to the salinity intrusion, larval fish would have encountered a freshwater plankton assemblage that was dominated by rotifers, Bosminidae, and cyclopoid copepods throughout the river. This assemblage was proportionally higher in EPA relative to DHA. During the salinity intrusion, the microzooplankton remained dominated by rotifers; however, the mesozooplankton community became dominated by the copepod *Acartia* spp.. Concurrently, the proportion of DHA increased and remained elevated into June, particularly in the middle and lower river. The intrusion of saline water increased the overall proportion of omega-3 fatty acids in the River, presumably due to the increased fraction of micro- and mesozooplankton feeding on a more marine-like phytoplankton based food web and this signal propagated through the food web. Additionally, we observed that FA appeared to be incorporated relatively unchanged in micro and mesozooplankton in terms of relative composition; however, MUFA and PUFA percent compositions increased in zooplankton relative to seston. This suggested that MUFA and PUFA are bioaccumulated at higher trophic levels, as seen in other studies (Persson and Verde 2006; Gladyshev et al. 2010; Ravett, Brett,

Arhonditsis 2010; Burns, Brett, and Schallenberg 2011). Overall, the FA composition of the food web indicated that the Chowan River is likely to provide nutrition in terms of FA composition for larval fish growth and development. This is based on the presence of higher chain (>20 carbons) PUFAs present in the mesozooplankton throughout the nursery.

The seston fatty acid composition consisted mainly of saturated fatty acids. Seston from freshwater and estuarine systems typically has a large percentage of SFA and this fraction has been attributed to detrital input, as opposed to originating from phytoplankton (Persson and Vrede 2006; Gladyshev et al. 2010; Ravett, Brett, Arhonditsis 2010; Burns, Brett, and Schallenberg 2011; Gonclaves et al 2012). Müller-Navarra et al. (2004) and Bec et al. (2010) analyzed seston and found phytoplankton only explained 27% of variance in FA composition and concluded that detritus and heterotrophic organisms also needed to be considered (Müller-Navarra et al. 2004; Bec et al. 2010). Bec et al (2010) therefore concluded that the seston can affect the fatty acid profiles of higher organisms, but may not relate individual groups of phytoplankton or microzooplankton. This agreed with our findings as seen in the reduced correlations between seston and the mesozooplankton.

We did not examine the seston composition directly by counting phytoplankton cells or examining pigment concentrations, thus we were unable to attribute the origin of particular fatty acids to phytoplankton or other sources. However, we were able to use the available literature to identify potential indicators of fatty acid origin. The top four fatty acids by percent composition varied by group, but 16:1 ω -7, 18:1 ω -9, 18:2 ω -6, and ALA were the most prevalent. Potential phytoplankton sources for these fatty acids may be diatoms, which have been shown to have increased 16:1 ω -7 and EPA in both freshwater and marine systems (Napolitano et al. 1997; Dalsgaard et al. 2003; Boschker et al. 2005; Arts et al. 2009; Bec et al. 2010) and we observed

this occurred in May and June coincident with the salinity increase. Green algae (Chlorophytes) have been shown to possess higher proportions of 18:2 ω -6, and ALA, and 18:4 ω -3 (Ahlgren et al. 1990; Dalsgaard et al. 2003; Boschker et al. 2005; Masclaux et al. 2012; Strandberg et al. 2015). Fatty acids corresponding to these phytoplankton groups were observed during April throughout the river and June in the middle and upper river. One other source of seston FA may have been present, pine pollen, which is found in large quantities during spring. Pine pollen is transported to freshwater systems via aerial deposition and floats at the surface, and the pine pollen fatty acid profile has a high percent composition of 18:1 ω -9 and 18:2 ω -6 (Masclaux et al. 2013), which can be observed in our samples. Obviously, seston FA are a mixture of multiple sources, thus the variability seen across the groups identified by the cluster analysis would be expected.

The microzooplankton fatty acid profiles were different throughout the sampling period with a change from decreased omega-3s to increased omega-3s in the system. This suggests a switch in microzooplankton diet had occurred over the sampling period and two pathways appear to be present during the study. The April microzooplankton fatty acid profiles for all river sections had a high percentage of 18:1 ω -9 and 18:2 ω -6 suggesting that the microzooplankton could be consuming either terrestrial material or chlorophytes during this time. Two sites in April had an increase in omega-3 fatty acids (ALA, 18:4 ω -3, EPA, DHA), and this would suggest a different dietary pathway that was reduced in SFA, perhaps consisting of either smaller microzooplankton such as ciliates or phytoplankton such as diatoms and/or dinoflagellates (Park and Marshall 2000; Gladyshev et al. 2010). The community was dominated by rotifers during this time and communities high in rotifer abundance have been shown to closely reflect the seston composition (Gladyshev et al. 2010). The microzooplankton fatty acid profiles in May

and June at all river locations had an increase in 16:1 ω -7, and omega-3s (ALA, EPA, and DHA). These changes can be correlated to the likely increase in diatoms and dinoflagellates from the saltwater intrusion event. The changes in June could be the increase in copepod nauplii of Calanoid copepods, and the presence of dinoflagellates and diatoms even when the system returned to freshwater. Our results are similar to systems where the phytoplankton composition was represented by diatoms and dinoflagellates by having increased 16:1 ω -7 and PUFAs (Müller-Navara et al. 2000; Dalsgaard et al. 2003; Gladyshev et al. 2010; Ravett, Brett, Arhonditsis 2010).

The mesozooplankton fatty acid profiles throughout the river in April and in the upper river in May and June were defined by higher percentages of 16:1 ω -7, 18:1 ω -9, ALA, EPA, and DHA, and mixed mesozooplankton community consisting of cladoceran and copepods. These fatty acids profiles are similar to those found in freshwater systems that have a mixed zooplankton composition (Persson and Verde 2006; Arts et al. 2009; Kainz et al. 2009; Gladyshev et al 2010; Burns et al. 2011; Masclaux et al. 2012). Cladocerans have low or no DHA compared to copepods and high EPA levels have been shown to correlate with the high somatic growth rates of cladoceran (Persson and Verde 2006). A saltwater intrusion changed the mesozooplankton species composition in May resulting in numerical dominance by *Acartia* spp. in the lower and middle sections of the river. *Acartia* spp. is the dominant copepod species in temperate, estuarine systems (Ambler et al. 1985; Orsi 1986; Cervetto et al. 1999; Mouny and Dauvin 2002; Kimmel and Roman 2004; Lawrence et al. 2004; Islam et al. 2005). The mesozooplankton fatty acid profiles in May were represented by 16:1 ω -7, EPA, and the highest observed percentages of DHA. This is clearly a reflection of the dominance of *Acartia* spp. in the system and a diet primarily consisting of marine algae higher in omega-3 FAs (Stottrup et al.

1999; Persson and Verde 2006; Arts et al. 2009; Kainz et al. 2009; Gladyshev et al 2010; Masclaux et al. 2012). Mesozooplankton fatty acid percent composition in June at the lower and middle site remained similar to that observed in May, despite the species composition having returned to a mix of cladocerans and copepods. This suggests that physical shifts in the system that result in seston changes may persist in the system despite shifts in zooplankton community composition.

The relevance of the food web fatty acid composition can be determined by examining the potential feeding behavior of larval fish within the Chowan River nursery. Alewife and blueback herring start feeding on smaller cladocerans and copepods at about 6 mm total length (Mullen et al. 1986). Binon (2011) reported that river herring at 6 mm notochord length had a maximum gape width of 400 μ m, and estimated maximum prey size of 200 μ m, which would result in mesozooplankton being an important food resource. In the Connecticut River, the diet for blueback herring were dominated by rotifers for fish 5-12 mm, Bosminidae for fish 12-16 mm, and cyclopoid copepods for fish >16mm in total length (Crecco and Blake 1983). Based on these dietary studies, the larval and juvenile river herring would be feeding across the size range of zooplankton prey that we sampled; however, fish would be consuming primarily microzooplankton early in the year (April) and mesozooplankton later in the season (May and June). In April, two pathways for FA propagation were present in the microzooplankton. Thus, fish feeding during this time may experience variability in the quality of the microzooplankton prey in terms of percentage of PUFAs. Larval fish need PUFAs (ALA, EPA and DHA) for growth, visual acuity, survival, and development of normal pigmentation (Bell et al. 1995; Bell and Sargent 1996; Rainuzzo et al. 1997; Sargent et al. 1999; Rossi et al. 2006). The shift to larger prey later in the year (May and June) resulted in a change in prey quality as the relative

percentage of EPA and DHA increased. This was the result of a salinity intrusion into the middle and lower reaches of the estuary that was associated with dominance of the cladoceran *Acartia* spp. and a significant increase in DHA and EPA. This could allow larval and juvenile fish to consume prey with a higher proportion of PUFAs. It is unknown if river herring can elongate precursor FA into PUFAs. Even if fish can convert precursor FA, larval fish could not receive all nutritional need for those fatty acids (Agaba et al. 2005). The larval fish would not have to use energy for the conversion, and could continue to put energy into growth (Wacker and von Elert 2001; Rossi et al 2006). This allows the larval fish to survive and grow past the critical period.

The fish nursery present in the lower Chowan River may undergo significant changes during the critical time of larval fish growth and our results demonstrate how changes in the seston community may propagate through the food web. The results also highlight that additional information concerning the fatty acid composition of the zooplankton prey base for larval fish can provide insight into habitat quality, our stated goal. Sheaves et al. (2004) pointed out the need to expand the nursery habitat concept to include relevant ecosystem processes, particularly resource dynamics. This research begins to explore the mechanisms that allow nursery habitat to function. We plan further research to investigate the linkage between the seston community fatty acid composition, the zooplankton community fatty acid composition, and larval fish to determine how lower food web variability relates to larval fish condition and survival.

Acknowledgements

I would like to thank S. Lichti, C. Krahforst, J. Osborne, A. Powell, M. Baker, and E. Diaddorio for help in the field collection. I would like to thank L. Stratton, C. Kolb, and R. Pattridge for help in Dr. Jacques Rincharde's lab in processing my fatty acid samples. I would like to thank Dr.

463 Ariane Peralta for help with the improved statistical analysis, and two anonymous reviewers for
 464 comments, which helped to improve the manuscript.

465 Work Cited

- 466 Able, K.W. (2005) A re-examination of fish estuarine dependence: Evidence for connectivity
467 between estuarine and ocean habitats. *Estuarine, Coastal, and Shelf Science* 64: 5-17.
- 468 Agaba, M.K., D.R. Tocher, X.Z. Zheng, C.A. Dickson, J.R. Dick, and A.J. Teale. (2005) Cloning
469 and functional characterization of polyunsaturated fatty acid elongases of marine and freshwater
470 teleost fish. *Journal of Comparative Physiology B* 142: 342-352.
- 471 Ahlgren, G., L. Lundstedt, M. Brett, and C. Forsberg. (1990) Lipid composition and food quality
472 of some freshwater phytoplankton for cladoceran zooplankters. *Journal of Plankton Research* 12:
473 809-818.
- 474 Ambler, J.W., J.E. Cloern, and A. Hutchinson. (1985) Seasonal cycles of zooplankton from San
475 Francisco Bay. *Hydrobiologia* 129: 177-197.
- 476 Anderson, M. J. (2001), A new method for non-parametric multivariate analysis of variance.
477 *Austral Ecology*, 26: 32–46. doi:10.1111/j.1442-9993.2001.01070.pp.x
- 478 Arts, M.T., M.T. Brett, and M.J. Kainz. (2009) *Lipids in aquatic ecosystems*. Springer. New
479 York, New York.
- 480 ASMFC (Atlantic States Marine Fisheries Commission). (2012) River herring benchmark stock
481 assessment volume 1. Stock Assessment Report No. 12-02, May 2012.
- 482 Beck, M.W., K.L. Heck Jr., K.W. Able, D.L. Childers, D.B. Eggleston, B.M. Gillanders, B.
483 Halpern, C.G. Hays, K. Hoshino, T. J. Minello, R.J. Orth, P.F. Sheridan, and M.P. Weinstein.
484 (2001) The identification, conservation, and management of estuarine and marine nurseries for
485 fish and invertebrates. *BioScience* 51: 633- 641.
- 486 Bell, M.V., R.S. Batty, J.R. Dick, K. Fretwell, J.C. Navarro and J.R. Sargent. (1995) Dietary
487 deficiency of docosahexanoic acid impairs vision at low light intensities in juvenile herring
488 (*Clupea harengus* L.). *Lipids* 30 (5): 443-449.
- 489 Bec, A., M.E. Perga, C. Desvillettes, and G. Bourdier. (2010) How well can the fatty acid content
490 of lake seston be predicted from its taxonomic composition? *Freshwater Biology* 55: 1985-1972.
- 491 Bell, M.V. and J.R. Sargent. (1996) Lipid nutrition and fish recruitment. *Marine Ecology*
492 *Progress Series* 134: 315-316.
- 493 Binion, S. (2011) Evaluating spatial and temporal overlap between larval alosines and potential
494 zooplankton prey in lower Roanoke River and Albemarle Sound, North Carolina. Thesis. East
495 Carolina University, Greenville, NC.
- 496 Boehlert, G.W. and B.C. Mundy. (1988) Role of behavioral and physical factors in larval and
497 juvenile fish recruitment to estuarine nursery areas. *American Fisheries Society Symposium*
498 3:51-67.
- 499 Boesch, D.F., and R.F. Turner. (1984) Dependence of fisheries species on salt marshes: the role
500 of food and refuge. *Estuaries and Coasts* 7: 460-468.
- 501 Boschker, H.T., J.C. Krombamp, and J.J. Middelburg. (2005) *Limnology and Oceanography* 50:
502 70-80.
- 503 Brett, M.T. and D.C. Müller-Navarra. (1997) The role of highly unsaturated fatty acids in aquatic
504 foodweb processes. *Freshwater Biology* 38: 483-499.
- 505 Brett, M.T., D.C. Müller-Navarra, A.P. Ballantyne, J.L. Ravet, C.R. Goldman. (2006) *Daphnia*
506 fatty acid composition reflects that of their diet. *Limnology and Oceanography* 51: 2428-2437.
- 507 Burns, C., M.T. Brett, and M. Schallenberg. (2010) A comparison of the trophic transfer of fatty
508 acids in freshwater plankton by cladocerans and calanoid copepods. *Freshwater Biology* 56:889-
509 903.

510 Cervetto, G., R. Gaudy, and M. Pagano. (1999) Influence of salinity on the distribution of
511 *Acartia tonsa* (Copepoda, Calanoida). Journal of Experimental Marine Biology and Ecology
512 239: 33-45.

513 Chambord, S., T. Maris, F. Colas, T. van Engeland, A.C. Sossu, F. Azemar, M. Le Coz, T. Cox,
514 L. Buisson, S. Souissi, P. Meire, and M. Tackx. (2016) Mesozooplankton affinities in a
515 recovering freshwater estuary. Estuarine, Coastal, and Shelf Science 177: 47-59.

516 Chick, J.H. and M.J. Van Den Avyle. (1999) Effects of zooplankton spatial variation on growth
517 of larval striped bass: an experimental approach. Transactions of the American Fisheries Society
518 128: 339-351.

519 Chick, J.H., A.P. Levchuk, K.A. Medley and J.H. Havel. (2010) Underestimation of rotifer
520 abundance a much greater problem than previously appreciated. Limnology and Oceanography:
521 Methods 8, 79-87.

522 Clarke, K.R. and R.N. Gorley. (2006) Primer V6: User Manual/Tutorial. Primer-E Ltd.
523 Plymouth, United Kingdom.

524 Cooper, J.E., R.A. Rulifson, J.J. Isely, S.E. Winslow. (1998) Food habits and growth of juvenile
525 striped bass, *Morone saxatilis*, in Albemarle Sound, North Carolina. Estuaries 21: 307-317.

526 Copeman, L.A., C.C. Parrish, J.A. Brown, M. Harel. (2002) Effects of docosahexaenoic,
527 eicosapentaenoic, and arachidonic acids on the early growth, survival, lipid composition and
528 pigmentation of yellowtail flounder (*Limanda ferruginea*): a live food enrichment experiment.
529 Aquaculture 210: 285-304.

530 Czesny, S. and K. Dabrowski. (1998) The effect of fatty acids concentration in wild and
531 domesticated walleye (*Stizostedion vitreum*) eggs on embryos' viability. Aquatic Living
532 Resources 11: 371-378.

533 Dalsgaard, J., M. St. John, G. Kattner, D. Müller-Navarra and W. Hagen. (2003) Fatty acids
534 trophic markers in the pelagic marine environment. Advances in Marine Biology 46: 225-340.

535 Desvillettes, C.H., G. Bourdier, C.H. Amblard and B. Barth. (1997) Use of fatty acids for the
536 assessment of zooplankton grazing on bacteria, protozoans and microalgae. Freshwater Biology
537 38, 629-637.

538 Farkas, T., and S. Herodek. (1964) The effect of environmental temperature on the fatty acid
539 composition of crustacean plankton. Journal of Lipid Research 5: 369-373.

540 Folch, J., M. Lees, and G.H. Sloane Stanley. (1957) A simple method for isolation and
541 purification of total lipids from animal tissues. Journal of Biological Chemistry 226: 497-509.

542 Fraser, A.J., J.R. Sargent, J.C. Gamble, and D.D. Seaton. (1989) Formation and transfer of fatty
543 acids in an enclosed marine food chain comprising phytoplankton, zooplankton, and herring
544 (*Clupea harengus* L.) larvae. Marine Chemistry 27: 1-18.

545 Gladyshev, M.I., N.N. Sushchik, O.N. Makhutova, O.P. Dubovskaya, E. S. Kravchuk, and E.B.
546 Khromechek. (2010) Correlations between fatty acid composition of seston and zooplankton and
547 effects of environmental parameters in a eutrophic Siberian reservoir. Limnologica 40: 343-357.

548 Goncalves, A.M.M., U.M. Azeiterio, M.A. Pardal and M. De Troch. (2012) Fatty acid profiling
549 reveals seasonal and spatial shifts in zooplankton diet in a temperate estuary. Estuarine, Coastal
550 and Shelf Science 109: 70-80.

551 Goulden, C.E. and A.R. Place. (1990) Fatty acid synthesis and accumulation rates in daphniids.
552 Journal of Experimental Zoology 256: 168-178.

553 Greene, K.E., J.L. Zimmerman, R.W. Laney, and J.C. Thomas-Blate. (2009) Atlantic coast
554 diadromous fish habitat: A review of utilization, threats, recommendations for conservation, and

research needs. Atlantic States Marine Fisheries Commission Habitat Management Series No. 9, Washington, D.C.

Gulati, R.D. and W.R. Demott. (1997) The role of food quality for zooplankton: remarks on the state-of-the-art, perspectives and priorities. *Freshwater Biology* 38: 753-768.

Haney, J.F. and D.J. Hall. (1973) Sugar-coated *Daphnia*: A preservation technique for Cladocera. *Limnology and Oceanography* 18: 331-333.

Hiatt, Ashley. (2013) Climate Summary: June 2013. State Climate Office of North Carolina. <http://nc-climate.ncsu.edu/climateblog?id=28>. Accessed 15 September 2013.

Hjort, J. (1914) Fluctuations in the great fisheries of northern Europe viewed in the light of biological research. *Rapp. P.-v. Reun. Cons. Perm. Int. Explor. Mer.* 29: 1-228.

Islam, M.S., H. Ueda, and M. Tanaka. (2005) Spatial distribution and trophic ecology of dominant copepods associated with turbidity maximum along the salinity gradient in a highly embayed estuarine system in Ariake Sea, Japan. *Journal of Experimental Marine Biology and Ecology* 316: 101-115.

Iverson, S.J., C. Field, W.D. Bowen, and W. Blanchard. (2004) Quantitative fatty acid signature analysis: a new method of estimating predator diets. *Ecological Monographs* 74: 211-235.

Johnson, W.S. and D.M. Allen. (2012) *Zooplankton of the Atlantic and Gulf coasts: a guide to their identification and ecology*, 2nd edition. John Hopkins University Press. Baltimore, Maryland.

Kainz, M., M.T. Arts and A. Mazumder. (2004) Essential fatty acids in the planktonic food web and their ecological role for higher trophic levels. *Limnology and Oceanography* 49: 1748-1793.

Kainz, M.J., M. Perga, M.T. Arts and A. Mazumder. (2009) Essential fatty acid concentration of different seston sizes and zooplankton: a field study of monomictic coastal lakes. *Journal of Plankton Research* 31: 635-645.

Kimmel D.G., and M.R. Roman. (2004) Long-term trends in mesozooplankton abundance in Chesapeake Bay USA: influence of freshwater input. *Marine Ecology Progress Series* 267: 71-83.

Kimmerer, W.J. (2002) Effects of freshwater flow on abundance of estuarine organisms: physical effects or trophic linkages? *Marine Ecology Progress Series* 243: 39-55.

Lawrence, D., I. Valiela, and G. Tomasky. (2004) Estuarine calanoid copepod abundance in relation to season, salinity, and land-derived nitrogen loading, Waquoit Bay, MA. *Estuarine, Coastal and Shelf Science* 61: 547-557.

Leech, D., S. Ensign and M. Piehler. (2009) Zooplankton Assessment Project (ZAP): Reassessing prey availability for river herring in the Chowan River Basin- Year 2. North Carolina Sea Grant Report. FRG 08-EP-06/ 09-EP-03.

Legendre P., Legendre L. (1998) *Numerical Ecology*. 2nd Ed. Amsterdam: Elsevier

Malzahn, A.M., N. Aberle, C. Clemmesen, and M. Boersma. (2007) Nutrient limitation of primary producers affects planktivorous fish condition. *Limnology and Oceanography* 52: 2062-2071.

Marques, S.C., U.M. Azeiteiro, J. C. Marques, J. M. Neto, and M.A. Pardal. 2006. Zooplankton and ichthyoplankton communities in a temperate estuary: spatial and temporal patterns. *Journal of Plankton Research* 28: 297-312.

Martino, E.J. and E.D. Houde. (2010) Recruitment of striped bass in Chesapeake Bay: spatial and temporal environmental variability and availability of zooplankton prey. *Marine Ecology Progress Series* 409: 213-228.

Masclaux, H., A. Bec, M.J. Kainz, F. Perriere, C. Desvillettes and G. Bourdier. (2012) Accumulation of polyunsaturated fatty acids by cladocerans: effects of taxonomy, temperature and food. *Freshwater Biology* 57: 696-703.

Masclaux, H., M.E. Perga, M. Kagami, C. Desvillettes, G. Bourdier, and A. Bec. (2013) How pollen organic matter enters freshwater food web. *Limnology and Oceanography* 58: 1185-1195.

McHugh, J.L. (1967) Estuarine nekton. American Association for the Advancement of Science Publications 83: 581-620.

Metcalfe, L.D and A.A. Schmitz. (1961) The rapid preparation of fatty acid esters for gas chromatographic analysis. *Analytical Chemistry* 33: 363-364.

Moderan, J., V. David, P. Bouvais, P. Richard and D. Fichet. (2012) Organic matter exploitation in a highly turbid environment: planktonic food web in the Charente estuary, France. *Estuarine, Coastal Shelf Science* 98: 126-137.

Mouny, P. and J.C. Dauvin. (2002) Environmental control of mesozooplankton community structure in the Seine estuary (English Channel). *Oceanologica Acta* 25: 13-22.

Mullen, D.M., C.W. Fay, and J.R. Moring. (1986) Species profiles: life history and environmental requirements of coastal fishes and invertebrates (North Atlantic) alewife/blueback herring. Biological Report 82. USFWS and Army Corps of Engineers.

Müller-Navarra, D.C., M.T. Brett, A.M. Liston and C.R. Goldman. (2000) A highly unsaturated fatty acid predicts carbon transfer between primary producers and consumers. *Nature* 403: 74-77.

Müller-Navarra, D.C., M.T. Brett, S. Park, S. Chandra, A. P. Ballantyne, E. Zorita, and C.R. Goldman. 2004. Unsaturated fatty acid content in seston and tropho-dynamic coupling in lakes. *Nature* 427: 69-72.

Müller-Navarra, D. (2008) Food web paradigms: the biochemical view on trophic interactions. *International Review Hydrobiology* 93: 489-505.

Napolitano, G.E., R.J. Pollero, A.M. Gayoso, B.A. MacDonald and R.J. Thompson. (1997) Fatty acid as trophic markers of phytoplankton blooms in the Bahia Blanca Estuary (Buenos Aires, Argentina) and in Trinity Bay (Newfoundland, Canada). *Biochemical Systematics and Ecology* 25: 739-755.

NCDENR (North Carolina Department of Environment and Natural Resources). (2006) Basinwide assessment report Chowan River Basin. NCDENR. Raleigh, North Carolina.

NCDMF (North Carolina Division of Marine Fisheries). (2007) North Carolina fisheries management plan amendment 1 river herring. NCDMF. Morehead City, North Carolina.

Newton, G.M. (1996) Estuarine ichthyoplankton ecology in relation to hydrology and zooplankton dynamics in a salt-wedge estuary. *Marine and Freshwater Research*. 47: 99-111.

Odum, E.P. (1980) The status of three ecosystem-level hypotheses regarding salt marsh estuaries: tidal subsidy, outwelling, and detritus-based food chains. In: Kennedy VS (ed) *Estuarine Perspectives*. Academic, New York.

Orsi, J.J. and W.L. Mecum. (1986) Zooplankton distribution and abundance in the Sacramento-San Joaquin delta in relation to environmental factors. *Estuaries* 9, 326-339.

Park, G.S., and H.G. Marshall. (2000) The trophic contributions of rotifers in tidal freshwater and estuarine habitats. *Estuarine, Coastal, and Shelf Science* 51: 729-742.

Paulsen, M., C. Hammer, A.M. Malzahn, P. Polte, C. von Dorrien, and C. Clemmesen. (2014) Nutritional situation for larval Atlantic herring (*Clupea harengus* L.) in two nursery areas in the western Baltic Sea. *ICES Journal of Marine Science* 71: 991-1000.

Persson, J., and T. Verde. (2006) Polyunsaturated fatty acids in zooplankton: variation due to taxonomy and trophic position. *Freshwater Biology* 51: 887-900.

646 Rainuzzo, J.R., K.I. Reitan, and Y. Olsen. (1997) The significance of lipids at early stages of
647 marine fish: a review. *Aquaculture* 155: 103-115.

648 Ravet, J.L., M.T. Brett, and G.B. Arhonditsis. (2010) The effects of seston lipids on zooplankton
649 fatty acid composition in Lake Washington, Washington, USA. *Ecology* 9: 180-190.

650 Rossi, S., A. Sabates, M. Latasa and E. Reyes. (2006) Lipid biomarkers and trophic linkages
651 between phytoplankton, zooplankton and anchovy (*Engraulis encrasicolus*) larvae in the NW
652 Mediterranean. *Journal of Plankton Research* 28: 551-562.

653 RStudio Team (2015). RStudio: Integrated Development for R. RStudio, Inc., Boston, MA URL
654 <http://www.rstudio.com/>.

655 Rulifson, R.A., J.E. Cooper, D.W. Stanley, M.E. Shepherd, S.F. Wood and D.A. Daniel. (1993)
656 Food and feeding of young finfish species in lower Roanoke River, Batchelor Bay, and western
657 Albemarle Sound, North Carolina, 1982-1988. Albemarle-Pamlico Estuarine Study APES 90-
658 16:21-27 to U.S. Environmental Protection Agency.

659 Sargent, J., L. McEvoy, A. Estevez, G. Bell, M. Bell, J. Henderson, D. Tocher. (1999) Lipid
660 nutrition of marine fish during early development: current status and future directions.

661 Sheaves, M., R. Baker, I. Nagelkerken, and R.M. Connolly. (2015) True value of estuarine and
662 coastal nurseries for fish: incorporating complexity and dynamics. *Estuaries Coast* 38: 401-414.

663 Sheaves, M. (2016) Simple processes drive unpredictable differences in estuarine fish
664 assemblages: Baselines for understanding site specific ecological and anthropogenic impacts.
665 *Estuarine, Coastal and Shelf Science* 170: 61-69.

666 Smyntek, P.M., M.A. Teece, K.L. Schulz, and A.J. Storch. (2008) Taxonomic differences in the
667 essential fatty acid composition of groups of freshwater zooplankton relate to reproductive
668 demands and generation time. *Freshwater Biology* 53: 1768-1782.

669 Stottrup, J.G., J.G. Bell and J.R. Sargent. (1999) The fate of lipids during development and cold
670 storage of eggs in the laboratory-reared calanoid copepod, *Acartia tonsa* Dana, and in response
671 to different algal diets. *Aquaculture* 176: 257-269.

672 Tackx, M.L., N. de Pauw, R. van Mieghem, F. Azemar, A. Hannouti, S. van Damme, F. Fiers, N.
673 Daro, and P. Meire. (2004) Zooplankton in the Schelde estuary, Belgium and the Netherlands.
674 Spatial and temporal patterns. *Journal of Plankton Research* 26: 133-141.

675 Taipale, S., P. Kankaala, K. Hamalainen, R.I. Jones. (2008) Seasonal shifts in the diet of the lake
676 zooplankton revealed by phospholipid fatty acid analysis. *Freshwater Biology* 54: 90-104.

677 Townsend, D.W. 1983. The relations between larval fishes and zooplankton in two inshore areas
678 of the Gulf of Maine. *Journal of Plankton Research* 5: 145-173.

679 Vargas, C.A., R. Escribano, and S. Poulet. (2006) Phytoplankton food quality determines time
680 windows for successful zooplankton reproductive pulses. *Ecology* 87: 2992-2999.

681 Wacker, A. and E. von Elert. (2001) Polyunsaturated fatty acids: evidence for non-substitutable
682 biochemical resources in *Daphnia galeata*. *Ecology* 82: 2507-2520.

683 Walsh, H.J., L.R. Settle, and D.S. Peters. (2005) Early life history of blueback herring and
684 alewife in the lower Roanoke River, North Carolina. *Transactions of the American Fisheries*
685 *Society* 134: 910-926.

686 Winder, M. and A.D. Jassby. (2011) Shifts in zooplankton community structure: Implications for
687 food web processes in the upper San Francisco Estuary. *Estuaries and Coasts* 34: 675-690.

688 Webster, C.D. and R.T. Lovell. (1990) Response of striped bass larvae fed brine shrimp from
689 different sources containing different fatty acid compositions. *Aquaculture* 90: 49-61.

690 Webster, K.E. and R.H. Peters. (1978) Some size-dependent inhibitions of larger cladoceran
691 filterers in filamentous suspensions. *Limnology and Oceanography* 23: 1238-1245.

692 Tables

693 Table 1: Results of SIMPER analysis for each group; all the zooplankton species that contributed
 694 community composition, and their corresponding values in % are given. Dash marks represent
 695 species that were not included in the contribution of 70%

	Microzooplankton		Mesozooplankton		
	Group 1	Group 2	Group 3	Group 4	Group 5
Overall Similarity	91.18	80.18	84.39	57.16	52.52
Species	Percent Composition		Percent Composition		
Bosminidae	-	-	17.26	34.25	12.20
<i>Leptodora</i> spp.	-	-	-	-	21.80
Calanoida	-	-	-	33.32	26.09
Cyclopoida	-	-	-	10.24	9.07
<i>Acartia</i> spp.	-	-	74.66	-	-
Copepod nauplii	10.60	41.61	-	-	-
Rotifer	86.47	50.46	-	-	-

696
 697
 698
 699
 700
 701
 702
 703
 704
 705
 706
 707
 708
 709
 710
 711
 712
 713
 714
 715
 716
 717
 718
 719
 720
 721
 722
 723
 724
 725

Table 2: Results of SIMPER analysis for each group; the fatty acids that contributed to the differences in fatty acid profile, and their corresponding values in % are given. Dash marks represent fatty acids that were not included in the contribution of 70%.
Group G had a sample size < 2.

	Seston			Microzooplankton			Mesozooplankton			
	Group A	Group B	Group C	Group D	Group E	Group F	Group G	Group H	Group I	Group J
Overall Similarity	66.40	72.32	64.34	84.88	92.49	81.05	N/A	81.42	89.42	84.89
Fatty Acids	Percent Composition			Percent Composition			Percent Composition			
16:1 ω -7	3.12	6.53	-	-	-	7.98	-	9.53	-	8.51
18:1 ω -9	7.04	2.27	2.42	31.63	13.46	-	17.01	10.58	-	-
18:2 ω -6	-	-	-	12.15	-	-	-	-	-	-
18:3 ω -3 (ALA)	2.37	-	3.33	-	8.43	6.80	-	6.14	10.65	-
18:4 ω -3	-	-	-	-	10.29	-	-	-	7.05	-
20:5 ω -3 (EPA)	-	1.55	2.44	-	11.72	9.91	-	11.24	13.84	16.40
22:6 ω -3 (DHA)	-	-	1.87	-	-	9.45	-	-	10.11	16.58

Figures




Figure 1. The overview of Albemarle Sound in North Carolina (a). The close up view of the location for two main tributaries (Chowan and Roanoke Rivers), and the Albemarle Sound in North Carolina (b). The three sections used to collect zooplankton on the Chowan River (c).




Figure 2. The two microzooplankton community composition groups from cluster analysis (a) at 65% similarity. The mean microzooplankton community composition (% \pm S.D.) for the two groups from cluster analysis (b).




Figure 3. The three mesozooplankton community composition groups from cluster analysis (a) at 50% similarity. The mean mesozooplankton community composition (% \pm S.D.) for the three groups from cluster analysis (b).




Figure 4. The mean (\pm S.D.) saturated (SFA), monounsaturated (MUFA), and polyunsaturated (PUFA) fatty acid composition (%) for the seston, microzooplankton, and mesozooplankton (a). The mean fatty acid composition (% \pm S.D.) for the seston, microzooplankton, and mesozooplankton (b).




Figure 5. The three seston fatty acid composition groups from cluster analysis (a) at 60% similarity. The mean seston fatty acid composition ($\%$, \pm S.D.) for the three groups from cluster analysis (b).




Figure 6. The three microzooplankton fatty acid composition groups from cluster analysis (a) at 60% similarity. The mean microzooplankton fatty acid composition (% \pm S.D.) for the three groups from cluster analysis (b).




Figure 7. The four mesozooplankton fatty acid composition groups from cluster analysis (a) at 60% similarity. The mean mesozooplankton fatty acid composition (% \pm S.D.) for the four groups from cluster analysis (b).

798

799 **Appendix**

800 Table A1: Mean fatty acid composition (\pm standard deviation) (percentage of total fatty acids
801 detected) of seston from the Chowan River by group. SFA: saturated fatty acids, MUFA:
802 monounsaturated fatty acids, and PUFA: polyunsaturated fatty acids.

	Seston Group		
	A (8)	B (5)	C (4)
14:0	6.6 \pm 1.8	11.2 \pm 5.1	7.6 \pm 1.0
15:0	1.5 \pm 1.5	1.3 \pm 0.4	1.7 \pm 0.3
16:0	47.7 \pm 5.5	54.0 \pm 5.1	50.8 \pm 4.4
17:0	2.3 \pm 0.4	2.7 \pm 0.7	2.6 \pm 0.2
18:0	14.2 \pm 5.6	9.7 \pm 1.2	11.4 \pm 2.5
20:0	0.4 \pm 0.3	0.3 \pm 0.1	0.4 \pm 0.2
Σ SFA	72.7	79.2	74.0
16:1 ω -9	1.2 \pm 1.0	1.3 \pm 0.6	1.6 \pm 0.7
16:1 ω -7	3.1 \pm 1.9	6.3 \pm 3.3	2.1 \pm 1.9
18:1 ω -9	7.0 \pm 1.8	2.3 \pm 0.8	2.4 \pm 0.3
18:1 ω -7	0.3 \pm 0.2	0.1 \pm 0.2	0.1 \pm 0.1
20:1	0.1 \pm 0.0	0.1 \pm 0.0	0.2 \pm 0.1
Σ MUFA	11.7	10.1	6.4
18:2 ω -6	2.9 \pm 4.2	0.6 \pm 0.6	0.8 \pm 1.2
18:3 ω -3	2.4 \pm 1.4	0.9 \pm 0.5	3.3 \pm 1.3
18:4 ω -3	0.7 \pm 1.1	0.1 \pm 0.1	1.7 \pm 1.4
20:2 ω -6	0.5 \pm 0.4	0.2 \pm 0.2	0.7 \pm 0.5
20:3 ω -6	0.4 \pm 0.1	0.4 \pm 0.3	0.3 \pm 0.2
20:4 ω -6	0.9 \pm 0.6	0.8 \pm 0.4	0.8 \pm 0.5
20:3 ω -3	0.6 \pm 0.3	0.4 \pm 0.3	1.1 \pm 0.6
20:4 ω -3	0.6 \pm 1.4	0.5 \pm 0.3	2.3 \pm 1.9
20:5 ω -3	1.6 \pm 1.1	1.6 \pm 0.4	2.4 \pm 1.7
22:5 ω -6	0.7 \pm 0.5	0.8 \pm 0.8	1.2 \pm 0.6
22:5 ω -3	0.8 \pm 0.6	0.7 \pm 0.6	1.2 \pm 0.7
22:6 ω -3	1.4 \pm 1.1	1.1 \pm 0.6	1.9 \pm 1.0
Σ PUFA	13.5	8.1	17.7

803

804

805

806

807

808

809

810

811

812

813

814 Table A2: Mean fatty acid composition (percentage of total fatty acids detected) of
815 microzooplankton (<60µm) from the Chowan River by groups. SFA: saturated fatty acids,
816 MUFA: monounsaturated fatty acids, and PUFA: polyunsaturated fatty acids.

	Microzooplankton Groups		
	D (5)	E (2)	F (8)
14:0	2.6 ± 0.5	3.7 ± 0.9	5.9 ± 1.1
15:0	0.3 ± 0.1	0.5 ± 0.2	1.1 ± 0.3
16:0	27.4 ± 3.0	24.5 ± 0.3	23.5 ± 3.4
17:0	0.4 ± 0.1	0.6 ± 0.0	1.2 ± 0.3
18:0	3.4 ± 0.8	3.4 ± 0.0	7.1 ± 0.3
20:0	1.8 ± 0.4	0.8 ± 0.1	0.2 ± 0.1
ΣSFA	35.9	33.5	39.0
16:1ω-9	0.3 ± 0.1	0.8 ± 0.3	0.9 ± 0.6
16:1ω-7	2.2 ± 1.6	1.4 ± 0.1	7.1 ± 2.0
18:1ω-9	31.6 ± 5.3	13.5 ± 0.5	5.3 ± 3.1
18:1ω-7	0.7 ± 0.8	0.8 ± 0.2	1.9 ± 0.5
20:1	0.6 ± 0.2	0.8 ± 0.1	1.4 ± 0.6
ΣMUFA	35.4	17.3	16.6
18:2ω-6	12.2 ± 1.6	6.9 ± 0.9	2.8 ± 1.2
18:3ω-3	2.9 ± 1.3	8.4 ± 1.3	7.0 ± 1.7
18:4ω-3	3.2 ± 2.3	10.3 ± 1.5	2.8 ± 0.7
20:2ω-6	0.4 ± 0.1	0.5 ± 0.1	0.3 ± 0.1
20:3ω-6	0.1 ± 0.1	0.1 ± 0.0	0.2 ± 0.1
20:4ω-6	0.6 ± 0.7	0.4 ± 0.1	3.2 ± 0.5
20:3ω-3	0.3 ± 0.1	1.0 ± 0.2	0.4 ± 0.1
20:4ω-3	1.3 ± 0.4	2.8 ± 0.1	2.1 ± 0.3
20:5ω-3	5.0 ± 2.0	11.7 ± 1.4	9.9 ± 2.0
22:5ω-6	0.2 ± 0.2	1.0 ± 0.1	2.3 ± 0.9
22:5ω-3	0.2 ± 0.2	0.2 ± 0.1	1.9 ± 1.1
22:6ω-3	2.1 ± 1.5	5.6 ± 0.3	9.8 ± 3.2
ΣPUFA	28.5	48.9	42.7

817
818
819
820
821
822
823
824
825
826
827
828
829

830 a: Mean fatty acid composition (\pm standard deviation) (percentage of total fatty acids detected) of
 831 mesozooplankton from the Chowan River by group. SFA: saturated fatty acids, MUFA:
 832 monounsaturated fatty acids, and PUFA: polyunsaturated fatty acids.

	Mesozooplankton Groups			
	G (1)	H (5)	I (5)	J (5)
14:0	4.5	5.4 \pm 0.9	5.2 \pm 1.1	5.1 \pm 1.1
15:0	1.1	1.1 \pm 0.4	0.7 \pm 0.0	0.9 \pm 0.1
16:0	29.7	24.4 \pm 2.6	19.5 \pm 1.3	22.0 \pm 1.5
17:0	1.3	1.4 \pm 0.4	1.0 \pm 0.1	1.4 \pm 0.3
18:0	11.2	7.7 \pm 1.2	5.1 \pm 0.3	6.8 \pm 0.7
20:0	0.4	0.2 \pm 0.0	0.2 \pm 0.1	0.1 \pm 0.1
Σ SFA	48.2	40.2	31.7	36.3
16:1 ω -9	0.3	0.9 \pm 0.2	0.9 \pm 0.4	0.8 \pm 0.4
16:1 ω -7	4.1	9.5 \pm 2.2	6.2 \pm 2.2	8.6 \pm 2.2
18:1 ω -9	17.0	10.6 \pm 4.3	6.4 \pm 1.2	4.0 \pm 1.4
18:1 ω -7	2.1	3.9 \pm 1.7	2.4 \pm 0.4	3.0 \pm 1.1
20:1	0.3	0.1 \pm 0.1	0.3 \pm 0.1	0.2 \pm 0.1
Σ MUFA	23.8	25.0	16.2	16.6
18:2 ω -6	4.2	3.0 \pm 0.9	5.1 \pm 0.6	2.2 \pm 0.7
18:3 ω -3	2.9	6.1 \pm 2.1	10.7 \pm 1.5	5.1 \pm 1.9
18:4 ω -3	1.4	2.4 \pm 0.8	7.1 \pm 1.2	2.2 \pm 0.4
20:2 ω -6	0.2	0.2 \pm 0.1	0.4 \pm 0.1	0.2 \pm 0.0
20:3 ω -6	0.1	0.1 \pm 0.0	0.1 \pm 0.0	0.1 \pm 0.0
20:4 ω -6	3.7	4.2 \pm 1.5	2.1 \pm 0.6	2.5 \pm 1.2
20:3 ω -3	0.2	0.2 \pm 0.1	0.5 \pm 0.3	0.2 \pm 0.1
20:4 ω -3	0.8	0.6 \pm 0.3	1.2 \pm 0.6	0.8 \pm 0.3
20:5 ω -3	5.9	11.2 \pm 2.4	13.8 \pm 0.9	16.2 \pm 0.6
22:5 ω -6	0.4	0.4 \pm 0.2	0.8 \pm 0.2	1.8 \pm 0.8
22:5 ω -3	0.4	0.3 \pm 0.2	0.4 \pm 0.2	0.6 \pm 0.3
22:6 ω -3	7.3	5.7 \pm 2.9	10.1 \pm 2.8	15.1 \pm 7.3
Σ PUFA	27.5	34.4	52.3	47.0

833