

# Comparison of chemical compounds associated with sclerites from healthy and diseased sea fan corals (*Gorgonia ventalina*)

Carlos Toledo-Hernández <sup>Corresp., 1</sup>, Claudia P Ruiz-Díaz <sup>2</sup>, Liz Díaz <sup>3</sup>, Vanessa Santiago-Cárdenas <sup>4</sup>, Derick N Rosario-Berrios <sup>3</sup>, Derek M Garcia-Almedina <sup>3</sup>, Loretta M Roberson <sup>5</sup>

<sup>1</sup> Sociedad Ambiente Marino SAM, San Juan, Puerto Rico

<sup>2</sup> Department of Environmental Science, University of Puerto Rico, San Juan, Puerto Rico

<sup>3</sup> Department of Chemistry, University of Puerto Rico, San Juan, Puerto Rico

<sup>4</sup> Department of Biology, University of Puerto Rico, San Juan, Puerto Rico

<sup>5</sup> Marine Resources Program, Marine Biological Laboratory, Falmouth, MA, United States

Corresponding Author: Carlos Toledo-Hernández

Email address: cgth0918@gmail.com

**Background:** The roles of gorgonian sclerites as structural components and predator deterrents have been widely studied. Yet, their role as barriers against microbes has only recently been investigated, and even less is known about the diversity and roles of the chemical compounds associated with sclerites. **Methods:** Here, we examine the semi-volatile organic compound fraction (SVOCs) associated with sclerites from healthy and diseased *Gorgonia ventalina* sea fan corals to understand their possible role as a stress response or in defense of infection. We also measured the oxidative potential of compounds from diseased and healthy *G. ventalina* colonies. **Results:** The results showed that sclerites harbor a great diversity of SVOCs. Overall, 70 compounds were identified, the majority of which are novel with unknown biological roles. The majority of SVOCs identified exhibit multiple immune-related roles including antimicrobial and radical scavenging functions. The free radical activity assays further confirmed the anti-oxidative potential of some these compounds. The anti-oxidative activity was, nonetheless, similar across sclerites regardless of the health condition of the colony, although sclerites from diseased sea fans display slightly higher anti-oxidative activity than the healthy ones. **Discussion:** Sclerites harbor great SVOCs diversity, the majority of which are novel to sea fans or any other corals. Yet the scientific literature consulted showed that the roles of compounds found in sclerites vary from antioxidant to antimicrobial compounds. However, this study fell short in determine the origin of the SVOCs identified, undermining our capacity to determine the biological roles of the SVOCs on sclerites and sea fans.

1 **Comparison of chemical compounds associated with sclerites from healthy and diseased sea**  
 2 **fan corals (*Gorgonia Ventalina*).**

3 Carlos Toledo-Hernández<sup>1</sup>, Claudia P. Ruiz-Díaz<sup>1,2</sup>, Liz M. Díaz-Vázquez<sup>3</sup>, Vanessa Santiago<sup>4</sup>,  
 4 Derick N. Rosario-Berrios<sup>3</sup>, Derek M. García-Almedina<sup>3</sup>, Loretta M. Roberson<sup>5</sup>.

5 1 Sociedad Ambiente Marino (SAM), San Juan, Puerto Rico

6 2 Department of Environmental Science, San Juan, Puerto Rico

7 3 Department of Chemistry, University of Puerto Rico, San Juan, Puerto Rico

8 4 Department of Biology, University of Puerto Rico San Juan, Puerto Rico

9 5 Marine Biological Laboratory, Woods Hole, MA USA

10 Corresponding author: Carlos Toledo-Hernández

11 Sociedad Ambiente Marino (SAM), PO Box 22158, San Juan, Puerto Rico 00931

12 e-mail address: cgth0918@gmail.com

13 **ABSTRACT**

**Background:** The roles of gorgonian sclerites as structural components and predator deterrents have been widely studied. Yet, their role as barriers against microbes has only recently been investigated, and even less is known about the diversity and roles of the chemical compounds associated with sclerites.

**Methods:** Here, we examine the semi-volatile organic compound fraction (SVOCs) associated with sclerites from healthy and diseased *Gorgonia ventalina* sea fan corals to understand their possible role as a stress response or in defense of infection. We also measured the oxidative potential of compounds from diseased and healthy *G. ventalina* colonies.

**Results:** The results showed that sclerites harbor a great diversity of SVOCs. Overall, 70 compounds were identified, the majority of which are novel with unknown biological roles. The majority of SVOCs identified exhibit multiple immune-related roles including antimicrobial and radical scavenging functions. The free radical activity assays further confirmed the anti-oxidative potential of some these compounds. The anti-oxidative activity was, nonetheless, similar across sclerites regardless of the health condition of the colony, although sclerites from diseased sea fans display slightly higher anti-oxidative activity than the healthy ones.

**Discussion:** Sclerites harbor great SVOCs diversity, the majority of which are novel to sea fans or any other corals. Yet the scientific literature consulted showed that the roles of compounds found in sclerites vary from antioxidant to antimicrobial compounds. However, this study fell short in determine the origin of the SVOCs identified, undermining our capacity to determine the biological roles of the SVOCs on sclerites and sea fans.

## INTRODUCTION

Sea fans (*Gorgonia* sp.) are one of the top competitors for space within Caribbean coral reefs ecosystems. Similar to their relatives, the scleractinian corals, sea fans have suffered several

environmental disturbances such as aspergillosis disease outbreaks (a fungal infection caused by the saprophytic fungus *Aspergillus sydowii*) that seriously compromised the viability of many populations across the Caribbean during the 1990s (Smith et al., 1996; Nagelkerken et al., 1997; Geiser et al., 1998; Kim et al., 2004). Yet, in contrast to the scleractinian, sea fans have overcome these disturbances and continue to thrive. In fact, gorgonians are the prevailing corals in many reefs formerly dominated by scleractinian corals. Their relative success is provided in part, by their strong immune defenses, which have allowed gorgonians to maintain their internal homeostasis and health (Ruiz-Diaz et al., 2013; Sabat & Toledo-Hernández, 2015).

Gorgonians have a rather diverse repertoire of defensive mechanisms to cope and respond to disturbances, both abiotic and biotic (Toledo-Hernández & Ruiz-Diaz 2014; Ruiz-Diaz et al., 2016). For instance, to respond to pathogen infection, gorgonians are equipped with chemical pathways such as the melanin cascade that prevent or reduce the pathogen dissemination throughout the host tissue (Petes et al., 2003; Mydlarz et al., 2008). Concomitant to this response, gorgonians have the ability to activate the production of peroxidase enzymes, which induce the production of reactive oxygen species with cytotoxic roles and cellular signaling (Mydlarz & Harvell, 2007). Furthermore, gorgonians are also characterized for their diverse production of secondary metabolites, many of which are thought to be immune-related compounds due to their anti-predation and microbial properties (Harvell, Fenical & Green, 1988; Kim, 1994; Jensen et al., 1996; Smith et al., 1996; Geiser et al., 1998; Kim et al., 2000a, 2000b, Alker et al., 2004).

Gorgonians also have sclerites, which are minute skeletal elements mainly composed of calcium carbonate surrounding an organic matrix made up primarily of glycoproteins, spread throughout the gorgonians' epidermis, mesoglea, and axial skeleton (Kingsely & Watabe, 1982, 1984; Harvell & Suchanek, 1987, Van Alstyne & Paul, 1992). Sclerites are thought to have multiple defensive roles. For instance, sclerites provide structural support to gorgonians by reducing the elasticity and stiffness of the axial skeleton against water motion (Koehl, 1982; Lewis & Willis, 1991). In addition, sclerites play a role in deterring predation from gastropods and reef fishes (Harvell, Fenical & Greene, 1988; Van Alstyne & Paul, 1992; Koh et al., 2000). Furthermore, the purpling of sclerites (i.e., darkening of sclerites through the accumulation of pigments; Leverette et al., 2008) has also been linked to defensive mechanisms against several biological threats (Petes et al., 2003; Mydlarz & Harvell, 2007; Mydlarz et al., 2008).

Yet, the role of sclerites as chemical barriers against microbes has only recently been investigated. For instance, Toledo-Hernández et al., (2016) showed that sea fan sclerites treated with acetone and placed on top of the surfaces of culture media as to block the entrance of microbes, were less effective at delaying the crossing and further colonization of fungi in the agar than sclerites not treated with any solvent. However, no screening of chemical compounds in sclerites was conducted in that study. In fact, studies devoted to surveying the chemical compounds found in gorgonian sclerites are rather scarce, with only a few of them describing the chemical composition of the core of sclerites (Kingsley & Watabe, 1982, 1984; Kingsley et al., 1990). Here we perform for the first time, a screening of the semi volatile organic compounds (SVOCs) fraction found in sclerites from healthy and diseased *G. ventalina* colonies. Furthermore, we performed a scientific literature survey of the SVOCs found in sclerites to determine their potential biological roles. Finally, we conducted radical scavenging activity assays to examine the oxidative potential of chemical compounds associated with sclerites from healthy and diseased fans.

## METHODS

### Tissue collection and sclerite isolation

Sclerites used for this study were isolated from ten healthy and ten diseased *Gorgonia ventalina* colonies located at a depth of 1.5-2.0 m from El Escambrón beach, San Juan, Puerto Rico (18°28'00"N, 66°05'12"W). The tissue samples were collected under permit 2012-IC-086 issued to Claudia P. Ruiz Diaz, University of Puerto Rico (UPR) Rio Piedras campus, given by the Puerto Rico Department of Natural Resources, Commonwealth of Puerto Rico. Healthy colonies showed neither lesions nor tissue purpling, whereas diseased colonies showed at least one lesion usually overgrown by algae and surrounded by a ring of purple tissue. Previous studies have used the aforementioned pathology to diagnose algal tumor affliction, sea fan aspergillosis, as well as competitive interactions between inter and conspecific individuals (Morse et al., 1977; Smith et al., 1996; Alker et al., 2004). However, since we performed no microbiological or histological analyses to determine the precise etiology of the injured tissue collected for this study, we avoided the use of any term that could be associated with the afflictions previously described for sea fans. One tissue fragment approximately 4 cm<sup>2</sup> was cut from the edge of each healthy fan (hereafter HH). In diseased fans, 2 cm<sup>2</sup> samples were cut from both healthy tissue at the edge of the fan (healthy-diseased or HD) as well as from the diseased area (diseased-diseased or DD). Fragments were placed individually in 50 mL centrifuge tubes and brought to the laboratory on

ice. Once in the laboratory, each fragment was further cut into smaller fragments of 2-3 cm<sup>2</sup> in size and placed in 20.0 mL-sterilized leveled vials filled with 16.0 mL of distilled water. Then, to disassociate sclerites from the soft tissue matrix, vials were vortexed at room temperature, sonicated for 5 min, and then allowed to settle for approximately 4 hrs. This procedure was repeated over a period of 5-7 days to minimize impacts on the structure of the sclerites. After disassociated, sclerites were rinsed several times with distilled water, individually transferred to clean 55 mm Pyrex petri dishes, and dried at 35 °C for 72 hrs.

# **Gas chromatography-mass spectrometry analysis**

From each dried sclerite sample, 0.10 g were grounded in a clean porcelain mortar and the homogenized powder was transferred to a 1.5 mL micro-test tube with 1 mL of acetone. Each tube was vortexed and centrifuged at 6,000 rpm for 5 min. Then, 1.5 mL of the resulting extract from each sample was filtered and transferred without introducing air bubbles to clean amber vials (32 x12 mm). Afterward, samples were purged with nitrogen to remove dissolve oxygen and then septum caps were fitted to each vial and these were inverted to ensure no headspace was visible and finally, vials were hermetically sealed and not opened until analyzed to preserve the solution. Prior to injecting the solvent used for the extraction, reagents, glassware and other samples processing hardware were blank tested to ensure these materials will not add artifacts to the sclerite extracts analyses. Then, vials with sclerite extracts were placed in an Agilent 7890A gas chromatograph (GC) coupled to an Agilent 5975 C mass-selective detector for splitless injection. Because removing a sample aliquot from the vials may compromise the integrity of the sample, multiple aliquots were injected to allow for the screening and re-analysis of the samples. The injection temperature was 250 °C. Separation of samples was performed using a fused silica capillary column (SLB®-5 ms 30 m × 0.32 mm × 0.25 µm film thickness; Supelco, Bellefonte, PA). The initial column temperature was held at 70 °C for 1 min, and then increased by 8 °C min<sup>-1</sup> until reaching 270 °C, where it was held for 3 min, for a total run time of 29 min per sample. Ultrapure helium at constant-flow of 1.2 mL min<sup>-1</sup> was used as the carrier gas. Analysis of the generated chromatograms was performed using the AMDIS version 2.7/NIST version 2.0 computer program and the criteria for selection in the analysis for the screening compounds were: (1) R- match >800; (2) S/N > 15; and (3) peak areas >1000. Compound identification was confirmed through the use of NIST library appropriated standards and retention time index (<http://www.nist.gov/srd/nist1a.cfm>). To determine the potential biological roles of each of the

detected compounds, a scientific literature review using the IUPAC names of each compound was conducted.

### Free radical scavenging assay

Free radical scavenging activity of sclerite extracts was estimated using 1,1,-diphenyl-2-picrylhydrazyl (DPPH) (MacDonald-Wisk, Wood & Garg, 2006). Briefly, 0.10 g of homogenized powder not previously used from each of the sclerite samples was placed into labeled and clean 50 mL centrifuge tubes filled with 5.0 mL methanol. Then, each tube was vortexed for 10 sec, placed in a water bath at 40 °C for 40 min, and sonicated afterward for 30 min. The samples were then centrifuged for 10 min (6,000 rpm) at 25 °C. Afterwards, 1.0 mL of the supernatant (“sclerite extract”) was transferred to a clean 50 mL centrifugal tube containing 5.0 mL methanol. Then, 1 µL of the sclerite extract and 39 µL of DPPH (0.09 mM) were added to a 96-well microplate and the absorbance of the each sample was recorded at 0, 1, 4 and 5 minutes and then every 5 minutes for a one-hour period at 560 nm in a Thermo Scientific Multiscan FC spectrophotometer. Only those samples with absorbances with absorbance < 2 were used in the analysis. Scavenging activity in this study was expressed as IC<sub>50</sub>, which represents the concentration of extracts (mg/mL) needed to inhibit 50% of the DPPH radicals. To calculate IC<sub>50</sub> for each sclerite sample, the percent inhibition of radical production (%I) was estimated using the following equation (Mishra et al., 2012):

$$\%I = (A_i - A_f) \cdot 100 / A_i ,$$

where A<sub>i</sub> and A<sub>f</sub> represent the initial and final absorbance, respectively. A linear regression analysis was performed with extract concentration as the independent variable and %I as the dependent variable. Finally, the resulting linear regression equations with a correlation coefficient R<sup>2</sup> > 80% were used to estimate the IC<sub>50</sub>.

### Statistical analyses

Nonmetric Multi Dimensional Scaling (nMDS) along with an Analysis of Similarity (ANOSIM) with Bray-Curtis distances measurement were utilized to compare the composition of chemical compounds among the sclerites from different tissues. We used nMDS ordination, achieved by the *metaMDS* wrapper function from the package *vegan* in R version 3.2. The ordination was applied such that the data was scaled down to two dimensions.

For the analysis of free radical scavenging, a Kruskal-Wallis was performed to detect differences between  $IC_{50}$  index from sclerites from different tissues. In this analysis,  $IC_{50}$  index was assigned as the dependent variable and the origin of the sclerite was assigned as the independent variable. Finally, a  $\chi^2$  test was conducted to determine statistical differences in the relative abundances of alcohols, alkanes and alkenes and sclerite origin. In all analyses, P-value was fixed at 0.05. Statistical analyses were performed using R version 3.2 (R Core Team, 2014). For the statistical analyses, HD and DD samples from the same colony were considered independent and analyzed accordingly. In modular organisms such as gorgonian corals, modules are semi-autonomous units. Consequently, any particular stressors (abiotic or biotic) may trigger or have a differential consequence in tissue subjected to different health states (Van Valen, 1978; Tuomi & Vuorisalo, 1989).

## RESULTS

### Semi-volatile organic compounds

A total of 70 SVOCs were found in the sampled sclerites; 37% of these were identified from DD sclerites, while 35% and 29% were identified from DH and HH sclerites, respectively. Yet, the nMDS and ANOMIS analyses failed to group the chemical compounds according to their origin (e.g., HH, HD and DD sclerites). This holds true even when compounds found only once or twice across the samples were eliminated from the analysis.

In general, three major groups were identified: alcohols, alkanes and alkenes. When data was organized based on sclerite origin, alcohols and secondarily alkanes were the most frequently identified SVOCs in all samples, although the number of identified alkenes was higher in HD sclerites (Figure 1). However, the  $\chi^2$  test analysis revealed no statistical differences between alcohols, alkanes and alkenes and the sclerites origin. Furthermore, DD sclerites exhibited the highest number of exclusive compounds with a total of 17 unique SVOCs, followed by HH and HD sclerites with 15 and 8 unique SVOCs, respectively (Table S1; Figure 2). The remaining SVOCs were either shared across all sclerites or between two distinct combinations of sclerites (Figure 2).

Evidence from the scientific literature concerning the possible biological roles of the SVOCs identified in sclerites was obtained for 44% the compounds. Most were associated with immune related roles such as antimicrobials and antioxidants. Less represented roles were related to



cytotoxic and apoptotic compounds, cell membrane components such as steroids and oxidized steroids derivatives, and intermediate signaling molecules.

### Free radical scavenging assay

Thirteen out of the 30 sclerites sampled tested had  $R^2 > 80\%$  and thus were used for the free radical scavenging assays. Of these, six samples were HH sclerites, whereas four and three were DD and HD sclerite samples, respectively. The free radical scavenging  $IC_{50}$  ranged from 0.21 mg dry extract/mL to 0.01 mg dry extract/mL DPPH. Observed differences in antioxidant activity among sclerites were not statistically different, however on average there was a trend for higher values in HH and lowest in DD sclerites (Figure 3).

## DISCUSSION

Overall, 70 SVOCs associated with sclerites were identified. The majority of compounds have not been reported previously in corals and consequently little is known of their biological roles (Table S1). Several of these SVOCs were common to all sclerites samples, whereas others were found in two or exclusively in one of sclerite types (Table S1; Figure 2). Yet, no clusters between the compounds and health states were formed, suggesting that the chemical composition of sclerites is similar across the samples independent of the health state of the colony. Nonetheless, the concentration of these compounds may vary among colonies with contrasting health state, but our data do not provide sufficient details to validate or reject this argument.

On the other hand, the scientific literature surveyed revealed potential roles for some of the compounds. For instance, some of the compounds identified in this study have been characterized as free radical scavengers (Table S1). The oxidative analysis assays performed in this study, which revealed that extracts from the three sclerite origins exhibit antioxidant activity, further confirms this fact.

Antioxidant production in gorgonians is thought to have mitigating roles during several immune responses including phagocytosis, peroxidase enzymatic activity, and melanization (Olano & Bigger, 2000; Midlarz & Harvell, 2007). These immune responses produce reactive oxygen species (ROS) and other free radical molecules, inducing oxidative stress in the coral. Consequently, to counteract this oxidative stress, sea fans have developed regulatory systems such as increased production of non-enzymatic antioxidant scavengers, which neutralize the

218 potential harmful impact of ROS and other free radical molecules (Yost, Jones & Mitchelmore,  
219 2010; Shahbudin et al., 2011).

220 Antioxidant capacity in corals has also been linked to thermal stress. Several studies have  
221 reported that corals under thermal stress exhibit higher antioxidant potential than those under  
222 normal temperature conditions (Downs et al., 2002; Griffin & Bhagooli, 2004). In fact, over-  
223 expression of genes involved in the oxidative-stress response has been reported to increase in  
224 acroporids after they have undergone thermal stress (Csász , Senaca & Oppen, 2009).  
225 Alternatively, microbes associated with sea fans, both prokaryotes and eukaryotes, may be  
226 producing the antioxidant compounds found in sclerites. In the case of eukaryotes such as  
227 zooxanthellae for instance, antioxidants may be synthesized as a mitigating mechanism against  
228 ROS produced during photosynthesis (Foyer and Shigeoka, 2011).

229 Compounds known to have antimicrobial properties such as alcohols and esters were also  
230 identified in sclerites (Table S1). Many of these compounds have been isolated from terrestrial  
231 plants. Nonetheless, the fact that gorgonians produce chemical compounds with antimicrobial  
232 activities is not surprising. Multiple culture-based studies using extracts from several gorgonians  
233 species have revealed the antimicrobial potential of these extracts (Kim1994, 2000; Jensen et al.,  
234 1996; Alker, Smith & Kim, 2001; Alker et al., 2004; Ward et al., 2007). Nonetheless, in this  
235 study, no relationship could be established between the origin of compounds (i.e., diseased or  
236 healthy corals) and their microbial roles, as the majority of the potential roles of the compounds  
237 could not be obtained from the scientific literature. Many of these compounds could be  
238 precursors or intermediaries of potentially biologically relevant molecules that were trapped  
239 within the sclerites during their formation. If indeed some of the compounds isolated from  
240 sclerites are precursors of biologically active molecules, as well as environmental cues or internal  
241 stressors, sclerites could be used as archives of metabolic pathways or past climatic and biotic  
242 stress events.

## 243 CONCLUSIONS

244 Sclerites harbor a great SVOC diversity, the majority of which are novel to sea fans or any other  
245 corals. However, a major drawback of the study is our inability to determine the origins of these  
246 SVOCs. For instance, it is uncertain which SVOCs, if any, were synthesized by microbes  
247 associated with sea fans. On the other hand, certain compounds may be derivatives from other

compounds. For instance, 1-undecene, 9 methyl, could be a derivative of other compounds such as alcohols which have undergone esterification reactions to form esters that are important fractions of lipids and steroids of living cells. In either case, not knowing the origin of the SVOCs hinders our capacity to understand the underlying mechanism of their biological function. However, as research progresses and our understanding of sclerite formation is improved, coupled with the identification of other chemical compounds using different extraction solvents (polar and non-polar), our understanding of the roles the chemical compounds from sclerites and the corals in general will improve. Yet, the present study represents a necessary first step in that process.

## Acknowledgements

We like to express our gratitude to Alberto M. Sabat for his friendly review.

## REFERENCES


- Alker P, Smith GW, Kim K. 2001. Characterization of *Aspergillus sydowii* (Thom et Church), a fungal pathogen of Caribbean sea fan corals. *Hydrobiology* 460:105-111.
- Alker AP, Kiho K, Dube DH, Harvell CD. 2004. Localized induction of a generalized response against multiple biotic agents in Caribbean sea fans. *Coral Reefs* 23:397-405.
- Császár NBM, Seneca FO, van Oppen MJH. 2009. Variation in antioxidant gene expression in the scleractinian coral *Acropora millepora* under laboratory thermal stress *Marine Ecology Progress Series* 392:93-102.
- Downs, CA, Fauth JE, Halas JC, Dustan P, Bemiss J, Woodley CM. 2002. Oxidative stress and seasonal coral bleaching. *Free Radical Biology and Medicine* 33:533-543.
- Foyer CH, Shigeoka S. 2011. Understanding oxidative stress and antioxidant function to enhance photosynthesis. *Plant Physiology* 155:93-100.
- Geiser DM, Taylor JW, Ritchie KB, Smith GW. 1998. Cause of sea fan death in the West Indies. *Nature* 394:137-138.
- Griffin PS, Bhagooli R. 2004. Measuring antioxidant potential in corals using the FRAP assay.

- 274 *Journal of Experimental Marine Biology and Ecology* 302:201-211.
- 275 Harvell CD, Suchanek TH. 1987. Partial predation on tropical gorgonians by *Cyphoma gibbosum*
- 276 (Gastropoda). *Marine Ecology Progress Series* 38:37- 44.
- 277 Harvell CD, Fenical W, Greene CH. 1988. Chemical and structural defenses of Caribbean
- 278 gorgonians (*Pseudopterogorgia* spp.). I. Development of an in situ feeding assay. *Marine*
- 279 *Ecology Progress Series* 49:287-294.
- 280 Jensen PR, Harvell CD, Wirtz K, Fenical W. 1996. Antimicrobial activity of extracts of
- 281 Caribbean gorgonian corals. *Marine Biology* 125:411-419.
- 282 Kim K. 1994. Antimicrobial activity in gorgonian corals (Coelenterata, Octocorallia). *Coral*
- 283 *Reef*. 13(2):75-80.
- 284 Kim K, Harvell CD, Smith GW, Merkel SM. 2000a. Fungal disease resistance of Caribbean sea
- 285 fan corals (*Gorgonia* spp.). *Marine Biology* 136:259-267.
- 286 Kim K, Kim PD, Alker AP, Harvell CD. 2000b. Chemical resistance of gorgonian corals against
- 287 fungal infections. *Marine Biology* 137:393-401.
- 288 Kim K and Harvell CD. 2004. The rise and fall of a six-year coral fungal epizootic. *American*
- 289 *Naturalist* 164:S52-S63.
- 290 Kingsley R, Watabe N. 1982. Ultrastructure invitation of spicule formation in the gorgonian
- 291 *Leptogorgia virgulata* (Lamarck) (Coelenterata: Gorgonacea). *Cell and Tissues Research* 223:
- 292 325-334.
- 293 Kingsley R, Watabe N. 1984. Synthesis and transport of the organic matrix of the spicules in the
- 294 gorgonian *Leptogorgia virgulata* (Coelenterata: Gorgonacea). An autoradiographic investigation.
- 295 *Cell and Tissues Research* 235:533-538.
- 296 Kingsley RJ, Tsuzak m, Watabe N, Gerald I, Mechanic GL. 1990. Collagen in the spicule organic
- 297 matrix of the gorgonian *Leptogorgia virgulata*. *Biological Bulletin* 179:207-213.


- 298 Koehl MAR. 1982. Mechanical design of spicule-reinforced connective tissue: stiffness. *Journal*  
299 *of Experimental Biology*. 98:239-267.
- 300 Koh LL, Goh NKC, Chou LM, Tan YW. 2000. Chemical and physical defenses of Singapore  
301 gorgonians (Octocorallia: Gorgonacea). *Journal of Experimental Marine Biology and Ecology*.  
302 251:103-115.
- 303 Lewis, JC, Wallis, EV. 1991. The function of surface sclerites in gorgonians (Coelenterata,  
304 Octocorallia). *Biological Bulletin*.18:275–288.
- 305 Leverette CL, Warren M, Smith MA, Smith GW. 2008. Determination of carotenoid as the purple  
306 pigment in *Gorgonia ventalina* sclerites using Raman microscopy. *Spectrochimica Acta Part A*.  
307 69:1058-1061.
- 308 MacDonald LK, Wood LG, Garg M. 2006. Methodology for the determination of biological  
309 antioxidant capacity in vitro: a review. *Science of Food and Agriculture*. 86:2046-2056.
- 310 Mishra K, Ojha H, Nabo Kumar Chaudhury NK. 2012. Estimation of antiradical properties of  
311 antioxidants using DPPH assay: A critical review and results. *Food Chemistry*. 130:1036-1043  
312 DOI10.1016/j.foodchem.2011.07.127
- 313 Morse DE, Morse ANC, Duncan H. 1997. Algal tumor in the Caribbean sea fan, *Gorgonia*  
314 *ventalina*. Proceeding of the 3th Third International Coral Reef Symposium, Miami Florida, 623-  
315 629.
- 316 Mydlarz LD and Harvell CD. 2007. Peroxidase activity and inducibility in the sea fan coral  
317 exposed to a fungal pathogen. *Comparative Biochemistry Physiology, Part A*. 146:54-62.
- 318 Mydlarz LD, Holthouse SF, Pters EC, Haverl CD. 2008. Cellular responses in sea fan corals:  
319 Granular amoebocytes react to pathogen and climate stressors. *PloS ONE*.  
320 10.1371/journal.pone.0001811.
- 321 Nagelkerken IK, Smith GW, Bonair K, Bush P, Garzón-Ferriera J, Botero L, Gayle P, Harvell  
322 CD, Heberer C, Kim K, Petrovic C, Pors L, Yoshioka P. 1997. Widespread disease in Caribbean  
323 sea fans: II. Patterns of infection and tissue loss. *Marine Ecology Progress Series* 160:255-263.
- 324 Olano CT and Bigger CH. 2000. Phagocytic activities of the gorgonian coral *Swiftia exserta*.

- 325 *Journal of Invertebrate Pathology* 76:176-84.
- 326 Petes LE, Harvell CD, Peters EC, Webb MAH, Mullen KM. 2003. Pathogens compromise  
327 reproduction and induce melanization in Caribbean sea fans. *Marine Ecology Progress Series*  
328 264:167-171.
- 329 Sabat AM and Toledo-Hernández C. 2015. Viability of sea fan population impacted by disease:  
330 Recruitment versus Incidence. *Journal of Marine Biology* [10.1155/2015/987060](https://doi.org/10.1155/2015/987060).
- 331 Shahbudin S, Deny S, Zakirun AMT, Haziyamin TAH, Akbar John B, Taher M. 2011.  
332 Antioxidant properties of soft coral *Dendronephthya* sp. *International Journal of Pharmacology*  
333 7:263-267.
- 334 Smith GW, Ives, LD, Nagelkerken IA, Ritchie KB. 1996. Caribbean sea-fan mortalities.  
335 *Nature*.383:487.
- 336 R Core Team. 2014. R: A language and environment for statistical computing. R Foundation for  
337 statistical Computing, Vienna, Austria. URL <http://www.R-project.org/>.
- 338 Ruiz-Diaz CP, Toledo-Hernández C, Sabat AM, Marcano M. 2013. Immune response to a  
339 pathogen in corals. *Journal of Theoretical Biology* 332:141-148.
- 340 Ruiz-Diaz CP, Toledo-Hernández C, Mercado-Molina A, Sabat AM. 2016. Scraping and  
341 extirpating: two strategies to induce recovery of diseased *Gorgonia ventalina* sea fans. *Marine*  
342 *Ecology* 37: 336-343.
- 343 Tuomi J and Vuorisalo T. 1989. What are the units of selection in modular organisms? *Oikos*  
344 54:227-233.
- 345 Toledo-Hernández C and Ruiz-Díaz CP. 2014. The immune response of corals. *Invertebrate*  
346 *Survival Journal* 11:319-328.
- 347 Toledo-Hernández C, Ruiz-Díaz CP, Paniagua R, Roberson L, Ramírez-Camejo L, Sabat AM,  
348 Bayman P. 2016. The role of sclerites in the defense against pathogens of the sea fan *Gorgonia*  
349 *ventalina* (Octocorallia). *Journal of Experimental Marine Biology and Ecology* 483:20-24.
- 350 Ward JR, Rypien KL, Bruno JF, Harvell, CD, Jordán-Dahlgre E, Mullen KM, Rodríguez-

- 351 Martínez RE, Sánchez J, Smith G. 2007. Coral diversity and disease in Mexico. *Diseases of*  
 352 *Aquatic Organisms* 69:23-31.
- 353 Yost DM, Jones RJ, Mitchelmore CL. 2010. Alterations in dimethylsulfoniopropionate (DMSP)  
 354 levels in the coral *Montastraea franksi* in response to copper exposure. *Aquatic Toxicology*  
 355 98:367-373.
- 356 Van Alstyne KL, Paul VJ. 1992. Chemical and structural defenses in the sea fan *Gorgonia*  
 357 *ventalina*: effects against generalist and specialist predators. *Coral Reefs* 11:155-159.
- 358 Van Valen L. 1978. Arborescent animals and other colonids. *Nature* 276:318.




359 **Figure 1.** Percentages of the main functional chemicals in sclerites from healthy sea fans (HH),  
 360 from healthy tissue from diseased sea fans (DH), and diseased tissue from diseased sea fans  
 361 (DD).



362 **Figure 2.** Number of volatile organic compounds (VOCs) in sclerites from healthy sea fans (HH),  
 363 healthy tissue from diseased sea fans (HD), and diseased tissue from diseased sea fans (DD).





**Figure 3.** Average of free radical scavenging analysis expressed as IC<sub>50</sub> of sclerites from healthy sea fans (HH), healthy tissue from diseased fans (HD) and diseased tissue from diseased fans (DD). Bars denote standard errors.