

Light enhanced calcification in *Stylophora pistillata*: effects of glucose, glycerol and oxygen

Zooxanthellate corals have long been known to calcify faster in the light than in the dark, however the mechanism underlying this process has been uncertain. Here we tested the effects of oxygen under controlled pCO₂ conditions and fixed carbon sources on calcification in zooxanthellate and bleached microcolonies of the branching coral *Stylophora pistillata*. In zooxanthellate microcolonies, oxygen increased dark calcification rates to levels comparable to those measured in the light. However in bleached microcolonies oxygen alone did not enhance calcification, but when combined with a fixed carbon source (glucose or glycerol), calcification increased. Respiration rates increased in response to oxygen with greater increases when oxygen is combined with fixed carbon. ATP content was largely unaffected by treatments, with the exception of glycerol which decreased ATP levels.

1 Light enhanced calcification in *Stylophora pistillata*: effects of glucose, glycerol and oxygen

2

3 Michael Holcomb*, Eric Tambutté, Denis Allemand, Sylvie Tambutté

4 Centre Scientifique de Monaco

5 8 Quai Antoine 1er

6 MC-98000 Monaco

7

8 *Corresponding author current address:

9 Michael Holcomb

10 ARC Centre of Excellence in Coral Reef Studies

11 School of Earth and Environment & Oceans Institute

12 The University of Western Australia

13 35 Stirling Highway M004

14 Crawley, WA 6009 Australia

15 email: mholcomb3051@gmail.com

16 Abstract

17 Zooxanthellate corals have long been known to calcify faster in the light than in the dark,
 18 however the mechanism underlying this process has been uncertain. Here we tested the effects of
 19 oxygen under controlled pCO₂ conditions and fixed carbon sources on calcification in
 20 zooxanthellate and bleached microcolonies of the branching coral *Stylophora pistillata*. In
 21 zooxanthellate microcolonies, oxygen increased dark calcification rates to levels comparable to
 22 those measured in the light. However in bleached microcolonies oxygen alone did not enhance
 23 calcification, but when combined with a fixed carbon source (glucose or glycerol), calcification
 24 increased. Respiration rates increased in response to oxygen with greater increases when oxygen
 25 is combined with fixed carbon. ATP content was largely unaffected by treatments, with the
 26 exception of glycerol which decreased ATP levels.

27 Introduction

28 Rising ocean temperatures and declining pH have received considerable attention in recent years
 29 (e.g. Hoegh-Guldberg et al., 2007; McCulloch et al., 2012). Among the major concerns are the
 30 impacts on coral reefs, particularly on corals themselves. As temperatures rise, bleaching events
 31 (the loss of symbiotic zooxanthellae) become more frequent and severe, reducing the ability of
 32 the coral to cope with other stressors, leading to the loss of coral reefs (e.g. McWilliams et al.,
 33 2005). Rising CO₂ levels further compound the problem by reducing aragonite saturation state
 34 and potentially calcification (e.g. Langdon and Atkinson 2005; Holcomb et al., 2010).

35 Central to our understanding of the response of corals to changing environmental conditions is an
 36 understanding of the role the symbiotic zooxanthellae play in coral calcification. It has long been
 37 established that calcification rates increase in zooxanthellate corals during periods in which
 38 photosynthesis is occurring (Yonge, 1931), a process known as light-enhanced calcification (e.g.
 39 Vandermeulen et al., 1972; Gattuso et al., 1999). Comparisons of facultatively symbiotic corals
 40 with and without zooxanthellae similarly show increased calcification in the presence of
 41 symbionts (Jacques et al., 1983). Yet, comparisons of zooxanthellate corals to azooxanthellate
 42 corals show that some azooxanthellate corals can grow just as rapidly (Marshall, 1996;
 43 Mortensen and Rapp, 1998), bleached corals can continue calcifying for some time at rates
 44 similar to those of unbleached corals (Rodrigues and Grottoli, 2006), in corals such as acroporids,
 45 the fastest growing regions are largely azooxanthellate (Pearse and Muscatine 1971), and light

does not necessarily stimulate calcification in zooxanthellate corals (e.g. Rinkevich and Loya, 1984). Thus, it is not clear that symbionts per se enhance calcification.

There are several mechanisms which could be responsible for the increase in calcification generally associated with symbiosis (for reviews see: Gattuso et al., 1999; Allemand et al., 2011; Tambutté et al., 2011). Calcification could increase due to the drawdown in CO₂ as a direct result of photosynthesis, leading to increased pH, shifting the dominate CO₂ species from bicarbonate to carbonate and increasing the saturation state (Kawaguti and Sakumoto, 1948). Other likely mechanisms by which calcification could be increased include the use of fixed carbon from photosynthesis as fuel to increase ion pumping and elevate saturation state (Goreau, 1959), or the use of fixed carbon for organic matrix synthesis, which may facilitate aragonite precipitation (Young et al., 1971). Oxygen produced by the zooxanthellae may relieve internal hypoxia and thus enhance calcification (Rinkevich and Loya, 1984; Rands et al., 1992, Nakamura et al., 2013). Or symbionts may remove compounds which inhibit calcification or metabolism (e.g. Goreau, 1961; Vandermuelen and Muscatine, 1974).

Though numerous studies have examined the production of fixed carbon by zooxanthellae and suggested that various forms of fixed carbon (glycerol (Muscatine, 1967), glucose (Burriesci et al., 2012), among other compounds (e.g. Whitehead and Douglas, 2003)) are translocated to the coral host, the role of fixed carbon specifically in calcification has received little attention. Various carbon compounds are known to be taken up by corals. Stephens (1960, 1962), and Lewis and Smith (1971) demonstrated that corals have the ability to take up fixed carbon compounds supplied exogenously. Stephens (1962) suggested that a glucose concentration of ~80 µM is sufficient for glucose uptake to balance respiration in *Fungia*. Oku et al. (2003a,b) claimed corals have the ability to incorporate a range of exogenous sugars into their tissues, and that supplying sugars exogenously can improve the survival of bleached corals. Thus, exogenous fixed carbon may substitute for that provided by zooxanthellae.

The effect of fixed carbon supplementation on calcification has been examined in five studies. Taylor (1977) showed that fixed carbon compounds supplied exogenously are transported to the regions of most rapid growth. Vandermeulen and Muscatine (1974) and Chalker (1975) exposed corals in the light and in the dark to glucose, glycerol and alanine. Calcification rates were either unchanged or declined as a result of exposure to fixed carbon sources at concentrations from 10 µM – 10mM. A similar result was found by Al-Horani et al. (2007) for both glucose and feeding

with *Artemia*. In contrast, Colombo-Pallotta et al. (2010) found that in bleached corals, glycerol could stimulate calcification, though not in zooxanthellate corals. Such results suggest that fixed carbon by itself is not generally limiting the calcification process.

In addition to fixed carbon, zooxanthellae also provide oxygen to the coral host. Several studies have documented the daily cycle in dissolved oxygen levels adjacent to the coral tissue, with hyperoxia persisting for much of the day, and hypoxia persisting for much of the night (e.g. Rands et al., 1992; Shashar et al., 1993; Marshall and Clode, 2003; Agostini et al., 2009). Respiration and calcification rates are linked to this cycle in oxygen saturation, with lower rates at reduced oxygen levels (Rinkevich and Loya, 1984; Shick, 1990; Al-Horani et al., 2007) and increased calcification rates under elevated oxygen (Colombo-Pallotta et al., 2010; Wijgerde et al., 2012, Nakamura et al., 2013). In both the studies of Vandermeulen and Muscatine (1974) and Chalker (1975), corals were incubated in unstirred, un-aerated containers, thus any potential benefit from the supplied carbon sources may have been masked by oxygen limitation. Under normal conditions, photosynthate is supplied and oxygen levels are elevated by photosynthesis, thus it may be the combination of oxygen and fixed carbon that allow the coral to respire and calcify faster.

A lack of, or insufficient, flow may be an important factor limiting the potential benefits of fixed carbon in early experiments. Corals, like many benthic organisms, depend upon water movement for their supply of nutrients, food, and waste removal. Water flow plays a major role in determining the diffusive boundary layers surrounding corals, affecting transport rates of O₂, CO₂, nutrients, etc. likely affecting all aspects of coral physiology (Dennison and Barnes 1988; Atkinson and Bilger 1992; Kuhl et al., 1995; Nakamura and van Woesik 2001). Respiration in particular may be limited by the diffusion of oxygen under stagnant or low-flow conditions (Shick 1990; Kuhl et al., 1995), which would in-turn limit energy availability to other processes, such as calcification. Thus adequate flow is likely critical if any benefit is to be seen to supplementation with fixed carbon.

Given that respiration and calcification rates respond to elevated oxygen levels, dark calcification may indeed be energy limited due to insufficient oxygen to maintain oxidative phosphorylation. Fixed carbon availability may also limit calcification if sufficient oxygen is available. Thus, elevated oxygen levels in combination with fixed carbon supplementation may be sufficient to increase dark calcification rates to levels comparable to light calcification rates. The primary

energy carrier used to convert energy derived from respiration into ion pumping and other energy requiring activities is adenosine triphosphate (ATP), thus energy limitation may be reflected in cellular ATP content. Here combinations of different fixed carbon sources and concentrations and elevated oxygen levels were tested on zooxanthellate and bleached microcolonies of the branching coral *Stylophora pistillata* to determine whether oxygen limitation or a combination of oxygen and fixed carbon limitation can explain light enhanced calcification. In addition to calcification, respiration and ATP were measured to provide insight into the potential underlying processes.

Materials & Methods

Experimental organisms

Colonies of the branching coral *Stylophora pistillata* were maintained in the laboratories of the Centre Scientifique de Monaco in flow-through aquaria receiving Mediterranean seawater (S=38) heated to 25 °C. Experimental microcolonies were prepared as described by Al-Moghrabi et al (1993) and Ferrier-Pagès et al (2003), composed of ~2 cm x ~1 cm terminal portions of branches on nylon monofilament line. Microcolonies were used after at least three weeks of recovery, allowing tissue to fully cover the cut skeleton. Lighting was provided by JBL Marin Day T5 bulbs providing an irradiance of ~195 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ at the position of the microcolonies with a 12 h light/dark cycle. To prepare bleached microcolonies, microcolonies were maintained in the dark for a minimum of six weeks prior to use, at which point microcolonies appeared pure white. Microcolonies were fed twice a week with newly hatched *Artemia sp.* A subset of microcolonies were weighed periodically to estimate growth rates (buoyant weight technique, Jokiel et al. 1978; Davies 1989).

Experimental incubations

133 Exposures to treatment conditions, calcification, and respiration rate measurements were carried
 134 out in 250 ml Erlenmeyer flasks (made of borosilicate glass). Flasks were sealed using
 135 transparent acrylic stoppers. A small hole in the center of the acrylic stopper allowed a
 136 microcolony to be suspended in the center of the flask; a rubber stopper was used to seal the hole
 137 once the microcolony had been placed level with the 200 ml mark (Supplemental Figure S1).
 138 Preliminary experiments showed oxygen influx into nitrogen sparged seawater and oxygen efflux
 139 from oxygen sparged seawater were negligible with this sealing system. Flasks were placed on a
 140 multiposition magnetic stir plate and stirred at 145 rpm with teflon coated stir bars. Different stir
 141 bar sizes (7.8 x 39.2 mm, 5.9 x 15 mm, and 4.5 x 11.95 mm – values are average measured
 142 diameter and length, the manufacturer's stated size was generally slightly larger) were tested
 143 initially (Fig. 1), different stirring speeds were tested as well (data not shown, see supplemental
 144 material). Flasks were held within a constant temperature bath set at 25°C. Incubations were
 145 carried out either in the dark or under $\sim 195 \mu\text{mol photons m}^{-2} \text{s}^{-1}$ provided by JBL Marin Day T5
 146 bulbs. Microcolonies were starved for at least two days prior to being used in experiments to
 147 reduce nutrient efflux. All dark incubations for zooxanthellate colonies were started in the
 148 morning such that microcolonies had been pre-conditioned to dark conditions for 12 hr prior to
 149 the start of experiments. Light incubations were carried out during normally lit hours. Corals
 150 were maintained under treatment conditions for at least 20 min prior to starting the experiment to
 151 allow the coral to acclimate to treatment conditions. All seawater used for incubations was
 152 passed through two 0.45 μm membrane filters prior to use. The seawater mass present in each
 153 incubation was estimated based on the mass of seawater held by each incubation chamber
 154 without a coral and the displacement volume of the coral skeleton calculated from skeletal dry
 155 weight and assuming a density of 2.9 g/cm^3 . For each experimental run, in addition to treatment
 156 incubations, seawater only and treatment seawater incubations were carried out to check for

background changes and corals not exposed to treatment conditions were included as well to check for day-to-day variations in growth/metabolic rates. Following incubations, water from a subset of flasks which had previously contained corals was pooled and used for an additional incubation (without coral) to assess the potential for microbial growth in the treatment seawater to impact the measured values. Dark incubations of corals generally resulted in a ~14% decline in oxygen and a ~4% decline in alkalinity by the end of the incubation. In some instances (for bleached microcolonies), the change in alkalinity was below the resolution of the measurement method, calcification data for such incubations were discarded.

Water chemistry

Oxygen

Experimental oxygen and CO₂ levels were set by bubbling seawater held at 25 °C with humidified air or mixed gas, the composition of the mixed gas was set using mass flow controllers to mix ambient air, oxygen, and CO₂ to achieve ~2x ambient pO₂ at ambient pCO₂. CO₂ levels were verified using a Licor 6262 CO₂ analyzer. Oxygen (% saturation) was measured using a fiber optic oxygen measurement system (PreSens) with Pst3 sensors mounted on the inside of each flask. Sensors were calibrated with air (100%) and N₂ (0%) sparged seawater, a linear response was assumed. Atmospheric pressure was recorded at the time each experiment was performed. Oxygen concentrations were calculated as follows. The oxygen concentration for air saturated (100%) seawater at the experimental salinity and pressure was calculated using data from Benson and Krause (1984) as fit by equation 8 of Garcia and Gordon (1992), the first A₃ term was ignored. Concentrations calculated in this manner are for 1 atm total pressure, values were corrected for the measured atmospheric pressure and used to convert percent saturation measurements to concentrations.

Carbonate chemistry

Alkalinity samples were collected at the start and end of each experiment in pre-cleaned 20 ml scintillation vials and stored at 4 °C till measured. Semi-automated titrations with 0.03 N HCl (containing 41g/L NaCl) were carried out on ~4 g replicated samples using a Metrohm Titrando 808 dosimat. Certified reference material (supplied by the laboratory of Andrew Dickson, Scripps Institute of Oceanography) and an internal seawater standard were run each time samples were measured, alkalinity was calculated via regression per Holcomb et al. (2012). Samples for pH measurement were collected in 20 ml syringes, capped, and kept in a 25 °C water bath until measured. An initial sample was injected into a 10 cm fiber optic flow cell for collecting the reference spectrum, then ~1 µl of a 1.47 mM meta Cresol Purple (Acros 199250050 lot: A0264321) solution was added per ml seawater in the syringe using a micrometer syringe (Gilmont) to inject and mix the mCP. The solution was then injected into the flow cell and the pH measured, calculations followed DOE (1994). Alkalinity, ammonia, and pH were then used to estimate calcification assuming calcification alters alkalinity by 2 moles per mole CaCO₃ produced, ammonia was assumed to increase alkalinity by an amount equal to its concentration. Carbonate chemistry parameters were calculated using CO2Sys (van Heuven et al., 2009), additional species contributing to alkalinity were included as appropriate.

Fixed carbon

All incubations with fixed carbon substrates included antibiotics (35 IU penicillin, 0.035 mg streptomycin per ml) and antibiotics were included in the associated control incubations. The effects of this antibiotic concentration on calcification and respiration were tested as well. Glucose or glycerol was added as a concentrated stock solution. Target concentrations were ~1

% (120 mM) glycerol, 1 mM glycerol, 1 mM glucose, and 0.02 mM glucose. Concentrations were chosen based on the works of Whitehead and Douglas (2003) and Colombo-Pallotta et al. (2010). Concentrations of each respiratory substrate were verified at the start and end of each incubation using one of the following kits, protocols were modified slightly to improve results in seawater matrices: d-glucose kit (Megazyme) for glucose, glycerol GK kit (Megazyme) for glycerol. Measured concentrations were: 110 - 130 mM and 1 - 1.4 mM for glycerol, 0.9-1 mM and 0.019 - 0.021 mM for glucose. Concentrations of all fixed carbon compounds changed by less than 10% during the incubations, except for the 0.02 mM glucose treatment for which concentration declined by ~40% by the end of the incubation, which is consistent with estimates of the uptake of fixed carbon substrates (Stephens 1960; Whitehead and Douglas 2003).

Nutrients

Ammonia and phosphate concentrations were measured at the start and end of a subset of incubations using a Proxima autoanalyzer (Alliance Instruments) following the manufactures recommended protocols. No nutrient release was detected for zooxanthellate microcolonies, only for bleached microcolonies were nutrient releases measureable and thus alkalinity data corrected for changes in nutrient concentrations.

ATP

ATP was extracted with sulfuric acid following a protocol modified from Fang et al. (1987; 1989). Microcolonies were placed in 0.6 N sulfuric acid (kept on water/ice prior to use) and placed immediately in a sonicator (Branson 200 Ultrasonic Cleaner) filled with water/ice. Samples were sonicated for 15 min, tubes with microcolonies re-weighed, and a sample of the extract transferred to a 1.5 ml polypropylene tube (Eppendorf) and stored at -80 °C until measured. See supplementary materials for a comparison of this method with other techniques.

231

232 ATP measurements were made using a luciferin/luciferase based ATP detection kit (Roche ATP
233 Bioluminescence Assay Kit HS II) with light emission measured using either a Lumat 9507 or
234 9508 luminometer (Berthold), and concentrations calculated based on standard additions. A
235 detailed protocol is included in the supplemental materials.

236

237 Normalizations:

238 Protein was extracted from the tissue following ATP extraction by adding NaOH to produce a ~1
239 N NaOH solution which was held at 90 °C for ~20 min. Protein was measured using a Pierce
240 BCA protein assay kit; protein concentrations were calculated using standard addition (using
241 BSA as the standard).

242 Surface area was determined via the wax method of Stimson and Kinzie (1991). Normalization
243 data are summarized in the supplementary materials (Table S2).

244

245 Statistics

246 To calculate total protein and ATP per coral it was assumed that the internal water volume of the
247 microcolony equilibrated with the surrounding solution, thus the extract volume consisted of the
248 volume of the extraction solution added plus the volume of seawater added with the microcolony
249 (approximated by the change in weight of the tube upon addition of the coral minus the dry
250 weight of the coral).

251 ATP extracted from corals was normalized to surface area, protein, and skeletal dry weight and
252 compared using a one-way ANOVA. Measurements of calcification and respiration under
253 treatment conditions were normalized to measurements on the same coral under ambient (dark)
254 conditions, to account for day to day variations in rates, all rates were normalized to untreated
255 controls measured at the same time (per Holcomb et al., 2012), data were subsequently compared

using ANOVA. When the overall ANOVA model showed a significant treatment effect, treatments significantly different from each other were detected using Dunnett's multiple comparison to compare treatments to controls. Residuals were plotted against predicted values and on normal probability plots to assess violations of ANOVA assumptions. The SAS software package was used for all statistical tests.

Results

Effects of stirring on physiological parameters

Several stirring rates and stir bar sizes were tested to establish appropriate flow rates for the experimental incubations (Fig. 1). Significant differences were detected in dark respiration rates among stirring treatments ($F_{2,13}=18.68$, $p=0.0002$), with smaller stir bars being associated with lower respiration rates. No significant differences in calcification were detected, though there was an apparent trend toward higher dark calcification rates with larger stir bar sizes. A stirring speed of 145 rpm with a 39.2 x 7.8 mm (manufacturers stated size: 40 x 8 mm) stir bar was chosen for the experiments since it allowed maximal respiration and calcification rates while maintaining normal polyp expansion (visually assessed, not quantified).

Effect of treatments on calcification

Dark calcification was significantly affected by treatment conditions ($F_{7,125} = 10.69$, $p<0.0001$ for zooxanthellate and $F_{6,38} = 3.99$, $p=0.0034$ for bleached microcolonies, Fig. 2A, C). Exposure to light or oxygen significantly ($p<0.05$) increased calcification of zooxanthellate microcolonies relative to dark controls while 1% glycerol with oxygen led to a significant decline in dark calcification (Fig. 2A). Treatment with glucose and oxygen or lower concentrations of glycerol with oxygen was not associated with a significant change in dark calcification for zooxanthellate microcolonies. In contrast, for bleached microcolonies light or oxygen alone had no effect but

treatment with oxygen combined with fixed carbon increased dark calcification, with a significant increase ($p=0.0035$) for the 1% glycerol plus oxygen treatment (Fig. 2C).

Effects of treatments on respiration

Respiration rates were significantly affected by treatment conditions ($F_{6,65} = 26.96$, $p<0.0001$ for zooxanthellate and $F_{6,43} = 23.48$, $p<0.0001$ for bleached microcolonies, Fig. 2B, D). All treatments (except antibiotics) were associated with higher oxygen consumption in the dark ($p<0.05$) for zooxanthellate microcolonies with the largest increases for corals treated with oxygen and fixed carbon (Fig. 2B) (incubation in the light led to oxygen production, data were not compared with dark incubations). Similarly in bleached microcolonies treatment with fixed carbon combined with oxygen had the largest effect ($p<0.001$) on respiration (Fig. 2D) (light had no effect on respiration in bleached colonies).

Effect of treatments on ATP

Treatments significantly affected ATP levels in zooxanthellate microcolonies regardless of the normalization procedure ($F_{6,32}=3.18$, $p = 0.0146$ for normalization to protein, $F_{6,32} = 6.41$, $p=0.0002$ for dry weight, $F_{6,32} = 5.71$, $p=0.0004$ for surface area), with reduced ATP levels being associated with glycerol (Fig. 3). For bleached microcolonies similar trends were observed, however only for normalization to surface area was the 1% glycerol with oxygen treatment significantly lower than control ($F_{5,26} = 3.84$, $p=0.0097$, pairwise 1% glycerol to control $p=0.011$).

Discussion

Stirring

Flow limitation may in-part explain the wide range of estimates for ‘light enhanced calcification’ (Gattuso 1999; Al-Horani et al., 2007). In the initial experiments with different sized stir bars

(Fig. 1) light calcification and oxygen production remained nearly constant, while smaller stir bars (and consequent reduced flow) were associated with declines in dark respiration and calcification. If such trends extend to still lower flow regimes, as may be suggested by the larger declines in dark calcification observed by Dennison and Barnes (1988) under unstirred conditions, large differences in light v dark calcification could be generated by incubating corals in static or poorly stirred containers. As suggested by experiments with elevated oxygen (Fig. 2), and experiments under aerated versus un-aerated conditions (Rinkevich and Loya 1984), oxygen limitation may be the primary factor responsible for reduced calcification in the dark. The effects of oxygen limitation are likely to be amplified under low flow conditions due to the increased thickness of the diffusive boundary layer (e.g. Kuhl et al., 1995), as has been established for respiration (e.g. Shick 1990).

Calcification

For corals with intact symbiosis (zooxanthellate microcolonies), oxygen appears to be the primary factor limiting dark calcification (Fig. 2A). A similar rate enhancement was seen with both light and oxygen in zooxanthellate microcolonies, suggesting oxygen production by the symbionts is the primary mechanism for light enhanced calcification. Similar observations have been made previously (Colombo-Pallotta et al., 2010; Wijgerde et al., 2012), however earlier studies have not monitored pCO₂ when adjusting pO₂, thus the possibility of changes in saturation state being responsible for the reported increase in calcification could not be ruled out. Here pCO₂ has been controlled during oxygen manipulation to ensure that the observed effect is a result of oxygen elevation and not concomitant degassing of CO₂. In contrast to Wijgerde et al (2012), we found no indication that oxygen tensions of ~200% (~400 µmol/kg sw) negatively affected coral growth. Indeed it seems surprising that such a modest increase in external pO₂

would be associated with substantial declines in calcification given that higher pO_2 is present at the surface of the coral tissue when photosynthesis is occurring (Shashar et al., 1993), and that such conditions are usually associated with enhanced calcification e.g. light enhanced calcification.

Oxygen alone is not sufficient to increase calcification, as seen in bleached microcolonies (Fig. 2C). A respiratory substrate must also be present to allow the coral to benefit from increased oxygen tension. The precise nature of the respiratory substrate appears to be of less importance than its concentration as suggested by an increase in calcification for both glucose and glycerol which appears to be enhanced at high substrate concentrations. However, high concentrations of fixed carbon in the seawater are not necessarily beneficial to calcification as glycerol concentrations which enhanced calcification in bleached microcolonies reduced dark calcification in zooxanthellate microcolonies (Fig 2; Colombo-Pallotta et al., 2010).

The similarity of the results of the current study and those of Colombo-Pallotta et al. (2010) suggest that oxygen limitation in the dark for zooxanthellate corals is a common phenomenon, affecting both branching (*S. pistillata*, current study), and massive corals (*Montastraea faveolata*, Colombo-Pallotta et al., 2010). However, although oxygen may generally be limiting calcification at night in zooxanthellate corals, there is likely a gradual depletion in respiratory substrates over the course of the night and eventually respiratory substrates will become limiting even for zooxanthellate corals. This may help to explain why declines in calcification with prolonged dark incubations have been observed in some studies (Al-Horani et al., 2007) but not others (Moya et al., 2006), and be partly responsible for the lack of an enhancement of calcification in earlier studies of fixed carbon supplementation (Vandermeulen and Muscatine 1974; Chalker 1975). For corals in the natural environment, there is likely to be a high degree of variability amongst corals as to whether fixed carbon or oxygen is limiting in the dark, depending upon such factors as flow, light history, photosynthetic rates, etc.

355

356 Respiration

357 Respiration was similarly affected by oxygen and fixed carbon for both zooxanthellate and
358 bleached microcolonies (Fig 2B, D), with increased rates at elevated pO_2 and further increases
359 with added fixed carbon. Despite increases in respiration, calcification did not necessarily
360 increase, with different patterns being observed in bleached versus zooxanthellate microcolonies
361 (Fig. 2). Since treatments had different effects on calcification but similar effects on respiration
362 for zooxanthellate v bleached microcolonies, there could be fundamentally different processes
363 occurring e.g. additional fixed carbon stressing the carbon replete zooxanthellate microcolonies,
364 yet benefiting the fixed carbon limited bleached colonies, or use of additional fixed carbon (and
365 energy) in biosynthetic pathways for zooxanthellate microcolonies and respiration for bleached
366 colonies. However it is also possible that the additional fixed carbon is having similar effects for
367 both zooxanthellate and bleached corals. Since the oral tissue layers (which do not carry out
368 calcification) are most exposed to the surrounding environment, they are likely to be more
369 affected by changes in the composition of the external seawater. So oxygen and fixed carbon in
370 the seawater are likely to have a greater effect on the oral tissue layers than the aboral tissue
371 layers, thus the changes in respiration may reflect primarily the oral tissue layers. For
372 zooxanthellate microcolonies with their high respiration rates (Table S2) and greater tissue
373 thicknesses (assumed from a higher protein content per unit surface area in zooxanthellate
374 microcolonies relative to bleached (Table S2), which may be associated with a greater diffusion
375 distance through the tissue), increases in respiration in the oral tissue layers in response to fixed
376 carbon may reduce oxygen availability to the calciblasts and thus reduce calcification
377 (alternatively the increased CO_2 produced by respiration could generate similar patterns). In
378 bleached microcolonies, respiration rates are lower and the tissue thickness (protein content) is
379 reduced, thus the increase in respiration by oral tissue layers may be insufficient to restrict the

oxygen supply to the calicoblasts, thus allowing the calicoblasts to benefit from the supply of fixed carbon. Zooxanthellae are likely to be more effective than external substrates in enhancing calcification due to the production of oxygen and fixed carbon within the coral tissue, and thus closer to the calicoblasts. Though even photosynthesis may not be sufficient to completely eliminate hypoxia in parts of the calicodermis, as suggested by low pO_2 values measured within some corals (Agostini et al., 2011), so hypoxia may still affect calcification even under light conditions.

Estimates of gross photosynthesis which have been made by combining net oxygen release in the light and oxygen consumption in the dark have long been known to be underestimates (Edmunds and Davies 1988; Shick 1990). Our results suggest that light respiration is stimulated by both oxygen and fixed carbon produced by photosynthesis, and, based on our observed stimulation, respiration rates are likely to be at least 20% higher in the light than in the dark. Even greater enhancements have been reported for short-term respiration measurements made immediately after light incubations (Edmunds and Davies 1988; Colombo-Pallotta et al., 2010). However, much of this enhancement of respiration may be due to respiration by the zooxanthellae, with a relatively small enhancement in the animal host (Agostini et al., 2013).

ATP

Despite different fixed carbon sources having similar effects on respiration and calcification, glucose and glycerol had markedly different effects on ATP (Fig. 3). Glycerol was associated with reduced ATP levels while little change was observed for other treatment conditions.

Although it is not clear why such a difference is observed, it could be associated with high concentrations of glycerol stressing the corals as reduced ATP levels have been associated with stress events (Fang et al., 1991). Or it may reflect different pathways being used for glycerol versus glucose (e.g. Whitehead and Douglas 2003), and the demands of those respective

pathways on cellular ATP. Light v dark conditions appear to make relatively little difference in ATP levels – here we measured slightly lower values in the light, while previously slightly higher levels have been measured in the light (Al-Horani et al., 2003), in neither case were the differences statistically significant, suggesting that corals generally regulate metabolic activities to maintain stable ATP levels, though glycerol may perturb this balance (Fig. 3). ATP levels have been suggested to be potentially useful indicators of coral health (Fang et al., 1991), however our results suggest that such an interpretation is far from clear – calcification, which is also used as an indicator of coral health, showed little relationship to ATP levels. However, ATP appears to be a potential indicator of coral biomass. ATP normalized to protein was similar for zooxanthellate and bleached microcolonies, but when normalized to either skeletal mass or surface area, ATP was lower in bleached microcolonies, suggesting that ATP is closely linked to biomass as has been observed in deep-sea corals (Hamoutene et al., 2008), and since tissue biomass is a potential indicator of coral health, ATP may be similarly useful.

Due to the nature of the measurements, we cannot, however, rule out a role for ATP levels in controlling calcification rates. Our measurements estimate total ATP and not the ATP content of the calicoblastic cells, nor the turn-over of ATP within the calicoblasts. Thus though the ATP content of the full tissue layer does not track calcification, we cannot assess whether ATP levels within different cell layers respond differently to treatments and what the ATP content of the calicoblasts may be.

Environmental context

Although aquarium conditions are far removed from natural environmental conditions in many respects – with corals often experiencing constant temperatures, flow regimes, and light intensities under aquarium conditions, many of the parameters measured agree well with measurements made on corals taken from natural environments. ATP levels for a variety of

corals collected from the wild were reported to range from 8.2 – 52.6 $\mu\text{g/g}$ (Fang et al., 1991), our measurements ($\sim 9.3 - 46.8 \mu\text{g/g}$) fall well within this range, however Fang et al. (1991) did not report values for *S. pistillata*. Our respiration rate of 12.5 $\mu\text{mol O}_2/\text{d}/\text{cm}^2$ (for zooxanthellate control corals) falls within the range measured in *S. pistillata* in the field: 0.5 – 16.6 $\mu\text{mol O}_2/\text{d}/\text{cm}^2$ (Mass et al., 2007). Calcification rates (23.7 $\mu\text{mol CaCO}_3/\text{d}/\text{cm}^2$ for zooxanthellate control corals), were however higher than recently reported in the field for *S. pistillata* -0.24 – 9.6 $\mu\text{mol CaCO}_3/\text{d}/\text{cm}^2$ (Mass et al., 2007). However, in other species, higher calcification rates have been reported during the light: 24 $\mu\text{mol CaCO}_3/\text{d}/\text{cm}^2$ (Schneider et al., 2009), 88 $\mu\text{mol CaCO}_3/\text{d}/\text{mg protein}$ (Dennison and Barnes 1988), and in the dark: 44 $\mu\text{mol CaCO}_3/\text{d}/\text{mg protein}$ (Dennison and Barnes 1988) versus our 10.2 $\mu\text{mol CaCO}_3/\text{d}/\text{mg protein}$. Despite the differences between aquarium and natural conditions, for the range of physiological parameters measured, values fall within the range encountered in the natural environment. Thus, results from the current laboratory experiment likely apply to the natural environment, though with the diversity of species, growth forms, and environmental conditions found in nature, it is difficult to assess how common oxygen or fixed carbon limitation may be in corals growing under natural conditions.

Conclusions

Oxygen limitation plays a role in explaining reduced calcification rates during dark periods for scleractinian corals. Factors such as flow rate and tissue structure likely contribute to the degree of oxygen limitation and thus influence ‘light enhanced calcification’. Fixed carbon can also be limiting, as was the case for bleached microcolonies. However changes in total ATP in the tissue did not reflect changes in calcification, but rather changed in response to the form of fixed

453 carbon. This suggests that any regulation of calcification by energy availability occurs via a
 454 pathway which is not responding directly to the total ATP level in the tissue.

455

456

Acknowledgments

We would like to thank our colleagues for assistance and fruitful discussions, particularly R. Grover, S. Sikorski, C. Rottier, C. Godinot, M. Naumann, N. Segonds, D. Desgre, A. Venn, N. Techer, C. Ferrier-Pagès, and E. Elia. We also thank G. Gaetani (Woods Hole Oceanographic Institution) and M. McCulloch (University of Western Australia) for the use of equipment.

References

- Agostini S, Suzuki Y, Higuchi T, Casareto BE, Yoshinaga K, Nakano Y, Fujimura H. 2011. Biological and chemical characteristics of the coral gastric cavity. *Coral Reefs* 31:147-156.
- Agostini S, Fujimura H, Fujita K, Suzuki Y, Nakano Y. 2013. Respiratory electron transport system activity in symbiotic corals and its link to calcification. *Aquatic Biology* 18:125-139.
- Al-Horani FA, Al-Moghrabi SM, de Beer D. 2003. The mechanism of calcification and its relation to photosynthesis and respiration in the scleractinian coral *Galaxea fascicularis*. *Marine Biology* 142: 419-426.
- Al-Horani FA, Tambutté É, Allemand D. 2007. Dark calcification and the daily rhythm of calcification in the scleractinian coral, *Galaxea fascicularis*. *Coral Reefs* 26:531-538.
- Al-Moghrabi S, Allemand D, Jaubert J. 1993. Valine uptake by the scleractinian coral *Galaxea fascicularis*: characterization and effect of light and nutritional status. *Journal Comparative Physiology B* 163: 355-362.
- Allemand D, Ferrier-Pagès C, Furla P, Houlbrèque F, Puverel S, Reynaud S, Tambutté É, Tambutté S, Zoccola D. 2011. Biomineralization in reef-building corals : from molecular mechanisms to environmental control. *CR Acad Sc Paris Palévol* 3:453-467.
- Benson BB, Krause JD. 1984. The concentration and isotopic fractionation of oxygen dissolved in freshwater and seawater in equilibrium with the atmosphere. *Limnology Oceanography* 29:620-632.
- Burriesci MS, Raab TK, Pringle JR. 2012. Evidence that glucose is the major transferred metabolite in dinoflagellate–cnidarian symbiosis. *Journal of Experimental Biology* 215:3467-3477.
- Chalker BE. 1975. Calcification, metabolism, and growth by the staghorn coral, *Acropora cervicornis* (Lamarck). Thesis; University of Miami.
- Chida J, Yamane K, Takei T, Kido H. 2012. An efficient extraction method for quantitation of adenosine triphosphate in mammalian tissues and cells. *Anal Chim Acta* 727:8-12.
- Cohen AL, Holcomb M. 2009. Why corals care about ocean acidification. *Oceanography* 22: 118-127.
- Colombo-Pallotta MF, Rodríguez-Román A, Iglesias-Prieto R. 2010. Calcification in bleached and unbleached *Montastraea faveolata*: evaluating the role of oxygen and glycerol. *Coral Reefs* 29:899-907.
- Crossland CJ, Barnes DJ. 1974. The role of metabolic nitrogen in coral calcification. *Marine Biology* 28: 325-332.
- Davies PS. 1989. Short-term growth measurements of corals using an accurate buoyant weighing technique. *Marine Biology* 101:389-395.

- 498 Dennison WC, Barnes DJ. 1988. Effect of water motion on coral photosynthesis and
499 calcification. *Journal Experimental Marine Biology Ecology* 115:67-77.
- 500 DOE. 1994. Handbook of methods for the analysis of the various parameters of the carbon
501 dioxide system in sea water version 2
- 502 Edmunds PJ. 2011. Zooplanktivory ameliorates the effects of ocean acidification on the reef coral
503 *Porites* spp. *Limnology Oceanography* 56: 2402-2410.
- 504 Edmunds PJ, Davies S. 1988. Post-illumination stimulation of respiration rate in the coral *Porites*
505 *porites*. *Coral Reefs* 7:7-9.
- 506 Falter JL, Atkinson MJ, Coimbra CFM. 2005. Effects of surface roughness and oscillatory flow
507 on the dissolution of plaster forms: Evidence for nutrient mass transfer to coral reef
508 communities. *Limnology Oceanography* 50: 246-254.
- 509 Fang L-S, Chen Y-W, Chen C-S. 1991. Feasibility of using ATP as an index for environmental
510 stress on hermatypic coral. *Marine Ecology Progress Series* 70: 257-262.
- 511 Fang L-S, Chen Y-WJ, Chen C-S. 1989. Why does the white tip of stony coral grow so fast
512 without zooxanthellae? *Marine Biology* 103: 359-363.
- 513 Fang L-S, Chen Y-WJ, Soong KY. 1987. Methodology and measurement of ATP in coral.
514 *Bulletin Marine Science* 41: 605-610.
- 515 Ferrier-Pages C, Witting J, Tambutte E, Sebens KP. 2003. Effect of natural zooplankton feeding
516 on the tissue and skeletal growth of the scleractinian coral *Stylophora pistillata*. *Coral Reefs*
517 22: 229-240.
- 518 Garcia HE, Gordon LI. 1992. Oxygen solubility in seawater: Better fitting equations. *Limnology*
519 *Oceanography* 37:1307-1312.
- 520 Gattuso JP, Allemand D, Frankignoulle M. 1999. Photosynthesis and calcification at cellular,
521 organismal and community levels in coral reefs: a review of interactions and control by
522 carbonate chemistry. *American Zoologist* 39: 160-183.
- 523 Gladfelter EH. 1982. Skeletal development in *Acropora cervicornis*: I. patterns of calcium
524 carbonate accretion in the axial corallite. *Coral Reefs* 1:45-51.
- 525 Goreau TF. 1959. The physiology of skeleton formation in corals I. a method for measuring the
526 rate of calcium deposition by corals under different conditions. *Biological Bulletin* 116:59-75.
- 527 Goreau TF. 1961. On the relation of calcification to primary productivity in reef building
528 organisms. In Lenhoff, HM, Loomis, WH ed. *The biology of hydra and some other*
529 *coelenterates*. 269-285.
- 530 Hamoutene D, Burt K, Samuelson S, Wareham V, Miller-Banoub J. 2008. Adenosine triphosphate
531 (ATP) and protein data in some species of deep-sea corals in Newfoundland and Labrador
532 Region (Northwest Atlantic Ocean). *Canadian Technical Report Fisheries Aquatic Science*
533 2801: 1-18.
- 534 Hand SC, Hardewig I. 1996. Downregulation of cellular metabolism during environmental stress:
535 mechanisms and implications. *Annual Review Physiology* 58: 539-563.
- 536 Hoegh-Guldberg O, Mumby PJ, Hooten AJ, Steneck RS, Greenfield P, Gomez E, Harvell CD,
537 Sale PF, Edwards AJ, Caldeira K, Knowlton N, Eakin CM, Iglesias-Prieto R, Muthiga N,
538 Bradbury RH, Dubi A, Hatziolos ME. 2007. Coral reefs under rapid climate change and
539 ocean acidification. *Science* 318:1737-1742
- 540 Holcomb M, Cloud JG, Woolsey J, Ingermann RL. 2004. Oxygen consumption in unfertilized
541 salmonid eggs: an indicator of egg quality? *Comparative Biochemistry Physiology A:*
542 *Comparative Physiology* 138: 349-354.
- 543 Holcomb M, McCorkle DC, Cohen AL. 2010. Long-term effects of nutrient and CO2 enrichment
544 on the temperate coral *Astrangia poculata* (Ellis and Solander, 1786). *Journal Experimental*
545 *Marine Biology Ecology* 386: 27-33.

- 546 Holcomb M, Cohen AL, McCorkle DC. .2012. An investigation of the calcification response of
547 the scleractinian coral *Astrangia poculata* to elevated pCO₂ and the effects of nutrients,
548 zooxanthellae and gender. *Biogeosciences* 9:29-39
- 549 Jacques TG, Marshall N, Pilson MEQ. 1983. Experimental ecology of the temperate scleractinian
550 coral *Astrangia danae* II. Effect of temperature, light intensity and symbiosis with
551 zooxanthellae on metabolic rate and calcification. *Marine Biology* 76: 135-148.
- 552 Jokiel PL, Maragos JE, Franzisket L. 1978. Coral growth: buoyant weight technique. In: Stoddart
553 DR, Johannes RE. eds), *Coral Reefs: Research Methods*. UNESCO monographs on
554 oceanographic methodology, Paris: 529–542.
- 555 Kardish N, Ronen R, Bubrick P, Garty J. 1987. The influence of air pollution on the concentration
556 of ATP and on chlorophyll degradation in the lichen, *Ramalina duriae* (de not.) Bagl. *New*
557 *Phytologist* 106: 697-706.
- 558 Kawaguti S, Sakumoto D. 1948. The effect of light on the calcium deposition of corals. *Bull*
559 *Oceanographica Inst Taiwan* 4:65-70.
- 560 Kuhl M, Cohen Y, Dalsgaard T, Jorgensen BB, Revsbech NP. 1995. Microenvironment and
561 photosynthesis of zooxanthellae in scleractinian corals studied with microsensors for O₂, pH
562 and light. *Marine Ecology Progress Series* 117: 159-172.
- 563 Langdon C, Atkinson MJ. 2005. Effect of elevated pCO₂ on photosynthesis and calcification of
564 corals and interactions with seasonal change in temperature/irradiance and nutrient
565 enrichment. *J Geophys Res* 110:C09S07
- 566 Lannig G, Eilers S, Portner HO, Sokolova IM, Bock C. 2010. Impact of ocean acidification on
567 energy metabolism of oyster, *Crassostrea gigas*--changes in metabolic pathways and thermal
568 response. *Marine Drugs* 8: 2318-2339.
- 569 Larsson C-M, Olsson T. 1979. Firefly assay of adenine nucleotides from algae: Comparison of
570 extraction methods. *Plant Cell Physiology* 20:145-155.
- 571 Lewis DH, Smith DC. 1971. The autotrophic nutrition of symbiotic marine coelenterates with
572 special reference to hermatypic corals. 1. Movement of photosynthetic products between
573 symbionts. *Proceedings of the Royal Society London (B)* 178:11 1-129.
- 574 Marshall AT, 1996. Calcification in hermatypic and ahermatypic corals. *Science* 271: 637-639.
- 575 Marshall AT, Clode PL. 2003. Light-regulated Ca²⁺ uptake and O₂ secretion at the surface of a
576 scleractinian coral *Galaxea fascicularis*. *Comparative Biochemistry and Physiology* 136A:
577 417-426.
- 578 Mass T, Einbinder S, Brokovich E, Shashar N, Vago R, Erez J, Dubinsky Z. 2007.
579 Photoacclimation of *Stylophora pistillata* to light extremes: metabolism and calcification.
580 *Marine Ecology Progress Series* 334: 93-102.
- 581 McCulloch M, Falter J, Trotter J, Montagna P. 2012. Coral resilience to ocean acidification and
582 global warming through pH up-regulation. *Nature Climate Change* 2:623-627
- 583 McWilliams, J.P., Cote, I.M., Gill, J.A., Sutherland, W.J., Watkinson, A.R., 2005. Accelerating
584 impacts of temperature-induced coral bleaching in the caribbean. *Ecology* 86: 2055-2060.
- 585 Mortensen PB, Rapp HT. 1998. Oxygen and carbon isotope ratios related to growth line patterns
586 in skeletons of *Lophelia pertusa* (L) (Anthozoa, scleractinia): implications for determination
587 of linear extension rates. *Sarsia* 83: 433-446.
- 588 Moya A, Tambutté S, Tambutté E, Zoccola D, Caminiti N, Allemand D. 2006. Study of
589 calcification during a daily cycle of the coral *Stylophora pistillata*: implications for ‘light-
590 enhanced calcification’. *Journal Experimental Biology* 209:3413-3419
- 591 Muscatine L. 1967. Glycerol excretion by symbiotic algae from corals and tridacna and its
592 control by the host. *Science* 156:516-519

- 593 Nakamura T, Nadaoka K, Watanabe A. 2013. A coral polyp model of photosynthesis, respiration
594 and calcification incorporating a transcellular ion transport mechanism. *Coral Reefs* 32:779-
595 794.
- 596 Oku H, Yamashiro H, Onaga K. 2003. Lipid biosynthesis from [14C]-glucose in the coral
597 *Montipora digitata*. *Fisheries Science* 69: 625-631.
- 598 Oku H, Yamashiro H. 2003. Effect of glucose addition on the coral *Montipora digitata*. Sangusho
599 ni kansuru Chosa Kenkyu Hokokusho Heisei 14 Nendo 40-45.
- 600 Pearse VB, Muscatine L. 1971. Role of symbiotic algae (Zooxanthellae) in coral calcification.
601 *Biological Bulletin* 141: 350-363.
- 602 Ries JB. 2011. A physicochemical framework for interpreting the biological calcification
603 response to CO₂-induced ocean acidification. *Geochimica Cosmochimica Acta* 75: 4053-
604 4064.
- 605 Rands ML, Douglas AE, Loughman BC, Ratcliffe RG. 1992. Avoidance of hypoxia in a cnidarian
606 symbiosis by algal photosynthetic oxygen. *Biological Bulletin* 182: 159-162.
- 607 Rinkevich B, Loya Y. 1984. Does light enhance calcification in hermatypic corals? *Marine*
608 *Biology* 80: 1-6.
- 609 Rodrigues LJ, Grottoli AG. 2006. Calcification rate and the stable carbon, oxygen, and nitrogen
610 isotopes in the skeleton, host tissue, and zooxanthellae of bleached and recovering Hawaiian
611 corals. *Geochimica Cosmochimica Acta* 70: 2781-2789.
- 612 Schneider K, Levy O, Dubinsky Z, Erez J. 2009. In situ diel cycles of photosynthesis and
613 calcification in hermatypic corals. *Limnol. Oceanogr.* 54: 1995-2002.
- 614 Shashar N, Cohen Y, Loya Y. 1993. Extreme diel fluctuations of oxygen in diffusive boundary
615 layers surrounding stony corals. *Biological Bulletin* 185: 455-461.
- 616 Shick JM, 1990. Diffusion limitation and hyperoxic enhancement of oxygen consumption in
617 zooxanthellate sea anemones, zoanthids, and corals. *Biological Bulletin* 179: 148-158.
- 618 Smith JE, Shaw M, Edwards RA, Obura D, Pantos O, Sala E, Sandin SA, Smriga S, Hatay M,
619 Rohwer FL. 2006. Indirect effects of algae on coral: algae-mediated, microbe-induced coral
620 mortality. *Ecology Letters* 9: 835-845.
- 621 Stephens GC. 1960. Uptake of glucose from solution by the solitary coral, *Fungia*. *Science* 131:
622 1532.
- 623 Stephens GC. 1962. Uptake of organic material by aquatic invertebrates. I. uptake of glucose by
624 the solitary coral, *Fungia scutaria*. *Biological Bulletin* 123: 648-659.
- 625 Stimson J, Kinzie RA III. 1991. The temporal pattern and rate of release of zooxanthellae from
626 the reef coral *Pocillopora damicornis* (Linnaeus) under nitrogen-enrichment and control
627 conditions. *Journal Experimental Marine Biology Ecology* 153: 63-7
- 628 Tambutté S, Holcomb M, Ferrier-Pagès C, Reynaud S, Tambutté É, Zoccola D, Allemand D.
629 2011. Coral biomineralization: From the gene to the environment. *Journal Experimental*
630 *Marine Biology Ecology* 408: 58-78
- 631 Taylor DL. 1977. Intra-colonial transport of organic compounds and calcium in some atlantic reef
632 corals. *Proceedings of the 3rd International Coral Reef Symposium*, 431-435.
- 633 van Heuven S, Pierrot D, Lewis E, Wallace DWR. 2009. MATLAB Program Developed for CO₂
634 System Calculations. ORNL/CDIAC-105b. Carbon Dioxide Information Analysis Center,
635 Oak Ridge National Laboratory, U.S. Department of Energy, Oak Ridge, Tennessee
- 636 Vandermeulen JH, Davis ND, Muscatine L. 1972. The effect of inhibitors of photosynthesis on
637 zooxanthellae in corals and other marine invertebrates. *Marine Biology* 16:185-191
- 638 Vandermeulen JH, Muscatine L. 1974. Influence of symbiotic algae on calcification in reef
639 corals: critique and progress report. In Vernberg WB, ed. *Symbiosis in the Sea* 1-19.

- 640 Whitehead LF, Douglas AE. 2003. Metabolite comparisons and the identity of nutrients
641 translocated from symbiotic algae to an animal host. *Journal of Experimental Biology* 206:
642 3149-3157.
- 643 Wijgerde T, Jurriaans S, Hoofd M, Verreth JAJ, Osinga R. 2012. Oxygen and Heterotrophy Affect
644 Calcification of the Scleractinian Coral *Galaxea fascicularis*. *PLoS ONE* 7(12): e52702.
645 doi:10.1371/journal.pone.0052702
- 646 Yonge CM. 1931. The significance of the relationship between corals and zooxanthellae. *Nature*
647 128:309-311
- 648 Young SD, O'Connor JD, Muscatine L. 1971. Organic material from scleractinian coral skeletons
649 - II. Incorporation of ¹⁴C into protein, chitin and lipid. *Comparative Biochemistry and*
650 *Physiology* 40B: 945-958.

652

Figure 1

Effect of stir bar size

Figure 1. Effect of stir bar size on relative calcification rates (A) and oxygen consumption (B) by zooxanthellate microcolonies maintained in the light or in the dark, the 40 mm stir bar was used as the control value for calculating relative rates. N=4 or greater for all treatments, symbols are mean, error bars are standard deviation.




Figure 2

Relative calcification and respiration rates

Figure 2. Relative calcification (A, C) and respiration (B, D) rates for zooxanthellate (A, B) and bleached (C, D) microcolonies maintained under each treatment condition. Treatments were: cont = control, ab = antibiotics, O2 = ~2x atmospheric pO2, glu20um = ~20 uM glucose with oxygen, glu1mm = ~1 mM glucose with oxygen, gly1mm = ~1 mM glycerol with oxygen, gly120mM = 120 mM glycerol with oxygen, light = light. With the exception of the light treatment, all incubations were conducted in the dark. N=4 or greater for all treatments, symbols are mean, error bars are standard deviation, * = treatment significantly different from control at $p = 0.05$.




Figure 3

ATP measurements

Figure 3. ATP contents (μg) normalized to: mg protein (up triangle, plotted against left axis), cm^2 surface area (filled circles, plotted against left axis), and g skeletal dry weight (squares, plotted against the right axis) for zooxanthellate (A) and bleached (B) microcolonies exposed to each treatment (per Fig. 2). N=2 or greater for all treatments, symbols are mean, error bars are standard deviation, * = treatment significantly different from control at $p = 0.05$.

