

***Trichoplax adhaerens* reveals an endocrine-like network sensitive to 9-*cis*-retinoic acid at the base of metazoan evolution (#6219)**

1

First submission

Please read the **Important notes** below, the **Review guidance** on page 2 and our **Standout reviewing tips** on page 3. When ready [submit online](#). The manuscript starts on page 4.

Important notes

Editor and deadline

Christopher Cooper / 7 May 2017

Files

8 Figure file(s)

7 Raw data file(s)

Please visit the overview page to [download and review](#) the files not included in this review PDF.

Declarations

One or more DNA sequences were reported.



Please read in full before you begin

How to review

When ready [submit your review online](#). The review form is divided into 5 sections. Please consider these when composing your review:

- 1. BASIC REPORTING**
- 2. EXPERIMENTAL DESIGN**
- 3. VALIDITY OF THE FINDINGS**
4. General comments
5. Confidential notes to the editor

 You can also annotate this PDF and upload it as part of your review

To finish, enter your editorial recommendation (accept, revise or reject) and submit.

BASIC REPORTING

-  Clear, unambiguous, professional English language used throughout.
-  Intro & background to show context. Literature well referenced & relevant.
-  Structure conforms to [PeerJ standards](#), discipline norm, or improved for clarity.
-  Figures are relevant, high quality, well labelled & described.
-  Raw data supplied (see [PeerJ policy](#)).

EXPERIMENTAL DESIGN

-  Original primary research within [Scope of the journal](#).
-  Research question well defined, relevant & meaningful. It is stated how the research fills an identified knowledge gap.
-  Rigorous investigation performed to a high technical & ethical standard.
-  Methods described with sufficient detail & information to replicate.

VALIDITY OF THE FINDINGS

-  Impact and novelty not assessed. Negative/inconclusive results accepted. *Meaningful* replication encouraged where rationale & benefit to literature is clearly stated.
-  Data is robust, statistically sound, & controlled.
-  Conclusions are well stated, linked to original research question & limited to supporting results.
-  Speculation is welcome, but should be identified as such.

The above is the editorial criteria summary. To view in full visit <https://peerj.com/about/editorial-criteria/>

7 Standout reviewing tips

3



The best reviewers use these techniques

Tip

Example

Support criticisms with evidence from the text or from other sources

Smith et al (J of Methodology, 2005, V3, pp 123) have shown that the analysis you use in Lines 241-250 is not the most appropriate for this situation. Please explain why you used this method.

Give specific suggestions on how to improve the manuscript

Your introduction needs more detail. I suggest that you improve the description at lines 57- 86 to provide more justification for your study (specifically, you should expand upon the knowledge gap being filled).

Comment on language and grammar issues

The English language should be improved to ensure that your international audience can clearly understand your text. I suggest that you have a native English speaking colleague review your manuscript. Some examples where the language could be improved include lines 23, 77, 121, 128 - the current phrasing makes comprehension difficult.

Organize by importance of the issues, and number your points

1. Your most important issue
2. The next most important item
3. ...
4. The least important points

Give specific suggestions on how to improve the manuscript

Line 56: Note that experimental data on sprawling animals needs to be updated. Line 66: Please consider exchanging "modern" with "cursorial".

Please provide constructive criticism, and avoid personal opinions

I thank you for providing the raw data, however your supplemental files need more descriptive metadata identifiers to be useful to future readers. Although your results are compelling, the data analysis should be improved in the following ways: AA, BB, CC

Comment on strengths (as well as weaknesses) of the manuscript

I commend the authors for their extensive data set, compiled over many years of detailed fieldwork. In addition, the manuscript is clearly written in professional, unambiguous language. If there is a weakness, it is in the statistical analysis (as I have noted above) which should be improved upon before Acceptance.

***Trichoplax adhaerens* reveals an endocrine-like network sensitive to 9-*cis*-retinoic acid at the base of metazoan evolution**

Jan Philipp Novotný^{1,2}, Ahmed Ali Chughtai¹, Markéta Kostrouchová^{1,3}, Veronika Kostrouchová¹, David Kostrouch¹, Filip Kaššák¹, Radek Kaňa⁴, Bernd Schierwater^{5,6}, Marta Kostrouchová¹, Zdenek Kostrouch

Corresp. 1

¹ Biocev, First Faculty of Medicine, Charles University in Prague, Vestec, Czech Republic

² Department of Medicine V., University of Heidelberg, Heidelberg, Germany

³ Department of Pathology, Third Faculty of Medicine, Charles University Prague, Prague, Czech Republic

⁴ Institute of Microbiology, Laboratory of Photosynthesis, Czech Academy of Sciences, Třeboň, Czech Republic

⁵ Institute for Animal Ecology and Cell Biology, University of Veterinary Medicine, Hannover, Germany

⁶ Department of Molecular, Cellular and Developmental Biology, Yale University, New Haven, CT, United States

Corresponding Author: Zdenek Kostrouch

Email address: Zdenek.kostrouch@lf1.cuni.cz

Trichoplax adhaerens, the only known species of Placozoa is likely to be closely related to an early metazoan that preceded branching of Cnidaria and Bilateria. This animal species is surprisingly well adapted to free life in the World Ocean inhabiting tidal coastal zones of oceans and seas with warm to moderate temperatures and shallow waters. The genome of *T. adhaerens* (sp. Grell) includes four nuclear receptors, namely homologues of RXR (NR2B), HNF4 (NR2A), COUP (NR2F) and ERR (NR3B) that show a high degree of similarity with human homologues. In the case of RXR, the sequence identity to human RXR alpha reaches 81% in the DNA binding domain and 70 % in the ligand binding domain. We show that *T. adhaerens* RXR (TaRXR) binds 9-*cis* retinoic acid (9-*cis*-RA) with high affinity, as well as high specificity and that exposure of *T. adhaerens* to 9-*cis*-RA regulates the expression of the putative *T. adhaerens* homologue of vertebrate L-malate-NADP+ oxidoreductase (EC 1.1.1.40) which in vertebrates is regulated by a heterodimer of RXR and thyroid hormone receptor. Treatment by 9-*cis*-RA alters the relative expression profile of *T. adhaerens* nuclear receptors, suggesting the existence of natural ligands. Keeping with this, algal food composition has profound effect on *T. adhaerens* growth and appearance. Our results uncover an endocrine-like network of nuclear receptors sensitive to 9-*cis*-RA in *T. adhaerens* and support the existence of a ligand-sensitive network of nuclear receptors at the base of metazoan evolution.

***Trichoplax adhaerens* reveals an endocrine-like network sensitive to 9-cis-retinoic acid at the base of metazoan evolution.**

Jan Philipp Novotný^{1,2}, Ahmed Chughtai¹, Markéta Kostrouchová^{1,3}, Veronika Kostrouchová¹, David Kostrouch¹, Filip Kaššák¹, Radek Kaňa⁴, Bernd Schierwater^{5,6}, Marta Kostrouchová¹, Zdeněk Kostrouch¹

1 Biocev, First Faculty of Medicine, Charles University, Vestec, Czech Republic

2 Department of Medicine V, University of Heidelberg, Heidelberg, Germany

3 Department of Pathology, Third Faculty of Medicine, Charles University, Ruská 87, 10000 Praha 10, Czech Republic

4 Laboratory of Photosynthesis, Centrum Algatech, Institute of Microbiology, Academy of Sciences Novohradska 237 - Opatovický mlýn, CZ 379 01 Třeboň, Czech Republic.

5 Division of Ecology and Evolution, Stiftung Tierärztliche Hochschule, Hannover Germany

6 Department of Molecular, Cellular and Developmental Biology, Yale University

New Haven, CT, USA

*Corresponding author:

Zdenek Kostrouch

Biocev, First Faculty of Medicine,

Charles University in Prague,

Průmyslová 595, 252 42 Vestec,

Czech Republic

email: zdenek.kostrouch@lf1.cuni.cz

ABSTRACT

Trichoplax adhaerens, the only known species of Placozoa is likely to be closely related to an early metazoan that preceded branching of Cnidaria and Bilateria. This animal species is surprisingly well adapted to free life in the World Ocean inhabiting tidal costal zones of oceans and seas with warm to moderate temperatures and shallow waters. The genome of *T. adhaerens* (sp. Grell) includes four nuclear receptors, namely homologues of RXR (NR2B), HNF4 (NR2A), COUP (NR2F) and ERR (NR3B) that show a high degree of similarity with human homologues. In the case of RXR, the sequence identity to human RXR alpha reaches 81% in the DNA binding domain and 70 % in the ligand binding domain. We show that *T. adhaerens* RXR (TaRXR) binds 9-*cis* retinoic acid (9-*cis*-RA) with high affinity, as well as high specificity and that exposure of *T. adhaerens* to 9-*cis*-RA regulates the expression of the putative *T. adhaerens* homologue of vertebrate L-malate-NADP⁺ oxidoreductase (EC 1.1.1.40) which in vertebrates is regulated by a

heterodimer of RXR and thyroid hormone receptor. Treatment by 9-*cis*-RA alters the relative expression profile of *T. adhaerens* nuclear receptors, suggesting the existence of natural ligands. Keeping with this, algal food composition has profound effect on *T. adhaerens* growth and appearance. Our results uncover an endocrine-like network of nuclear receptors sensitive to 9-*cis*-RA in *T. adhaerens* and support the existence of a ligand-sensitive network of nuclear receptors at the base of metazoan evolution.

INTRODUCTION

Life on Earth began 4.1 to 3.5 billion years ago (Bell et al. 2015) with the appearance of the first unicellular prokaryotic organisms that subsequently evolved, in part, to multicellular lifeforms forming the kingdom Metazoa that have specialized tissues for digestion, regulation of homeostasis, locomotion, perception, analysis of the environment and reproduction.

In contrast to unicellular organisms, metazoans are in need of regulatory mechanisms that provide the means of coordination between various tissues in a tight arrangement with cellular homeostasis. This coordination on the level of humoral signaling includes regulation by nuclear receptors (NRs), which respond to small, mostly hydrophobic molecules, including hormones produced by specific tissues, metabolites or even molecules present in the environment and transfer these signals to the nucleus, and thus leading to a dynamically changing but adaptive gene expression (Escriva et al. 2004).

NRs therefore play an important role in maintaining intra- and inter-cellular functions in multicellular organisms. Their overall structure is conserved in most nuclear receptors and consists

of an A/B (N-terminal) domain, the DNA binding domain (DBD), a hinge region, the ligand binding domain (LBD) and the C-terminal domain (Kumar & Thompson 1999; Robinson-Rechavi et al. 2003). The DBD and LBD of NRs exhibit an especially high degree of conservation and the changes that were acquired during evolution allow for classification of the NR protein family into six subfamilies (Laudet 1997; Escriva et al. 1998). NRs that evolved within these subfamilies show functional connections that include specialization of regulatory functions in time or cell type restriction, fortification of ancestral functions or their specific inhibition by newly evolved NRs (Escriva et al. 2004; Kostrouchova & Kostrouch 2015).

With the overall structure maintained across metazoan species, nuclear receptors show significant heterogeneity regarding their quantity and function, many of which have not yet been explored in e.g. *Caenorhabditis elegans* with over 280 nuclear receptors (reviewed in (Kostrouchova & Kostrouch 2015)).

The evolutionary changes accumulated in diversified NRs allow functional subspecialization at the level of specific sequence binding within gene promoters (response elements), protein-protein interactions with functionally linked receptor interactors and adoption of new ligands as specific hormonal regulators. The evolution of hormonal ligands acquired by different species during evolution is well documented and indicates the potential of NRs to adopt new ligands as regulators (Escriva et al. 2000; Markov & Laudet 2011). This is accompanied by two essential questions: to what degree is the plasticity of ligand selection a fundamental property of NRs; and what might be the origin of specific ligand binding by NRs. It has been suggested that the original NR, which is the ancestral NR possessing gene regulatory capacity, may have been an unliganded molecular regulator. More recently, it was proposed that the ligand

binding and ligand-dependent regulatory potential of NRs is an inherent feature of the evolution of NRs (Bridgham et al. 2010).

A search for NRs that may be closely related to an ancient ancestor of the NR family led to the discovery of an RXR homologue in Cnidaria (Kostrouch et al. 1998). Surprisingly, this receptor showed not only extremely high degree of sequence homology with vertebrate RXRs, far surpassing the degree of conservation observed in insects, but also by its ability to bind the same ligand as vertebrate RXRs, *9-cis*-retinoic acid (*9-cis*-RA), with an affinity close to that reported for vertebrate RXRs. Similarly, as vertebrate RXRs, the jellyfish RXR showed specific binding preference for *9-cis* RA over all-*trans*-retinoic acid (AT-RA) and was able to heterodimerize with vertebrate thyroid hormone receptor alpha. Recent genome sequencing projects confirmed the existence of highly conserved RXR across several metazoan species including insects (*Locusta migratoria*, (Nowickyj et al. 2008)) that are evolutionarily older than species with a more diversified RXR homologue such as Usp found in *Drosophila* (reviewed in (Gutierrez-Mazariegos et al. 2014)).

To date, the nuclear receptor network has mainly been studied in complex organisms already in possession of an extensive endocrine network. Genome analysis of the basal metazoan *Trichoplax adhaerens* by whole genome sequencing revealed four highly conserved nuclear receptors, namely homologues of HNF4 (NR2A), RXR (NR2B), ERR (NR3) and COUP (NR2F) (Srivastava et al. 2008). Surprisingly, the degree of conservation of the predicted placozoan NRs with known vertebrate NRs is not only very high at the level of the predicted secondary structure, as can be expected for true NRs, but also at the level of the primary amino acid sequence. Especially the similarity of the placozoan RXR (TaRXR) to its vertebrate homologues is high, as it is in the case of the cubomedusan RXR. *T. adhaerens*, which shows characteristics of a basal

metazoan with only few cell types (Smith et al. 2014) and a relatively simple 4 member NR complement, offers a unique model that may shed light on the evolution of gene regulation by NRs.

In this presented work, we attempted to study the placozoan RXR homologue functionally. Our results show that *T. adhaerens* RXR binds 9-*cis*-RA with an affinity comparable to that of vertebrate and jellyfish RXRs and *T. adhaerens* responds to nanomolar concentrations of 9-*cis*-RA with a transcriptional upregulation of the putative homologue of malic enzyme that is regulated by a heterodimer formed by liganded thyroid hormone receptor and RXR in vertebrates. We also show that 9-*cis*-RA affects the relative expression of the four NRs present in *T. adhaerens* genome suggesting that these NRs may form a regulatory network capable of responding to possible ligands present in these animals or in their environment. In line with this, growth, multiplication and appearance of *T. adhaerens* are strongly affected by food composition, especially by red pigment containing algae suggesting that specific food components or their metabolites may be ligands involved in the ancestral regulatory network of NRs.

METHODS

Bioinformatics and cloning of RXR

The predicted RXR gene models on jgi (<http://jgi.doe.gov/>) (Nordberg et al. 2014) were screened for the characteristic molecular signature of the DNA binding domain (C-X2-C-X13-C-X2-C-X15-C-X5-C-X9-C-X2-C-X4-C-X4-M) (Kostrouch et al. 1995) and the appropriate predicted gene model (protein ID 53515) was selected for further use.

The alignment of different RXRs was performed by Clustal Omega (<http://www.ebi.ac.uk/Tools/msa/clustalo/>) (Sievers et al. 2011) and adjusted/exported as an image file using Jalview (<http://www.jalview.org>). Protein domain characterization was performed with SMART (Schultz et al. 1998; Letunic et al. 2015). Analysis of HNF4, ERR and COUP was done similarly. Phylogenetic analysis was performed on RXR ClustalO alignment using PhyMLv3.1 (Guindon et al. 2010) using SeaView v4.6.1 with a 100 bootstrap analysis and SPR distance computation. The tree was then visualized using FigTree v1.4.3.

T. adhaerens total RNA was obtained from 50-100 animals and extracted using TRIZOL® reagent (Invitrogen, Carlsbad, CA, USA) according to the protocol supplied by the manufacturer.

Subsequently, cDNA was prepared with random hexamers and SuperScript III (Invitrogen™) according to the manufacturer's protocol.

Several RXR transcripts were then amplified by PCR with primers covering the starting sequence ((GCGGATCC)ATGGAGGACAGATCGTTTAAAAAA), starting at 32 bp 5' of ATG (TCTACCAATGTTTATCGCATCGGTTA) and starting at 97 bp 5' of ATG (TTAAGGCTTAACTGATGATGTTGTGAATG) with a common reverse primer covering the last 24 bp of the predicted gene sequence ((CGGAATTC)TTAAGAACTGCCTGTTTCCAGCAT).

Each PCR product was then ligated into pCR®2.1-TOPO® or pCR®4-TOPO® vector with the classic TA Cloning Kit and TOPO TA Cloning Kit (Invitrogen™), respectively. The ligated products were then transformed using One Shot® TOP10 Chemically Competent *E. coli* and cultured on LB Agar plates containing 100 µg/ml ampicillin. Plasmid DNA was extracted from obtained colonies and screened for mutations by sequencing using vector specific M13 forward and reverse primers. Only non-mutated sequences were used in subsequent experiments.

The RXR fragments were then restricted and inserted into pGEX-2T vector system for bacterial expression (Addgene, Cambridge, MA, USA). Proper insertion was verified by sequencing.

Protein expression

BL21 pLysS bacteria were transformed with previously described RXR mRNA inserted into pGEX-2T vector. Stocks of transformed bacteria were stored in 8% glycerol according to the Novagen pET System Manual (11th edition) (https://www.google.cz/search?q=Novagen+pET+System+Manual+&ie=utf-8&oe=utf-8&client=firefox-b&gfe_rd=cr&ei=T9z1WMHJDsni8AfpmoGoCQ). For protein expression, bacteria were scraped from stock and incubated in Liquid Broth (LB) with ampicillin (100 µg/ml) and chloramphenicol (34 µg/ml) overnight. The culture was then used to inoculate 100 ml of LB + antibiotics and grown to OD600 = 0.6-0.8 at 37 °C, then induced with 100 µl 1M IPTG (isopropyl-D-thiogalactopyranoside) (Sigma-Aldrich, St. Louis, MO, USA). and moved to 25 °C (RT) for 5 h. The culture was then spun at 9000 ~~x~~g for 15 min and the supernatant discarded. The bacterial pellet was resuspended in 10 ml GST binding buffer (25 mM Tris pH 7.5, 150 mM NaCl, 1 mM EDTA + protease inhibitor (S8820 Sigma Fast, Sigma-Aldrich, St. Louis, MO, USA or cOmplete™, EDTA-free Protease Inhibitor Cocktail, Roche, Basel, Switzerland)). Bacteria were then lysed by 6 x 20 s ultrasonication on ice (50 watts, 30kHz, highest setting – 100%) (Ultrasonic Processor UP50H, (Hielscher Ultrasonics GmbH, Teltow, Germany) and subsequently incubated with 15-20 mg glutathione agarose beads (Sigma-Aldrich®) prepared according to manufacturer's instructions. Incubation took place at 4 °C for about 10 h and beads were washed according to instructions, resuspended in regeneration buffer (50mM Tris-HCl pH7.4, 1mM EDTA, 120 mM KCl, 5 mM DTT, 8% glycerol (v/v)) or 50mM TRIS–HCl pH 7.4 + 9% (v/v) glycerol for

subsequent thrombin (bovine plasma, Sigma-Aldrich®) cleavage, if performed, and then adjusted for regeneration buffer conditions.

The size of the GST-RXR fusion protein was checked by SDS-PAGE. Thrombin cleavage was performed at RT for 4 h.

Radioactive *9-cis-R* binding assay

Radioactive *9-cis*-RA and AT-RA were purchased from PerkinElmer (Waltham, MA, USA). Binding was performed in 100 µl binding buffer (50mM Tris-HCl pH7.4, 1mM EDTA, 120 mM KCl, 5 mM DTT, 8% glycerol (v/v), 0.3% to 0.5% (w/v) CHAPS (3-[(3-Cholamidopropyl)dimethylammonio]-1-propanesulfonate hydrate, Sigma-Aldrich) for 2 h on wet ice in a dark environment. The protein used for binding was either GST-TaRXR fusion protein on beads with about 375 ng/assay or thrombin-cleaved TaRXR. For estimation of specific binding, 200x excess of either *9-cis*-RA or AT-RA (Sigma-Aldrich) was used. In case of GST-RXR fusion protein, 50 µl of the supernatant was removed after 30 s centrifugation at 1300 ~~x~~g and washed 3x with 1000 µl wash buffer (50 mM Tris-HCl pH7.4, 1 mM EDTA, 120 mM KCl, 5 mM DTT, 8% (v/v) glycerol, 0.5% (w/v) CHAPS) removing 900 µl after each wash. For cleaved RXR protein 10 µl hydroxyapatite slurry suspended in binding buffer (12.7 mg/100µl) was added to the assay and mixed twice, collecting the apatite slurry by centrifugation (15 s at 600 xg). 95 µl of the supernatant was removed and the slurry washed twice with 1 ml of wash buffer, removing 900 µl after each wash. Work with retinoids was done under indirect illumination with 60W, 120V yellow light bulb (BugLite, General Electric Co, Nela Parc, Cleveland Oh, USA) as described (Cahnmann 1995).

201 **Culture of *T. adhaerens* and algae**

202 *Trichoplax adhaerens* was cultured in Petri dishes containing filtered artificial seawater
 203 (Instant Ocean, Spectrum Brands, Blacksburg, VA, USA) with a salinity of approx. 38-40 ppt.
 204 *Rhodomonas salina* (strain CCAP 978/27), *Chlorella sp.*, *Porphyridium cruentum* (UTEX B637)
 205 and other non-classified algae, as well as aquarium milieu established in the laboratory by mixing
 206 salt water obtained from a local aquarium shop were used to maintain the stock. The cultures were
 207 kept at approx. 23°C and an automated illumination for 12 h/day was used with a conventional
 208 lightbulb on a daylight background from late spring to mid-summer in the laboratory located at
 209 50.0703122N, 14.4293456E with laboratory windows oriented eastward. The natural illumination
 210 included almost direct morning light from 8 AM to 10.30 AM, indirect sunlight for most of the
 211 daytime and sunlight reflected from a building across the street from 1 PM to 6 PM. Algae were
 212 maintained as described (Kana et al. 2012; Kana et al. 2014). The experiments were performed
 213 predominantly during sunny weather.

214

215 **Treatment of *T. adhaerens* with retinoic acids**

216 Incubation of the animals was done overnight in the absence of light. Each batch within an
 217 experiment was derived from similar cultures and fed with similar amounts and composition of
 218 algae. All experiments were started in a dark room with indirect yellow light illumination
 219 (similarly as in case of the ligand binding studies) and further incubations were done in the dark.

220

221 **Quantitative PCR**

222 Droplet digital PCR was performed on a QX100 Droplet Digital PCR System (Bio-Rad
 223 Laboratories, Hercules, CA, USA). For this, *T. adhaerens* was cultured according to culture

224 conditions described and 4-10 animals were removed per 100µl TRIzol reagent. RNA purification
225 was performed according to manufacturer's instructions. Total RNA was measured by a UV
226 spectrophotometer and used as a reference for normalization.

227 Reverse transcription was performed with SuperScript III Reverse Transcriptase according
228 to manufacturer's instructions. The cDNA was then mixed with ddPCR Supermix (Bio-Rad)
229 according to the manufacturer's instructions (corresponding to 100-500 ng of RNA) and analyzed.
230 PCR primers were designed using the UPL online ProbeFinder (Roche) software and were as
231 follows:

232 TaRXR – left:tctgcaagttggtatgaagca, right: agttggtgtgctattctttacgc

233 TaHNF4 ([ref|XM_002115774.1|](#)):

234 left: ggaatgatttgattttacctcgac, right: tacgacaagcgatacgagca

235 TaCOUP ([ref|XM_002109770.1|](#)):

236 left: attttgaatgctgccaatg , right: ttactggttgaggatggaac

237 TaSoxB1 ([ref|XM_002111308.1|](#)):

238 left: tgtcagatgcggataaacga , right: ggatgttcctcatgtgtaatgc

239 TaTrox-2 ([ref|XM_002118165.1|](#)):

240 left: gcctatagtcgacctgccata , right: ttggtgatgatggtgtcca

241 TaPaxB1 ([gb|DQ022561.1|](#)):

242 left: tcaaacgggttctgtagcc , right: ggtgtgccaccttaggc

243 TaERR (nuclear receptor 3, [gb|KC261632.1|](#)):

244 left: ttacgcatgtgatggttatgg , right: agcgtgcctatttatttctct

245

Results were subsequently analyzed using the Bio-Rad ddPCR software. Manual correction of the cut off was performed when automated analysis was not possible. To visualize changes in nuclear receptor expression in the absence of a reliable housekeeping gene as reference, we considered the absolute quantity of each nuclear receptor as a percentage of the overall nuclear receptor expression and subsequently visualized the change of receptor expression by subtraction of the percentage of the control experiment. Absolute copy numbers of the proposed malic enzyme homologue in *T. adhaerens* have been normalized to overall RNA quantity for expressional analysis.

Experiments with quantification by qRT-PCR were performed on a Roche LightCycler II with OneTaq polymerase and the same probes as for ddPCR.

For the estimation of the relative expression of NRs in small (< 0.5 mm) versus big animals (>1 mm), 20 to 30 animals from the same culture were used for each paired experiment.

Identification of *T. adhaerens* homologue of L-malate-NADP+ oxidoreductase (EC 1.1.1.40)

P48163 (MAOX_HUMAN) protein sequence was used as query sequence and searched against *T. adhaerens* database with BLASTP on <http://blast.ncbi.nlm.nih.gov/Blast.cgi> using standard algorithm parameters. The best hit was a hypothetical protein TRIADDRAFT_50795 with a sequence identity of 57% and a query coverage of 93% and was assumed to be *T. adhaerens* closest homologue of vertebrate L-malate-NADP+ oxidoreductase.

Microscopy and image analysis

Observation of *T. adhaerens* was done with an Olympus SZX10 microscope equipped with DF Plan 2x objective and Olympus DP 73 camera operated by CellSens Dimension computer

program (kindly provided by Olympus, Prague, Czech Republic) or Olympus CKX41 or SZX7 with Olympus E-410 camera and QuickPhoto Micro 3.1 program.

Circularity was calculated by establishing the area (A) and perimeter (p) of *T. adhaerens* using ImageJ (<https://imagej.nih.gov/ij/>) and then calculated with the isoperimetric quotient $Q = \frac{4\pi A}{p^2}$, (A – Area, p – perimeter). GraphPad Prism 5 (or higher) was used for graphical representation and calculations of the confidence intervals with $p = 0.05$.

RESULTS

***T. adhaerens* retinoid X receptor shows high cross-species sequence identity.**

By using the ab initio model of the JGI *Trichoplax* database as reference we screened the *Trichoplax* JGI database for RXR homologues with a complete DBD and LBD sequence and were able to obtain, as well as verify a full length RXR transcript previously not annotated as the ‘best model’. Blastp analysis showed a high sequence similarity to human, as well as mouse RXR with 66% overall sequence identity to human RXRalpha.

SMART analysis of the proposed TaRXR sequence showed a zinc finger DNA binding domain (amino acid residues 16-87) and a ligand binding domain (amino acid residues 155-342) with E values $<10^{-40}$. Blast analysis of the zinc finger DNA binding and ligand binding domains revealed a sequence identity of 81% and 70% to human RXR alpha, respectively. Both domains contained the predicted molecular pattern characteristic for each domain. The heptad repeat LLLRLPAL proposed for dimerization activity (Forman & Samuels 1990b; Forman & Samuels 1990a; Kiefer 2006) as well as the LBD signature for 9-*cis*-RA binding Q-x(33)-L-x(3)-F-x(2)-R-

x(9)-L-x(44)-R-x(63)-H were present (**Fig. 1**) and also amino acid residues shown as critical for binding 9-*cis*-retinoic acid (Egea et al. 2000) . Out of 11 amino acid residues critical for 9-*cis*-RA binding (A271, A272, Q 275, L 309, F 313, R 316, L 326, A 327, R 371, C 432, H 435) 9 are conserved (A271, A272, Q 275, L 309, F 313, R 316, L 326, A 327, R 371, C 432, H 435), while the remaining two amino acids are substituted [A327S, and C432T (C432A in *Tripedalia cystophora*)]. Due to the high sequence identity, we propose a 9-*cis*-retinoic acid binding capability of the hypothesized TaRXR sequence, as well as DNA binding capability.

Phylogenetic analysis using PhyML algorithm indicates that TaRXR is likely to precede branching of RXRs in cubomedusae and scyphomedusae and clusters with RXRs in Bilateria (**Fig 2**).

The remaining three NRs identified in the *T. adhaerens* genome show also very high overall sequence identity with vertebrate homologues (**Table 1**). Alignments of *T. adhaerens* HNF4, COUP and ERR with homologues from selected species can be found in **Supplementary File S2**.

Table 1. BLASTP comparison of *T. adhaerens* NRs with the human closest homologues and cross-species conservation of RXR homologues

<i>T. adhaerens</i> homologue	Percent identity (<i>H. sapiens</i> homologue)
TaRXR	66% (RXR α)
TaCOUP	43% (COUP1)
TaHNF4	67% (HNF4 α)
TaERR	46% (ERR3)

<i>Drosophila melanogaster</i> homologue	Percent identity (<i>H. sapiens</i> homologue)
USP	47% (RXR α)

310

<i>T. adhaerens</i> homologue	Cnidaria (<i>T. cysotophora</i> homologue)
TaRXR	35% (jRXR)

311

Cnidaria (<i>T. cysotophora</i> homologue)	Percent identity (<i>H. sapiens</i> homologue)
jRXR	48% (RXR α)

312

313

314 **TaRXR shows preferential binding affinity to 9-*cis* retinoic acid over all-*trans*-**
 315 **retinoic acid**

316 Consistent with the bioinformatically high conservation of the LBD shown, we were able
 317 to show that TaRXR binds 9-*cis*-retinoic acid with high affinity and specificity (**Fig. 3A**). A 9-*cis*-
 318 RA binding assay to determine the dissociation constant of TaRXR indicated high affinity binding
 319 with a saturation plateau from 5 nM to 10 nM. This was clearly observed in repeated experiments
 320 (**Fig. 3B**). In contrast, TaRXR does not show high affinity displaceable binding of all-*trans*-
 321 retinoic acid. GST-RXR and Thrombin-cleaved RXR showed similar 9-*cis*-RA binding capability
 322 (not shown).

323

324 **9-*cis*-retinoic acid induces malic enzyme gene expression at nanomolar**
 325 **concentrations**

Next, we searched whether 9-*cis*-RA has observable biological effects on *T. adharens* at nanomolar concentrations. We hypothesized that TaRXR is likely to be involved in the regulation of metabolic events. In vertebrates, RXR is a dimerization partner of TR and together, these two NRs are regulating a wide range of metabolic pathways. We therefore searched for an orthologue of vertebrate L-malate-NADP⁺ oxidoreductase (EC 1.1.1.40) in *T. adhaerens* genome since this enzyme is an established reporter of the state of thyroid hormone dependent regulation (see Discussion).

The sequence of the *T. adhaerens* likely orthologue of vertebrate L-malate-NADP⁺ oxidoreductase was retrieved from the *Trichoplax* genomic database together with its presumed promoter based on the predicted sequence (**Supplementary File S3**).

Droplet digital PCR showed an increased transcription of the malic enzyme gene after incubation of *T. adhaerens* with 9-*cis*-RA, but not with all-*trans*-RA (**Fig 4**). In repeated experiments, we observed that the level of induction was higher at 9-*cis*-RA concentrations in the range of 1 to 10 nM, than above 10 nM. We also noticed that the level of the induction slightly varied based on the actual *T. adhaerens* cultures and the algal food composition of the *T. adhaerens* cultures.

Changes in the culture environment alter the expression pattern of the nuclear receptor complement in *T. adhaerens*

From the experience we gained by culturing *T. adhaerens*, as well as from the previous experiments we knew that the culture conditions can dramatically influence phenotype. Having

the possible developmental functions of the ancestral NRs in mind, we raised the question whether the expression patterns of the NRs reflect changes in phenotype.

Firstly, we assayed the relative expression of RXR against all three other NRs in small versus big animals. The relative proportion of the RXR expression compared to the remaining NRs was found to be higher in big animals (33%) than in small animals (24%). The treatment by 3.3 nM 9-*cis*-RA led to a dramatic increase of the relative expression of RXR in comparison to the rest of the NR complement (51%), indicating that phenotypic changes are connected with differential expression of NRs and that 9-*cis*-RA affects the expression of RXR.

In order to see the effect of 9-*cis*-RA on all NRs, we sampled and extracted RNA from cultures containing the same number of big and small animals treated with different concentrations of 9-*cis*-RA. The experimental cultures were started from the same original cultures and during incubation were fed with *Chlorella* sp. Only, since this algal food showed to have the least effect on *T. adhaerens* cultures. All four *T. adhaerens* NRs were quantified by either qRT-PCR or ddPCR.

Analysis of NR expression pattern in animals incubated with different concentrations of 9-*cis*-RA, revealed a relative increase in RXR expression at low nanomolar concentrations (<10 nM) in repeated experiments. In contrast, further increase of 9-*cis*-RA resulted in smaller changes compared to the expression pattern of NRs in control animals or even reverted the values observed in low nanomolar conditions (**Fig 5**).

Food composition dramatically changes the phenotype and the reproduction rate of *T. adhaerens*

T. adhaerens retrieved from laboratory aquariums used for the stock cultures were relatively similar in appearance and included small round animals containing approximately 50 cells and grew to animals with an approximate diameter of 0.2 mm and rarely were bigger. Their rate of multiplication when transferred to Petri dishes was doubling in one month or even one week, depending on whether the glass was covered by microbial and algal films established during culturing in aquariums. We attempted to use several defined algae as artificial food. They included *Pyrrenomonas helgolandii*, *Picocystis salinarium*, *Tetraselmis subcoriformis*, *Rhodomonas salina*, *Phaeodactylum tricornutum*, *Porphyridium cruentum* and *Chlorella sp.* Individual subcultures of *T. adhaerens* differed in the rate of propagation and appearance as well as colors that were varying from greenish to brown and reddish taints depending on the food that was used as singular species food or mixtures (**Fig 6**). Also, contaminants from the original algal food, which prevailed in some cultures, had an influence on *T. adhaerens* growth and behavior. In controlled experiments, it became clear that some food components or food compositions are ~~more affecting~~ the growth and appearance of *T. adhaerens* than food availability. When *T. adhaerens* were fed with equal amounts of algal cells (although they differed in size and expected digestibility), the addition of algae containing red pigments - Cryptophytes (*Pyrrenomonas helgolandii* and *Rhodomonas salina*) or Rhodophyta (*Porphyridium cruentum*) - had a strong positive effect on *T. adhaerens* growth, especially in combination with the green algae *Chlorella sp.* (**Fig 7**).

Furthermore, the addition of *Porphyridium cruentum* to *Chlorella sp.* resulted in significant change in circularity, while feeding *T. adhaerens* with triple food containing *Chlorella*, *Rhodomonas* and *Porphyridium* showed the most pronounced effect. Culturing *T. adhaerens* on either single food showed similar isoperimetric values (**Fig 8**).

DISCUSSION

T. adhaerens is probably the closest living species to basal metazoans with only four NRs

Trichoplax adhaerens is an especially interesting species from an evolutionary perspective. It shows the most primitive metazoan planar body arrangement with a simple dorsal-ventral polarity, the establishment of which is one of the most ancient events in the evolution of animal symmetries (Smith et al. 1995; Stein & Stevens 2014). The Placozoa dispose with only a few (probably six) morphologically recognizable cell types (Jakob et al. 2004; Smith et al. 2014).

In strong contrast to this, the *T. adhaerens* genome shows larger blocks of conserved synteny relative to the human genome than flies or nematodes (Srivastava et al. 2008). Genome analyses indicate that Placozoa are basal relative to Bilateria as well as all other diploblast phyla (Schierwater et al. 2009), but all kinds of different views are also discussed (reviewed in (Schierwater et al. 2016)).

In concordance with this, its genome contains four (Srivastava et al. 2008) rather than 17 NRs, which can be found in the cnidarian *Nematostella vectensis* (Reitzel & Tarrant 2009). Even though it has been proposed that Placozoa lost representatives of NR6 (SF1/GCNF), TR2/TR4 of the NR2 subfamily and invertebrate specific nuclear receptors (INR, clade of invertebrate-only nuclear receptors with no standard nomenclature) NR1/NR4 (Bridgham et al. 2010), reasoning in this direction depends on the assumed phylogenic position of the phylum Placozoa.

The four NRs found in the genome of *T. adhaerens* are relatively closely related to their vertebrate counterparts, RXR (NR2B), HNF4 (NR2A), COUP (NR2F) and ERR (NR3) (Srivastava et al. 2008). Among them, *T. adhaerens* RXR and HNF4 show the highest degree of identity in protein sequence and the relatedness of *T. adhaerens* RXR (TaRXR) to human RXR is similar to that of *Tripedalia cystophora* RXR (jRXR) (Kostrouch et al. 1998), which has also been shown to bind 9cis-RA at nanomolar concentrations. These results suggest that TaRXR is structurally and also functionally very closely related to its vertebrate counterparts, most probably representing the most basal liganded NR of all Metazoa known today.



***T. adhaerens* RXR binds preferentially 9-cis-RA**

By using a radioactively labelled ligand, we could demonstrate that the RXR homologue in *T. adhaerens* binds 9-cis-RA with very high affinity and shows a strong binding preference to 9-cis-RA over all-trans-RA similarly, as is the case in vertebrate RXRs (Allenby et al. 1993) and the cnidarian TcRXR (Kostrouch et al. 1998).

9-cis-RA affects the expression of a *T. adhaerens* homologue of a conserved metabolically active enzyme, L-malate-NADP⁺ oxidoreductase

A biological role of 9-cis-RA binding with high affinity to the placozoan RXR receptor is supported by our *in vivo* experiments. In search for genes that may be under the regulation of

TaRXR, we identified a putative homologue of vertebrate L-malate-NADP⁺ oxidoreductase (EC 1.1.1.40) and analyzed the effect of 9-*cis*-RA or all-*trans*-RA on its expression. In agreement with our binding experiments, we observed induction of this enzyme's expression at low nanomolar concentrations of 9-*cis*-RA (1 to 10 nM). Interestingly, higher concentrations of 9-*cis*-RA (30 nM) had a smaller effect on expression and all-*trans*-RA had no effect up to 30 nM concentrations. A plausible explanation for this could be that 9-*cis*-RA also acts as a ligand for other *T. adhaerens* NRs which may have an opposite effect on the expression of L-malate-NADP⁺ oxidoreductase. Furthermore, 9-*cis*-RA could act as a 'supranatural' ligand and the continuous occupation of TaRXR by this high affinity ligand may interfere with the normal function of the receptor which has been demonstrated in the case of the glucocorticoid receptor (Schaaf & Cidlowski 2003; George et al. 2009).

In mammals regulation of malic enzyme expression is mediated by a thyroid hormone receptor (TR) – RXR heterodimer (Dozin et al. 1985a; Dozin et al. 1985b; Petty et al. 1989; Petty et al. 1990). By showing a 9-*cis*-RA dependent change in expression of the likely placozoan malic enzyme homologue *in vivo*, we provide indirect evidence of a conserved RXR mediated regulation of gene expression. Although the expression of L-malate-NADP⁺ oxidoreductase in mammals is usually used as a factor reflecting regulation by thyroid hormone (Dozin et al. 1986), it has also been shown that its cell type associated differences depend on the expression level of RXR alpha (Hillgartner et al. 1992; Fang & Hillgartner 2000) suggesting that regulation by RXR has been conserved throughout metazoan evolution while additional regulation via thyroid hormone represents an innovation of Bilateria (Wu et al. 2007).

NRs form a network responding to 9-*cis*-RA

Since autoregulation and cross-regulation of NRs by their specific ligands is well documented for a large number of nuclear receptors (Tata 1994), we searched if 9-*cis*-RA affects the expression of TaRXR mRNA relative to the other *T. adhaerens* NRs. Our *in vivo* experiments showed not only effects on specific gene expression in response to very low concentrations of 9-*cis*-RA (at 1 or 3 nM), but also an additional dose-dependent reverse effect of higher concentrations. This is likely to be in line with our binding experiments that suggested the possibility of an additional binding site or sites with higher capacity and lower affinity. We also cannot rule out that higher concentrations of 9-*cis*-RA affect some of the three remaining *T. adhaerens* NRs. Nevertheless, an inhibitory effect of 9-*cis*-RA on the expression of its cognate receptor at the protein level (through protein degradation) was reported (Nomura et al. 1999).

Although it is not clear if 9-*cis*-RA is the natural ligand for RXRs (Wolf 2006; Ruhl et al. 2015) conserved in all metazoan phyla studied to date, we show not only that 9-*cis*-RA binds TaRXR with nanomolar affinity but also positively regulates its expression, which resembles auto-activation of several NRs in vertebrates [e.g. ER and TR (Tata 1994; Bagamasbad & Denver 2011)]. Furthermore, three out of four NRs constituting the NR complement in *T. adhaerens* respond to treatment by 9-*cis*-RA at transcriptional level. Two NRs, RXR itself and ERR respond positively to nanomolar concentrations of 9-*cis*-RA, while COUP, which often acts as an inhibitor of specific gene expression (Tran et al. 1992), is regulated negatively by 9-*cis*-RA. COUP was recently shown to be inactivated by small hydrophobic molecules (Le Guevel et al. 2017). The regulatory connections of *T. adhaerens* NRs places the auto-regulation and cross-regulation of NRs to the base of metazoan evolution.

Food composition rather than quantity affects phenotype of *T. adhaerens*

At first glance, *T. adhaerens* seems to benefit from any source of biological material on surfaces that can be digested and absorbed by its digestive system (e.g. aquarium microorganisms and detritus). Feeding with certain live microorganisms in laboratory cultures, however, dramatically changes the dynamics of *T. adhaerens* cultures, such as shape, size, color, body transparency, growth and divisions of the animal. For example, we observed poor growth and reproduction rates of *T. adhaerens* fed solely on *Chlorella sp.* even at a relatively high density. In contrast, cultures fed with red pigment containing *Rhodomonas salina* showed much faster proliferation and led, in part, to the formation of giant animals, seeming to halt their division. Despite *Porphyridium cruentum* containing similar pigments as *Rhodomonas*, such as phycoerythrin, cultures grown with *Porphyridium* as the main nutrient source did not show phenotypical abnormalities but addition of it to a culture with *Chlorella* and *Rhodomonas* resulted in an additive effect on reproduction rate.

Even though growth of *T. adhaerens* seems to follow a simple program, it is likely to require strict regulatory mechanisms. Formation of specific cellular types is connected with phenotypic appearance of animals possessing larger proportions of certain cells, e.g. upper epithelium in balloon like animals or larger proportion of peripheral regions containing stem-cell like cells in narrow or prolonged animals. Analysis of circularity as a measure of location specific cellular proliferation is in concordance with the observed culture characteristics and shows that lower isoperimetric values (less ‘roundness’) indicate higher reproduction rates.

Our experiments provide evidence that food composition is more important for *T. adhaerens* growth and propagation than its quantity, which is in line with the recent finding of phosphate and nitrate playing important roles determining distribution of placozoans around the globe (Paknia & Schierwater 2015). It indicates that food constituents, especially those present in

the algae containing phycobilin based red pigments like *Rhodomonas salina* and *Porphyridium cruentum* might possess hormone-like molecules or molecules resulting in hormone-like metabolites in *T. adhaerens* that act through the NR complement and, indeed, analysis of NRs in differently sized animals indicates impact of food composition on NR expression.

When viewed together, our work shows that the presence of 9-*cis*-RA binding RXR in Placozoa argues for the existence of ligand regulated NRs at the base of metazoan evolution. This work also suggests that an endocrine-like network was present in early metazoans, likely to be an ancestral regulatory network linking nutritional or environmental factors with the regulation of gene expression that is mediated by NRs. This also suggests that the regulation by liganded NRs evolved as an evolutionary need connected with heterotrophy and multicellularity.

In fact, despite fragments of NR domains being found in prokaryotes, no single full sized NR has been discovered in bacteria or archaea and the closest known relatives to metazoans, unicellular and colonial Choanoflagellates, lack nuclear receptors, as well as genes of several other regulatory pathways (King et al. 2008). On the other hand in fungi, the sister group of Holozoa, (Shalchian-Tabrizi et al. 2008) transcription factors surprisingly similar to metazoan NRs evolved independently possibly for the regulation of metabolism and response to xenobiotics (Thakur et al. 2008; Naar & Thakur 2009). Thus, the evolution of NRs seems to be associated with two key evolutionary features of metazoans: multicellularity and heterotrophy.

Ctenophores, a possible sister phylum to *Cnidaria* do not contain classical NRs featuring both mechanistically critical domains of NRs, the DNA binding and ligand binding domains. Nevertheless, the ctenophore *Mnemiopsis* contains two homologues of NR2A (HNF4) that lack DNA binding domain (Reitzel et al. 2011).

In conclusion, the *T. adhaerens* endocrine-like network supports the hypothesis of a basic regulatory mechanism by NRs, which may have been subspecialized with the appearance of new NRs in order to cope with new environmental and behavioral challenges during the course of early metazoan evolution.

Acknowledgements

The authors thank NCBI for accessibility of data and bioinformatics support and Joint Genome Institute of United States Department of Energy (<http://jgi.doe.gov/>) (Nordberg et al. 2014).

REFERENCES

- Allenby G, Bocquel MT, Saunders M, Kazmer S, Speck J, Rosenberger M, Lovey A, Kastner P, Grippo JF, Chambon P, and et al. 1993. Retinoic acid receptors and retinoid X receptors: interactions with endogenous retinoic acids. *Proc Natl Acad Sci U S A* 90:30-34.
- Bagamasbad P, and Denver RJ. 2011. Mechanisms and significance of nuclear receptor auto- and cross-regulation. *Gen Comp Endocrinol* 170:3-17. 10.1016/j.ygcen.2010.03.013
- Bell EA, Boehnke P, Harrison TM, and Mao WL. 2015. Potentially biogenic carbon preserved in a 4.1 billion-year-old zircon. *Proc Natl Acad Sci U S A* 112:14518-14521. 10.1073/pnas.1517557112
- Bridgham JT, Eick GN, Larroux C, Deshpande K, Harms MJ, Gauthier ME, Ortlund EA, Degnan BM, and Thornton JW. 2010. Protein evolution by molecular tinkering: diversification

of the nuclear receptor superfamily from a ligand-dependent ancestor. *PLoS Biol* 8.
10.1371/journal.pbio.1000497

Cahnmann HJ. 1995. A fast photoisomerization method for the preparation of tritium-labeled
9-cis-retinoic acid of high specific activity. *Anal Biochem* 227:49-53.
10.1006/abio.1995.1251

Dozin B, Cahnmann HJ, and Nikodem VM. 1985a. Identification of thyroid hormone receptors
in rat liver nuclei by photoaffinity labeling with L-thyroxine and triiodo-L-thyronine.
Biochemistry 24:5197-5202.

Dozin B, Magnuson MA, and Nikodem VM. 1985b. Tissue-specific regulation of two functional
malic enzyme mRNAs by triiodothyronine. *Biochemistry* 24:5581-5586.

Dozin B, Magnuson MA, and Nikodem VM. 1986. Thyroid hormone regulation of malic
enzyme synthesis. Dual tissue-specific control. *J Biol Chem* 261:10290-10292.

Egea PF, Klaholz BP, and Moras D. 2000. Ligand-protein interactions in nuclear receptors of
hormones. *FEBS letters* 476:62-67.

Escriva H, Bertrand S, and Laudet V. 2004. The evolution of the nuclear receptor superfamily.
Essays Biochem 40:11-26.

Escriva H, Delaunay F, and Laudet V. 2000. Ligand binding and nuclear receptor evolution.
Bioessays 22:717-727. 10.1002/1521-1878(200008)22:8<717::AID-BIES5>3.0.CO;2-
I

Escriva H, Langlois MC, Mendonca RL, Pierce R, and Laudet V. 1998. Evolution and
diversification of the nuclear receptor superfamily. *Ann N Y Acad Sci* 839:143-146.

Fang X, and Hillgartner FB. 2000. Alterations in retinoid X receptor-alpha expression
contribute to cell-type dependent differences in thyroid hormone regulation of malic
enzyme transcription. *Mol Cell Endocrinol* 164:41-52.

Forman BM, and Samuels HH. 1990a. Dimerization among nuclear hormone receptors. *New
Biol* 2:587-594.

Forman BM, and Samuels HH. 1990b. Interactions among a subfamily of nuclear hormone
receptors: the regulatory zipper model. *Mol Endocrinol* 4:1293-1301. 10.1210/mend-
4-9-1293

George AA, Schiltz RL, and Hager GL. 2009. Dynamic access of the glucocorticoid receptor to
response elements in chromatin. *Int J Biochem Cell Biol* 41:214-224.
10.1016/j.biocel.2008.09.019

Guindon S, Dufayard JF, Lefort V, Anisimova M, Hordijk W, and Gascuel O. 2010. New
algorithms and methods to estimate maximum-likelihood phylogenies: assessing the
performance of PhyML 3.0. *Syst Biol* 59:307-321. 10.1093/sysbio/syq010

Gutierrez-Mazariegos J, Schubert M, and Laudet V. 2014. Evolution of retinoic acid receptors
and retinoic acid signaling. *Subcell Biochem* 70:55-73. 10.1007/978-94-017-9050-
5_4

Hillgartner FB, Chen W, and Goodridge AG. 1992. Overexpression of the alpha-thyroid
hormone receptor in avian cell lines. Effects on expression of the malic enzyme gene
are selective and cell-specific. *J Biol Chem* 267:12299-12306.

Jakob W, Sagasser S, Dellaporta S, Holland P, Kuhn K, and Schierwater B. 2004. The Trox-2
Hox/ParaHox gene of Trichoplax (Placozoa) marks an epithelial boundary. *Dev Genes
Evol* 214:170-175.

601 Kana R, Kotabova E, Lukes M, Papacek S, Matonoha C, Liu LN, Prasil O, and Mullineaux CW.
602 2014. Phycobilisome Mobility and Its Role in the Regulation of Light Harvesting in
603 Red Algae. *Plant Physiol* 165:1618-1631. 10.1104/pp.114.236075

604 Kana R, Kotabova E, Sobotka R, and Prasil O. 2012. Non-photochemical quenching in
605 cryptophyte alga *Rhodomonas salina* is located in chlorophyll a/c antennae. *PLoS One*
606 7:e29700.

607 Kiefer JC. 2006. Emerging developmental model systems. *Dev Dyn* 235:2895-2899.

608 King N, Westbrook MJ, Young SL, Kuo A, Abedin M, Chapman J, Fairclough S, Hellsten U, Isogai
609 Y, Letunic I, Marr M, Pincus D, Putnam N, Rokas A, Wright KJ, Zuzow R, Dirks W, Good
610 M, Goodstein D, Lemons D, Li W, Lyons JB, Morris A, Nichols S, Richter DJ, Salamov A,
611 Sequencing JG, Bork P, Lim WA, Manning G, Miller WT, McGinnis W, Shapiro H, Tjian
612 R, Grigoriev IV, and Rokhsar D. 2008. The genome of the choanoflagellate *Monosiga*
613 *brevicollis* and the origin of metazoans. *Nature* 451:783-788. 10.1038/nature06617

614 Kostrouch Z, Kostrouchova M, Love W, Jannini E, Piatigorsky J, and Rall JE. 1998. Retinoic
615 acid X receptor in the diploblast, *Tripedalia cystophora*. *Proc Natl Acad Sci U S A*
616 95:13442-13447.

617 Kostrouch Z, Kostrouchova M, and Rall JE. 1995. Steroid/thyroid hormone receptor genes in
618 *Caenorhabditis elegans*. *Proc Natl Acad Sci U S A* 92:156-159.

619 Kostrouchova M, and Kostrouch Z. 2015. Nuclear receptors in nematode development:
620 Natural experiments made by a phylum. *Biochim Biophys Acta* 1849:224-237.
621 10.1016/j.bbagr.2014.06.016

622 Kumar R, and Thompson EB. 1999. The structure of the nuclear hormone receptors. *Steroids*
623 64:310-319.

624 Laudet V. 1997. Evolution of the nuclear receptor superfamily: early diversification from an
625 ancestral orphan receptor. *J Mol Endocrinol* 19:207-226.

626 Le Guevel R, Oger F, Martinez-Jimenez CP, Bizot M, Gheeraert C, Firmin F, Ploton M, Kretova
627 M, Palierne G, Staels B, Barath P, Talianidis I, Lefebvre P, Eeckhoutte J, and Salbert G.
628 2017. Inactivation of the Nuclear Orphan Receptor COUP-TFII by Small Chemicals.
629 *ACS Chem Biol*. 10.1021/acscchembio.6b00593

630 Letunic I, Doerks T, and Bork P. 2015. SMART: recent updates, new developments and status
631 in 2015. *Nucleic Acids Res* 43:D257-260. 10.1093/nar/gku949

632 Markov GV, and Laudet V. 2011. Origin and evolution of the ligand-binding ability of nuclear
633 receptors. *Mol Cell Endocrinol* 334:21-30. 10.1016/j.mce.2010.10.017

634 Naar AM, and Thakur JK. 2009. Nuclear receptor-like transcription factors in fungi. *Genes Dev*
635 23:419-432. 10.1101/gad.1743009

636 Nomura Y, Nagaya T, Hayashi Y, Kambe F, and Seo H. 1999. 9-cis-retinoic acid decreases the
637 level of its cognate receptor, retinoid X receptor, through acceleration of the turnover.
638 *Biochem Biophys Res Commun* 260:729-733. 10.1006/bbrc.1999.0969

639 Nordberg H, Cantor M, Dusheyko S, Hua S, Poliakov A, Shabalov I, Smirnova T, Grigoriev IV,
640 and Dubchak I. 2014. The genome portal of the Department of Energy Joint Genome
641 Institute: 2014 updates. *Nucleic Acids Res* 42:D26-31. 10.1093/nar/gkt1069

642 Nowickij SM, Chithalen JV, Cameron D, Tyshenko MG, Petkovich M, Wyatt GR, Jones G, and
643 Walker VK. 2008. Locust retinoid X receptors: 9-Cis-retinoic acid in embryos from a
644 primitive insect. *Proc Natl Acad Sci U S A* 105:9540-9545. 10.1073/pnas.0712132105

- Paknia O, and Schierwater B. 2015. Global Habitat Suitability and Ecological Niche Separation in the Phylum Placozoa. *PLoS One* 10:e0140162. 10.1371/journal.pone.0140162
- Petty KJ, Desvergne B, Mitsushashi T, and Nikodem VM. 1990. Identification of a thyroid hormone response element in the malic enzyme gene. *J Biol Chem* 265:7395-7400.
- Petty KJ, Morioka H, Mitsushashi T, and Nikodem VM. 1989. Thyroid hormone regulation of transcription factors involved in malic enzyme gene expression. *J Biol Chem* 264:11483-11490.
- Reitzel AM, Pang K, Ryan JF, Mullikin JC, Martindale MQ, Baxevanis AD, and Tarrant AM. 2011. Nuclear receptors from the ctenophore Mnemiopsis leidyi lack a zinc-finger DNA-binding domain: lineage-specific loss or ancestral condition in the emergence of the nuclear receptor superfamily? *Evodevo* 2:3. 10.1186/2041-9139-2-3
- Reitzel AM, and Tarrant AM. 2009. Nuclear receptor complement of the cnidarian Nematostella vectensis: phylogenetic relationships and developmental expression patterns. *BMC Evol Biol* 9:230. 10.1186/1471-2148-9-230
- Robinson-Rechavi M, Escriva Garcia H, and Laudet V. 2003. The nuclear receptor superfamily. *J Cell Sci* 116:585-586.
- Ruhl R, Krzyzosiak A, Niewiadomska-Cimicka A, Rochel N, Szeles L, Vaz B, Wietrzyk-Schindler M, Alvarez S, Szklenar M, Nagy L, de Lera AR, and Krezel W. 2015. 9-cis-13,14-Dihydroretinoic Acid Is an Endogenous Retinoid Acting as RXR Ligand in Mice. *PLoS Genet* 11:e1005213. 10.1371/journal.pgen.1005213
- Schaaf MJ, and Cidlowski JA. 2003. Molecular determinants of glucocorticoid receptor mobility in living cells: the importance of ligand affinity. *Mol Cell Biol* 23:1922-1934.
- Schierwater, Holland PWH, Miller DJ, Stadler PF, Wiegmann BM, Worheide G, Wray GA, and R D. 2016. Never Ending Analysis of a Century Old Evolutionary Debate: "Unringing" the Urmetazoon Bell. *Front Ecol Evol*, 4:5. 10.3389/fevo.2016.00005
- Schierwater B, Kolokotronis SO, Eitel M, and DeSalle R. 2009. The Diploblast-Bilateria Sister hypothesis: parallel revolution of a nervous systems may have been a simple step. *Commun Integr Biol* 2:403-405.
- Schultz J, Milpetz F, Bork P, and Ponting CP. 1998. SMART, a simple modular architecture research tool: identification of signaling domains. *Proc Natl Acad Sci U S A* 95:5857-5864.
- Shalchian-Tabrizi K, Minge MA, Espelund M, Orr R, Ruden T, Jakobsen KS, and Cavalier-Smith T. 2008. Multigene phylogeny of choanozoa and the origin of animals. *PLoS One* 3:e2098. 10.1371/journal.pone.0002098
- Sievers F, Wilm A, Dineen D, Gibson TJ, Karplus K, Li W, Lopez R, McWilliam H, Remmert M, Soding J, Thompson JD, and Higgins DG. 2011. Fast, scalable generation of high-quality protein multiple sequence alignments using Clustal Omega. *Mol Syst Biol* 7:539. 10.1038/msb.2011.75
- Smith CL, Giordano H, Schwartz M, and DeLotto R. 1995. Spatial regulation of Drosophila snake protease activity in the generation of dorsal-ventral polarity. *Development* 121:4127-4135.
- Smith CL, Varoqueaux F, Kittelmann M, Azzam RN, Cooper B, Winters CA, Eitel M, Fasshauer D, and Reese TS. 2014. Novel cell types, neurosecretory cells, and body plan of the early-diverging metazoan Trichoplax adhaerens. *Curr Biol* 24:1565-1572. 10.1016/j.cub.2014.05.046

691 Srivastava M, Begovic E, Chapman J, Putnam NH, Hellsten U, Kawashima T, Kuo A, Mitros T,
692 Salamov A, Carpenter ML, Signorovitch AY, Moreno MA, Kamm K, Grimwood J,
693 Schmutz J, Shapiro H, Grigoriev IV, Buss LW, Schierwater B, Dellaporta SL, and
694 Rokhsar DS. 2008. The Trichoplax genome and the nature of placozoans. *Nature*
695 454:955-960.

696 Stein DS, and Stevens LM. 2014. Maternal control of the Drosophila dorsal-ventral body axis.
697 *Wiley Interdiscip Rev Dev Biol* 3:301-330. 10.1002/wdev.138

698 Tata JR. 1994. Autoregulation and crossregulation of nuclear receptor genes. *Trends*
699 *Endocrinol Metab* 5:283-290.

700 Thakur JK, Arthanari H, Yang F, Pan SJ, Fan X, Breger J, Frueh DP, Gulshan K, Li DK, Mylonakis
701 E, Struhl K, Moye-Rowley WS, Cormack BP, Wagner G, and Naar AM. 2008. A nuclear
702 receptor-like pathway regulating multidrug resistance in fungi. *Nature* 452:604-609.
703 10.1038/nature06836

704 Tran P, Zhang XK, Salbert G, Hermann T, Lehmann JM, and Pfahl M. 1992. COUP orphan
705 receptors are negative regulators of retinoic acid response pathways. *Mol Cell Biol*
706 12:4666-4676.

707 Wolf G. 2006. Is 9-cis-retinoic acid the endogenous ligand for the retinoic acid-X receptor?
708 *Nutr Rev* 64:532-538.

709 Wu W, Niles EG, and LoVerde PT. 2007. Thyroid hormone receptor orthologues from
710 invertebrate species with emphasis on Schistosoma mansoni. *BMC Evol Biol* 7:150.
711 10.1186/1471-2148-7-150
712

Figure 1(on next page)

Multiple sequence alignment of selected metazoan homologues of RXR compared with TaRXR.

Aligned with ClustalO, amino acid residue types colored according to Clustal scheme in Jalview, * indicates DBD footprint residues, # LBD footprint residues. Black box shows the DBD, red box represents the LBD. Sequences from top to bottom (organism, identifier): *Trichoplax adhaerens*, TaRXR ID 53515; *Homo sapiens*, sp|P19793|RXRA_HUMAN; *Homo sapiens*, sp|P28702|RXRB_HUMAN; *Homo sapiens*, sp|P48443|RXRG_HUMAN; *Xenopus laevis*, RXR alpha, sp|P51128|RXRA_XENLA; *Aurelia aurita*, RXR, tr|S5ZWR0|S5ZWR0_AURA Retinoid X receptor; *Tripedalia cystophora*, RXR, tr|O96562|O96562_TRICY Retinoic acid X receptor; *Drosophila melanogaster*, USP, sp|P20153|USP_DROME. Readers with specific color preferences may download the compared sequences (**Supplementary File S1**) and create the Clustal scheme with different color specifications using the Jalview program (<http://www.jalview.org/>).

PeerJ

Manuscript to be reviewed

17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848

[illegible][illegible][illegible][illegible][illegible]

AXRXR	207	--	D	G	L	L	S	G	H	Y	I	T	T	S	A	N	A	G	V	G	A	F	O	R	I	L	T	E	L	V	N	M	Y	K	M	D	E	L	G	C	L	A	I	V	L	F	N	P	D	V	S	G	O	L	S	A	D	R	V	K	E	L	V	G	283	
RXRA_HUMAN	322	--	D	G	L	L	S	G	H	Y	I	T	T	S	A	N	A	G	V	G	A	F	O	R	I	L	T	E	L	V	N	M	Y	K	M	D	E	L	G	C	L	A	I	V	L	F	N	P	D	V	S	G	O	L	S	A	D	R	V	K	E	L	V	G	398	
RXRB_HUMAN	393	--	D	G	L	L	S	G	H	Y	I	T	T	S	A	N	A	G	V	G	A	F	O	R	I	L	T	E	L	V	N	M	Y	K	M	D	E	L	G	C	L	A	I	V	L	F	N	P	D	V	S	G	O	L	S	A	D	R	V	K	E	L	V	G	469	
RXRG_HUMAN	323	--	D	G	L	L	S	G	H	Y	I	T	T	S	A	N	A	G	V	G	A	F	O	R	I	L	T	E	L	V	N	M	Y	K	M	D	E	L	G	C	L	A	I	V	L	F	N	P	D	V	S	G	O	L	S	A	D	R	V	K	E	L	V	G	399	
RXRA_XENLA	348	--	D	G	L	L	S	G	H	Y	I	T	T	S	A	N	A	G	V	G	A	F	O	R	I	L	T	E	L	V	N	M	Y	K	M	D	E	L	G	C	L	A	I	V	L	F	N	P	D	V	S	G	O	L	S	A	D	R	V	K	E	L	V	G	424	
SSZWRO_AURAU	288	--	D	G	L	L	S	G	H	Y	I	T	T	S	A	N	A	G	V	G	A	F	O	R	I	L	T	E	L	V	N	M	Y	K	M	D	E	L	G	C	L	A	I	V	L	F	N	P	D	V	S	G	O	L	S	A	D	R	V	K	E	L	V	G	364	
O96562_TRICY	297	--	D	G	L	L	S	G	H	Y	I	T	T	S	A	N	A	G	V	G	A	F	O	R	I	L	T	E	L	V	N	M	Y	K	M	D	E	L	G	C	L	A	I	V	L	F	N	P	D	V	S	G	O	L	S	A	D	R	V	K	E	L	V	G	373	
USP_DROME	358	L	P	Q	E	L	N	S	S	E	S	Y	N	S	A	I	K	A	G	V	S	A	F	O	R	I	L	T	E	L	V	N	M	Y	K	M	D	E	L	G	C	L	A	I	V	L	F	N	P	D	V	S	G	O	L	S	A	D	R	V	K	E	L	V	G	445

[illegible]

Figure 2

Phylogenetic tree of sequences shown in Fig. 1.

Phylogenetic tree constructed with PhyML algorithm with a 100 bootstrap and SPR distance computation visualized by FigTree of selected metazoan RXR sequences shown in Fig. 1 (and listed in **Supplementary File S1**). Bootstrap values are shown next to nodes.




Figure 3

Binding of retinoic acids to TaRXR.

A – Single point analysis of binding preference of *T. adhaerens* RXR to 9-*cis*-RA over all-*trans*-RA. Radioactive 9-*cis*-RA binds at a concentration of 4 nM to 200 micrograms of *T. adhaerens* RXR (dark column). 200x excess of unlabeled 9-*cis*-RA displaces more than 80 % labeled 9-*cis*-RA from binding to *T. adhaerens* RXR (second column) while the same molar excess of all-*trans*-RA (marked as AT-RA) (which is likely to contain approximately 1% spontaneously isomerized 9-*cis*-RA) competes away less than 50 % of bound labeled 9-*cis*-RA (column 3 from left). Radioactive all-*trans*-RA at identical conditions binds only slightly more than observed the non-specific binding with 9-*cis*-RA (column 4 from left) and this interaction is not displaced by the excess of non-labeled 9-*cis*-RA nor non-labeled all-*trans*-RA. B – Kinetic analysis of binding of 9-*cis*-RA to *T. adhaerens* RXR. The plateau is reached at around 3 to 5 $\times 10^{-9}$ M.




Figure 4

The effect of 9-*cis*-RA on the expression of the *T. adhaerens* closest putative homologue and likely orthologue of L-malate-NADP+ oxidoreductase (EC1.1.1.40).

Ten to fifteen animals were cultured in the dark overnight with indicated ligands or in medium containing only the solvent used for ligand solutions. Total RNA and cDNA were prepared using identical conditions and diluted to working concentrations suitable for ddPCR. In repeated experiments, incubation with 3 nM 9-*cis*-RA induced expression of the putative *T. adhaerens* L-malate-NADP+ oxidoreductase more than 4 times. Incubation with 30 nM 9-*cis*-RA induced enzyme expression also, but to a lesser extent and 30 nM all-*trans*-RA (AT-RA) did not upregulate the expression of the predicted L-malate-NADP+ oxidoreductase.




Figure 5

The effect of 9-*cis*-RA on the expression of *T. adhaerens* NRs.

A representative experiment of the expression of *T. adhaerens* NRs in animals exposed to various concentrations of 9-*cis*-RA expressed as change of the ratio compared to the control. One and 3 nM 9-*cis*-RA upregulate RXR and ERR, while downregulating COUP. The expression of *T. adhaerens* HNF4 is not changed. Similarly, as in the case of the putative L-malate-NADP+ oxidoreductase homologue shown in **Fig 4**, the effect is stronger in the case of 1 nM 9-*cis*-RA compared to 3 nM, 30 nM, as well as 50 nM concentrations reverse the effect on RXR and COUP, but do not influence the expression of ERR. The data suggest that a network sensitive to nanomolar concentrations of 9-*cis*-RA at an expressional level is formed by RXR, COUP, HNF4 and ERR.




Figure 6

Phenotypes of *T. adhaerens* change at various feeding conditions.

T. adhaerens acquires various body shapes in individual cultures dependent on food availability and composition. At conditions maintained in stable and biologically equilibrated stock aquariums, *T. adhaerens* is usually small and pale with diameter varying from 50 μm to 400 μm while cultures with added algae contain large flat animals with diameter reaching up to 1 mm (A and B). In some cultures, animals grow as long stretching structures, reaching a length exceeding one or even several centimeters (C). The algal food makes the animals greenish, reddish, rusty or brown with variable proportion of prominent dark cells. Animal shapes also vary from flat and round with smooth circumference, to curved or ruffled circumference or animals with long projections. Bars represent 1mm in A, B, I, 1cm in C, 250 μm in D, 500 μm in E and H, 200 μm in F, and 100 μm in G.




Figure 7 (on next page)

Propagation of *T. adhaerens* depends on algal food composition.

Three cultures of 5 large animals in each were established and fed with the same number of algal cells consisting of *Chlorella sp.*, *Chlorella sp.* and *Rhodomonas salina* and *Chlorella sp.*, *Rhodomonas salina* and *Porphyridium cruentum*. While the culture fed with *Chlorella sp.* only doubled in the number of animals within a period of one week, cultures with red pigment containing algae multiplied more than 5 times and 10 times within the same time period.




Figure 8

The effect of algal food composition on *T. adhaerens* growth and appearance.

T. adhaerens was cultured similarly as shown in Fig. 6 and all animals photographed and analyzed using ImageJ program for their number (panel A), mean area (panel B), mean perimeter (panel C), Perimeter/Area ratio (panel D) and isometric quotient (panel E) after one-week. Ch – stands for feeding with *Chlorella sp.*, Rh – *Rhodomonas salina*, Po – *Porphyridium cruentum*, and their combinations. 3Rh stands for a culture with three times higher concentration of *Rhodomonas salina* and 3Ch for three times higher concentration of *Chlorella sp.* Panel A shows that addition of *Rhodomonas salina* (Ch+Rh) greatly increases the number of animals observed after one week of culture. This effect is even more pronounced in cultures containing all three algae, while three times bigger concentration of only one type of algae (Ch and Rh) has little or no effect. This is even more pronounced when the area and perimeter are determined (panels B and C). Determination of the isoperimetric quotient in individual cultures indicates that cultures with *Rhodomonas* have a significantly smaller ratio, suggesting higher proliferative rate of structures at the animal circumference (panel E). Bars represent 95% confidence interval. Raw data are provided as Supplementary **File S5** and **S6**.

