

Anti-apoptotic properties of carbon monoxide in porcine oocyte during *in vitro* aging

David Němeček ^{Corresp., 1}, Markéta Dvořáková ¹, Ivona Heroutová ¹, Eva Chmelíková ¹, Markéta Sedmíková ¹

¹ Department of Veterinary Sciences, Czech University of Life Sciences, Prague, Czech Republic

Corresponding Author: David Němeček

Email address: nemecekd@af.czu.cz

If fertilization of matured oocyte does not occur, unfertilized oocyte undergoes aging, resulting in a time-dependent reduction of the oocyte's quality. The aging of porcine oocytes can lead to apoptosis. Carbon monoxide (CO), a signal molecule produced by the heme oxygenase (HO), possesses cytoprotective and anti-apoptotic effects that have been described in somatic cells. However, the effects of CO in oocytes have yet to be investigated. By immunocytochemistry method we detected that both isoforms of heme oxygenase (HO-1 and HO-2) are present in the porcine oocytes. Based on the morphological signs of oocyte aging, it was found that the inhibition of both HO isoforms by Zn-protoporphyrin IX (Zn-PP IX) leads to an increase in the number of apoptotic oocytes and decrease in the number of intact oocytes during aging. Contrarily, the presence of CO donors (CORM-2 or CORM-A1) significantly decrease the number of apoptotic oocytes while increasing the number of intact oocytes. We also determined that CO donors significantly decrease the caspase-3 (CAS-3) activity. Our results suggest that HO/CO contributes to the sustaining viability through regulation of apoptosis during *in vitro* aging of porcine oocytes

Anti-apoptotic Properties of Carbon Monoxide in Porcine Oocyte during *in Vitro* Aging

David Němeček¹, Markéta Dvořáková¹, Ivona Heroutová¹, Eva Chmelíková¹, Markéta Sedmíková¹

¹ Department of Veterinary Sciences, Czech University of Life Sciences, Prague, Czech Republic

Corresponding Author:

David Němeček

Kamýcká 129, Praha 6, 165 00, Czech Republic

Email address: nemecekd@af.czu.cz

Abstract

If fertilization of matured oocyte does not occur, unfertilized oocyte undergoes aging, resulting in a time-dependent reduction of the oocyte's quality. The aging of porcine oocytes can lead to apoptosis. Carbon monoxide (CO), a signal molecule produced by the heme oxygenase (HO), possesses cytoprotective and anti-apoptotic effects that have been described in somatic cells. However, the effects of CO in oocytes have yet to be investigated. By immunocytochemistry method we detected that both isoforms of heme oxygenase (HO-1 and HO-2) are present in the porcine oocytes. Based on the morphological signs of oocyte aging, it was found that the inhibition of both HO isoforms by Zn-protoporphyrin IX (Zn-PP IX) leads to an increase in the number of apoptotic oocytes and decrease in the number of intact oocytes during aging. Contrarily, the presence of CO donors (CORM-2 or CORM-A1) significantly decrease the number of apoptotic oocytes while increasing the number of intact oocytes. We also determined that CO donors significantly decrease the caspase-3 (CAS-3) activity. Our results suggest that HO/CO contributes to the sustaining viability through regulation of apoptosis during *in vitro* aging of porcine oocytes.

Introduction

In most mammals, a mature oocyte is in the stage of the metaphase of the second meiotic division, when it awaits fertilization. If the fertilization does not occur within the time referred to as a ‘temporal window for optimal fertilization’ (Fissore et al., 2002; Goud et al., 2005), a time-dependent decrease in oocyte quality takes place, which is also referred to as post-ovulatory oocyte aging (Fissore et al., 2002; Miao et al., 2009; Lord & Aitken, 2013). In pigs, as well as in other mammals, the aging process takes place in both *in vivo* and *in vitro* conditions (Petrová et al., 2004; Petrová et al., 2009). Oocyte aging is one of the factors limiting various assisted reproductive technologies (ART) outcome in several mammalian species (Miao et al., 2009). Many changes of cellular functions occur during oocyte ageing, as well as morphological changes of the cytoskeleton and cellular organelles. The negative effects of aging include premature exocytosis of cortical granules (Szollosi, 1971), structural changes of the *zona pellucida* (Xu et al., 1997), the decrease of the fertilizing capability (Lanman, 1968), increase of polyspermy (Badenas et al., 1989), parthenogenesis (Blandau, 1952) and chromosomal aberrations (Szollosi, 1971).

Several mechanisms contribute to the formation of negative effects of the oocyte aging. Aging process leads to the progressive increase in ROS production and the concomitant depletion of antioxidant protection and as a consequence the post ovulatory aged oocyte experiences a state of oxidative stress (Lord & Aitken, 2013). This relates to the disruption in functions of the mitochondria and Ca^{2+} signaling (Liu et al, 2000; Lord & Aitken, 2013). Changes in the activity of the M-phase promotion factor (MPF) and the mitogen-activated protein kinase (MAPK) that maintain the meiotic arrest in metaphase II also occurs during aging (Kikuchi et al., 1995; Miao et al., 2009; Jiang et al., 2011). The decrease of the MPF activity causes parthenogenetic activation in the aged oocytes and consequently cellular death. The increased MAPK activity also contributes to the triggering of cellular death (Sadler et al., 2004, Jeseta et al., 2008, Miao et al., 2009). Aging process finally leads to lytic or, more often, apoptotic cell death of aged oocytes (Fissore et al., 2002; Miao et al., 2009; Petrová et al., 2009; Lord & Aitken, 2013).

Programmed cell death is characterized by the activation of caspases (aspartate-

specific cysteine proteases) that are activated upon the receipt of either an extrinsic or intrinsic death signal. Both signals induce the execution phase of the apoptotic pathway characterized by the activation of executioner caspases that subsequently activate cytoplasmic endonucleases and proteases. Their activation leads to characteristic morphological and biochemical changes observed during apoptosis (Salvesen & Dixit, 1997; Slee, Adrai & Martin, 2001; Taylor, Cullen & Martin, 2008). Caspase-3 (CAS-3) is one of the most important executioner caspases. The CAS-3 activity is often used as a marker of apoptotic cell death, regardless of whether the apoptosis was triggered through an extrinsic or intrinsic pathway (Elmore, 2007). Similarly, as in somatic cells, also in aged oocytes is CAS-3 activated during apoptotic cell death (Zhu et al., 2015; Zhu et al., 2016).

Carbon monoxide (CO), endogenously produced by heme oxygenase (HO) or exogenously delivered by CO gas or CO-releasing molecules (CORMs) (Motterlini et al., 2003) is one of the known factors that can modulate apoptotic pathway in various types of somatic cells (Brouard et al., 2000; Petrache et al., 2000; Wu & Wang, 2005; Ryter, Alam & Choi, 2006; Kim et al., 2011), but the effect of CO in oocytes is unknown. HO enzyme catalyzes oxidative cleavage of heme producing ferrous iron, biliverdin-IX α and CO (Tenhunen et al., 1968; Tenhunen et al., 1969). HO exists in two active isoforms, HO-1 and HO-2. HO-1 is an inducible isoform activated by different kinds of stresses (e.g. oxidative stress) (Biswas et al., 2014; Ryter & Choi, 2016), while the constitutive isoform HO-2 is responsible for the HO basal activity (Turkseven et al., 2007; Muñoz-Sánchez & Cháñez-Cárdenas, 2014).

CO influences a variety of signalling pathways and generally has cytoprotective, anti-apoptotic and anti-inflammatory properties (Motterlini & Otterbein, 2010). In murine endothelial cells CO suppresses apoptosis through activation of the p38 MAPK (Brouard et al., 2000; 2002) and in rat endothelial cells CO prevents the initiation of apoptosis through increasing the expression of anti-apoptotic factor Bcl-2 (Zhang et al., 2003a) and decreasing the expression/activation of pro-apoptotic factors Bid and Bax (Zhang et al., 2003; Wang et al., 2007a). It further prevents the release of cytochrome c from mitochondrial matrix (Zhang et al., 2003a; Wang et al., 2007a). Overall, CO decreases caspases activation (CAS-3, CAS-8, CAS-9), which was proven in

endothelial cells of mice and rats (Zhang et al., 2003a; Zhang et al., 2003b; Wang et al., 2007a; Wang et al., 2011), in murine astrocytes (Almeida et al., 2012), rat ganglion cells (Schallner et al., 2012) and porcine lung tissue (Goebel et al., 2008).

The anti-apoptotic effect of the HO/CO system was investigated in somatic cells, but in the case of oocytes the effect of CO is so far unknown. While it was proven that HO-1 deficiency in mice causes lower fertilization capability of oocytes (Zenclussen et al., 2011), knowledge of the HO/CO significance in oocytes is still insufficient. We assumed that, similarly as in somatic cells, CO could also prevent the apoptotic pathway in oocytes. We hypothesized that CO could improve the viability of porcine oocytes. The aim of our work is to identify the effect of CO on the course of the porcine oocyte aging.

Materials and Methods

Collection, Cultivation and *in vitro* Aging of Porcine Oocytes

Porcine ovaries were obtained from local slaughterhouses from gilts during an unknown stage of the oestrous cycle. Porcine oocytes were obtained from the ovaries through aspiration of the follicular fluid from follicles (2 – 5 mm). Only oocytes with intact cytoplasm and compact cumuli were chosen for the experiments. Oocytes were cultivated in the modified M199 medium (Gibco-BRL, Life Technologies, Paisley, Scotland) containing sodium bicarbonate (32.5 mM), calcium L-lactate (2.75 mM), gentamicin (0.025 mg/ml), HEPES (6.3 mM), 13.5 IU eCG: 6.6 IU hCG/ml (P.G.600; Intervet, Boxmeer, Holland) and 10% (v/v) fetal calf serum (GibcoBRL). Oocytes were cultivated for 48 hours up to the metaphase stage of the second meiotic division (MII) in 4-well Petri dishes (Nunc, Fisher Scientific, USA; 1 ml of cultivation medium; 39°C; 5,0 % CO₂). Meiotically matured oocytes were denuded and used for *in vitro* aging (24, 48 and 72 hours) under the same conditions in the cultivation medium without P.G. 600.

Evaluation of *in vitro* Aged Porcine Oocytes

Upon completion of the *in vitro* aging, the oocytes were fixed in a mixture of ethanol and acetic acid (3:1, w/v, 48 hours). Fixed oocytes were stained using 1.0% (w/v) orcein. The oocytes were classified into four groups according to the morphological signs of aging: intact oocytes (oocytes in the stages: metaphase II, anaphase II or telophase II), parthenogenetically activated oocytes (embryos and oocytes containing pronuclei), apoptotic oocytes (called fragmented oocytes; oocytes containing apoptotic vesicles under the *zona pellucida*) and lysed oocytes (oocytes with loss of integrity and rupture of cytoplasmic membrane) (Petrová et al., 2004).

Immunocytochemical Detection of Heme Oxygenase and Cleaved Caspase-

3

Upon completion of the oocyte cultivation period, *zona pellucida* was removed (0.1% pronase). Subsequently, the oocytes were fixed in 2.5% (w/v) paraformaldehyde in Phosphate-buffered saline (PBS). The oocyte membrane was permeabilized by 0.5% (w/v) Triton X in PBS with 0.01% (w/v) BSA. After rinsing in PBS with 0.1% (w/v) Tween 20 followed incubation with the primary antibody anti-heme oxygenase 1 or anti-heme oxygenase 2 (Abnova Corporation, Taipei, Taiwan; 1:200) or anti-cleaved caspase-3

(Asp175) (Cell Signaling Technology, Danvers, USA; 1:400). The incubation period took over 14-16 hours in moisture at a temperature of 4°C in 0.1 % (w/v) BSA and 0.01 % (w/v) Tween 20 in PBS. After incubation, the oocytes were rinsed with 0.1% (w/v) Tween 20 in PBS and cultivated with secondary anti-mouse IgG antibody conjugated with fluorescein-5-isothiocyanate (FITC, Sigma–Aldrich GmbH, Munich, Germany; 1:100) or anti-rabbit IgG conjugated with FITC (ThermoFisher Scientific, Rockford, USA; 1:500). The incubation with the secondary antibody took place at laboratory temperature in 0.1% (w/v) BSA and 0.01% (w/v) Tween 20 in PBS over the course of one hour in darkness. After incubation, the oocytes were rinsed in 0.1% (w/v) Tween 20 in PBS.

The chromatin was stained using 4',6-diamidino-2-phenylindole (DAPI, Sigma–Aldrich GmbH, Munich, Germany). To exclude non-specific secondary antibody binding the oocytes in the control group were treated in the same way as the experimental group, except that primary antibody incubation was not performed. Oocytes were mounted on slides and images were acquired using confocal scanning microscope (Zeiss, Germany). The images were analyzed in NIS Elements AR Software (NIKON, Japan). The data was expressed as mean signal intensity of the FITC fluorescence, reduced by a basal signal intensity of appropriate negative control. Each experiment was repeated at least three times at a minimal amount of 15 oocytes in each experimental group.

Western Blot

Western blot was performed for validation of the antibodies specificity in accordance to Tůmová et al., 2013. Briefly, MII oocytes were lysed and separated by SDS-PAGE. After blotting the membranes were incubated with primary antibodies - anti-heme oxygenase-1, anti-heme oxygenase-2 (Abnova Corporation, Taipei, Taiwan; 1:1000) or cleaved caspase-3 (Asp175) (Cell Signaling Technology, Danvers, USA; 1:1000) and then with secondary antibodies - Mouse IgG (Amersham GE Healthcare, Life Sciences, United Kingdom; 1:30,000) or rabbit IgG (Amersham GE Healthcare, Life Sciences, United Kingdom; 1:120,000). Proteins were detected by ECL Advanced Western blotting detection kit (Amersham GE Healthcare, Life Sciences, United Kingdom). Each western blot was repeated at least three times with a minimal amount of 200 oocytes.

***In Vitro* Cultivation of Aging Oocytes with Exogenous CO Donors or Heme Oxygenase Inhibitor**

Oocytes were cultivated in a cultivation media with the CO donors: CORM-2 (tricarbonyl dichlororuthenium (II) dimer; Sigma–Aldrich GmbH, Munich, Germany) at concentrations of 5, 25, 50 and 100 μ M, dissolved in dimethyl sulfoxide (DMSO) or CORM-A1 (sodium boranocarbonate; Sigma–Aldrich GmbH, Munich, Germany) at concentrations of 25, 50 and 100 μ M, dissolved in H₂O. CORM-2 and CORM-A1 both have distinct rate of CO release, wherein CORM-2 is a fast CO releaser, whereas CORM-A1 shows a slow and gradual CO release (Motterlini et al., 2002; 2005). In the case of HO inhibition, the oocytes were cultivated in a cultivation medium with HO inhibitor Zn-protoporphyrin IX (Zn-PP IX; Sigma–Aldrich GmbH, Munich, Germany), at concentrations of 2.5, 5 and 25 μ M dissolved in DMSO. In order to eliminate the effect of DMSO on the aging of porcine oocytes, a control group was incubated in a cultivation medium containing only DMSO. In order to confirm the effect of CO, oocytes were cultivated in cultivation medium with ruthenium (III) chloride (inactive CORM2; iCORM-2; Sigma–Aldrich GmbH, Munich, Germany) or inactive CORM-A1 (iCORM-A1). iCORM-A1 consisted of CORM-A1 prepared in 0.1 M HCl bubbled with N₂ gas for 10 min to dissipate all of the CO and then the pH of the solution was adjusted to 7.4. The effect of the iCORMs at comparable concentrations was not significant compared to oocytes cultivated in pure cultivation medium (supplemental information tab. 1). Upon completion of the given cultivation period, the oocytes were morphologically evaluated or prepared for immunocytochemical technique. Each experiment was repeated at least three times and at least 80 oocytes were used for each experimental group.

Statistical Analysis

Data from all experiments were subjected to statistical analysis. All experiments were repeated at least three times. The SAS 9.0 software (SAS Institute Inc., USA) was used for statistical analysis. Significant differences between groups were determined using analysis of variance (ANOVA) followed by Scheffe's method. A p-value of less than 0.05 was considered significant. Significant differences among different groups of oocytes are indicated by different symbols.

Results

Heme Oxygenase Isoforms Are Present in Porcine Oocyte during *in vitro* Aging

The objective of the experiment was to localize HO-1 and HO-2 in meiotically mature porcine oocytes (MII) and oocytes exposed to *in vitro* aging for the period of 24, 48 and 72 hours. HO isoforms are present in porcine oocyte (Fig. 1, Fig 2), and their expression gradually increase during *in vitro* aging. In case of HO-1, the signal was predominant in the nucleus/perichromosomal area (Fig. 2). On the contrary, the HO-2 signal was mainly observed in the oocyte cytoplasm (Fig. 3).

During the *in vitro* aging, the expression of both isoforms increased primarily in the oocyte cytoplasm. In comparison with meiotically matured oocytes, in oocytes aged 24 hours the expression of HO-1 increased by 2.2 ± 0.2 times, aged 48 hours increased by 3.5 ± 0.3 times and aged 72 hours increased by 6.7 ± 1.3 times (Fig. 4A). In the case of HO-2, the expression in oocytes aged 24 hours increased by 1.3 ± 0.1 times, aged 48 hours increased by 1.7 ± 0.14 times, and in oocytes aged 72 hours increased by 3.1 ± 0.6 times (Fig. 4B).

The Inhibition of Heme Oxygenase Increases the Ratio of Apoptotic Porcine Oocytes during the *in Vitro* Aging

This experiment focused on the effect of HO inhibition by HO inhibitor Zn-protoporphyrin IX (Zn-PP IX) on *in vitro* aging of porcine oocytes. After 24 hours of *in vitro* aging, the effect was significant only when the highest concentration of Zn-PP IX (25 μ M) was used, where the amount of intact oocytes in comparison to the control group decreased by 10.6% (94.0 ± 1.5 vs. $83.4 \pm 2.1\%$ for control and HO inhibitor, respectively) (Fig. 5A). In other concentrations of Zn-PP IX, the effect on the morphological signs of aging was not significant.

After 48 hours of *in vitro* aging in the presence of Zn-PP IX, the effect was significant in all concentrations, causing decreases in the ratio of intact oocytes and increases in the ratio of apoptotic oocytes. In oocytes treated with Zn-PP IX, the ratio of intact oocytes decreased by 11.1-12.9% (67.1 ± 1.6 vs. 54.2 ± 2.1 - $56.0 \pm 1.7\%$ for control and HO inhibitor, respectively) while the ratio of apoptotic oocytes increased by 5.6-9.4% (21.5 ± 2.3 vs. 27.0 ± 1.1 - $30.8 \pm 2.4\%$ for control and HO inhibitor, respectively).

Moreover, the highest concentration of inhibitor (25 μ M) also increased the ratio of lytic oocytes by 8.3% (1.1 ± 1.5 vs. $9.4 \pm 1.6\%$ for control and HO inhibitor, respectively). The effects of different concentrations of inhibitor did not differ significantly (Fig. 5B).

The most obvious manifestation of negative signs of aging was observed in oocytes aged 72 hours, wherein the ratio of apoptotic oocytes reached 60.4 ± 3.7 – $67.5 \pm 3.4\%$ in all groups (control and experimental). However, the effect of the HO inhibitor was not significant (Fig. 5C).

CO donors Suppress Apoptosis in Porcine Oocytes during the *in Vitro* Aging

The objective of this experiment was to identify the effect of CO on the course of *in vitro* aging of porcine oocytes evaluated based on morphological signs of aging. As the CO source, we used two CO donors (CORM-2 or CORM-A1) each with different CO release kinetic. The control group of oocytes was cultivated in the presence of inactive compounds (iCORM-2 or iCORM-A1) to eliminate its effect. It was found that CO delivered by CORMs causing decreases in the ratio of apoptotic oocytes and increases in the ratio of intact oocytes.

The effect of the CO donors on *in vitro* aging became apparent after 48 hours of *in vitro* aging. The effect of CORM-A1 was significant at the concentrations 25 and 50 μ M, which decreased the ratio of apoptotic oocytes by 10.2-14.4% (30.3 ± 4.3 vs. 15.6 ± 5.8 – 19.8 ± 4.6 for control and CORM-A1, respectively) and simultaneously increased the ration of intact oocytes by 12.6-16.0% (59.8 ± 3.6 vs. 72.3 ± 3.8 – 75.7 ± 3.7 for control and CORM-A1, respectively) (Fig. 6A, 6B). The difference between control group and group cultivated in the CORM-A1 at the concentration 100 μ M was not significant. The effect of CORM-2 was significant in all concentrations. CORM-2 decreased the ratio of apoptotic oocytes by 6.7-9.9% (21.5 ± 1.3 vs. 11.5 ± 1.5 – $14.8 \pm 0.3\%$ for control and CORM-2, respectively), while simultaneously increased the ratio of intact oocytes by 8.6-13.5% (67.1 ± 1.6 vs. 75.7 ± 1.4 – $80.6 \pm 1.4\%$ for control and CORM-2, respectively). The effects of different concentrations of CORM-2 did not differ significantly (Fig. 7A, 7B).

Both CO donors significantly suppressed apoptosis even in oocytes aged 72 hours. Effect of CORM-A1 was significant at the concentrations 50 and 100 μ M, which

decreased the ratio of apoptotic oocytes by 12.3-14.3% (59.5 ± 3.6 vs. 45.2 ± 7.3 – 47.2 ± 3.2 for control and CORM-A1, respectively) and simultaneously increased the ratio of intact oocytes by 11.2-12.5% (28.8 ± 3.7 vs. 40.0 ± 7.3 – 41.3 ± 6.0 for control and CORM-A1, respectively) (Fig. 6C). The difference between control group and group cultivated in the CORM-A1 concentration 25 μ M was not significant. As with oocytes aged 48 hours, also in oocytes aged 72 hours the effect of CORM-2 was significant in all concentrations. CORM-2 decreased the ratio of apoptotic oocytes by 8.9-17.4% (60.4 ± 2.7 vs. 43.0 ± 3.6 – $51.5 \pm 1.5\%$ for control and CORM-2, respectively) while increased the ratio of intact oocytes by 12.6-21.6% (17.5 ± 1.7 vs. 30.1 ± 2.4 – $39.1 \pm 2.2\%$ for control and CORM-2, respectively). The effects of different concentrations of CORM-2 did not differ significantly (Fig. 7C).

CO Donors Decrease the Activity of Caspase-3 in Porcine Oocytes during the *in Vitro* Aging

This experiment focused on the effect of CO donors CORM-2 or CORM-A1 on the fluorescence intensity of activated CAS-3 (cleaved CAS-3; cCAS-3) as a marker of apoptosis. It was found that the CO donors decrease the fluorescence intensity of cCAS-3 during *in vitro* aging (Fig. 8, Fig. 9, Fig. 10).

The effect of both CO donors on the activity of CAS-3 was significant in all groups, i.e. in oocytes aged 24, 48 and 72 hours. In the case of CORM-2, the effect of different concentrations did not differ significantly, whereas the effect of CORM-A1 differs depending to the concentration. We observed the most significant effect of both CO donors in oocytes aged 24 hours, wherein CORM-A1 causing decreases of cCAS-3 fluorescence intensity by 40.9-48.5% (100.0 ± 6.0 vs. 51.6 ± 6.2 – $59.2 \pm 5.9\%$ for control and CORM-A1 at the concentrations 25 and 50 μ M, respectively). The effect of CORM-A1 at the concentration 100 μ M was less significant and decreased cCAS-3 fluorescence intensity by 16.5% (100.0 ± 6.0 vs. $83.5 \pm 5.2\%$ for control and CORM-A1 at the concentrations 100 μ M, respectively) In the case of CORM-2, the cCAS-3 fluorescence intensity decreased by 55.6-63.7% (100.0 ± 8.4 vs. 36.3 ± 4.5 – $44.4 \pm 2.7\%$ for control and CORM-2, respectively).

After 48 and 72 hours of *in vitro* aging, the cCAS-3 fluorescence intensity was significantly lower in comparison with oocytes aged 24 hours, however, the cultivation of

oocytes with the CO donors also lead to the significant decrease of cCAS-3 fluorescence intensity. In group of oocytes aged 48 hours with CORM-A1 the cCAS-3 fluorescence intensity decreased by 16.7-23.5% (65.6 ± 10.3 vs. $42.1 \pm 4.5 - 48.9 \pm 4.9$ for control and CORM-A1 at the concentrations 25 and 50 μM , respectively) The effect of CORM-A1 at the concentration 100 μM was not significant (Fig. 11). In the case of CORM-2 the cCAS-3 fluorescence intensity decreased by 13.1-22.4% (35.3 ± 2.9 vs. $13.1 \pm 1.8 - 22.4 \pm 1.8$ for control and CORM-2, respectively) in oocytes aged 48 hours (Fig. 12). In group of oocytes aged 72 hours with CORM-A1 the cCAS-3 fluorescence intensity decreased by 13.4-22.9% (69.0 ± 7.6 vs. $46.1 \pm 8.7 - 55.6 \pm 5.2$ for control and CORM-A1 at the concentrations 25 and 50 μM , respectively). The effect of CORM-A1 at the concentration 100 μM was not significant. In the case of CORM-2 the cCAS-3 fluorescence intensity decreased by 10.8-14.1% (24.9 ± 5.2 vs. $10.8 \pm 3.8 - 14.1 \pm 2.0$ for control and CORM-2, respectively) in oocytes aged 72 hours.

Discussion

Post-ovulatory oocyte aging causes progressive decrease in oocyte quality and ultimately leads to cell death. Heme oxygenase (HO) is essential enzyme which cytoprotective properties being caused by the production of an important signalling molecule, carbon monoxide (CO) (Wu & Wang, 2005). Together with hydrogen sulfide (H₂S) and nitric oxide (NO), CO belongs to gasotransmitters, which are a subfamily of endogenous molecules of gases or gaseous signalling molecules. The role of H₂S and NO as signal molecules in porcine oocytes was identified (Bu et al., 2003, Goud et al., 2005; Nevoral et al., 2014; Krejčová et al., 2015), but information regarding the function of the HO/CO system is yet to be unavailable. Our work proves the presence of both HO isoforms in porcine oocytes and characterizes the effect of the HO inhibitor and CO donors on the course of *in vitro* aging of porcine oocytes.

We have shown that both HO isoforms (HO-1 and HO-2) are localized in meiotically matured and aged porcine oocytes and that over the course of *in vitro* aging, the expression of both HO isoforms increase. The HO-1 is a member of the superfamily of stress proteins, the expression of which increase in reaction to a wide spectrum of inducers (for example oxidative stress) (Dennerly, 2000; Ryter, Alam & Choi, 2006). The increased expression of the HO-1 isoform and higher production of CO decrease the production of reactive oxygen species (ROS) and suppress apoptosis (Pileggi et al, 2001; Li et al., 2016). We also assumed that HO-1/CO has the anti-apoptotic effect in oocytes. The increased expression of HO-1 during the *in vitro* aging of porcine oocytes, which we described, is likely induced by an increased production of ROS. High level of ROS is considered to be a major factor responsible for negative signs of aging in aged oocytes (Lord & Aitken, 2013). We also observed an increased expression of the HO-2 isoform. HO-2 is generally considered to be a constitutively expressed isoform responsible for a stable production of CO, whose level of expression does not respond to the effect of stress factors (Turkseven et al., 2007; Muñoz-Sánchez & Cháñez-Cárdenas, 2014). However, some authors have reported that in response to the stress factors the expression of HO-2 can also increase, which has cytoprotective effect (Kim et al., 2008; Ding et al., 2011). We suggest that both HO-1 and HO-2 can modulate ROS levels and suppress aging process in porcine oocytes.

The HO isoforms differed in their intracellular localization. It was found that the HO-1 was localized in meiotically mature and aged porcine oocytes predominantly in the area of chromosomes, and during the *in vitro* aging the HO-1 expression increased primarily in the cytoplasm. On the other hand, HO-2 was primarily localized in the cytoplasm. In general, HO-1 and HO-2 are considered to be proteins bound to the membrane of endoplasmic reticulum (Yoshida & Kikuchi, 1978; Ma et al., 2004), however, they may also be localized in other cellular compartments. HO-1 may be localized in caveolae, mitochondria and the nucleus (Dunn et al., 2014). Nuclear translocation of HO-1 requires cleavage of the membrane bound domain yielding ~28 kDa protein fragment (Lin et al., 2007; Linnenbaum et al., 2012). The transfer of HO-1 into the nucleus takes place in the reaction to stress factors, and the nuclear HO-1 protects cells from the effect of oxidative stress (Lin et al., 2007). We assumed that this isoform may have similar function also in porcine oocytes. Meiotic maturation of oocytes under *in vitro* conditions is endangered by increased oxidative stress in comparison with *in vivo* conditions (Dvořáková et al., 2016). Because HO-1 is present in the perichromosomal area in matured oocytes, we assumed that the transfer of HO-1 could have already taken place during the meiotic maturation, due to possible oxidative stress.

We had proved that in porcine oocytes, the HO/CO contributes to sustaining their viability and affects the regulation of apoptosis. HO inhibition using Zn-PP IX leads to apoptosis in somatic cells (Hirai et al., 2007), which we also observed in our experiments. Porcine oocyte aging in the presence of the HO inhibitor worsened the effect of aging and increased the ratio of apoptotic oocytes. In the oocytes aged 24 hours, the effect of the inhibitor was observable in the group cultivated with the highest concentration of the inhibitor (25 μ M). In the oocyte aged 48 hours, the effect of the inhibitor was significant in all concentrations; however the differences between the individual concentrations on proportion of apoptotic, parthenogenetically activated and intact oocytes were not significant. Given that the Zn-PP IX, particularly in higher concentrations, may also inhibit nitric oxide synthase (NOS) and soluble guanylyl cyclase (sGC) (Luo & Vincent, 1994; Grundemar & Ny, 1997), the possibility that this effect also appeared in our experiments cannot be excluded. However, Appleton et al.

(1999) state that at the low concentrations of Zn-PP IX (up to 5 μ M) there is selective inhibition of HO activity with minor effect on sGC and NOS. Therefore we assume that the non-selective inhibition could be seen only at the highest concentration (25 μ M) used by us. Although the inhibition of sGC leads to the increased occurrence of negative signs of oocyte aging (Goud et al., 2005), the NOS inhibition, on the contrary, decreases the ratio of apoptotic oocytes (Nevoral et al., 2013). The increased production of NO through inducible-NOS (i-NOS) isoform in the cell contributes to the formation of oxidative stress (Wang et al., 2007b). It is therefore possible that while the inhibitor Zn-PP IX suppresses the protective effect of HO/CO, it also simultaneously decreases oxidative stress through iNOS inhibition.

This work is the first that proved the cytoprotective effect of CO on oocyte aging. As a source of CO we used CORM-2 and CORM-A1. CORM-2 and CORM-A1 both have distinct rates of CO release (Motterlini et al., 2002; 2005). Both CO donors increased the ratio of intact oocytes and decreased the ratio of apoptotic oocytes during *in vitro* aging. In somatic cells, it was proved that HO/CO affects the viability of cells and apoptosis through the regulation of activity of pro-apoptotic and anti-apoptotic proteins. CO decreases the level of apoptosis, for example, through up-regulation of expression of the anti-apoptotic factor Bcl-2 and down-regulation of the activity of caspase-3 (Zhang et al., 2003a, Cepinskas et al., 2007). Caspase-3 activation also occurs during oocyte aging (Zhu et al., 2015, Zhu et al., 2016). As apparent from the results obtained, the CAS-3 attains highest activity in oocytes during the first day of aging, while over the course of the following days the activity of CAS-3 is significantly lower. Suppressing its activity through the CO donors affected the entire course of cultivation of the aging oocytes; but it was most significant, precisely during the first 24 hours of aging. However, the fact that the effect also lasted through the following days of aging could be a consequence of the fact that CO has the ability of pre-conditioning, where a short exposure to CO subsequently leads to increased resistance of cells against stressors and thus to decreased level of apoptosis (Queiroga et al., 2012; Andreadou et al., 2015). Both CO donors suppress apoptosis even each of them has slightly different effect. The effect of CORM-2 on the ratio of apoptotic oocytes and the CAS-3 activity was not dose-dependent. However, in the case of CORM-A1, we observed the effect of

different concentrations. We attribute it to the different kinetics of CO release from CORM-A1 and CORM-2 (Motterlini & Foresti, 2017).

The mechanism through which CO affects CAS-3 activity is apparently complex, because CO may regulate the apoptotic pathway by multiple mechanisms. Beside the modulation of ROS levels, CO can modulate activity of c-jun terminal kinase (JNK), a member of the MAPK family. In somatic cells, CO decreases the activity of JNK (Morse et al., 2003) while in aging of porcine oocytes, it leads to the decrease of its activity and suppression of apoptosis (Sedmíková et al., 2013). Additionally, CO may suppress apoptosis through the modulation of the level of Ca^{2+} . In somatic cells, CO may decrease the Ca^{2+} level (Lin & McGrath, 1988; Gende, 2004; Lim et al., 2005), while in aging oocytes the release of Ca^{2+} can trigger the cell death (Gordo et al., 2002; Zhu et al., 2016). Another molecular target for CO includes pro-apoptotic and anti-apoptotic factors. In somatic cells, CO increases the expression of the anti-apoptotic factor Bcl-2 and decreases the expression of the pro-apoptotic factors Bid and Bax (Zhang et al., 2003; Wang et al., 2007a). It is possible that CO may also act in a similar manner in oocytes.

Conclusions

These experiments have shown that the CO donors suppress negative signs of aging in porcine oocytes and inhibit apoptosis through reduction of CAS-3 activity. For these reasons, it can be assumed that the HO/CO system is functional in porcine oocytes and that it contributes to sustaining the viability of oocytes and regulates the programmed cell death of oocytes.

References

- Almeida AS, Queiroga CS, Sousa MF, P. M. Alves PM, Viera HL. 2012. Carbon monoxide modulates apoptosis by reinforcing oxidative metabolism in astrocytes: role of Bcl-2. *Journal of Biological Chemistry* 287(14):10761-10770. DOI 10.1074/jbc.M111.306738.
- Andreadou I, Efstathios K, Iliodromitis EK, Rassaf T, Schulz R, Papapetropoulos A, Ferdinandy P. 2015. The role of gasotransmitters NO, H₂S and CO in myocardial ischaemia/reperfusion injury and cardioprotection by preconditioning, postconditioning and remote conditioning. *British Journal of Pharmacology* 172(6):1587-1606. DOI 10.1111/bph.12811.
- Appleton SD, Chretien ML, McLaughlin BE, Vreman HJ, Stevenson DK, Brien JF, Nakatsu K, Maurice DH, Marks GS. 1999. Selective inhibition of heme oxygenase, without of nitric oxide synthase or soluble guanylyl cyclase, by metalloporphyrins at low concentrations. *Drug Metabolism & Disposition* 27(10): 1214-1219.
- Badenas J, Santaló J, Calafell JM, Estop AM, Egozcue J. 1989. Effect of the degree of maturation of mouse oocytes at fertilization: A source of chromosome imbalance. *Gamete Research* 24(2):205-218. DOI 10.1002/mrd.1120240208.
- Biswas C, Shah N, Muthu M, La P, Fernando AP, Sengupta S, Yang G, Dennery PA. 2014. Nuclear heme oxygenase-1 (HO-1) modulates subcellular distribution and activation of Nrf2, impacting metabolic and anti-oxidant defenses. *Journal of Biological Chemistry* 289(39):26882-26894. DOI 10.1074/jbc.M114.567685.
- Blandau RJ. 1952. The female factor in fertility and infertility. I. Effects of delayed fertilization on the development of the pronuclei in rat ova. *Fertility and Sterility* 3(5):349-365.
- Brouard S, Otterbein LE, Anrather J, Tobiasch E, Bach FH, Choi AK, Soares MP. 2000. Carbon monoxide generated by heme oxygenase 1 suppresses endothelial cell apoptosis. *The Journal of Experimental Medicine*. 192 (7): 1015-1026.
- Brouard S, Berberat PO, Tobiasch E, Seldon MP, Bach FH, Soares MP. 2002. Heme oxygenase-1-derived carbon monoxide requires the activation of transcription factor NF-kappa B to protect endothelial cells from tumor necrosis factor-alpha-mediated apoptosis. *The Journal of Biological Chemistry*. 277(20): 17950-17961. DOI

10.1074/jbc.M108317200.

Bu S, Xia G, Tao Y, Lei L, Zhou B. 2003. Dual effects of nitric oxide on meiotic maturation of mouse cumulus cell-enclosed oocytes in vitro. *Molecular and Cellular Endocrinology* 207(1-2):21-30. DOI 10.1016/S0303-7207(03)00213-2.

Cepinskas G, Katada K, Bihari A, Potter RF. 2007. Carbon monoxide liberated from carbon monoxide-releasing molecule CORM-2 attenuates inflammation in the liver of septic mice. *The American Journal of Physiology: Gastrointestinal and Liver Physiology* 294(1):G184-G191. DOI 10.1152/ajpgi.00348.2007.

Dennery PA. 2000. Regulation and role of heme oxygenase in oxidative injury. *Current Topics in Cellular Regulation* 1(36):181. DOI 10.1016/S0070-2137(01)80008-X.

Ding B, Gibbs PEM, Brookes PS, Maines MD. 2011. The coordinated increased expression of biliverdin reductase and heme oxygenase-2 promotes cardiomyocyte survival: a reductase-based peptide counters β -adrenergic receptor ligand-mediated cardiac dysfunction. *The FASEB Journal* 25(1):301-313. DOI 10.1096/fj.10-166454.

Dunn LL, Midwinter RG, Ni J, Hamid HA, Parish CHR, Stocker R. 2014. New insights into intracellular locations and functions of heme oxygenase-1. *Antioxidants* 20(11):1723-1742. DOI 10.1089/ars.2013.5675.

Dvořáková M, Heroutová I, Němeček D, Adámková K, Krejčová T, Nevoral J, Kučerová-Chrpová V, Petr J, Sedmíková M. 2016. The antioxidative properties of S-allyl cysteine not only influence somatic cells but also improve early embryo cleavage in pigs. *PeerJ* 4:e2280. DOI 10.7717/peerj.2280.

Elmore S. 2007. Apoptosis: A review of programmed cell death. *Toxicologic Pathology* 35(4):495-516. DOI 10.1080/01926230701320337.

Fissore R, Kurokawa M, Knott J, Zhang M, Smyth J. 2002. Mechanisms underlying oocyte activation and postovulatory ageing. *Reproduction* 124(6):745-754. DOI 10.1530/rep.0.1240745.

Gende OA. 2004. Carbon monoxide inhibits capacitative calcium entry in human platelets. *Thrombosis Research* 114(2):113-119. DOI 10.1016/j.thromres.2004.04.015.

Goebel U, Siepe M, Mecklenburg A, Stein P, Roesslein M, Schwer CI, Schmidt R, Doenst T, Geiger KK, Pahl HL, Schlensak C, Loop T. 2008. Carbon monoxide inhalation reduces pulmonary inflammatory response during cardiopulmonary bypass in

pigs. *Anesthesiology* 108(6):1025-1036. DOI 10.1097/ALN.0b013e3181733115.

Gordo AC, Rodrigues P, Kurokawa M, Jellerette T, Ellerette EE, Warner C, Fissore R. 2002. Intracellular calcium oscillations signal apoptosis rather than activation in in vitro aged mouse eggs. *Biology of Reproduction* 66(6):1828-1837. DOI 10.1095/biolreprod66.6.1828.

Goud AP, Goud PT, Diamond MP, Abu-Soud HM. 2005. Nitric oxide delays oocyte aging. *Biochemistry* 44(34):11361–11368. DOI 10.1021/bi050711f.

Grundemar L, Ny L. 1997. Pitfalls using metalloporphyrins in carbon monoxide research. *Trends in Pharmacological Sciences* 18(6):193-195. DOI 10.1016/S0165-6147(97)01065-1.

Hirai K, Sasahira T, Ohmori H, Fuji K, Kuniyasu H. 2007. Inhibition of heme oxygenase-1 by zinc protoporphyrin IX reduces tumor growth of LL/2 lung cancer in C57BL mice. *International Journal of Cancer* 120(3):500-505. DOI 10.1002/ijc.22287.

Jeřeta, M, Petr J, Krejčová T, Chmelíková E, Jílek F. 2008. In vitro ageing of pig oocytes: effects of the histone deacetylase inhibitor trichostatin A. *Zygote* 16(02):145-152. DOI 10.1017/S0967199408004668.

Jiang GJ, Wang K, Miao DQ, Guo L, Hou Y, Schatten H, Sun QY, Singh SR. 2011. Protein profile changes during porcine oocyte aging and effects of caffeine on protein expression patterns. *PLoS ONE* 6(12):e28996. DOI 10.1371/journal.pone.0028996.

Kang MK, Han SJ. 2011. Post-transcriptional and post-translational regulation during mouse oocyte maturation. *BMB reports* 44(3): 147-157. DOI 10.5483/BMBRep.2011.44.3.147.

Kikuchi K, Izaike Y, Noguchi J, Furukawa T, Daen FP, Naito K, Toyoda Y. 1995. Decrease of histone H1 kinase activity in relation to parthenogenetic activation of pig follicular oocytes matured and aged in vitro. *Journal of Reproduction and Fertility* 105(2):325-330.

Kim YM, Choi BM, Kim YS, Kwon YG, Kibbe MR, Billiar TR, Tzeng E. 2008. Protective effect of p53 in vascular smooth muscle cells against nitric oxide-induced apoptosis is mediated by up-regulation of heme oxygenase-2. *BMB Reports* 41(2):164-169. DOI 10.5483/BMBRep.2008.41.2.164.

Kim YM, Pae HO, Park JE, Lee YC, Woo JM, Kim NH, Choi YK, Lee BS, Kim SR, Chung HT. 2011. Heme oxygenase in the regulation of vascular biology: from molecular mechanisms to therapeutic opportunities. *Antioxidants & Redox Signaling* 14(1):137-167. DOI 10.1089/ars.2010.3153.

Krejčová T, Šmelcová M, Petr J, Bodart JF, Sedmíková M, Nevoral J, Dvořáková M, Vyskočilová A, Weingartova I, Kučerová-Chrpová V, Chmelíková E, Tůmová L, Jílek F. 2015. Hydrogen sulfide donor protects porcine oocytes against aging and improves the developmental potential of aged porcine oocytes. *PLoS ONE* 10(1):e0116964. DOI 10.1371/journal.pone.0116964.

Lanman JT. 1968. Delays during reproduction and their effects on the embryo and fetus. *New England Journal of Medicine* 278(18):993-999. DOI 10.1056/NEJM196805022781806.

Li C, Zhang C, Wang T, Xuan J, Su C, Wang Y. 2016. Heme oxygenase 1 induction protects myocardiac cells against hypoxia/reoxygenation-induced apoptosis. *Herz* 41(8):715-724. DOI 10.1007/s00059-016-4424-6.

Lim I, Gibbons SJ, Lyford GL, Miller SM, Strege PR, Sarr MG, Chatterjee S, Szurszewski JH, Shah VH, Farrugia G. 2005. Carbon monoxide activates human intestinal smooth muscle L-type Ca^{2+} channels through a nitric oxide dependent mechanism. *American Journal of Physiology-Gastrointestinal and liver physiology* 288(1):G7-G14. DOI 10.1152/ajpgi.00205.2004.

Lin H, McGrath JJ. 1988. Carbon monoxide effects on calcium levels in vascular smooth muscle. *Life Sciences* 43(22):1813-1816. DOI 10.1016/0024-3205(88)90280-9.

Lin Q, Weis S, Yang G, Weng YH, Helston R, Rish K, Smith A, Bordner J, Polte T, Gaunitz F, Dennerly PA. 2007. Heme oxygenase-1 protein localizes to the nucleus and activates transcription factors important in oxidative stress. *Journal of Biological Chemistry* 282(28):20621-20633. DOI 10.1074/jbc.M607954200.

Linnenbaum M, Busker M, Kraehling JR, Behrends S. 2012. Heme oxygenase isoforms differ in their subcellular trafficking during hypoxia and are differentially modulated by cytochrome P450 reductase. *PLoS ONE* 7(4): e35483. DOI 10.1371/journal.pone.0035483.

Liu L, Trimarchi JR, Keefe DL. 2000. Involvement of mitochondria in oxidative

stress-induced cell death in mouse zygotes. *Biology of Reproduction* 62 (6):1745–1753.
DOI 10.1095/biolreprod62.6.1745.

Lord T, Aitken RJ. 2013. Oxidative stress and ageing of the post-ovulatory oocyte. *Reproduction* 146(6):R217-R227. DOI 10.1530/REP-13-0111.

Luo D, Vincent SR. 1994. Metalloporphyrins inhibit nitric oxide-dependent cGMP formation in vivo. *European Journal of Pharmacology: Molecular Pharmacology* 267(3):263-267. DOI 10.1016/0922-4106(94)90149-X.

Ma N, Ding X, Doi M, Izumi N, Semba R. 2004. Cellular and subcellular localization of heme oxygenase-2 in monkey retina. *Journal of Neurocytology* 33(4):407-415. DOI 10.1023/B:NEUR.0000046571.90786.6e.

Miao YL, Kikuchi K, Sun QY, Schatten H. 2009. Oocyte aging: cellular and molecular changes, developmental potential and reversal possibility. *Human Reproduction Update* 15(5):573-585. DOI 10.1093/humupd/dmp014.

Morse D, Pischke SE, Zhou Z, Davis RJ, Flavell RA, Loop T, Otterbein SL, Otterbein LE, Choi AM. 2003. Suppression of inflammatory cytokine production by carbon monoxide involves the JNK pathway and AP-1. *Journal of Biological Chemistry* 278(39):36993-36998. DOI 10.1074/jbc.M302942200.

Motterlini R, Clark JE, Foresti R, Sarathchandra P, Mann BE, Green CJ. 2002. Carbon monoxide-releasing molecules: characterization of biochemical and vascular activities. *Circulation Research* 90(2): E17-24. DOI 10.1161/hh0202.104530.

Motterlini R, Mann BE, Johnson TR, Clark JE, Foresti R, Green CJ. 2003. Bioactivity and pharmacological actions of carbon monoxide-releasing molecules. *Current pharmaceutical design*. 9(30): 2525-2539.

Motterlini R, Mann BE, Foresti R. 2005. Therapeutic applications of carbon monoxide-releasing molecules. *Expert Opinion on Investigational Drugs* 14(11):1305-1318. DOI 10.1517/13543784.14.11.1305.

Motterlini R, Otterbein LE. 2010. The therapeutical potential of carbon monoxide. *Nature reviews. Drug discovery*. 9(9): 728-743. DOI 10.1038/nrd3228.

Motterline R, Foresti R. 2017. Biological signaling by carbon monoxide and carbon monoxide-releasing molecules. *American Journal of Physiology. Cell Physiology* 312(3): C302-C313. DOI 10.1152/ajpcell.00360.2016.

Muñoz-Sánchez J, Chánez-Cárdenas ME. 2014. A Review on hemeoxygenase-2: focus on cellular protection and oxygen response. *Oxidative Medicine and Cellular Longevity* 2014:1-16. DOI 10.1155/2014/604981.

Nevoral J, Petr J, Gelaude A, Bodart JF, Kučerová-Chrpová V, Sedmíková M, Krejčová T, Kolbabová T, Dvořáková M, Vyskočilová A, Weingartová I, Křivohlávková L, Žalmanová T, Jílek F. 2014. Dual effects of hydrogen sulfide donor on meiosis and cumulus expansion of porcine cumulus-oocyte complexes. *PLoS ONE* 9(7):e99613. DOI 10.1371/journal.pone.0099613.

Otterbein LE, Hedblom A, Harris C, Csizmadia E, Gallo D, Wegiel B. 2011. Heme oxygenase-1 and carbon monoxide modulate DNA repair through ataxia-telangiectasia mutated (ATM) protein. *PNAS* 108(35): 14491-14496. DOI 10.1073/pnas.1102295108.

Petrache I, Otterbein LE, Alam J, Weigand GW, Choi AMK. 2000. Heme oxygenase-1 inhibits TNF-alpha induced apoptosis in cultured fibroblasts. *American Journal of Physiology-Lung Cellular and Molecular Physiology*. 278(3): 312-319.

Petrová I, Sedmíková M, Chmelíková E, Svestková D, Rajmon R. 2004. In vitro ageing of porcine oocytes. *Czech Journal of Animal Science* 49:93–98.

Petrová I, Sedmíková M, Petr J, Vodková Z, Pytloun P, Chmelíková E, Reháček D, Čtrnáctá A, Rajmon R, Jílek F. 2009. The roles of c-Jun N-terminal kinase (JNK) and p38 mitogen-activated protein kinase (p38 MAPK) in aged pig oocytes. *Journal of Reproduction and Development* 55(1):75-82. DOI 10.1262/jrd.20061.

Pileggi A, Molano RD, Berney T, Cattani P, Vizzardelli C, Oliver R, Fraker C, Ricordi C, Pastori RL, Bach FH, Inverardi L. 2001. Heme oxygenase-1 induction in islet cells results in protection from apoptosis and improved in vivo function after transplantation. *Diabetes* 50(9):1983-1991. DOI 10.2337/diabetes.50.9.1983.

Queiroga CS, Tomasi S, Widerøe M, Alves PM, Vercelli A, Vieira HL. 2012. Preconditioning triggered by carbon monoxide (CO) provides neuronal protection following perinatal hypoxia-ischemia. *PLoS ONE* 7(8):e42632. DOI 10.1371/journal.pone.0042632.

Ryter SW, Alam J, Choi AM. 2006. Heme oxygenase-1/carbon monoxide: from basic science to therapeutic applications. *Physiological Reviews* 86(2):583-650. DOI 10.1152/physrev.00011.2005.

Ryter SW, Choi AM. 2016. Targeting heme oxygenase-1 and carbon monoxide for therapeutic modulation of inflammation. *Translational Research* 167(1):7-34. DOI 10.1016/j.trsl.2015.06.011.

Sadler KC, Yüce O, Hamaratoglu F, Vergé V, Peaucellier G, Picard A. 2004. MAP kinases regulate unfertilized egg apoptosis and fertilization suppresses death via Ca^{2+} signaling. *Molecular Reproduction and Development* 67(3):366-383. DOI 10.1002/mrd.20023.

Salvesen GS, Dixit VM. 1997. Caspases: intracellular signaling by proteolysis. *Cell* 91(4):443-446. DOI 10.1016/S0092-8674(00)80430-4.

Sedmíková M, Petr J, Dörflerová A, Nevoral J, Novotná B, Krejčová T, Chmelníková E, Tůmová L. 2013. Inhibition of c-Jun N-terminal kinase (JNK) suppresses porcine oocyte ageing in vitro. *Czech Journal of Animal Science* 58:535-545.

Schallner N, Fuchs M, Schwer CI, Loop T, Buerkle H, Lagrèze WA, van Oterendorp C, Biermann J, Goebel U. 2012. Postconditioning with inhaled carbon monoxide counteracts apoptosis and neuroinflammation in the ischemic rat retina. *PLoS ONE* 7(9):e46479. DOI 10.1371/journal.pone.0046479.

Slee EA, Adrain C, Martin SJ. 2001. Executioner caspase-3, -6, and -7 perform distinct, non-redundant roles during the demolition phase of apoptosis. *Journal of Biological Chemistry* 276(10):7320-7326. DOI 10.1074/jbc.M008363200.

Szollosi D. 1971. Morphological changes in mouse eggs due to aging in the fallopian tube. *American Journal of Anatomy* 130(2):209-225. DOI 10.1002/aja.1001300207.

Suttner DM, Dennery PA. 1999. Reversal of HO-1 related cytoprotection with increased expression is due to reactive iron. *FASEB Journal* 13(13):1800-1809.

Taylor RC, Cullen SP, Martin SJ. 2008. Apoptosis: controlled demolition at the cellular level. *Nature Reviews Molecular Cell Biology* 9(3):231-241. DOI 10.1038/nrm2312.

Tenhunen R, Marver HS, Schmid R. 1969. Microsomal heme oxygenase. Characterization of the enzyme. *The Journal of Biological Chemistry* 23(244):6388-6394.

Tenhunen R, Marver HS, Schmid R. 1968. The enzymatic conversion of heme to

bilirubin by microsomal heme oxygenase. *Proceedings of the National Academy of Sciences USA* 2(61):748–755.

Tůmová L, Petr J, Žalmanová T, Chmelíková E, Kott T, Tichovská H, Kučerová-Chrpová V, Hošková K, Jílek F. 2013. Calcineurin expression and localisation during porcine oocyte growth and meiotic maturation. *Animal Reproduction Science* 141(3-4): 154-163. DOI 10.1016/j.anireprosci.2013.07.011.

Turkseven S, Drummond G, Rezzani R, Rodella L, Quan S, Ikehara S, Abraham NG. 2007. Impact of silencing HO-2 on EC-SOD and the mitochondrial signaling pathway. *Journal of Cellular Biochemistry* 100(4):815-823. DOI 10.1002/jcb.21138.

Wang X, Wang Y, Kim HP, Nakahira K, Ryter SW, Choi AM. 2007a. Carbon monoxide protects against hyperoxia-induced endothelial cell apoptosis by inhibiting reactive oxygen species formation. *Journal of Biological Chemistry* 282(3):1718-1726. DOI 10.1074/jbc.M607610200.

Wang XL, Liu HR, Tao L, Liang F, Yan L, Zhao RR, Lopez BL, Christopher TA, Ma XL. 2007b. Role of iNOS derived reactive nitrogen species and resultant nitrative stress in leukocytes induced cardiomyocyte apoptosis after myocardial ischemia/reperfusion. *Apoptosis* 12(7):1209-1217. DOI 10.1007/s10495-007-0055-y.

Wang X, Wang Y, Lee SJ, Kim HP, Choi AM, Ryter SW. 2011. Carbon monoxide inhibits Fas activating antibody-induced apoptosis in endothelial cells. *Medical Gas Research* 1(1):8. DOI 10.1186/2045-9912-1-8.

Wilson JL, Bouillaud F, Almeida AS, Vieira HL, Ouidja MO, Dubois-Randé JL, Foresti R, Motterlini R. 2017. Carbon monoxide reverses the metabolic adaptation of microglia cells to an inflammatory stimulus. *Free Radical Biology & Medicine* 104, 311-323. DOI 10.1016/j.freeradbiomed.2017.01.022.

Wu L, Wang R. 2005. Carbon monoxide: endogenous production, physiological functions, and pharmacological applications. *Pharmacological Reviews* 57(4):585-630. DOI 10.1124/pr.57.4.3.

Xu Z, Abbott A, Kopf GS, Schultz RM, Ducibella T. 1997. Spontaneous activation of ovulated mouse eggs: time-dependent effects on M-phase exit, cortical granule exocytosis, maternal messenger ribonucleic acid recruitment, and inositol 1,4,5-trisphosphate sensitivity. *Biology of Reproduction* 57(4):743–750. Doi

10.1095/biolreprod57.4.743.

Yoshida T, Kikuchi G. 1978. Purification and properties of heme oxygenase from pig spleen microsomes. *The Journal of Biological Chemistry* 253(12):4224–4229.

Zenclussen ML, Jensen F, Rebelo S, El-Mousleh T, Casalis PA, Zenclussen AC. 2012. Heme oxygenase-1 expression in the ovary dictates a proper oocyte ovulation, fertilization, and corpora lutea maintenance. *American Journal of Reproductive Immunology* 67(5):376-382. DOI 10.1111/j.1600-0897.2011.01096.x.


Zhang X, Shan P, Alam J, David RJ, Flavell RA, Lee PJ. 2003a. Carbon monoxide modulates Fas/Fas ligand, caspases, and Bcl-2 family proteins via the p38 mitogen-activated protein kinase pathway during ischemia-reperfusion lung injury. *Journal of Biological Chemistry* 278(24):22061-22070. DOI 10.1074/jbc.M301858200.

Zhang X, Shan P, Otterbein LE, Alam J, Flavell RA, David RJ, Choi AM, Lee PJ. 2003b. Carbon monoxide inhibition of apoptosis during ischemia-reperfusion lung injury is dependent on the p38 mitogen-activated protein kinase pathway and involves caspase 3. *Journal of Biological Chemistry* 278(2):1248-1258. DOI 10.1074/jbc.M208419200.

Zhu J, Zhang J, Li H, Wang TY, Zhang CHX, Luo MJ, Tan JH. 2015. Cumulus cells accelerate oocyte aging by releasing soluble Fas Ligand in mice. *Scientific Reports* 5:8683. DOI 10.1038/srep08683.


Zhu J, Lin FH, Zhang J, Lin J, Li H, Li YW, Tan XW, Tan JH. 2016. The signaling pathways by which the Fas/FasL system accelerates oocyte aging. *Aging* 8(2):291-303. DOI 10.18632/aging.100893.

686 **Figures and Tables**




687 *Figure 1: Immunoblotting of HO isoforms (HO-2, MW~35,7 kDa; HO-1, MW~33/28 kDa) in*
 688 *porcine oocytes. Oocytes were matured to the second metaphase stage (MII stage). Proteins were*
 689 *separated by SDS-PAGE, transferred to a nitrocellulose membrane and then incubated with HO specific*
 690 *antibodies (anti-HO1, anti-HO2; both 1:1000). One sample contained proteins from 200 oocytes. HO1*
 691 *and HO2 and their truncated forms were detected by specific antibodies. The arrows indicate bands*
 692 *corresponding to the molecular weight of the HO protein.*

693
694




695 *Fig. 2: Localization of HO-1 in meiotically matured porcine oocytes (MII) (A) and in oocytes*
 696 *exposed to in vitro aging for 24 (B), 48 (C) and 72 (D) hours. HO-1 is shown in green (FITC), chromatin is*
 697 *shown in blue (DAPI), magnified 400x.*

698
699




700 *Fig. 3: Localization of HO-2 in meiotically mature porcine oocytes (MII) (A) and in oocytes*
701 *exposed to in vitro aging for 24 (B), 48 (C) and 72 (D) hours. HO-2 is shown in green (FITC), chromatin is*
702 *shown in blue (DAPI), magnified 400x.*

703



704 Fig. 4: Expression of HO-1 (A) and HO-2 (B) in meiotically matured (MI I) porcine oocytes and
 705 oocytes exposed to in vitro aging for 24, 48 and 72 hours. The level of expression of HO-1 and HO-2 was
 706 determined as the mean intensity of the given isoform's signal in porcine oocytes and related to the
 707 average signal intensity of the given HO isoform in meiotically matured (MI I) oocytes. Bars show the
 708 mean \pm SEM. *#+ indicates significant differences in the HO-1 and HO-2 signal intensity ($P<0.05$).

709



710 Fig. 5: The effect of the HO inhibitor zinc protoporphyrin IX (Zn-PP IX) on the porcine oocytes
 711 during in vitro aging for 24 (A), 48 (B) and 72 (C) hours. Control group was cultivated in a medium
 712 containing DMSO. The experimental group was cultivated in a medium containing Zn-PP IX at the
 713 concentrations of 2.5, 5 and 25 μ M. Data is expressed in a relative manner. The stages of in vitro aging
 714 were morphologically evaluated as: intact (MII), lytic (L), parthenogenetically activated (PA) and apoptotic
 715 (A). * indicates significant difference in the ratio of intact (MII) oocytes between different concentrations of
 716 inhibitor and control group ($P < 0.05$). # indicates significant difference in ratio of apoptotic (A) oocytes
 717 between different concentrations of inhibitor and control group ($P < 0.05$). + indicates significant difference
 718 in ratio of lytic (L) oocytes between different concentrations of inhibitor and control group. No significant

719 *difference was found in the ration of parthenogenetically activated (PA) oocytes between different*
 720 *concentrations of inhibitor and control group.*

721





Fig. 6: The effect of the CO donor CORM-A1 on the porcine oocytes during in vitro aging for 24 (A), 48 (B) and 72 (C) hours. The control group was cultivated in a medium containing an inactive form of CO donor (iCORM-A1; 100 μM). The experimental group was cultivated in a medium containing CORM-A1 at the concentrations of 25, 50 and 100 μM. The data is expressed in a relative manner. Stages of in vitro aging were determined morphologically as: intact (MII), lytic (L), parthenogenetically activated (PA) and apoptotic (A). * indicates significant difference in the ratio of intact (MII) oocytes between different concentrations of CORM-A1 and control group ($P < 0.05$). # indicates significant difference in the ratio of apoptotic (A) oocytes between different concentrations of CORM-A1 and control group ($P < 0.05$). No significant difference was found in the ration of parthenogenetically activated (PA) and lytic (L) oocytes

731 *between different concentrations of CORM-A1 and control group.*

732



733

734

735

736

737

738

739

740

741

*Fig. 7: The effect of the CO donor CORM-2 on the porcine oocytes during in vitro aging for 24 (A), 48 (B) and 72 (C) hours. The control group (C) was cultivated in a medium containing an inactive form of CO donor (iCORM-2; 100 μM). The experimental group was cultivated in a medium containing CORM-2 at the concentrations of 5, 25, 50 and 100 μM. The data is expressed in a relative manner. Stages of in vitro aging were determined morphologically as: intact (MII), lytic (L), parthenogenetically activated (PA) and apoptotic (A). * indicates significant difference in the ratio of intact (MII) oocytes between different concentrations of CORM-A1 and control group ($P<0.05$). # indicates significant difference in the ratio of apoptotic (A) oocytes between different concentrations of CORM-A1 and control group ($P<0.05$). No significant difference was found in the ration of parthenogenetically activated (PA) and lytic (L) oocytes*

742 *between different concentrations of CORM-2 and control group.*

743




Fig. 8: Typical expression pattern of activated CAS-3 (cleaved CAS-3) in porcine oocytes during in vitro aging for 24 (A), 48 (B) and 72 (C) hours in the presence of CO donor CORM-A1 at the concentrations of 25 and 50 μM. The control group represents oocytes cultivated with an inactive form of the CO donor (iCORM-A1). The experimental group represents oocytes cultivated with CORM-A1. The negative control group (NC) represents oocytes treated only with secondary antibody. Cleaved CAS-3 is shown in green (FITC), chromatin is shown in blue (DAPI), magnified 400x.





Fig. 9: Typical


expression pattern of activated CAS-3 (cleaved CAS-3) in porcine oocytes during in vitro aging for 24 (A), 48 (B) and 72 (C) hours in the presence of CO donor CORM-2. The control group represents oocytes cultivated with an inactive form of the CO donor (iCORM-2). The experimental group represents oocytes cultivated with CORM-2. The negative control group (NC) represents oocytes treated only with secondary antibody. Cleaved CAS-3 is shown in green (FITC), chromatin is shown in blue (DAPI), magnified 400x.

759




760 *Figure 10: Immunoblotting of cleaved caspase 3 (cCAS-3, MW~17/19 kDa) in porcine oocytes.*
 761 *Oocytes were meiotically matured in the second metaphase (MII stage). Proteins were separated on*
 762 *SDS-PAGE, transferred to a nitrocellulose membrane and then incubated with CAS-3 specific antibodies*
 763 *(anti-cleaved caspase-3 (Asp175). One sample contained proteins from 400 oocytes.*

764



765 *Fig. 11: The effect of the CO donor CORM-A1 on the expression of activated CAS-3 during in*
 766 *vitro* aging of porcine oocytes for 24, 48 and 72 hours. The data of expression of the activated CAS-3 are
 767 related to the data of expression of the activated CAS-3 in oocytes in the control group, exposed to in
 768 vitro aging for 24 hours. The control group represents oocytes cultivated in a medium containing an
 769 inactive form of the CO donor (iCORM-A1; 100 µM). The experimental group was cultivated in a medium
 770 containing CORM-A1 at the concentrations of 25, 50 and 100 µM. Bars show mean ± SEM. *# indicate
 771 significant differences in the level of expression of the activated CAS-3 between different concentrations
 772 of CORM-A1 or control group for each aging time separately ($P < 0.05$).

773



774 *Fig. 12: The effect of the CO donor CORM-2 on the expression of activated CAS-3 during in vitro*
 775 *aging of porcine oocytes for 24, 48 and 72 hours. The data of expression of the activated CAS-3 are*
 776 *related to the data of expression of the activated CAS-3 in oocytes in the control group exposed to in vitro*
 777 *aging for 24 hours. The control group represents oocytes cultivated in a medium containing an inactive*
 778 *form of the CO donor (iCORM-2; 100 µM). The experimental group was cultivated in a medium containing*
 779 *CORM-2 at the concentrations of 25, 50 and 100 µM. Bars show mean ± SEM. * indicates significant*
 780 *differences in the level of expression of the activated CAS-3 between different concentrations of CORM-2*
 781 *and control group for each aging time separately (P<0.05).*