

New species of the endemic Neotropical caddisfly genus *Contulma* from the Andes of Ecuador (Trichoptera: Anomalopsychidae)

Ralph Holzenthal ^{Corresp., 1}, Blanca Rios-Touma ², Ernesto Rázuri-Gonzales ¹

¹ Department of Entomology, University of Minnesota - Twin Cities Campus, Saint Paul, Minnesota, United States

² Facultad de Ingenierías y Ciencias Agropecuarias. Ingeniería Ambiental, Universidad de las Americas, Quito, Ecuador

Corresponding Author: Ralph Holzenthal

Email address: holze001@umn.edu

Adults of 3 new species of *Contulma* Flint (Trichoptera: Anomalopsychidae) are described and illustrated from the Andes of Ecuador, *Contulma lina*, new species, *Contulma quito*, new species, and *Contulma sangay*, new species. These species are similar to previously described species from the region, including *C. paluguillensis*, *C. nevada*, and *C. lancelolata*. New provincial records are provided for *C. bacula*, *C. cataracta*, and *C. echinata*. *Contulma duffi* Oláh, 2016 is considered a junior, subjective synonym of *C. penai*, Holzenthal and Flint, 1995, **new synonym**. Also, we provide an identification key to males of the 30 *Contulma* species now known. The genus is composed mostly of regionally endemic species occurring above 2000 m, with a few more widespread species and some that are found at lower elevations.

New species of the endemic Neotropical caddisfly genus *Contulma* from the Andes of Ecuador (Trichoptera: Anomalopsychidae)

Ralph W. Holzenthal^{1*} Blanca Ríos-Touma², Ernesto Rázuri-Gonzales¹

¹ Department of Entomology, University of Minnesota, St. Paul, Minnesota, USA

² Universidad de las Américas, Facultad de Ingenierías y Ciencias Agropecuarias, Ingeniería Ambiental, Campus Queri, Quito, Ecuador

* Corresponding author = Ralph Holzenthal, holze001@umn.edu

14

15 **Abstract.** Adults of 3 new species of *Contulma* Flint (Trichoptera: Anomalopsychidae) are
 16 described and illustrated from the Andes of Ecuador, *Contulma lina*, new species, *Contulma*
 17 *quito*, new species, and *Contulma sangay*, new species. These species are similar to previously
 18 described species from the region, including *C. paluguillensis*, *C. nevada*, and *C. lanceolata*.
 19 New provincial records are provided for *C. bacula*, *C. cataracta*, and *C. echinata*. *Contulma*
 20 *duffi* Oláh, 2016 is considered a junior, subjective synonym of *C. penai*, Holzenthal and Flint,
 21 1995, **new synonym**. Also, we provide an identification key to males of the 30 *Contulma* species
 22 now known. The genus is composed mostly of regionally endemic species occurring above 2000
 23 m, with a few more widespread species and some that are found at lower elevations.

24

Introduction

Neotropical Trichoptera currently includes more than 3,200 described species representing 155 genera and 25 families (Holzenthal and Calor, 2017) occurring in Mexico, the Caribbean, Central, and South America. Remarkably, 115 genera, or ca.75% of the total, are endemic to the region, making the fauna the second most diverse in the world for endemic genera after the Australasian (de Moore and Ivanov 2008). Moreover, several Neotropical genera are highly endemic regionally at the species level such as *Amphoropsycha* (Holzenthal 1985, Holzenthal and Rázuri-Gonzales 2011), *Atananolica* (Holzenthal 1988, Henriques-Oliveira and Santos, 2014), and *Contulma* (Holzenthal and Robertson 2006), the latter the main subject of this paper. Nevertheless, the Neotropical caddisfly fauna is incompletely known, mainly because there are regions in the Neotropics where the aquatic ecosystems are far from being well studied (e.g., Ríos-Touma et al., 2017). There is also a lack of regional researchers studying the order, especially in the Andean countries of Venezuela, Colombia, Ecuador, Peru, and Bolivia, which undoubtedly harbor 100s of species. For example, co-author Ríos-Touma is the first Ecuadorian to describe new species of caddisflies and Rázuri-Gonzales only the second Peruvian to do so.

A comprehensive revision of the endemic Neotropical genus *Contulma* was completed by Holzenthal and Flint (1995) and included 21 species, 18 described as new. Since then, 7 new species have been described, including 1 we synonymize here. Species in the genus are known from Costa Rica, the Andes of Colombia to Chile, and in the mountains of southeastern Brazil. This genus seems to display a high degree of local endemism among its species (Holzenthal and Robertson 2006), which are rarely collected using standard light trap techniques. Hand netting during the day, especially at high elevations, or the use of Malaise traps is generally more effective (Holzenthal and Ríos-Touma, 2012). The infrequency of collection does not mean the species are rare in nature, but most of the species have been described from only 1-5 individuals. The habitats of these species are small waterfalls, seeps, and small streams in lush forested mountainous areas as well as high elevation *páramo* streams above the tree line in the Andes (Holzenthal and Flint 1995).

Of the 30 species now known in the genus, including 3 new species described here, 19 occur in the tropical Andean countries (Ecuador, Colombia, Peru, Bolivia) and all occur above 2000 m, except for 2 of these species also found in lower elevations (Holzenthal and Flint 1995,

Holzenthall and Robertson 2006). Nine species occur in Ecuador, 5 endemic and 3 present also in Colombia. The 3 new species described here are from localities where no caddisfly collecting occurred previously, confirming the pattern of high endemism of species in the genus.

Materials & Methods

We used the methods described by Blahnik and Holzenthall (2004) to prepare adult specimens for taxonomic study. Genitalia were cleared in 85% lactic acid heated to 125°C for 20 min (Blahnik et al. 2007). An Olympus BX41 compound microscope outfitted with a drawing tube was used to observe specimens. Genitalic structures were drawn with pencil on paper and final illustrations were rendered in Adobe Illustrator. Morphological terminology follows that of Holzenthall and Flint (1995). Descriptions of species and generation of the identification key were accomplished using the software packages DELTA and INTKEY (Dallwitz 1980, Dallwitz et al. 1999).

Types of the new species are deposited in the collections of the Museo Ecuatoriano de Ciencias Naturales, Quito, Ecuador (MECN) and the University of Minnesota Insect Collection, St. Paul, Minnesota, USA (UMSP). All specimens examined in this study were affixed with a barcode label containing a unique 9 digit numeric code starting with the prefix UMSP. These codes are provided here for holotypes only. All associated specimen data are stored in the UMSP database. This study was performed under the Environmental Ministry of Ecuador study permits 36-2010-IC-FLO/FAU-DPA-MA and 005-15-IC-FAU-FLO-DNB/MA.

The electronic version of this article in Portable Document Format (PDF) will represent a published work according to the International Commission on Zoological Nomenclature (ICZN), and hence the new names contained in the electronic version are effectively published under that Code from the electronic edition alone. This published work and the nomenclatural acts it contains have been registered in ZooBank, the online registration system for the ICZN. The ZooBank LSIDs (Life Science Identifiers) can be resolved and the associated information viewed through any standard web browser by appending the LSID to the prefix <http://zoobank.org/>. The LSID for this publication is: urn:lsid:zoobank.org:pub:54BC56DC-5CC1-4DA0-82E4-AF69599C2F5D. The online version of this work is archived and available from the following digital repositories: PeerJ, PubMed Central and CLOCKSS.

Results

Species descriptions

Contulma lina, new species, Holzenthal, Ríos-Touma, Rázuri-Gonzales

LSID urn:lsid:zoobank.org:act:310DDC54-0008-4535-A385-BD71BB630F21

Figures 1A-E, 4

Diagnosis: The discoveries of *C. lina*, n. sp., as well as *C. quito*, n. sp., and the recently described *C. paluguillensis*, all from the high Andes of Ecuador, are beginning to fill the morphological gaps found between some of the species described by Holzenthal and Flint (1995). These 3 species, in addition to *C. bacula*, *C. caldensis*, *C. cataracta*, *C. echinata*, *C. nevada*, *C. papallacta*, and perhaps *C. spinosa*, all from either Ecuador or Colombia, share several features. All have some degree of development of a dorsolateral process on segment IX in the male genitalia and a development of setae or a setose process on the mesal face of the same segment below the dorsolateral process. In some species, one or the other of these characters may be more or less developed or absent, but the combination of characters shared by these species indicate that they may have a common origin. All are known from only a handful of specimens, most from only the type and a few paratypes, and the species each occur from only one or a few high altitude localities spread across a vast expanse of the northern Andes. With the collection of additional specimens, it may prove that some of these species are synonyms. On the other hand, the phallic structures seem to be unique to each species and attest to their distinctiveness. We predict that additional collecting will discover yet more new species in this radiation of Andean Trichoptera biodiversity. *Contulma lina* is distinguished from the above mentioned species, all illustrated by Holzenthal and Flint (1995) and Holzenthal and Ríos-Touma (2012), by the following combination of characters: mesal surface of segment IX bearing a sinuous band of setae from below dorsolateral process and continuing to near sternum IX; mesal process of sternum IX with prominent posteromesal, heavily sclerotized, strongly emarginate projection, without excavation; and phallus without spine-like setae, but instead with the apex bearing a lightly sclerotized scale-like structure, shallowly excavate at apex.

Description: *Male:* Forewing length 5.5 mm (n=1). Forewing color gray brown, immaculate, vestiture rubbed (specimen was netted during the day in light rain and mist). Male genitalia: Segment IX very short dorsally, narrow, deeply excavate mesally; in lateral view, IX quadrate, extended anterolaterally; posteriorly with short, broad, dorsolateral, spatulate process; posterior margin of IX produced medially to form broad, prominent, quadrate, heavily setose, paired lateral lobe; lobes widely separated ventrally; mesal face of segment IX bearing sinuous band of setae from below spatulate process continuing to near sternum IX; sternum IX with prominent posteromesal, heavily sclerotized, spatulate projection. Inferior appendages short, subtriangular, apices broadly rounded and bearing apical setae; inferior appendages apparently fused to base of IXth sternal projection, together forming highly complex structure as in Figs. 1A, C. Processes of subphallic membranes present, membranous, mound-like, setose. Segment X entirely membranous, apex entire, extending beyond apices of dorsolateral processes. Phallus complex; phallobase tubular, elongate, slender, sclerotized; phallicata very lightly sclerotized, dorsally with paired, very lightly sclerotized, semi-membranous lobes; apicoventral phallic membranes with paired, broad, lightly sclerotized, dorsolateral plates, membranes inflated anteroventrally, apically with lightly sclerotized scale-like structure, shallowly excavate at apex; phallotremal sclerite present, tubular, curved.

Female: Forewing length 6.5 mm (n=1). Color and vestiture as in male. Vaginal apparatus in ventral view elongate, hourglass shaped, narrowest in middle; base widely emarginate, subtriangular; apex trident shaped with wide, paired, sclerotized, slightly sinuate midlateral processes, their apices truncate; single medial process elongate, narrow, sclerotized, about same length as midlateral processes; medial membranes highly convoluted, as approximated in Fig. 4; apical membranes highly convoluted, with heavily sclerotized, rounded, beaklike apical sclerite.

Holotype male: ECUADOR: Napo: Reserva Ecológica Cayambe-Coca, waterfall, rd. to Oyacachi, 0.32621°S, 78.15049°W, el. 3690 m, 16.x.2011, Holzenthal, Ríos Touma, Pita (UMSP000098532) (UMSP). **Paratype:** same data as holotype, 1 female (MECN).

Etymology: This species is named for Lina Pita, lifelong friend of Blanca Ríos-Touma and one of the collectors of this species.

***Contulma quito*, new species, Holzenthal, Ríos-Touma, Rázuri-Gonzales**

148 LSID urn:lsid:zoobank.org:act:97969526-CF58-48A8-8548-8817326D04A0

149

150 Figures 2A-F

151

152 **Diagnosis:** This new species is very similar to *C. nevada* of the group of species mentioned
 153 above (*C. bacula*, *C. caldensis*, *C. cataracta*, *C. echinata*, *C. lina*, *C. nevada*, *C. paluguillensis*,
 154 *C. papallacta*, *C. spinosa*). *Contulma quito*, n. sp., and *C. nevada* have similar short dorsolateral
 155 processes on segment IX, a small setal patch or setal bearing process below the dorsolateral
 156 process, emarginate sternal projection of sternum IX, and short spines in the phallus. The major
 157 differences between *C. quito* and *C. nevada* are the broader excavation of the IXth sternal
 158 projection and the many more spine-like setae in the phallus of *C. quito*.

159 **Description:** *Male:* Forewing length 5.5 mm (n=3). Forewing color brown, with small patch
 160 of cream colored hairs at apex of subcosta and arculus, vestiture intact. Male genitalia: Segment
 161 IX very short dorsally, narrow, deeply excavate mesally; in lateral view, IX quadrate, slightly
 162 extended anterolaterally; posteriorly with very short, narrow, dorsolateral, spatulate process;
 163 posterior margin of IX produced medially to form broad, prominent, broadly rounded, heavily
 164 setose, paired lateral lobe; lobes widely separated ventrally; mesal face of segment IX with short,
 165 sclerotized, setose process below spatulate process. [*Variation:* in male paratype
 166 UMSP000148995, the spatulate processes and the short setose processes below them are each
 167 fused on both sides of the specimen to form a thin, crecentic shelf (Fig. 2F). In male paratype
 168 UMSP000148996, the spatulate processes and the setose processes below them are variously
 169 slightly longer or shorter on either the right or left sides than they are in the holotype]. Sternum
 170 IX with prominent posteromesal, heavily sclerotized, strongly emarginate projection. Inferior
 171 appendages short, crescentic, apices rounded and bearing apical setae; inferior appendages
 172 apparently fused to base of IXth sternal projection, together forming highly complex structure as
 173 in Figs. 2A, C. Processes of subphallic membranes present, membranous, mound-like, setose.
 174 Segment X entirely membranous, apex divided, with lightly sclerotized lateral flanges. Phallus
 175 complex; phallobase tubular, elongate, slender, sclerotized; phallicata very lightly sclerotized,
 176 dorsally with paired, very lightly sclerotized, semi-membranous lobes; apicoventral phallic
 177 membranes with paired, broad, lightly sclerotized, dorsolateral plates forming broadly rounded
 178 trough, apicoventrally with paired patches of numerous (ca. 50) very short spine-like setae;

179 phallotremal sclerite very lightly sclerotized, difficult to discern.

180 *Female*: Unknown.

181 **Holotype male: ECUADOR: Pichincha:** Distrito Metropolitano de Quito, Quebrada
182 Guapalito, 0.40113°S, 78.38378°W, el. 2807 m, 26.vii.2015, Rázuri, Ríos-Touma, Amigo
183 (UMSP000148997) (UMSP). **Paratypes: ECUADOR: Pichincha:** Distrito Metropolitano de
184 Quito, Quebrada Convalecencia, 0.40060°S, 78.38256°W, el. 2813 m, 26.vii.2015, Morabowen,
185 Hernández, 2 males (UMSP, MECN).

186 **Etymology:** Named for Quito, the capital of Ecuador, where the species was collected and
187 where the surrounding Andes seem to harbor a multitude of species in the genus.

188

189 ***Contulma sangay*, new species, Holzenthal, Ríos-Touma, Rázuri-Gonzales**
190 LSID urn:lsid:zoobank.org:act:38C17810-F31C-45A5-BFAC-3D3C5E68AD47

191

192 Figures 3A-D, 5

193

194 **Diagnosis:** This new species is similar to *C. lanceolata*, also from Ecuador. Both share a
195 similar shape of segment IX, which is strongly extended anterolaterally and with the posterior
196 portion excavated medially in both species. In addition, both have an elongate dorsolateral
197 lanceolate process on segment IX, but in *C. sangay* the process is divided apically into a pair of
198 narrow, terete lobes. Most distinctively, the new species lacks the ventrolateral process of
199 segment IX seen in *C. lanceolata* (Holzenthal and Flint, 1995, fig. 70).

200 **Description:** *Male*: Forewing length 4.5 mm (n=1). Forewing color brown, immaculate,
201 vestiture intact. Male genitalia: Segment IX short dorsally, narrow; in lateral view, IX
202 trapezoidal, strongly extended anterolaterally; posteriorly with elongate, dorsolateral, lanceolate
203 processs, apically divided into pair of terete, narrow, closely appressed lobes; process bearing
204 minute setae dorsally along length; posterior margin of IX excavated medially, produced
205 ventrally, to form prominent, semiquadrate, setose, paired lateral lobes; lobes close together
206 ventrally, forming acute separation; mesal face of segment IX without setae or processes;
207 sternum IX short, subquadrate projection, its surface rugose and with very small, mesal
208 sclerotized spur. Inferior appendages very short, crescentric, apices subacute, directed ventrally,

and bearing apical setae; inferior appendages apparently fused to base of IXth sternal projection, together forming highly complex structure as in Figs. 3A, C. Processes of subphallic membranes absent (or not apparent). Segment X entirely membranous, apex cleft, extending to apices of dorsolateral processes. Phallus complex; phallobase tubular, elongate, slender, sclerotized; phallicata very lightly sclerotized, dorsally with prominent membranous lobe; apicoventral phallic membranes with paired, broad, lightly sclerotized, ventrolateral plates, fused ventrally and forming apparent pore, apex with pair of small papillate lobes; phallotremal sclerite present, small, subquadrate.

Female: Forewing length 4.5 mm (n=1). Color as in male, but vestiture rubbed. Vaginal apparatus in ventral view short, oval, widest in middle; base rounded, urn-shaped; apex trident shaped with narrow, paired, lightly sclerotized, indistinct midlateral processes, their apices acute; single medial process elongate, narrow, sclerotized, longer than midlateral processes; medial membranes highly convoluted, as approximated in Fig. 5; apical membranes highly convoluted, with lightly sclerotized, broad, thin, shelf-like apical sclerite.

Holotype male: ECUADOR: Morona-Santiago: Río Salado, Highway E46 (via Riobamba-Macas), 2.24253°S, 78.27791°W, el. 1646 m, 26.i.2015, Holzenthal, Huisman, Ríos-Touma, Amigo (UMSP000147014) (UMSP). **Paratype:** same data as holotype, 1 female (MECN).

Etymology: Named for the type locality, Sangay National Park, where the species was collected from the Río Salado.

New Provincial Records

Contulma bacula Holzenthal and Flint, 1995:11 [Type locality: Ecuador, Napo, 1 mi E of Pa-pallacta; type depository: NMNH; holotype ♂]. —Medellín et al., 2004:201 [distribution; biology]. —Holzenthal and Calor, 2017:21 [catalog].

Distribution. Colombia, Ecuador.

New provincial record: ECUADOR: Morona-Santiago: Río Tinguichaca; Highway E46 (via Riobamba-Macas), 02.21474°S 078.44218°W, el. 2772 m, 25.i.2015, Holzenthal, Huisman, Ríos-Touma, Amigo, 1 male, 2 females (UMSP).

238

239 *Contulma cataracta* Holzenthal and Flint, 1995:12 [Type locality: Ecuador, Napo, Río Maspa
240 Chico, 2 km W Cuyuja; type depository: NMNH; holotype ♂]. —Holzenthal and Calor,
241 2017:22 [catalog].

242 **Distribution.** Ecuador.

243 **New provincial record: ECUADOR: Morona-Santiago:** Sangay National Park, waterfall 2,
244 2.18111°S, 78.5062°W, el. 3516 m, 12.xi.2015, Ríos-Touma, Thomson, Amigo, 3 males, 1
245 female (UMSP).

246

247 *Contulma echinata* Holzenthal and Flint, 1995:15 [Type locality: Colombia, Caldas, 5 km W
248 Termales de Ruíz; type depository: NMNH; holotype ♂; ♀]. —Muñoz-Quesada, 2000:274
249 [checklist]. —Holzenthal and Calor, 2017:22 [catalog].

250 **Distribution.** Colombia, Ecuador.

251 **New provincial record: ECUADOR: Napo:** Reserva Ecológica Cayambe-Coca, waterfall, rd.
252 to Oyacachi, 0.32621°S, 78.15049°W, el. 3690 m, 16.x.2011, Holzenthal, Ríos-Touma, Pita, 1
253 male (UMSP); same, except 26.ii.2012, Ríos-Touma & Pita, 1 male (UMSP).

254

255 **New synonymy**

256

257 *Contulma penai* Holzenthal and Flint, 1995:18 [Type locality: Ecuador, Zamora-Chinchipe, 30
258 km E Loja; type depository: NMNH; holotype ♂; ♀; larva]. —Muñoz-Quesada, 2000:274
259 [checklist]. —Holzenthal and Calor, 2017:23 [catalog].

260 —*Contulma duffi* Oláh, 2016:169 [Type locality: Colombia, Antioquia, Dusky Starfrontlet Bird
261 Reserve, Cordillera Occidental, Urrao, 6°25'N, 75°05'W, 9.ii.2014, caught by hand, leg. A.G.
262 Duff; type depository: private collection of J. Oláh; holotype ♂] **NEW SYNONYM**

263 **Distribution.** Colombia, Ecuador.

264

265 *Contulma duffi*, described from Antioquia, Colombia, fits clearly within the variation we have
266 seen in the species *C. penai*, which also occurs in Antioquia, Colombia, and Ecuador, where it is
267 rather common compared to other species in the genus. There are no features illustrated or
268 diagnosed in the original description that distinguish the species from *C. penai*. Most of the

characters discussed by Oláh are those common to the genus as a whole. The membranous structure of tergum X is identical in the 2 species. The dorsolateral processes of segment IX are illustrated as slightly curved in *C. duffi* compared to those illustrated for *C. penai*, but this slight difference is variable, and the inferior appendages are identical in the 2 species. In Oláh's diagnosis, the species is said to be most similar to *C. bacula*, but it shares little in common with that species. The inferior appendages, the posteromesal process of sternum IX, and the phallus are completely different between the 2 species. Further, the species is known from the holotype alone and is housed in the private collection of the author, making it unavailable to the scientific community.

Key to Males of *Contulma* Species¹

1. Posterior margin of segment IX with dorsolateral processes (HF 24, 44, 58, 79, 90, 107) (these processes are short, but present in *C. quito*, Fig. 2A, B)..... 2
- Posterior margin of segment IX without dorsolateral processes, although there may be small setose projections or patches of setae (HF 40, 49, 86, 95, 112)..... 21
- 2(1). Dorsolateral processes of segment IX very long, slender, strongly down curved, their apices rugose (HF 107, HR 2A, 4A)..... 3
- Dorsolateral processes of segment IX shorter and/or differently shaped, their apices not rugose..... 6
- 3(2). Phallus with pair of large, highly membranous convoluted dorsolateral lobes (in some specimens these lobes may not evert during the clearing process) (HR 2D-F, 4D-E) 4
- Phallus without highly membranous convoluted lobes or with much smaller, less well-developed lobes (HF 110, JN 5) 5
- 4(3). Segment IX extended anterodorsally (HR 4A); setose lobe of posterior margin of

¹ Citations of previously published illustrations referenced in the key are abbreviated as follows: HF=Holzenthal and Flint 1995, HR= Holzenthal and Roberston 2006, JN=Jardim and Nessimian 2011, HRT=Holzenthal and Ríos-Touma 2012.

294		segment IX situated close to middle of segment (HR 4A).....	<i>Contulma tripui</i>
295		Segment IX only slightly extended anterodorsally, if at all (HR 2A); setose lobe of	
296		posterior margin of segment IX situated close to ventral margin of segment (HR 2A) ..	
297		<i>Contulma fluminensis</i>
298	5(3).	Posterior margin of segment IX extended into a long, narrow, acute, setose lobe (JN	
299		1).....	<i>Contulma sana</i>
300		Posterior margin of segment IX extended into a much shorter, subtriangular, setose	
301		lobe (HF 107)	<i>Contulma tijuca</i>
302	6(2).	Phallus with pair of large, highly membranous convoluted dorsolateral lobes (in some	
303		specimens these lobes may not evert during the clearing process) (HF 28, 33, 56, 93) ..	
304		7
305		Phallus without highly membranous convoluted lobes or with much smaller, less	
306		well-developed lobes	10
307	7(6).	Segment IX with pair of elongate, heavily setose, ventrolateral lobes (HF 24, 26)	
308		<i>Contulma adamsae</i>
309		Segment IX without such lobes	8
310	8(7).	Dorsolateral processes of segment IX, in lateral view, long, linear, directed ventrad	
311		along entire length (sinuous and crossing apically in dorsal view) (HF 52-53);	
312		segment IX very short dorsolaterally (HF 52); segment X membranous, almost	
313		obliterated (HF 53).....	<i>Contulma cranifer</i>
314		Dorsolateral processes of segment IX, in lateral view, much shorter, linear to lorate,	
315		directed posteriad, apex only slightly to strongly directed ventrad (straight and not	
316		crossing apically in dorsal view); segment IX long dorsolaterally; segment X	
317		membranous, but well developed	9
318	9(8).	Dorsolateral processes of segment IX, narrow, terete, apex slightly directed ventrad	

319	(HF 30-31).....	<i>Contulma bacula</i>
320	Dorsolateral processes of segment IX lorate (strap-shaped with apex flexed), apex	
321	strongly directed ventrad (HF 90).....	<i>Contulma spinosa</i>
322	10(6). Segment IX with mesolateral patch of long or short spine-like setae or with	
323	mesolateral setose protuberances (HF 57, 75, 79; HRT 1B; Figs 1A, B, 2B)	11
324	Segment IX without mesolateral setae or setose protuberances	15
325	11(10). Apex of phallus with pair of large, tooth-like structures and/or with smaller	
326	apicoventral spines (HF 60, 81, Fig. 2D).....	12
327	Apex of phallus without such spines, although small setae or papillae may be present .	
328	14
329	12(11). Apex of phallus with pair of large, tooth-like structures and smaller apicoventral	
330	spines (HF 60)	<i>Contulma echinata</i>
331	Apex of phallus with pair of large, tooth-like structures only (HF 81)	
332	<i>Contulma papallacta</i>
333	Apex of phallus with smaller apicoventral spines only (HF 77; Figs 2D, E)	13
334	13(12). Projection of sternum IX subtriangular in ventral view, with broad, angulate mesal	
335	excavation (Fig 2C).....	<i>Contulma quito, new species</i>
336	Projection of sternum IX quadrate in ventral view, with narrow mesal excavation (HF	
337	76).....	<i>Contulma nevada</i>
338	14(11). Mesolateral setae of segment IX forming sinuous band from below dorsolateral	
339	process continuing to near sternum IX (Figs 1A, B); projection of sternum IX	
340	rounded (entire) apically (Fig. 1C).....	<i>Contulma lina, new species</i>
341	Mesolateral setae of segment IX restricted to patch below dorsolateral process (HRT	
342	1A, B); projection of sternum IX emarginate apically (HRT 1C)	

343	<i>Contulma paluguillensis</i>
344	15(10). Segment IX extended anterodorsally (HF 44, 61, 70, 82; HR 1A; Fig. 3A); inferior	
345	appendage, in lateral view, crescentic.....	16
346	Segment IX only slightly extended anterodorsally, if at all (HF 66); inferior	
347	appendage, in lateral view, subtriangular or quadrate (HF 66).....	<i>Contulma inornata</i>
348	16(15). Apex of phallus with pair of large, tooth-like structures (HF 64, 65); tergum IX with	
349	prominent, porsteromesal extension (HF 61, 62).....	<i>Contulma ecuadorensis</i>
350	Apex of phallus without such spines, although small setae or papillae may be present;	
351	tergum IX without posteromesal extension	17
352	17(16). Dorsolateral processes of segment IX clothed with short, fine setae (HF 44, 71, 82;	
353	Figs 3A, B).....	18
354	Dorsolateral processes of segment IX lacking vestiture of fine setae (HR 1A, B).....	
355	<i>Contulma boliviensis</i>
356	18(17). Dorsolateral processes of segment IX curved mesally, widely separated, subtriangular	
357	(HF 44-46).....	<i>Contulma colombiensis</i>
358	Dorsolateral processes of segment IX straight, parallel, directed downward, close	
359	together, lanceolate	19
360	19(18). Segment IX posteriorly with only single long dorsolateral or lateral setose process or	
361	none at all.....	20
362	Segment IX posteriorly with both long dorsolateral setose process and long, slender,	
363	sharply pointed, ventrolateral process (HF 70-72).....	<i>Contulma lanceolata</i>
364	20(19). Apex of dorsolateral processes of segment IX entire (HF 82, 83).....	<i>Contulma penai</i>
365	Apex of dorsolateral processes of segment IX divided into pair of terete, narrow,	
366	closely appressed lobes (Figs 3A, B)	<i>Contulma sangay, new species</i>

367	21(1).	Segment X with lightly sclerotized lateral regions that apparently articulate basally	
368		with sclerotized projections of dorsolateral corners of segment IX (HF 48-49, 103-	
369		104).....	22
370		Segment X without lightly sclerotized lateral regions that apparently articulate basally	
371		with sclerotized projections of dorsolateral corners of segment IX	23
372	22(21).	Segment IX with mesolateral patch of short spine-like setae (HF 48-49); projection of	
373		sternum IX truncate (HF 50); parameres long, slender (HF 51).....	
374		<i>Contulma costaricensis</i>
375		Segment IX without mesolateral setae (HF 104); projection of sternum IX acute	
376		apically (HF105); parameres short (HF 106)	<i>Contulma tica</i>
377	23(21).	Phallus with pair of large, highly membranous convoluted dorsolateral lobes (in some	
378		specimens these lobes may not evert during the clearing process) (HF 42-43, 89, 114-	
379		115).....	24
380		Phallus without highly membranous convoluted lobes or with much smaller, less	
381		well-developed lobes	26
382	24(23).	Segment IX with mesolateral patch of long (HF 39-40) or short (HF 86) spine-like	
383		setae.....	25
384		Segment IX without mesolateral setae or setoseprotuberances (HF 111-112)	
385		<i>Contulma valverdei</i>
386	25(24).	Dorsolateral membranous lobes of phallus each ending in sclerotized, scale-like	
387		process (HF 42-43); pojection of sternum IX long, rounded (entire) apically (HF 41)...	
388		<i>Contulma cataracta</i>
389		Dorsolateral membranous lobes of phallus entirely membranous (HF 89); projection	
390		of sternum IX short, slightly emarginate apically.....	<i>Contulma sancta</i>
391	26(23).	Segment IX with mesolateral patch of long or shorter spine-like setae (HF 95, 100) or	

392	with mesolateral setose protuberances (HF 36); inferior appendage, in lateral view,	
393	crescentic	27
394	Segment IX without mesolateral setae or setose protuberances (HR 3A, B); inferior	
395	appendage, in lateral view, subtriangular (HR 3A)	<i>Contulma meloi</i> ²
396	27(26). Apex of phallus with large, apicoventral spines (HF 97-98); parameres absent	
397	<i>Contulma talamanca</i>
398	Apex of phallus without such spines; parameres present, long, slender (HF 38, 102) 28	
399	28(27). Segment IX with patch of long, stout, spine-like mesolateral setae (HF 100);	
400	phallicata with parameres and lateral flanges (HF 102)	<i>Contulma tapanti</i>
401	Segment IX without patch of long, stout, spine-like mesolateral setae, although	
402	smaller setae may be present (HF 36); phallicata with parameres only, lateral flanges	
403	absent (HF 38)	<i>Contulma caldensis</i>

404 Discussion

405 *Contulma* species appear to be highly regionally endemic (Holzenthal and Flint 1995, Holzenthal
 406 and Robertson 2006). We confirm this pattern in our collections in Ecuador. Patterns of high
 407 endemicity in macroinvertebrate benthic larvae are well known in high altitude glacial streams;
 408 these habitats are highly threatened by climate change (Jacobsen et al. 2012). On the other hand,
 409 we added records to the Ecuadorian fauna of species that were previously known only from

² *Contulma meloi* is a member of the group of species including *C. fluminensis*, *C. sana*, *C. tijuca*, and *C. tripui* from southeastern Brazil. However, it lacks the very long, slender, strongly downcurved, apically rugose dorsolateral processes of segment IX. The species is known only from the male holotype and a male paratype. In the holotype, there are no dorsolateral processes (HR 3A), but in the paratype, a rudimentary process occurs (HR 3G). However, it shares all other diagnostic features of this group, including the posterior, setose extension of segment IX, the broad, shelf-like structure of sternum IX, including the flat, tooth-like, apical setae, and the complex, membranous lobes of the phallus. It appears artificially in couplet 26, separated from related species because it lacks the strongly downcurved, apically rugose dorsolateral processes indicated in couplet 2. The collection of additional specimens may prove that the absence of the dorsolateral process is an aberration; if so, the specimens would lead to *C. fluminensis* in the key presented above.

Colombia, demonstrating that some species may have spread across the Andes. Despite this distribution pattern, aquatic insect species, especially mayflies, have shown genetic isolation in the various ranges of the Ecuadorian Andes, even with relatively short distances between populations and with recent volcanic eruption history (Finn et al. 2016). No similar study has yet been performed with any Andean caddisfly species, but we expect a similar pattern of isolation among populations. This isolation could become intensified by the intense land use changes and water pollution occurring throughout the inter-Andean valleys (Ríos-Touma et al., 2014) potentially preventing dispersion among mountain populations and also local extinctions.

Acknowledgements

This study was supported by Universidad de Las Americas project AMB.BRT.17.005 "Diversidad y Distribucion de Trichoptera de Ecuador" and Minnesota Agricultural Experiment Station projects MIN17-017 and 17-029. ERG was funded through a doctoral fellowship from Cienciactiva, Consejo Nacional de Ciencia, Tecnologia, e Innovacion Tecnologica Peru (contract 277-2015- FONCECYT). The funders had no role in the study design, data collection and analysis, decision to publish, or preparation of the manuscript.

References

- Blahnik RJ, and Holzenthal RW. 2004. Collection and curation of Trichoptera, with an emphasis on pinned material. *Nectopsyche, Neotropical Trichoptera Newsletter* 1:8-20.
- Blahnik RJ, Holzenthal RW, and Prather AL. 2007. The lactic acid method for clearing Trichoptera genitalia. In: Bueno-Soria J, Barba-Álvarez R, and Armitage BJ, eds. *Proceedings of the 12th International Symposium on Trichoptera*. Columbus, Ohio: The Caddis Press, 9-14.
- Dallwitz MJ. 1980. A general system for coding taxonomic descriptions. *Taxon* 29:41-46.
- Dallwitz MJ, Paine TA, and Zurcher EJ. 1999. User's guide to the DELTA editor. Available from: <http://delta-intkey.com/>.
- de Moor FC, and Ivanov VD. 2008. Global diversity of caddisflies (Trichoptera: Insecta) in freshwater. *Hydrobiologia* 595:393-407.
- Finn DS, Encalada AC, and Hampel H. 2016. Genetic isolation among mountains but not between stream types in a tropical high-altitude mayfly. *Freshwater Biology* 61:702-714. 10.1111/fwb.12740
- Henriques-Oliveira AL, and Santos APM. 2014. Two new species of *Atanatolica* Mosely 1936 (Trichoptera: Leptoceridae) from Peru and Northeastern Brazil. *Zootaxa* 3869:537-547.
- Holzenthal RW. 1985. Studies in Neotropical Leptoceridae (Trichoptera) II: *Amphoropsyche*, a new genus and species of Leptocerinae from northern South America. *International Journal of Entomology* 27:255-269.
- Holzenthal RW. 1988. Studies in Neotropical Leptoceridae (Trichoptera), VIII: the genera *Atanatolica* Mosely and *Grumichella* Müller (Triplectidinae: Grumichellini). *Transactions of the American Entomological Society* 114:71-128.
- Holzenthal RW, and Calor AR. 2017. Catalog of the Neotropical Trichoptera (Caddisflies). *ZooKeys* 654:1-566. 10.3897/zookeys.654.9516
- Holzenthal RW, and Flint OS, Jr. 1995. Studies of Neotropical caddisflies, LI: systematics of the Neotropical caddisfly genus *Contulma* (Trichoptera: Anomalopsychidae). *Smithsonian Contributions to Zoology* 575:1-59.
- Holzenthal RW, and Rázuri-Gonzales LE. 2011. A new species of *Amphoropsyche* (Trichoptera, Leptoceridae) from Ecuador, with a key to the species in the genus. *ZooKeys* 211:59-65.

458 Holzenthal RW, and Ríos-Touma B. 2012. *Contulma paluguillensis* (Trichoptera:
459 Anomalopsychidae), a new caddisfly from the high Andes of Ecuador, and its natural
460 history. *Freshwater Science* 31:442-450. 10.1899/11-067.1

461 Holzenthal RW, and Robertson DR. 2006. Four new species of *Contulma* from South America
462 (Trichoptera: Anomalopsychidae). *Zootaxa* 1355:49-59.

463 Jacobsen D, Milner AM, Brown LE, and Dangles O. 2012. Biodiversity under threat in glacier-
464 fed river systems. *Nature Climate Change* 2:361-364.

465 Jardim GA, and Nessimian JL. 2011. A new species of *Contulma* Flint (Trichoptera,
466 Anomalopsychidae) from southeastern Brazil. *Revista Brasileira de Entomologia* 55:226-
467 228.

468 Medellín C F, Ramírez O M, and Rincón ME. 2004. Trichoptera del Santuario de Iguaque
469 (Boyacá, Colombia) y su relación con la calidad del agua. *Revista Colombiana de*
470 *Entomología* 30:197-203.

471 Muñoz-Quesada F. 2000. Especies del orden Trichoptera (Insecta) en Colombia [Colombian
472 species of the order Trichoptera (Insecta)]. *Biota Colombiana* 1:267-288.

473 Oláh J. 2016. New species and records of Trichoptera collected by Mr. A. G. Duff. More
474 phenomics and less genomics! *Opuscula Zoologica (Budapest)* 47:155-171.
475 <http://doi.org/10.5281/zenodo.167120>

476 Ríos-Touma B, Acosta R, and Prat N. 2014. The Andean Biotic Index (ABI): revised tolerance
477 to pollution values for macroinvertebrate families and index performance evaluation.
478 *Revista de Biología Tropical* 62 (suppl 2):249-273.

479 Ríos-Touma B, Holzenthal RW, Huisman J, Thomson R, and Rázuri-Gonzales E. 2017.
480 Diversity and distribution of the Caddisflies (Insecta: Trichoptera) of Ecuador. *PeerJ*
481 5:e2851. 10.7717/peerj.2851

Figure 1(on next page)

Contulma lina, figure 1

Catalog of the Neotropical Trichoptera (Caddisflies) Figure 1. **Male genitalia of *Contulma lina*, new species.** (A) segments IX and X, lateral (base of phallus indicated in crosshatch). (B) segments IX and X, dorsal. (C) segment IX, ventral. (D) phallus, lateral. (E) phallus apex, dorsal. Abbreviations: IX = abdominal segment IX; X = abdominal segment X.




Figure 2(on next page)

Contulma quito, figure 2

Catalog of the Neotropical Trichoptera (Caddisflies) Figure 2. **Male genitalia of *Contulma quito*, new species.** (A) segments IX and X, lateral (base of phallus indicated in crosshatch). (B) segments XI and X, dorsal. (C) segment IX, ventral. (D) phallus, lateral. (E) phallus apex, dorsal. (F) segments IX and X, dorsal, variation in paratype specimen UMSP000148995.




Figure 3(on next page)

Contulma sangay, figure 3

Catalog of the Neotropical Trichoptera (Caddisflies) Figure 3. **Male genitalia of *Contulma sangay*, new species.** (A) segments IX and X, lateral (base of phallus indicated in crosshatch); inset: apex of inferior appendage, caudal. (B) segments XI and X, dorsal. (C) segment IX, ventral. (D) phallus, lateral; insets: details of phallic structures.




Figure 4(on next page)

Contulma females, figures 4-5

Catalog of the Neotropical Trichoptera (Caddisflies) Figures 4 and 5. **Female vaginal apparatus of *Contulma*, new species, ventral.** (4) *C. lina*, new species. (5) *C. sangay*, new species.

