

Unanticipated discovery of two rare gastropod molluscs from recently located hydrothermally influenced areas in the Okinawa Trough

Chong Chen ^{Corresp., 1}, Hiromi Kayama Watanabe ^{2, 3, 4}, Junichi Miyazaki ^{1, 3, 4}, Shinsuke Kawagucci ^{1, 3, 4}

- ¹ Department of Subsurface Geobiological Analysis and Research, Japan Agency for Marine-Earth Science and Technology, Yokosuka, Kanagawa, Japan
- ² Department of Marine Biodiversity Research, Japan Agency for Marine-Earth Science and Technology, Yokosuka, Kanagawa, Japan
- ³ Research and Development Center for Submarine Resources, Japan Agency for Marine-Earth Science and Technology, Yokosuka, Kanagawa, Japan
- ⁴ Project Team for Development of New-generation Research Protocol for Submarine Resources, Japan Agency for Marine-Earth Science and Technology, Yokosuka, Kanagawa, Japan

Corresponding Author: Chong Chen
Email address: cchen@jamstec.go.jp

Background. Deep-sea hydrothermal vent is one of the most ‘extreme’ environments in the marine realm. Few species are capable of inhabiting such ecosystems, despite extremely high productivity there supported by microbial chemosynthesis, leading to high biomass and low species richness. Although gastropod molluscs are one of the main constituents of megafaunal communities at vent ecosystems, most species belong to several typical families (e.g., Provannidae, Peltospiridae, Lepetodrilidae) specialised and adapted to life at vents.

Methods. During recent surveys of Okinawa Trough hydrothermal vent systems, two snails atypical of vent ecosystems were unexpectedly found in newly discovered hydrothermally influenced areas. Shell and radular characteristics were used to identify the gastropods morphologically.

Results. One species was a vetigastropod, the calliostomatid *Tristichotrochus ikukoe* (Sakurai, 1994); and the other was a caenogastropod, the muricid *Abyssotrophon soyoae* (Okutani, 1959). Both gastropods were previously only known from regular non-chemosynthetic deep-sea and very rare – only two published records exist for each species. The radula formula of *Tristichotrochus ikukoe* is accurately reported for the first time and based on that it is returned to genus *Otukaia*. For both species, barcode sequences of the cytochrome c oxidase I (COI) gene were obtained and deposited for future references.

Discussion. These new records represent the second record of calliostomatids from vents (third from chemosynthetic ecosystems) and the third record of muricids from vents (tenth from chemosynthetic ecosystems), and extend the distribution of both species to the southwest. Neither family has been recorded at chemosynthetic ecosystems in the western Pacific. Both were from weakly diffuse flow areas not subject to high temperature venting but were nevertheless associated with typical vent-reliant taxa such as *Lamellibrachia* tubeworms and *Bathymodiolus* mussels. These new records show that these species are capable of tolerating environmental stress associated with weak hydrothermally influenced areas, despite not being vent endemic species, adding to the list of known vent/non-vent species intersections. This signify that such weakly influenced areas may provide key habitats for them, and that such areas may play a role in the evolution of biological adaptations to ‘extreme’ chemosynthetic ecosystems.

Research Article

Unanticipated discovery of two rare gastropod molluscs from recently located hydrothermally influenced areas in the Okinawa Trough

Chong Chen^{1*}, Hiromi Kayama Watanabe^{2,3,4}, Junichi Miyazaki^{1,3,4}, Shinsuke Kawagucci^{1,3,4}

¹Department of Subsurface Geobiological Analysis and Research (D-SUGAR), Japan Agency for Marine-Earth Science and Technology (JAMSTEC), 2-15 Natsushima, Yokosuka 237-0061, Japan

²Department of Marine Biodiversity Research (BIO-DIVE), Japan Agency for Marine-Earth Science and Technology (JAMSTEC), 2-15 Natsushima-cho, 237-0061 Yokosuka, Japan

³Research and Development Center for Submarine Resources, Japan Agency for Marine-Earth Science and Technology (JAMSTEC), 2-15 Natsushima-cho, 237-0061 Yokosuka, Japan

⁴Project Team for Development of New-generation Research Protocol for Submarine Resources, Japan Agency for Marine-Earth Science and Technology (JAMSTEC), 2-15 Natsushima-cho, 237-0061 Yokosuka, Japan

*Corresponding author. Tel: +81-(0)46-867-9717; E-mail: cchen@jamstec.go.jp

Short title: Two gastropods newly recorded from vents

17

Abstract

18 **Background.** Deep-sea hydrothermal vent is one of the most ‘extreme’ environments in the
19 marine realm. Few species are capable of inhabiting such ecosystems, despite extremely high
20 productivity there supported by microbial chemosynthesis, leading to high biomass and low
21 species richness. Although gastropod molluscs are one of the main constituents of megafaunal
22 communities at vent ecosystems, most species belong to several typical families (e.g.,
23 Provannidae, Peltospiridae, Lepetodrilidae) specialised and adapted to life at vents.

24 **Methods.** During recent surveys of Okinawa Trough hydrothermal vent systems, two snails
25 atypical of vent ecosystems were unexpectedly found in newly discovered hydrothermally
26 influenced areas. Shell and radular characteristics were used to identify the gastropods
27 morphologically.

28 **Results.** One species was a vetigastropod, the calliostomatid *Tristichotrochus ikukoe* (Sakurai,
29 1994); and the other was a caenogastropod, the muricid *Abyssotrophon soyoe* (Okutani, 1959).
30 Both gastropods were previously only known from regular non-chemosynthetic deep-sea and
31 very rare – only two published records exist for each species. The radula formula of
32 *Tristichotrochus ikukoe* is accurately reported for the first time and based on that it is returned to
33 genus *Otukaia*. For both species, barcode sequences of the cytochrome *c* oxidase I (COI) gene
34 were obtained and deposited for future references.

35 **Discussion.** These new records represent the second record of calliostomatids from vents (third
36 from chemosynthetic ecosystems) and the third record of muricids from vents (tenth from
37 chemosynthetic ecosystems), and extend the distribution of both species to the southwest. Neither
38 family has been recorded at chemosynthetic ecosystems in the western Pacific. Both were from
39 weakly diffuse flow areas not subject to high temperature venting but were nevertheless
40 associated with typical vent-reliant taxa such as *Lamellibrachia* tubeworms and *Bathymodiolus*
41 mussels. These new records show that these species are capable of tolerating environmental stress
42 associated with weak hydrothermally influenced areas, despite not being vent endemic species,
43 adding to the list of known vent/non-vent species intersections. This signify that such weakly
44 influenced areas may provide key habitats for them, and that such areas may play a role in the
45 evolution of biological adaptations to ‘extreme’ chemosynthetic ecosystems.

46

Introduction

Deep-sea hydrothermal vents exhibit increased water temperature, higher heavy metal concentrations, and raised concentrations of substances acting as energy sources for chemosynthesis (such as H₂, CH₄, H₂S) compared to the surrounding water (Van Dover, 2000; Petersen *et al.*, 2011). Due to such 'extreme' environmental conditions, few species are capable of inhabiting such ecosystems, despite extremely high productivity there for the deep-sea supported by microbial chemosynthesis (Grassle, 1987). As a result, although vent ecosystems support biomass comparable even to that of shallow water coral reefs, the macro- and megafaunal communities there are characterised by a low species richness (Lutz & Kennish, 1993; Van Dover & Trask, 2000).

Although gastropod molluscs are one of the main constituents of megafaunal communities at hydrothermal vent ecosystems, species present mostly to belong to one of several typical families (e.g., Provannidae, Peltospiridae, Lepetodrilidae) specialised and adapted to life at vent and other chemosynthetic ecosystems such as hydrocarbon seeps and organic falls (Sasaki *et al.*, 2010). Here, we report new records of two gastropod species, *Tristichotrochus ikukoe* (Sakurai, 1994) (Vetigastropoda: Calliostomatidae) and *Abyssotrophon soyoae* (Okutani, 1959) (Caenogastropoda: Muricidae), from two new diffuse flow hydrothermal sites discovered using the multibeam echo-sounding (MBES) method (as outlined in Nakamura *et al.*, 2015). Both species are exceedingly rare endemics of Japan, with only two published records each.

Calliostomatidae is an exclusively marine family of usually grazing gastropods which comprises about 300 extant species (Marshall, 1995; Vilvens, 2014; Marshall, 2016). Only two species have been recorded from chemosynthetic ecosystems, one from vent and one from seep (Sasaki *et al.*, 2010). Muricidae is a large family of carnivorous and generally predatory marine caenogastropods comprising about 1600 living species (Merle *et al.*, 2011). A total of seven muricids have been recorded from hydrocarbon seeps, but none from vents (Sasaki *et al.*, 2010) and two species (*Enixotrophon carduelis* (Watson, 1882) and *E. obtusus* (Marshall & Houart, 2011)) were recorded from Rumble V Volcano which is a hydrothermally influenced area on the Kermadec Ridge (Marshall & Houart, 2011). The present study, therefore, presents the second record of Calliostomatidae and the third record of Muricidae from hydrothermal vent ecosystems.

The two species newly recorded from vents herein are unlikely to be vent endemics but are clearly interacting with vent environment and other vent endemic fauna. These records add to the intersection of vent endemic and non-vent endemic species in the peripheral diffuse flow environment, where non-endemic species have often been recorded due to the much less stressful

environment compared to focused flow venting (reviewed in Levin *et al.*, 2016).

Materials & Methods

Potential signals of hydrothermal activity were detected in the Okinawa Trough by detecting acoustic water column anomalies, probably derived from CO₂ bubbles discharging from vents, using an EM122 (Kongsberg Maritime) MBES system on R/V *Yokosuka* (cruises YK14-16 and YK16-07), as outlined in Nakamura *et al.* (2015). The signals were then ground-truthed with Remotely Operated Vehicle (ROV) *KAIKO* (with vehicle *Mk-IV*) dives on-board R/V *Kairei* during cruises KR15-16 (Kawagucci *et al.*, 2015; Makabe *et al.*, 2016) and KR16-16 (Miyazaki *et al.*, 2016; Makabe *et al.*, 2017). A suction sampler equipped on the ROV *KAIKO* was used to collect the gastropods. A digital camera mounted on the same ROV was used to shoot the *in situ* observation images.

Upon recovery on to the research vessel, the specimens were fixed and stored in 99% ethanol for subsequent studies. Amplification and sequencing of the cytochrome *c* oxidase I (COI) barcode gene was done using a piece of foot using the universal primer pair LCO1490 and HCO2198 (Folmer *et al.*, 1994). Molecular methods follow that outlined in Chen *et al.* (2016). After taking tissue snips, soft parts were extracted from the specimens after rehydrating in MilliQ. The radula was dissected out from each specimen under a dissecting microscope (Olympus SZX1) and cleaned by washing in diluted commercial bleach followed by two washes, in MilliQ water and 99% ethanol each. The cleaned radulae were mounted on SEM stubs with carbon tapes and examined uncoated at 15kV with a table top scanning electron microscope (SEM; Hitachi TM-3000).

Results and Discussion

Detection and discovery of hydrothermal sites

One acoustic anomaly (28° 26.1' N, 128° 11.5' E; Fig. 1) was detected in the northern Okinawa Trough near Tokara Islands, in the 'Higashi-Ensei' area (Makabe *et al.*, 2017) east-northeast of Minami-Ensei Knoll (Hashimoto *et al.*, 1995). This site was explored on dive #724 of ROV *Kaiko*, which discovered a weakly active hydrothermal vent site mostly with diffuse flow areas. This site was 955–1180 m deep, dominated by dense *Lamellibrachia* tubeworm bushes grown

over by the barnacles *Ashinkailepas seepiophila* Yamaguchi, Newman & Hashimoto, 2004 and *Leucolepas longa* Southward & Jones, 2003, and was named ‘Fukai’ (Fig. 2A; for details see Miyazaki *et al.*, 2016; Makabe *et al.*, 2017). Other fauna typical of Okinawa Trough hydrothermal vents (Watanabe & Kojima, 2015), such as the squat lobster *Shinkaia crosnieri* Baba & Williams, 1998, the vesicomid clam *Akebiconcha kawamurai* Kuroda, 1943, and the mussel *Bathymodiolus platifrons* Hashimoto & Okutani, 1994 were also present in this site. Another signal (25° 4.5’ N, 124° 31.0’ E; Fig. 1) was located about 1950–1990 m deep on the northern slope of Tarama Hill in the southern Okinawa Trough, and was surveyed during ROV *KAIKO* dive #669 (for details see Makabe *et al.*, 2016). A small weakly hydrothermally influenced site dominated by white branching Poecilosclerida sponges (*sensu* Ise, 2012), *Alaysia* sp. and *Lamellibrachia* sp. tubeworms, and *Bathymodiolus aduloideus* Hashimoto & Okutani, 1994 mussels was found and named ‘Crane’ (Fig. 2C; Makabe *et al.*, 2017).

***Tristichotrochus ikukoe* (Sakurai, 1994)**

One specimen of a calliostomatid gastropod was collected from Fukai site, Higashi-Ensei, Okinawa Trough (28° 26.1064’ N, 124° 11.4889’ E, 962 m deep, R/V *Kairei* cruise KR16-16, ROV *KAIKO* Dive #724, 2016/xii/04). Based on shell (diameter 22.6 mm, height 21.2 mm; Fig. 3A–D) and radular (Fig. 4A) characteristics this specimen was clearly identifiable as *Tristichotrochus ikukoe* (Sakurai, 1994). To the knowledge of the authors, this is only the third time it has been formally reported in the literature (Higo *et al.*, 1999; Hasegawa, 2009). A total of three individuals were sighted at the same site (Fig. 2B), living attached on *Lamellibrachia* sp. tubeworm tubes or hard surrounding substrata.

The taxonomic history of *Tristichotrochus ikukoe* is rather perplexing. It was initially described from Kochi Prefecture, Japan as *Otukaia ikukoe* Sakurai, 1994. Sakurai compared it to *Otukaia kiheiziebisu* (Otuka, 1939), commenting that *O. ikukoe* is ‘smaller in size, and has obtuser ribs with larger granules’ as well as ‘has only 21 basal ribs in comparison’ to *O. kiheiziebisu* (Sakurai, 1994). Later, Okutani & Iwahori (1992) reported *O. kiheiziebisu* from Tosa Bay, Kochi Prefecture, Japan which extended its distribution range from Sagami Bay. They reported that their specimens were variable in spiral rib development, with some specimens exhibiting five equally strong ribs. Radulae of such ‘unusual’ specimens were illustrated (Okutani & Iwahori, 1992: figs. 7, 10–12), and were actually virtually identical to the holotype of *O. ikukoe*. Following this, Hasegawa & Saito (1995) considered *O. ikukoe* to be a junior synonym of *O. kiheiziebisu*, and Hasegawa (2009) formally synonymised the two species. More recently, however, Marshall (2016) figured the radula of a typical *O. kiheiziebisu* and found it to be clearly distinct from that

figured by Okutani & Iwahori (1992). *Otukaia ikukoe* was reinstated as a valid species and transferred to *Tristichotrochus* based on its fewer lateral teeth compared to *Otukaia* (four to six compared to nine in *O. kiheiziebisu*).

Though the radula of *T. ikukoe* was reported to have only four pairs of lateral teeth (Okutani & Iwahori, 1992) compared to nine in *O. kiheiziebisu* (Marshall, 2016), this was questioned by Marshall (2016) who reasoned that there may actually be as many as five to six pairs. The specimen of *T. ikukoe* investigated in the present study has eight pairs of laterals (Fig. 4A), which is more numerous than Marshall (2016) predicted and makes the radula formula $(n+8+1+8+n)$. Due to only a single specimen being collected it is not possible to comment on the intraspecific variation in the radula of *T. ikukoe* at this point, but this is the most reliable account of the radula characteristics of this rare calliostomatid. Having as many as eight pairs of laterals means the radula of *T. ikukoe* actually fits well in genus *Otukaia*, and it is therefore returned to that genus herein, as *Otukaia ikukoe* Sakurai, 1994.

It should be noted that *O. ikukoe* and *O. kiheiziebisu* are in fact easily distinguished by shell characteristics as well, by the following two points: 1. *O. ikukoe* has five to six strong ribs on the body whorl whereas *O. kiheiziebisu* only has two to three (including the base keel); 2. The ribs of *O. ikukoe* are less sharply raised but more strongly beaded than those of *O. kiheiziebisu*. To assist future identification of these two species based on molecular barcodes, a COI barcode of the present specimen was sequenced and deposited in DNA Data Bank of Japan (DDBJ; accession number LC333137). A sequence of the same gene is already available on GenBank for *O. kiheiziebisu* (from Williams, 2010) under the accession number AB505274, the identity of the individual sequenced has been ascertained and confirmed as *O. kiheiziebisu* (T. Nakano pers. comm., in Marshall, 2016). Based on comparison between these two sequences, the pairwise difference between the two species is 5.8%, which is higher than the average of 3–4% between closely related species in marine gastropods (Meyer & Pauley, 2005).

The size of the present specimen (shell height 21.2 mm) is just slightly smaller than previous reports as shell height of the holotype is 23.9 mm (Sakurai, 1994) and height of specimens definitively referable to *O. ikukoe* reported by Okutani & Iwahori (1992) range between 23.5 mm–26.7 mm.

Since *O. ikukoe* has not been recorded from outside Kochi Prefecture (Higo *et al.*, 1999; Hasegawa, 2009), the present new record extends its distribution range to the southwest by about 700 km. The collection depth of 962 m is deeper than the Tosa Bay record (700 m; Okutani &

Iwahori, 1992), and since the depth that the type specimen was collected from is not known (Sakurai, 1994), this represents the deepest record for the species.

Abyssotrophon soyoae (Okutani, 1959)

Two specimens of trophonine muricids were collected from the Crane site, Tarama Hill, Okinawa Trough (25° 4.5388' N, 124° 31.0103' E, 1973 m deep, R/V *Kairei* cruise KR15-16, ROV *KAIKO* Dive #669, 2015/x/29). A total of five individuals was sighted at this site (Fig. 2D). The morphology of the two specimens matched best, in terms of shell (specimen #1, height 22.2 mm, width 11.0 mm, Fig. 3E–H; specimen #1, height 22.5 mm, width 10.5 mm, Fig. 3I, J) and radular (specimen #1, Fig. 4B) characters, with *Abyssotrophon soyoae* (Okutani, 1959). Originally described from off Miki-Saki, Kumano-nada, Kii Peninsula, Japan (33° 48.8' N, 136° 34.8' E) as *Trophonopsis soyoae* Okutani, 1959, this is only the third time it has been recorded formally (Okutani & Iwahori, 1992; Higo *et al.*, 1999; Tsuchiya, 2017). Furthermore, three individuals that appeared to belong to the same species were seen in Fukai site (Fig. 2B, the same location as *O. ikukoe* collecting data above), but unfortunately none were collected.

Conchologically *A. soyoae* is easily recognised by two strong keels combined with very crowded axial ribs (Tsuchiya, 2017). One significant difference between the present material and previously reported specimens is that the axial ribs are less crowded (about 35 vs about 70; Okutani & Iwahori, 1992). However, there appears to be some variability in rib frequency across life stages (see Fig. 3J; Okutani & Iwahori, 1992: fig. 28) and the authors consider this difference to be intraspecific, rather than interspecific, variability until further evidence becomes available proving otherwise. The specimens examined herein also have higher spires than the holotype (Okutani, 1959), but they match well with specimens previously reported from Tosa Bay, Kochi Prefecture, Japan (Okutani & Iwahori, 1992). The shell height to width ratios were 2.02 and 2.14, which gives an impression that they are more stout than the previous specimens which range between 2.11 – 2.69 (2.11 in the holotype; Okutani & Iwahori, 1992) but only because the earlier whorls are much more corroded in the present specimens, likely due to the acidic hydrothermal environment. The radula is very much the same as imaged by Okutani & Iwahori (1992), including the formula (1+1+1) and in having a single middle cusp on the central tooth. Although the soft parts of the specimens investigated were not of good quality, eyes were present in both individuals and the smaller individual (specimen #1, male) possessed a very large penis as previously reported (Okutani & Iwahori, 1992). To facilitate future identification and genetic comparison, the COI barcode sequence was obtained from specimen #1 and deposited in DDBJ

under accession number LC333138. The closest match available on GenBank is a sequence of *Boreotrophon truncatus* (Strøm, 1768), accession number HQ919183 (from Layton *et al.*, 2014) with 9% pairwise difference.

Since this species has only been reported from the type locality and Tosa Bay (Higo *et al.*, 1999; Tsuchiya, 2017), the present record extends its distribution range over 1250 km to the southwest. The collection depth of the present record (1973 m) is within the bathymetric range between the type locality (2050 m; Okutani, 1959) and the Tosa Bay record (700 – 750 m; Okutani & Iwahori, 1992).

Significance of the new records from vent ecosystems

Trochoid gastropods are commonly found in many hydrothermal vent sites around the globe (e.g., *Itheyaspira*, *Cantrainea*) but calliostomatids have only been recorded twice (Sasaki *et al.*, 2010). These include *Falsimargarita nauduri* Warén and Bouchet 2001 from the 17°S hydrothermal vent site, East Pacific Rise (Warén & Bouchet, 2001) and *Maurea chilena* (Rehder, 1971) from Concepción methane seep off Central Chile (Sellanes *et al.*, 2008). Although Sellanes *et al.* (2008) also reported *Otukaia crustulum* Vilvens & Sellanes, 2006 from the Concepción methane seep, this species has been moved to Calliotropidae, as *Calliotropis crustulum* (Marshall, 2016). The present report of *Otukaia ikukoe* is therefore the second record of calliostomatids from vent ecosystems and the third record from chemosynthetic ecosystems.

Muricids are found from intertidal waters down to over 7000 m deep, and *Abyssotrophon* is one of the deepest-living genera occurring even over 7200 m deep in the case of *Abyssotrophon hadalis* (Sysoev, 1992) (Egorov, 1993). Five species of Muricidae are known from Concepción methane seep off Central Chile (Sellanes *et al.*, 2008) and two are known from methane seeps in Barbados Prism (Warén & Bouchet, 2001), making a total of seven records from hydrocarbon seeps. Except *Pagodula concepcionensis* Houart & Sellanes, 2006 which is in the subfamily Pagodulinae (Houart & Sellanes, 2006), all other six species recorded are in the subfamily Trophoninae (genera *Trophon* and *Coronium*; Sasaki *et al.*, 2010). The present record of *Abyssotrophon soyoae* is the third record of this family from vent ecosystems and the tenth from chemosynthetic ecosystems. Both families were previously unrecorded from chemosynthetic ecosystems of not only the Okinawa Trough but the western Pacific as a whole (Sasaki *et al.*, 2010; Watanabe & Kojima, 2015).

Although the two species reported herein were found in hydrothermally influenced settings, they

are also known from regular deep-sea bottom (Okutani & Iwahori, 1992) and thus not endemic to chemosynthetic ecosystems. Neither new hydrothermally influenced site reported herein is vigorously active, and is instead more like ‘hydrothermal seeps’ (Levin *et al.*, 2012) than conventional high-temperature fluid venting systems. The presence *O. ikukoe* and *A. soyoe* in weakly diffuse flow areas indicate that they are able to tolerate some levels of environmental stresses associated with hydrothermal ecosystems (e.g., increased acidity, concentration of toxic substances and heavy metals), allowing them to invade and benefit from the high productivity of hydrothermal vent ecosystems.

While feeding behaviour was not observed in the field, the ecological roles the two species play may be inferred as follows. Deep-water calliostomatids are mostly carnivorous grazers feeding on cnidarians or sponges (Marshall, 1995), and from the fact that they were observed living on tubeworm tubes *O. ikukoe* in Fukai site probably graze on sponges and/or biofilm growing on the tubes. Muricids are carnivores feeding on other marine invertebrates (Merle *et al.*, 2011), and *A. soyoe* is unlikely to be an exception. In both localities the *Abyssotrophon* individuals were seen in close proximity to other molluscs (e.g., *Bathymodiolus aduloide*s at Crane site) and barnacles (e.g., *Ashinkailepas seepiophila* at Fukai site) which the *Abyssotrophon* likely prey upon. In interacting with vent endemic species in such ways, *O. ikukoe* and *A. soyoe* may contribute to trophic transfer of chemosynthetically derived production from vents to surrounding ambient environment, which some other predators such as seastar, fish, and octopus have been suggested to play a role in (Levin *et al.*, 2016 and references therein). The apparent absence of these two species from other Okinawa Trough vents is possibly an artefact caused by the relatively little effort put into exploring the peripheral areas.

In fact, non-vent endemic deep-water predatory gastropods have been recorded from vent periphery in several occasions, for example *Buccinum viridum* Dall, 1889 on the Juan de Fuca Ridge (Tunnicliffe & Fontaine, 1987; Sarrazin *et al.*, 1997), *Enixotrophon carduelis* and *E. obtusus* on the Kermadec Ridge (Marshall & Houart, 2011), and *Neptunea robusta* Okutani, 1964 in the Okinawa Trough (Watanabe & Kojima, 2015). Their success may be a combined effect of the ability to tolerate moderate environmental stress from vent periphery and the fact that few vent endemic gastropod species are predators, meaning there is an available niche for these predatory species given the high productivity. In the Okinawa Trough vents, for instance, only one vent endemic predatory gastropod, *Thermosipho desbruyeresi* (Okutani & Ohta, 1993), is known (Watanabe & Kojima, 2015).

The present records signify that outskirts of hydrothermally influenced areas likely provide

important habitats and energy resources for some non-endemic species (Levin *et al.*, 2016). These peripheral areas may be important for the evolution of biological adaptations to chemosynthetic ecosystems by providing the intermediate environment between ‘extreme’ vents and the regular surrounding seafloor.

Conclusions

Exploration of two new hydrothermally influenced ecosystems in the Okinawa Trough led to the collection of two gastropod snails previously unknown from chemosynthetic ecosystems. One was *Otukaia ikukoe*, the second calliostomatid ever recorded from vents and the third record of the family for chemosynthetic ecosystems overall. The other was the muricid *Abyssotrophon soyoae*, marking the third time Muricidae has been found at hydrothermal vents and the tenth record at chemosynthetic ecosystems. Both species are extremely rare and this report represents the third record for each species in the literature as well as the first time they are seen in their natural habitat. Neither family has been recorded at chemosynthetic ecosystems in the western Pacific. Since both species are also known from regular non-chemosynthetic deep-sea and they were recovered from areas with only weak diffuse flow venting, they have most likely invaded vent environment from regular sea bottom and play a role in trophic transfer of chemosynthetic production to the surrounding seafloor. The fact that such species are able to tolerate (some) influence from hydrothermal vents signify that such ‘outskirt’ areas may provide key habitats for them, and that these areas may play a role in the evolution of biological adaptations to ‘extreme’ chemosynthetic ecosystems.

Additional Information and Declarations

Data Availability

Specimens studies in this study are deposited and accessioned at JAMSTEC marine biological sample collection; *Otukaia ikukoe*: JAMSTEC No. 1160052681; *Abyssotrophon soyoae*: JAMSTEC No. 1150046767. The newly generated COI barcode sequences are deposited in DNA Data Bank of Japan with accession numbers LC333137 (*O. ikukoe*) and LC333138 (*A. soyoae*).

Acknowledgements

The authors would like to express their sincere gratitude to the Captain and crews of R/Vs *Kairei* and *Yokosuka*, as well as the operation team of the ROV *KAIKO*, for their untiring support during the relevant cruises (YK14-16, KR15-16, YK16-07, KR16-16). Kyoko Okino (AORI, the University of Tokyo) is gratefully acknowledged for leading the YK14-16 cruise, and Kentaro Nakamura (the University of Tokyo) for analysing the MBES data which led to the discovery of the new hydrothermal sites. Roland Houart and Christopher Moe are thanked for their valuable aid in identifying the *Abyssotrophon*. Kazuya Kitada (JAMSTEC) kindly provided the base map for Fig. 1 used herein. Bruce Marshall (Museum of New Zealand Te Papa Tongarewa) and Verena Tunnicliffe (University of Victoria) are thanked for providing commenting on an earlier version which significantly improved the manuscript. The Fukai site was named in reference to a poisonous jungle of the same name in the film “Nausicaä of the Valley of the Wind” (Studio Ghibli, 1984); the director Hayao Miyazaki and Studio Ghibli are gratefully acknowledged for being a great source of inspiration.

References

- Chen C, Watanabe HK, Ohara Y. 2016.** A very deep *Provanna* (Gastropoda: Aabysochrysoidea) discovered from the Shinkai Seep Field, Southern Mariana Forearc. *Journal of the Marine Biological Association of the United Kingdom*. DOI: 10.1017/S0025315416001648.
- Egorov R. 1993.** Trophoninae (Muricidae) of Russian and adjacent seas. *Ruthenica Supplement 1*: 1–48.
- Folmer O, Black M, Hoeh W, Lutz R, Vrijenhoek R. 1994.** DNA primers for amplification of mitochondrial cytochrome c oxidase subunit I from diverse metazoan invertebrates. *Molecular Marine Biology and Biotechnology* **3**: 294–299.
- Grassle JF. 1987.** The ecology of deep-sea hydrothermal vent communities. In: Blaxter JHS and Southward AJ, eds. *Advances in Marine Biology*: Academic Press. 301–362.
- Hasegawa K. 2009.** Upper bathyal gastropods of the Pacific coast of northern Honshu, Japan, chiefly collected by R/V Wakataka-maru. In: Fujita T, ed. *Deep-sea fauna and pollutants off Pacific Coast of northern Japan*. Tokyo: National Museum of Nature and Science Monographs 39. 225–383.
- Hasegawa K, Saito H. 1995.** *Illustrations and annotated checklist of the molluscan type specimens contained in the Sakurai collection in the National Science Museum, Tokyo*. National Science Museum, Tokyo: Tokyo.
- Hashimoto J, Ohta S, Fujikura K, Miura T. 1995.** Microdistribution pattern and biogeography of the hydrothermal vent communities of the Minami-Ensei Knoll in the mid-Okinawa trough, Western Pacific. *Deep Sea Research Part I: Oceanographic Research Papers* **42**: 577–598.
- Higo S, Callomon P, Goto Y. 1999.** *Catalogue and bibliography of the marine shell-bearing Mollusca of Japan*. Elle Scientific Publications: Osaka.
- Higo S, Callomon P, Goto Y. 2001.** *Catalogue and bibliography of the marine shell-bearing mollusca of Japan: Type figures*. Elle Scientific Publications: Osaka.
- Houart R, Sellanes J. 2006.** New data on recently described Chilean trophonines (Gastropoda: Muricidae), with the description of a new species and notes of their occurrence at a cold-seep site. *Zootaxa* **1222**: 53–68.
- Ise Y. 2012.** Chapter 5: Porifera. In: Fujikura K, Okutani T and Maruyama T, eds. *Deep-sea life - Biological observations using research submersibles (2nd Edition)*. Tokyo: Tokai University Press.
- Kawagucci S, Kumagai H, Nakamura K, Miyazaki J, Kitada K, Torimoto J, Makabe A, Chen C, Kitahashi T, Fujiwara T, Tsu Tsumi S, Iwamoto H. 2015.** KR15-16 cruise report. ROV survey of new hydrothermal vent(s) at mid- and southern-Okinawa Trough, as a part of SIP. In: JAMSTEC, ed. *R/V Kairei Cruise Report*. Yokosuka: Japan Agency for Marine-Earth Science and

- Technology (JAMSTEC), available from:
http://www.godac.jamstec.go.jp/catalog/data/doc_catalog/media/KR15-16_all.pdf.
- Layton KKS, Martel AL, Hebert PDN. 2014.** Patterns of DNA barcode variation in Canadian marine molluscs. *PLOS ONE* **9**: e95003.
- Levin LA, Baco AR, Bowden DA, Colaco A, Cordes EE, Cunha MR, Demopoulos AWJ, Gobin J, Grupe BM, Le J, Metaxas A, Netburn AN, Rouse GW, Thurber AR, Tunnicliffe V, Van Dover CL, Vanreusel A, and Watling L. 2016.** Hydrothermal vents and methane seeps: Rethinking the sphere of influence. *Frontiers in Marine Science* **3**: 72.
- Levin LA, Orphan VJ, Rouse GW, Rathburn AE, Ussler W, Cook GS, Goffredi SK, Perez EM, Warren A, Grupe BM, Chadwick G, Strickrott B. 2012.** A hydrothermal seep on the Costa Rica margin: middle ground in a continuum of reducing ecosystems. *Proceedings of the Royal Society B: Biological Sciences* **279**: 2580–2588.
- Lutz RA, Kennish MJ. 1993.** Ecology of deep-sea hydrothermal vent communities: A review. *Reviews of Geophysics* **31**: 211–242.
- Makabe A, Tasumi E, Matsui Y, Horai S, Sato S, Chen C, Shibuya T, Kitada K, Takahashi A, Miyazaki J, Nakamura K, Kawagucci S, Kumagai H. 2017.** Geochemical and biological features of hydrothermal vent fields newly discovered in the Okinawa Trough. *Goldschmidt Conference Abstracts 2017*: Poster 2331, available from: <https://goldschmidt.info/2017/abstracts/abstractView?id=2017005519>.
- Makabe A, Tsutsumi S, Chen C, Torimoto J, Matsui Y, Shibuya T, Miyazaki J, Kitada K, Kawagucci S. 2016.** Discovery of new hydrothermal vent fields in the mid- and southern-Okinawa Trough. *Goldschmidt Conference Abstracts 2016*: 1945, available from: <https://goldschmidt.info/2016/uploads/abstracts/finalPDFs/1945.pdf>.
- Marshall BA. 1995.** A revision of the recent Calliostomatidae of New Zealand (Mollusca: Gastropoda: Trochoidea). *The Nautilus* **108**: 83–126.
- Marshall BA. 2016.** New species of Venustatrochus Powell, 1951 from New Zealand, and new species of *Falsimargarita* Powell, 1951 and a new genus of the Calliostomatidae from the southwest Pacific, with comments on some other calliostomatid genera (Mollusca: Gastropoda). *Molluscan Research* **36**: 119–141.
- Merle D, Garrigues B, Pointier J-P. 2011.** *Fossil and recent Muricidae of the world: Part Muricinae*. ConchBooks: Hackenheim.
- Meyer CP, Paulay G. 2005.** DNA barcoding: Error rates based on comprehensive sampling. *PLOS Biology* **3**: e422.
- Miyazaki J, Makabe A, Nakamura K, Chen C, Tasumi E, Fujiwara T, Horai S, Sato S, Inoue S, Hashimoto Y. 2016.** KR16-16 cruise report. Cross-ministerial Strategic Innovation Promotion Program (SIP) Next-Generation Technology for Ocean Resources Exploration: Seeking and

- 378 Hunting the Hydrothermal Activities in Southern Okinawa Trough In: JAMSTEC, ed. *R/V Kairei*
379 *Cruise Report*. Yokosuka: Japan Agency for Marine-Earth Science and Technology (JAMSTEC),
380 available from: http://www.godac.jamstec.go.jp/catalog/data/doc_catalog/media/KR16-16_all.pdf.
- 381 **Nakamura K, Kawagucci S, Kitada K, Kumagai H, Takai K, Okino K. 2015.** Water column imaging
382 with multibeam echo-sounding in the mid-Okinawa Trough: Implications for distribution of deep-
383 sea hydrothermal vent sites and the cause of acoustic water column anomaly. *Geochemical*
384 *Journal* **49**: 579–596.
- 385 **Okutani T. 1959.** A new muricid gastropod, *Trophonopsis soyoae*, sp. nov. from a deep water in Kumano-
386 Nada, Japan. *Venus* **20**: 350–352.
- 387 **Okutani T, Iwahori A. 1992.** Noteworthy gastropods collected from bathyal zone in Tosa Bay by the R/V
388 Kotaka-Maru in 1987 and 1988. *Venus* **51**: 235–268.
- 389 **Petersen JM, Zielinski FU, Pape T, Seifert R, Moraru C, Amann R, Hourdez S, Girguis PR, Wankel**
390 **SD, Barbe V, Pelletier E, Fink D, Borowski C, Bach W, Dubilier N. 2011.** Hydrogen is an
391 energy source for hydrothermal vent symbioses. *Nature* **476**: 176–180.
- 392 **Sakurai K. 1996.** Eight new species of trochid genera, *Tristichotrochus*, *Kombologion* and *Otukaia*
393 (*Calliostomatinae*) from Japan and adjacent waters. *Venus* **53**: 287–296.
- 394 **Sarrazin J, Robigou V, Juniper SK, and Delaney JR. 1997.** Biological and geological dynamics over
395 four years on a high-temperature sulfide structure at the Juan de Fuca Ridge hydrothermal
396 observatory. *Marine Ecology Progress Series* 153:5-24. 10.3354/meps153005
- 397 **Sasaki T, Warén A, Kano Y, Okutani T, Fujikura K, Kiel S. 2010.** Gastropods from recent hot vents
398 and cold seeps: systematics, diversity and life strategies. In: Kiel S, ed. *The vent and seep biota*.
399 Netherlands: Springer 169–254.
- 400 **Sellanes J, Quiroga E, Neira C. 2008.** Megafauna community structure and trophic relationships at the
401 recently discovered Concepción Methane Seep Area, Chile, ~36°S. *ICES Journal of Marine*
402 *Science* **65**: 1102–1111.
- 403 **Tsuchiya K. 2017.** Muricidae. In: Okutani T, ed. *Marine mollusks in Japan (2nd Editon)*. Tokyo: Tokai
404 University Press. 282–310.
- 405 **Tunnicliffe V, and Fontaine AR. 1987.** Faunal composition and organic surface encrustations at
406 hydrothermal vents on the southern Juan De Fuca Ridge. *Journal of Geophysical Research: Solid*
407 *Earth* 92:11303-11314.
- 408 **Van Dover CL. 2000.** *The ecology of deep-sea hydrothermal vents*. Princeton University Press: Princeton.
- 409 **Van Dover CL, and Trask JL. 2000.** Diversity at deep-sea hydrothermal vent and intertidal mussel beds.
410 *Marine Ecology Progress Series* **195**:169-178.
- 411 **Vilvens C. 2014.** New species and new records of Calliostomatidae (Gastropoda: Trochoidea) from
412 eastern and central Indo-Pacific. *Novapex* **15**: 37–48.

- 413 **Warén A, Bouchet P. 2001.** Gastropoda and Monoplacophora from hydrothermal vents and seeps new
414 taxa and records. *The Veliger* **44**: 116–231.
- 415 **Watanabe H, Kojima S. 2015.** Vent fauna in the Okinawa Trough. In: Ishibashi J-i, Okino K and
416 Sunamura M, eds. *Subseafloor Biosphere Linked to Hydrothermal Systems: TAIGA Concept*.
417 Tokyo: Springer Japan. 449–459.
- 418 **Williams ST, Donald KM, Spencer HG, Nakano T. 2010.** Molecular systematics of the marine
419 gastropod families Trochidae and Calliostomatidae (Mollusca: Superfamily Trochoidea).
420 *Molecular Phylogenetics and Evolution* **54**: 783–809.

Figure Captions

Figure 1. Map of the Okinawa Trough and surrounding areas showing locations of the two relevant hydrothermally influenced sites. The blue star indicates Fukai site, Higashi-Ensei and the red dot indicates Crane site, Tarama Hill. Black dots indicate known vent sites in the Okinawa Trough (following Watanabe & Kojima, 2015). Site name abbreviations are as follows: Cr = Crane Site, Ht = Hatoma Knoll, Ih = Iheya Ridge, Ik = Iheya North Knoll, Ir = Irabu Knoll, Iz = Izena Hole, Me = Minami-Ensei Knoll, Yg = Daiyon-Yonaguni Knoll, Yr = Yoron Hole. Original 500 m mesh data originally obtained by Japan Coast Guard and publicly available from Japan Oceanographic Data Center (http://jdoss1.jodc.go.jp/vpage/depth500_file_j.html).

Figure 2. *In situ* photographs of the two hydrothermally influenced sites. **A–B:** Fukai site, Higashi-Ensei; **A:** Overview, **B:** Close-up; the black arrowhead indicates *Otukaia ikukoe* and the white arrowhead indicates *Abyssotrophon* cf. *soyoe*. **C–D:** Crane site, Tarama Hill; **C:** Overview, **D:** Close-up; the white arrowhead indicates *Abyssotrophon soyoe*. The cut-out photographs are magnified images of gastropods seen at each site.

Figure 3. Specimens of the two gastropods collected from hydrothermally influenced areas. **A–D:** *Otukaia ikukoe* from Fukai site, Higashi-Ensei. **E–J:** *Abyssotrophon soyoe* from Crane site, Tarama Hill; **E–H** = Specimen #1, **I, J** = Specimen #2. Scale bars: **A–D** = 1 cm, **E–J** = 0.5 cm.

Figure 4. Scanning electron micrographs of radulae. **A:** *Otukaia ikukoe* from Fukai site, Higashi-Ensei. **B:** *Abyssotrophon soyoe* from Crane site, Tarama Hill. Scale bars: **A** = 500 μ m, **B** = 50 μ m.

Figure 1

Map of the Okinawa Trough and surrounding areas showing locations of the two relevant hydrothermally influenced site.

The blue star indicates Fukai site, Higashi-Ensei and the red dot indicates Crane site, Tarama Hill. Black dots indicate known vent sites in the Okinawa Trough (following Watanabe & Kojima, 2015). Site name abbreviations are as follows: Cr = Crane Site, Ht = Hatoma Knoll, Ih = Iheya Ridge, Ik = Iheya North Knoll, Ir = Irabu Knoll, Iz = Izena Hole, Me = Minami-Ensei Knoll, Yg = Daiyon-Yonaguni Knoll, Yr = Yoron Hole. Original 500 m mesh data originally obtained by Japan Coast Guard and publicly available from Japan Oceanographic Data Center (http://jdoss1.jodc.go.jp/vpage/depth500_file_j.html).




Figure 2

In situ photographs of the two hydrothermally influenced sites.

A-B: Fukai site, Higashi-Ensei; A: Overview, B: Close-up; the black arrowhead indicates *Tristichotrochus ikukoeae* and the white arrowhead indicates *Abyssotrophon* cf. *soyoae*. C-D: Crane site, Tarama Hill; C: Overview, D: Close-up; the white arrowhead indicates *Abyssotrophon soyoae*. The cut-out photographs are magnified images of gastropods seen at each site.




Figure 3

Specimens of the two gastropods collected from hydrothermally influenced areas.

A-D: *Tristichotrochus ikukoeae* from Fukai site, Higashi-Ensei. **E-J:** *Abyssotrophon soyoae* from Crane site, Tarama Hill; **E-H** = Specimen #1, **I, J** = Specimen #2. Scale bars: **A-D** = 1 cm, **E-J** = 0.5 cm.




Figure 4

Scanning electron micrographs of radulae.

A: *Tristichotrochus ikukoeae* from Fukai site, Higashi-Ensei. **B:** *Abyssotrophon soyoae* from Crane site, Tarama Hill. Scale bars: **A** = 500 μm , **B** = 50 μm .

**Note: Auto Gamma Correction was used for the image. This only affects the reviewing manuscript. See original source image if needed for review.*

