

Domestic carnivore interactions with wildlife in the Cape Horn Biosphere Reserve, Chile: husbandry and perceptions of impacts from a community perspective (#17813)

1

First revision

Editor guidance

Please submit by **30 Oct 2017** for the benefit of the authors (and your \$200 publishing discount).



Structure and Criteria

Please read the 'Structure and Criteria' page for general guidance.



Custom checks

Make sure you include the custom checks shown below, in your review.



Raw data check

Review the raw data. Download from the [materials page](#).



Image check

Check that figures and images have not been inappropriately manipulated.

Privacy reminder: If uploading an annotated PDF, remove identifiable information to remain anonymous.

Files

Download and review all files from the [materials page](#).

- 1 Tracked changes manuscript(s)
- 1 Rebuttal letter(s)
- 5 Figure file(s)
- 4 Table file(s)
- 1 Other file(s)

! Custom checks

Human participant/human tissue checks

- ! Have you checked the authors [ethical approval statement](#)?
- ! Does the study meet our [article requirements](#)?
- ! Has identifiable info been removed from all files?
- ! Were the experiments necessary and ethical?



Structure your review

The review form is divided into 5 sections.

Please consider these when composing your review:

1. BASIC REPORTING

2. EXPERIMENTAL DESIGN

3. VALIDITY OF THE FINDINGS

4. General comments

5. Confidential notes to the editor

 You can also annotate this PDF and upload it as part of your review

When ready [submit online](#).

Editorial Criteria

Use these criteria points to structure your review. The full detailed editorial criteria is on your [guidance page](#).

BASIC REPORTING

-  Clear, unambiguous, professional English language used throughout.
-  Intro & background to show context. Literature well referenced & relevant.
-  Structure conforms to [PeerJ standards](#), discipline norm, or improved for clarity.
-  Figures are relevant, high quality, well labelled & described.
-  Raw data supplied (see [PeerJ policy](#)).

EXPERIMENTAL DESIGN

-  Original primary research within [Scope of the journal](#).
-  Research question well defined, relevant & meaningful. It is stated how the research fills an identified knowledge gap.
-  Rigorous investigation performed to a high technical & ethical standard.
-  Methods described with sufficient detail & information to replicate.

VALIDITY OF THE FINDINGS

-  Impact and novelty not assessed. Negative/inconclusive results accepted. *Meaningful* replication encouraged where rationale & benefit to literature is clearly stated.
-  Data is robust, statistically sound, & controlled.
-  Conclusions are well stated, linked to original research question & limited to supporting results.
-  Speculation is welcome, but should be identified as such.

Standout reviewing tips

3



The best reviewers use these techniques

Tip

Support criticisms with evidence from the text or from other sources

Example

Smith et al (J of Methodology, 2005, V3, pp 123) have shown that the analysis you use in Lines 241-250 is not the most appropriate for this situation. Please explain why you used this method.

Give specific suggestions on how to improve the manuscript

Your introduction needs more detail. I suggest that you improve the description at lines 57- 86 to provide more justification for your study (specifically, you should expand upon the knowledge gap being filled).

Comment on language and grammar issues

The English language should be improved to ensure that an international audience can clearly understand your text. Some examples where the language could be improved include lines 23, 77, 121, 128 – the current phrasing makes comprehension difficult.

Organize by importance of the issues, and number your points

1. Your most important issue
2. The next most important item
3. ...
4. The least important points

Please provide constructive criticism, and avoid personal opinions

I thank you for providing the raw data, however your supplemental files need more descriptive metadata identifiers to be useful to future readers. Although your results are compelling, the data analysis should be improved in the following ways: AA, BB, CC

Comment on strengths (as well as weaknesses) of the manuscript

I commend the authors for their extensive data set, compiled over many years of detailed fieldwork. In addition, the manuscript is clearly written in professional, unambiguous language. If there is a weakness, it is in the statistical analysis (as I have noted above) which should be improved upon before Acceptance.

Domestic carnivore interactions with wildlife in the Cape Horn Biosphere Reserve, Chile: husbandry and perceptions of impacts from a community perspective

Elke Schüttler^{Corresp., 1}, Lorena Saavedra-Aracena¹, Jaime E Jiménez^{1, 2, 3, 4}

¹ Sub-Antarctic Biocultural Conservation Program, Universidad de Magallanes, Punta Arenas, Región de Magallanes y Antártica Chilena, Chile

² Department of Biological Sciences, University of North Texas, Denton, Texas, United States

³ Department of Philosophy and Religion, University of North Texas, Denton, Texas, United States

⁴ Institute of Ecology and Biodiversity (IEB), Santiago, Chile

Corresponding Author: Elke Schüttler
Email address: elkeschuttler@gmail.com

Background. Hundreds of millions of domestic carnivores worldwide have diverse positive affiliations with humans, but can provoke serious socio-ecological impacts when free-roaming. Unconfined dogs (*Canis familiaris*) and cats (*Felis catus*) interact with wildlife as predators, competitors, and disease-transmitters; their access to wildlife depends on husbandry, perceptions, attitudes, and behaviors of pet owners and non-owners.

Methods. To better understand husbandry and perceptions of impacts by unconfined, domestic carnivores, we administered questionnaires (n=244) to pet owners and non-owners living in one of the last wilderness areas of the world, the Cape Horn Biosphere Reserve, located in southern Chile. We used descriptive statistics to provide demographic pet and husbandry information, quantify free-roaming dogs and cats, map their sightings in nature, and report experiences and perceptions of the impact of free-roaming dogs and cats on wildlife. We corroborated our results with an analysis of prey remains in dog feces (n=53). With generalized linear models, we examined which factors (i.e., food provisioning, reproductive state, rural/village households, sex, and size) predicted that owned dogs and cats bring wildlife prey home.

Results. Thirty-one percent of village dogs (n=121) and 60% of dogs in rural areas (n=47) roamed freely day and/or night. We found free-roaming dog packs frequently observed (64% of participants) in the wild, including a feral dog population on Navarino Island. Dogs (31 of 168) brought invasive muskrats (*Ondatra zibethicus*) and avian prey home, and over half of all cats (27 of 51) mainly brought birds. Birds were also the most harassed group by one third of all dogs and cats. Despite these facts, dog-wildlife conflicts were hardly perceived (<9% of observed conflicts and suspected problems) and only 34% of the participants thought that cats might impact birds. Diet analysis revealed that dogs consumed livestock (64% of 59 prey occurrences), beavers (*Castor canadensis*, 14%), and birds (10%). The probability that dogs brought prey to owners' homes was higher in rural and larger dogs. There was also evidence that cats from rural households and with an inadequate food supply brought more prey home than village cats.

Discussion. Despite that muskrat, beavers, and birds were brought home, harassed, or found in dog feces, free-roaming dogs and, to a lesser extent, cats are perceived predominantly in an anthropogenic context (i.e., as pets) and not as carnivores interacting with wildlife. Therefore, technical and legal

measures should be applied to encourage neutering, increase confinement, particularly in rural areas, and stimulate social change via environmental education that draws attention to the possibility and consequences of unconfined pet interaction with wildlife in the southernmost protected area of the globe.

**Domestic carnivore interactions with wildlife in the Cape Horn Biosphere Reserve, Chile:
husbandry and perceptions of impacts from a community perspective**

Elke Schüttler¹, Lorena Saavedra-Aracena¹ and Jaime E. Jiménez^{1,2,3,4}

¹ Sub-Antarctic Biocultural Conservation Program, Universidad de Magallanes, Punta Arenas,
Región de Magallanes y Antártica Chilena, Chile

² Department of Biological Sciences, University of North Texas, Denton, Texas 76201, USA

³ Department of Philosophy and Religion, University of North Texas, Denton, Texas 76201, USA

⁴ Institute of Ecology and Biodiversity (IEB), Santiago, Chile

Corresponding Author:

Elke Schüttler¹

Teniente Muñoz 396, Puerto Williams, Región de Magallanes y Antártica Chilena, Chile

Email address: elkeschuttler@gmail.com

Abstract

Background. Hundreds of millions of domestic carnivores worldwide have diverse positive affiliations with humans, but can provoke serious socio-ecological impacts when free-roaming. Unconfined dogs (*Canis familiaris*) and cats (*Felis catus*) interact with wildlife as predators, competitors, and disease-transmitters; their access to wildlife depends on husbandry, perceptions, attitudes, and behaviors of pet owners and non-owners.

Methods. To better understand husbandry and perceptions of impacts by unconfined, domestic carnivores, we administered questionnaires (n=244) to pet owners and non-owners living in one of the last wilderness areas of the world, the Cape Horn Biosphere Reserve, located in southern Chile. We used descriptive statistics to provide demographic pet and husbandry information, quantify free-roaming dogs and cats, map their sightings in nature, and report experiences and perceptions of the impact of free-roaming dogs and cats on wildlife. We corroborated our results with an analysis of prey remains in dog feces (n=53). With generalized linear models, we examined which factors (i.e., food provisioning, reproductive state, rural/village households, sex, and size) predicted that owned dogs and cats bring wildlife prey home.

Results. Thirty-one percent of village dogs (n=121) and 60% of dogs in rural areas (n=47) roamed freely day and/or night. We found free-roaming dog packs frequently observed (64% of participants) in the wild, including a feral dog population on Naverino Island. Dogs (31 of 168) brought invasive muskrats (*Ondatra zibethicus*) and avian prey home, and over half of all cats (27 of 51) mainly brought birds. Birds were also the most harassed group by one third of all dogs and cats. Despite these facts, dog-wildlife conflicts were hardly perceived (<9% of observed conflicts and suspected problems) only 34% of the participants thought that cats might impact birds. Diet analysis revealed that dogs consumed livestock (64% of 59 prey occurrences), beavers (*Castor canadensis*, 14%) and birds (10%). The probability that dogs brought prey to owners' homes was higher in rural and larger dogs. There was also evidence that cats from rural households and with an inadequate food supply brought more prey home than village cats.

Discussion. Despite that muskrat, beavers, and birds were brought home, harassed, or found in dog feces, free-roaming dogs and, to a lesser extent, cats are perceived predominantly in an anthropogenic context (i.e., as pets) and not as carnivores interacting with wildlife. Therefore, technical and legal measures should be applied to encourage neutering, increase confinement, particularly in rural areas, and stimulate social change via environmental education that draws

attention to the possibility and consequences of unconfined pet interaction with wildlife in the southernmost protected area of the globe.

Introduction

In parallel to human population growth, the number of companion animals is constantly increasing as well. Pet feral dogs (*Canis familiaris*) have reached population estimates of 900 million and cats (*Felis catus*) of 600 million (O'Brien & Johnson, 2007; Gompper, 2014a), existing on all continents except Antarctica (Hughes et al., 2015). Since their domestication thousands of years ago, domestic dogs have had profound roles in human lives. These include companionship, livestock guarding, rescue, hunting, tourism, service animals, and wildlife management (Hart & Yamamoto, 2017). As dogs are part of a diversity of human cultures, their various roles, husbandry, and people's attitudes towards them have different implications for human-dog-wildlife interactions (Miller, Ritchie & Weston, 2014).

The possibility to interact with wildlife depends on dogs' and cats' husbandry, particularly on their confinement. This ranges from complete mobility restriction, for leashed and/or indoor dogs and cats, to feral domestic carnivores that survive independently of supplemental provisioning from humans (Kays & DeWan, 2004; Vanak & Gompper, 2009). In between these extremes, there exists a range of free-roaming animals that are owned or unowned and are, to some extent, subsidized by humans. As subsidized predators, domestic carnivores can reach higher population densities than wild carnivore populations (Gompper, 2014b), leading to complex socio-ecological consequences.

The impacts of free-roaming subsidized and feral domestic dogs include loss of livestock (Baker et al., 2008; Echegaray & Vilà, 2010), aggression towards humans (Schalamon et al., 2006), disease transmission (Matter & Daniels, 2010), and wildlife interference (reviewed in Young et al., 2011; Hughes & Macdonald, 2013). Dogs prey upon (Butler, du Toit & Bingham, 2004; Manor & Saltz, 2004), compete with (Mitchell & Banks, 2005; Vanak, Thaker & Gompper, 2009), infect (Acosta-Jamett, 2009), and disturb (Silva-Rodríguez, Ortega-Solís & Jiménez, 2010; Silva-Rodríguez & Sieving, 2012) wild animals. Suburban cats are successful small vertebrate predators (Woods, McDonald & Harris, 2003; Loyd et al., 2013). On islands, Medina et al. (2011) demonstrated that feral cats were responsible for at least 14% of global avian, mammalian, and

reptilian extinctions (see also Nogales et al., 2013). Finally, both dogs and cats may hybridize with their wild relatives (Randi, 2008).

While the biology of domestic carnivore-wildlife interactions is often the focus of research, studies on social dimensions are still in their infancy (Miller, Ritchie & Weston, 2014). Understanding the perceptions of free-roaming pet carnivores is indispensable to inform conservation action, as causes and solutions to this problem directly depend upon humans' relationships with them (Gramza et al., 2016). Conflicts between dogs/cats and wildlife could thus be minimized by a better understanding of how husbandry, attitudes, and behaviors of pet owners influence dogs and cats in their access to and interaction with wild prey or carnivores, particularly when close to protected areas. For example, dog owners felt more obliged to leash their dogs when they thought their dog would harm beach-nesting birds or people (Williams et al., 2009). Recent studies have shown that a more adequate diet (i.e., commercial or prepared food instead of e.g., household scraps) for dogs and cats decreases the probability of preying upon wild animals (Silva-Rodríguez & Sieving 2012, Sepúlveda et al., 2014). Additionally, in the absence of biological data or for logistical and cost reasons, a community-based assessment may provide suitable information on species' impacts on other species (White et al., 2005).

Here, we focus on understanding the interactions with wildlife by a population of free-roaming dogs and cats in a sensitive conservation area of southern Chile, the Cape Horn Biosphere Reserve, from a community perspective. To date, the only substantial human impacts in the biosphere reserve are biological invasions, particularly of wild and domestic exotic mammals that outnumber their native counterparts (Anderson et al., 2006). Additionally, native terrestrial predators are absent on many of the islands of the reserve, including Navarino Island, where the only village of the reserve is located (Anderson et al., 2006). This indicates that impacts on native and often naïve prey generated by introduced predators may cause extensive population reductions (discussed in Sih et al., 2010). Although free-roaming and feral dogs are commonly observed by locals on Navarino Island, few observations of their impacts exist. Dogs have been reported to prey upon the southernmost population of guanacos (*Lama guanicoe*), which is virtually unstudied and considered in danger of local extinction (Cunazza, 1991; González, 2005). Evidence also exists of dogs preying upon nests of solitary nesting waterfowl, such as the Flightless Steamer Duck (*Tachyeres pteneres*), a species endemic to Patagonia, and nesting colonies of the South

American Tern (*Sterna hirundinacea*, Schüttler et al., 2009). There are no published accounts of local cat impacts.

We administered questionnaires to pet owners and non-owners to address the following objectives: (1) provide demographic and husbandry information relevant for future dog and cat management in one of the last wilderness areas of the globe, (2) quantify free-roaming dogs and cats and map their locations in nature, (3) examine experiences and perceptions of the impact of free-roaming dogs and cats on wildlife and corroborate those with an analysis of prey remains of dog feces, and (4) predict which factors best explain when owned dogs and cats bring wildlife prey home. With regard to the paucity of ecological data on domestic carnivore-wildlife interactions in this sub-Antarctic wilderness area, this study allows a first insight into this conservation problem.

Materials & Methods

Ethics statement

We obtained prior informed consent from each participant by reading a printed statement explaining the project aims, the lack of risks in participating, the possibility to omit questions, information about use of and access to the results, and that the interview was anonymous and voluntary. The participants agreed to participate by signing a copy of the informed consent; they were also given a copy. Paper and digital questionnaires were stored anonymously. The Scientific Ethical Committee of the University of Magallanes, Chile, certified ethical approval of the instrument.

Study area

We carried out this study in the Cape Horn Biosphere Reserve (CHBR) (19,172 km² terrestrial surface), Chile, located at the extreme southern tip of South America (Fig. 1). The biosphere reserve exists within the Magellanic Sub-Antarctic forest ecoregion, one of the remaining 24 wilderness areas of the world (Mittermeier et al., 2003). The dominant habitats within this ecoregion are unfragmented evergreen and deciduous southern beech (*Nothofagus* spp.) and Winter's bark (*Drimys winteri*) forests, Magellanic peat bogs (predominantly *Sphagnum* spp.), high-Andean habitats, and glaciers (Pisano, 1977). The human population in the CHBR is of mixed cultural and ethnic origin (i.e., Yaghan indigenous people, Chilean Navy, fishermen, and Chilean

and foreign short- and long-term settlers) and is concentrated in the only town within the reserve, Puerto Williams (2,800 inhabitants) on Navarino Island. A small fishing village, Puerto Toro, exists on the eastern coast of Navarino as well as eight rural farm settlements. In the remainder of the biosphere reserve, there are only 11 Chilean Navy posts operated by a new family each year, and one farm on Hoste Island (Fig. 1). The principal economic activities on Navarino Island include fishing, tourism, and small-scale livestock farming. The infrastructure is limited to a dirt road along the northern coast of Navarino; public maritime transport within the reserve does not exist.

[please insert Figure 1 here]

Survey

From May 2015 to April 2016, we interviewed 215 households in Puerto Williams, seven farm owners on Navarino Island, and 22 Chilean Navy families living for one year on the 11 Navy posts on different islands in the CHBR (n=244 total interviews). To test the questionnaire design and adapt the questions, we conducted a pilot study with four trial informants who were later excluded from the dataset.

We calculated a representative 95% probability sample of 215 interviewees in Puerto Williams based on a census of households applied by ES in May 2015 (490 houses). We used a confidence interval of 5% and applied the finite population correction for smaller populations (Bernard, 2006: 183). We randomly chose 280 households from a map of numbered houses in town (adding 30% to the sample size of 215 to correct for non-responses). When an adult was not present, we left a written message explaining the motivation for our visit and our contact information. We visited each household up to three times before it was replaced. The questionnaires were administered at different times in a face-to-face interview approach at the participant's home and took 10-30 min. Two interviewers conducted the interviews in Spanish (n=92 by ES and n=152 by LS).

To meet objective 1, we collected information on dog and cat demographics (i.e., number, age, sex, origin, purpose of pets, number of pups/kittens in previous year, elimination, and temporary and definite pet loss) as well as on pet care/husbandry (i.e., veterinary treatments, food type, in how many households pets eat, and whether owned or unowned street dogs are fed). We

asked participants whether they restricted the movements of their dogs (i.e., day and/or night and if free-ranging, why), whether they saw unaccompanied dogs and cats (adults, pups/kittens) outside of town and from where they thought the animals came (objective 2). We asked participants about their observations of problematic dog situations in and out of town, perceptions of possible impacts of free-roaming dogs and cats, and their personal experiences regarding the pet's interaction(s) with other animals (objective 3). To better understand predictors (i.e., food provisioning, reproductive state, rural/village households, sex, size) of dog and cat interactions with wildlife, we asked owners whether their pets brought wildlife prey home (objective 4). We finished the questionnaire by asking for suggestions for reducing the number of free-roaming dogs, personal data (i.e., age, sex, education, residence time), and the owner's permission to take a photo of their pet. From this photo, we then classified dog sizes (i.e., small, medium, large) and calculated the mean of two independent estimations by ES and LSA. For farm owners, we added questions on their experience in losing domestic animals such as cattle (*Bos taurus*) or sheep (*Ovis aries*) to dog attacks. For participants without pets, non-relevant questions were not asked. Some questions were focused only on dogs because dogs produce visible socio-ecological conflicts in the study area. The complete questionnaire is provided as Supplementary Material S1.

Dog diet

Dog feces of owned and feral dogs (n=70) were opportunistically collected within one km of the 11 Navy posts and during trekking events on Navarino Island (June 2015 to April 2016). Each sample was dried, rehydrated, and degreased with detergent, and then grouped into hair, bones, feathers, and rubbish. Using a microscope, we examined the medullary and cuticular patterns of guard hairs to identify up to species level using gelatin as the printing medium (similar to González-Esteban, Villate & Irizar, 1996). As reference collections, we used those provided by the Instituto de la Patagonia, Universidad de Magallanes, complemented by our own additions, and local keys (Chehébar & Martín, 1989).

Statistics

We first calculated descriptive statistics to analyze basic demographic and husbandry information from the interviews to (i) quantify free-roaming dogs and cats, (ii) report experienced problems associated with dogs, (iii) perceptions of the potential impact of free-roaming dogs and



cats on wildlife, (iv) report percentages of prey and harassed animals by dogs and cats, and (v) percentages of prey groups in dog feces. We then used generalized linear models (GLMs) to examine predictors of interactions with wildlife of owned dogs and cats (Table 1). Our response variable for the dog and the cat model was whether pets brought wildlife home, defined here as any exotic or native wild mammal or bird (PREY). As covariates, we considered diet (dinner leftovers and/or commercial pet food) provided by the animal's owner (FOOD), sex (SEX), sterility (STERILIZED), owners' household location (LOCATION, rural or village), and pet size (SIZE, only for dogs). We focused on those covariates as they might identify pet characteristics or levels of care which can then inform recommendations for pet management for the benefit of wildlife. The location was included because rural households are imbedded in wilderness settings and access to wildlife is immediate. Response and predictor variables are explained in detail in Table 2.

As the response variable of the two models was binomial, we fitted generalized linear models (GLMs) with binomial error structure and logit link. The models were parameterized with all possible covariate combinations. Prior to analysis, we explored these data following Zuur et al. (2010). Collinearity between ordinal covariates was assessed with Spearman correlation coefficients (no coefficients were $>|0.4|$). We tested the independence of categorical variables using contingency tables (Chi-square and Fisher's exact tests, $p < 0.05$). We removed SEX and STERILIZED from the dog model for being significantly associated with LOCATION; similarly, we removed STERILIZED from the cat model for being significantly associated with LOCATION and SEX. FOOD was maintained here, despite its collinearity with LOCATION, to test the same variables in the dog and cat models (the conclusions for models tested with and without FOOD were identical, see Mundry, 2014, for treating collinearity). For model selection, we used Akaike's Information Criterion corrected for small sample size (AIC_c). We tested for an interviewer effect by including interviewer as a random effect in the models (generalized linear mixed models, GLMMs), but did not detect any ($AIC_{GLMMs} > AIC_{GLMs}$ of the global models, respectively). We accounted for model selection uncertainty (model weights ω_i were < 0.9) using full-model averaging (Symonds & Moussalli, 2011). To rank predictor variables in terms of importance, we summed Akaike weights for each model in which the variable under consideration appeared (Burnham & Anderson, 2002). We explored the direction of predictor impacts on the response

variable by calculating log odds ratios of the averaged estimates with 95% confidence intervals. Statistical modelling was conducted in R (R Core Team, 2016).

[Please insert Table 1 here]

[Please insert Table 2 here]

Results

We conducted 215 interviews in Puerto Williams, seven in rural households, and 22 with Navy post families. Only five people in Puerto Williams refused to participate. Of the 244 participants, 61.5% were female, mean participant's age was 39.5 years (SD 11.6, range 18-76 years), and mean residence time in the biosphere reserve was 11.6 years (SD 14.2, range one month-66 years).

The Puerto Williams participants owned 121 dogs and 36 cats, predominantly for companionship. The seven farm households owned 30 dogs and 15 cats, mainly kept as working dogs and for rodent control, respectively. The 22 Navy families owned 17 dogs primarily for companionship and two cats for rodent control. Two dogs and two cats stayed at the Navy post when families were exchanged after one year ("inherited animals"), the others left with their families. Reproductive control was moderate to high in Puerto Williams (41.7% of dogs and 19.4% of cats not sterilized), but almost absent in rural areas (83.3% dogs, 93.3% cats) and Navy posts (86.7% dogs, 100% cats, but 9 of 17 dogs were ≤ 1 -year, Table 3). Additionally, four participants had eliminated unwanted dog pups.

In Puerto Williams, over half of the dogs and around one third of the cats were vaccinated against rabies (55.4% and 33.3%, respectively) and treated for parasites (60.3% and 36.1%, respectively). Pet owners in rural households did not vaccinate against rabies, but treated them for parasites (100% dogs, 40.0% cats). At Navy posts, only dogs were vaccinated and dewormed (64.7%). Pets in Puerto Williams, rural dogs, and dogs at Navy posts were provided mainly with commercial food and/or meat ($>77.7\%$). However, 35 village dogs (28.9%) were fed in more than one household, and 74 interviewees in town (34.4%) reported feeding street dogs on a regular basis (71.8% at least once a week), mostly with leftovers (73.5% of 83 mentioned food items).

Sixty-five of 168 dogs (38.7%) roamed freely day or night and 39 dogs (23.2%) were always unrestricted. Using an extrapolated number of dogs for the 490 households in Puerto

Williams (n=275.8 with 1.4 dogs/household), we estimate that 84 dogs (30.6%) roam freely in town during day or night. The most common method of dog restriction (69.4% in town, 53.3% rural, 17.6% Navy posts) was keeping dogs in the house (63.6% of 121 responses), fewer were kept free in the yard (18.2%) or leashed (18.2%). Reasons mentioned for allowing unrestricted movement of dogs in town and rural environments were (i) the owner claimed animal freedom, (ii) leashing may increase aggressiveness, (iii) acclimation to free-roaming and releasing energy, (iv) the dog is not dangerous, and (v) unsuitable facilities (together 77.2% of 70 explanations). Also, 44.0% of 91 non-pet owners thought that street dogs enjoyed their freedom, and most dog owners (87.2% of 86) thought that street dogs roamed into the forest. Finally, 22 of 168 dogs (13.1%) went missing between 12-24 hours during the last year (2014/15), among which 13 dogs had disappeared up to one week before returning home. Cats disappeared more frequently (n=18, 34.0%); 14 cats for 2-7 days. Over the last 10 years, 35 pets (23 dogs, 12 cats) never returned.

[Please insert Table 3 here]

Free-roaming dogs not accompanied by people outside of town were frequently observed on Navarino Island (63.9% of 244 participants), whereas cat sightings in the wild were almost absent (6.1%, Fig. 2). The greatest distance of sighted dogs and cats was 19.4 km and 5.2 km from the northern settled coast, respectively. Neither dogs nor cats were seen roaming around Navy posts, except near the two posts on Navarino Island. Dogs were mostly observed in packs, with a median pack size of four dogs (mean 6.6, SD 7.5, range 2-60, n=172 sightings), while only 9.2% of the sightings were single dogs. Dog pups (abandoned or feral) outside Puerto Williams were sighted by 52 participants (21.3%) with a mean litter size of 4.0 (SD 2.3, range 1-12). Four participants observed pups and kittens (n=17) abandoned in cardboard boxes outside town. Apparently, the landfill, at an approximately 500 m from Puerto Williams, was a point of attraction for dogs, as 12 participants observed dog packs with a median size of eight (mean 10.75, SD 8.0, range 1-25, Fig. 2) in this area.

[Please insert Figure 2 here]

Over half of the participants (55.9%, n=222) on Navarino Island had experienced problems with dogs in Puerto Williams during 2009-2015 (83.9% had occurred during the last year, 2014/15), whereas 41 participants (18.5%) reported problems outside the town (61.4% during 2014/2015). Predominant experienced dog problems in town were conflicts with people (biting, attacking, frightening, disease transmitting, accidents, and stealing; 40.6% of 143 problems, 24.1% concerned children) and free-ranging domestic animals (cows, horses, and their offspring) in town, mostly foals (37.1%, Fig. 3). Outside of town, people experienced conflicts between dogs and domestic animals, particularly involving cattle (77.3% of 44 problems), whereas only two people saw dogs feeding on wildfowl eggs (4.5%).

Beyond personal experiences, most participants thought that free-roaming dogs caused problems both in and outside of town (91.9% and 89.2%, respectively). In town, suspected dog problems mainly involved people (68.8% of 221 problems), whereas outside of town concerns involved domestic animals and people (Fig. 3). Dog-wildlife conflicts (e.g., involving guanacos) were only mentioned 19 times (9.4% of 202 problems). However, when asked directly whether feral dogs could have negative impacts on wildlife and what kind of, most participants said yes (80.8%, n=239) regarding birds (67.3% of 349 problems). Guanacos were only mentioned 16 times among the affected wildlife (4.6%).

Regarding suspected cat problems, only one third of participants (33.8% of 240) associated problems with them outside of town, particularly with cats harassing and eating wild birds and their eggs (67.5% of 77 problems, Fig. 3).

Five of seven farm owners reported losing domestic animals due to unrestricted dogs of Puerto Williams or feral dogs during 2014/15. The estimated total loss during these events were 62 calves, 25 cows, and 30 sheep, while 30 sheep and 2 calves were found injured. The losses corresponded to 3.3%, 11.5%, 16.7%, 18.8%, 100% (cows), and 75.0% (sheep) of owned livestock.

[Please insert Figure 3 here]

Thirty-one village and rural dogs (18.5%) brought prey home, mainly invasive muskrats (32.5%) and birds (27.5%, Fig. 4). Among avian prey were songbirds (Austral Blackbird *Curaeus curaeus*, Austral Thrush *Turdus falcklandii*, Rufous-collared Sparrow *Zonotrichia capensis*),



shorebirds (terns, Upland Goose *Chloephaga picta*), and raptors (Chimango Caracara *Milvago chimango*). Over one third of all dogs (n=64) were observed harassing (but not killing) other animals, particularly birds (38.0% of 79 items mentioned: Chimango Caracara, Patagonian Sierra-Finch *Phrygilus patagonicus*, Upland Goose, and ducks such as Flightless Steamer Ducks), other dogs (16.5%), and horses (15.2%); three dogs harassed native mammals (i.e., foxes and seals). For half of all village and rural cats (n=27) brought prey home (birds were 57.6% of 33 prey items): among songbirds, the Austral Thrush, Fired-eyed Duquenois *Xolmis pyrope*, House Sparrow *Passer domesticus*, Patagonian Sierra-Finch, and among shorebirds, ducks and terns. Birds were also the most commonly-harassed prey group (72.0%) by 18 cats (only one species referred to here: Patagonian Sierra-Finch). The 244 participants mentioned diverse food items they thought feral dogs would eat (Fig. 4). Domestic livestock was the most important group mentioned (39.9%), whereas native birds and guanacos were less perceived (15.2% and 2.2%, respectively).

[Please insert Figure 4 here]



we could not identify prey remains of seven feces of owned and feral dogs (n=70) in rural zones of the CHBR and 10 feces were excluded for only containing dog hair. The subsequent diet analysis revealed that the main food item (64.4% of 59 prey occurrences) found in 53 feces was ungulates (i.e., horses and cows, which could not be distinguished here), followed by beavers (13.6%), birds (10.2%), mice (5.1%), rubbish (5.1%), and Fuegian red fox (*Pseudalopex culpaeus*, 1.7%).

Three models best explained whether dogs would bring prey home (Table 4). The most important variable with the highest summed Akaike weights (ω , upper limit=1.0) was LOCATION (ω =0.99); FOOD (ω =0.54) and SEX (ω =0.30) had less importance. The averaged estimates indicated that dogs in rural areas were more likely to bring prey home (Fig. 5A), whereas an adequate diet and the dog's sex had little influence (their confidence intervals overlapped the odds ratio at one). LOCATION (ω =0.99) and FOOD (ω =0.99) were the most important variables in the cat model, whereas sex played a minor role (ω =0.24, Table 4). Based on the averaged model estimates (Fig. 5B), there is evidence that rural cats that received more leftovers brought more prey home than village cats, whereas the cat's sex was a poor predictor. From the collinear variables

removed before modelling, only SIZE had a significant association with LOCATION and PREY, respectively (Fisher's exact tests, $p < 0.001$), indicating that not only rural, but also larger dogs were more prone to preying upon wildlife.

[Please insert Table 4 here]

[Please insert Figure 5 here]

Importantly, participants suggested several measures to diminish the number of street dogs, including reproductive control (18.9%, $n=534$ suggestions), registration (14.6%), responsible pet husbandry (11.2%), establishing an animal shelter (10.5%), education and adoption campaigns (9.2%), controls and penalties (8.4%), movement restriction (8.1%), animal health (7.3%), limiting the number of dogs per family or not abandoning dogs (7.1%), and euthanasia (4.7%).

Discussion

In the absence of biological studies, this survey provides a first understanding of the interactions of domestic carnivores with wildlife in a sub-Antarctic wilderness setting. A representative sample of the local population in the Cape Horn Biosphere Reserve gave insight into pet husbandry and perceptions of impacts of free-roaming dogs and cats. We found that free-roaming dog packs were frequently observed (64% of participants) in natural areas on Navarino Island. Many of these individuals may be owned, as many participants indicated their dogs were not confined (65 of 168 dogs roamed freely day or night, and 22 dogs had even disappeared for 24 h or more). Further, travel distances of free-roaming owned rural dogs vary, with reports of 4 km (Sepúlveda et al., 2015) or up to 8-30 km (Meek, 1999). However, such large foray distances are an exception. Finding a village dog at a distance >1 km from its home had a 10% chance in a study of dogs scavenging sea turtle nests (Ruiz-Izaguirre et al., 2014) and most rural dogs even stayed 95% of their time within <200 m from their households (Sepúlveda et al., 2015).

Our findings indicate that there is evidence of a feral dog population on Navarino Island. The participants reported sightings of unaccompanied dog packs in remote parts of the island (up to 19.4 km from the northern settled coast, Fig. 2), and sightings of dog pups (feral or abandoned) outside town (52 participants). They declared 52% of the 172 sightings as feral dogs. This may be

an over-estimate, but given that free-roaming village dogs are easily recognized in the small town of Puerto Williams, it is likely that participants could distinguish between owned and feral dogs. However, it is not clear whether this presumably feral population has achieved long-term human independence, as for example the dogs eradicated from Isabela Island, Galápagos (Reponen et al., 2014). The reported population of abandoned dog pups and missing dogs may have been recruited into feral dog packs (e.g., Boitani et al., 2017). The importance of the local landfill (12 sightings of dog packs of 8-11 animals on average) as a food subsidy warrants further investigation, as these novel ecosystems can produce a variety of positive and negative impacts on vertebrate species exploiting them (Boitani et al., 2017; Plaza & Lambertucci, 2017). For cats, the few sightings (5% of participants) in natural areas were along the northern-settled coast, except for one cat sighted 5.2 km south of the coast. Further phenotypical, genetic, and ecological research is needed to better understand the feral dog and possible feral cat population and their impacts on Navarino Island.

Although there were 227 sightings during 2014/2015, dog-wildlife conflicts of free-roaming dogs were hardly perceived (4.5% of 44 observed problems, Fig. 3). The direct observation of dog-wildlife interactions is probably a rare situation, particularly with mammals, as the mammalian community on Navarino Island is limited (Anderson et al., 2006). Only five terrestrial native species exist: two species each of bats and mice, and the vulnerable guanaco. Among exotic mammals, there are three elusive wild species (North American beaver, American mink *Neovison vison*, and muskrat) and free-ranging domestic mammals such as cows, horses, sheep, and pigs. Dog interactions with exotic mammals may not have been considered as true wildlife-conflict by the participants as the population reduction of exotic mammals such as mink and beavers was a general consensus in a community survey (Schüttler et al., 2011).

Guanacos have not been sighted along the northern coast for many years (González, Zapata & Marín, 2002) and their densities were as low as 0.14 individuals/km² for the northeastern coast of Navarino Island during 2002-2005 (González, 2005). Thus, it is almost impossible to see depredation or harassment of guanacos by dogs (one piece of photographic evidence was taken by Denis Chevally in 2002). However, individual dog attacks on rare species may impact their persistence substantially (e.g., pudus, Silva-Rodríguez & Sieving, 2012; mountain gazelles *Gazella gazella*, Manor & Saltz, 2004). Therefore, future studies on dog impacts on the southernmost isolated population of guanacos are an urgent need.

The likelihood of interactions among dogs and avian species **should** be much higher, as birds, among them many seabirds, are the most diverse and abundant group among vertebrates in the CHBR (Rozzi et al., 2006). Indeed, eleven dogs brought bird prey home and 30 dogs were observed by their owners to harass birds (songbirds, **shorebirds**, and raptors, Fig. 4). However, these experiences were not translated into the context of a possible “dog-wildlife” conflict: only 9% of 202 suspected dog problems outside town were dog-wildlife problems (Fig. 3); most were dog-domestic animal (54%) or dog-people conflicts (35%). On the one hand, this may be due to a lack of knowledge of the local fauna by short-term residents; further, Cape Horn biocultural identity is missing in the classrooms. Rozzi et al. (2008) reported an absence of native fauna in the minds of local short-term residents who primarily mentioned cosmopolitan roses and apple trees as local plant species. On the other hand, the absence of dog-wildlife interactions in the participants’ minds might indicate that dogs are mainly perceived as domestic animals that act in a human-dominated context and not as carnivores in a natural ecosystem. Personal safety was also the most common public concern regarding free-roaming dogs and cats in central Italy (Slater et al., 2008). This perception may be attributed to the historical attachment bonding between the dog-human dyad believed to be similar to a child-parent relationship (review in Payne, Bennett & McGreevy, 2015).

For cats, awareness of possible cat-wildlife problems, particularly with birds, was higher (68% of 77 problems, Fig. 3). On the one hand, these problems might be more visible, at least for cat owners, whose cats brought birds home (37% of cats in this study, Fig. 4). This number, however, probably clearly underestimates true capture rates, as Loyd et al. (2013) demonstrated with animal-borne video cameras worn by urban cats in the United States (see also Kays & DeWan, 2004). On the other hand, Arahori et al. (2017) showed that owners’ views of their cats and dogs differed; for example, cat owners had a weaker tendency to regard their pets as family members than dog owners. This perception may also influence their view on how cats behave outside their homes (i.e., as carnivorous species).

Besides prey brought home by dogs and cats, our diet analysis of dog feces showed that dogs indeed preyed upon domestic animals such as cows and horses (64%); but beavers (14%) and birds (10%) were present in feces as well. This finding is consistent with cow losses reported by Navarino’s farm owners. However, more biological methods, such as analyzing cat diet or

observations of what free-roaming dogs and cats actually do, are needed to further validate and complement the self-reported observations on wildlife interactions in this survey.

Finally, with GLMs, we showed that rural provenience, large dog size, and an adequately food supply for cats played a significant role as predictors for bringing wildlife prey home. When rural households are imbedded in wilderness settings and spatial and behavioral barriers between domestic and wild animals lack, the apparent consequence are higher depredation rates of (larger) dogs and cats on wildlife. Increasing the confinement of those pets should thus be part of management strategies. Unlike other studies, an inadequate food supply (i.e., higher percentage of leftovers) was not associated with dogs preying upon wildlife. This may be due to methodological differences. Silva-Rodríguez & Sieving (2011) and Ruiz-Izaguirre et al. (2014) considered body condition score and metabolic energy intake, whereas we only relied on the participants' statements. To some extent, the social desirability bias (where the participants wish to appear socially or morally worthy, Maccoby & Maccoby, 1954) might underlie these differences by biasing results that dogs are fed a commercial food diet. While restricting boundaries for farm dogs and cats is probably a difficult task due to their roles as working dogs or for rodent control, the necessity of improving nutrition could be more assumable by owners, not only for lowering wildlife depredation, but also for pet welfare.

Conclusions

Unconfined dogs and cats in the Cape Horn Biosphere Reserve interact with wildlife, although this passes almost unperceived by the local community, particularly with dogs. To guarantee the future intactness of this wilderness area, it is essential to put the possible impacts of free-roaming pet carnivores on wildlife into perspective. This should be done using an integrative approach that respects the many dimensions of pet carnivores in their beneficial and problematic interactions with their human, conspecific, and natural environment: (1) Although over half of pets in Puerto Williams were sterilized, encouraging neutering, particularly in rural areas, could reduce pet density and avoid elimination or abandonment of unwanted pups/kittens. Moreover, sterilized dogs were described to show lower rates of escaping from home and less roaming behavior (Neilson, Eckstein & Hart, 1997; Spain, Scarlett & Houpt, 2004, but see Garde et al., 2015), which might lower their access to wildlife. (2) Despite its probable unpopularity (e.g., Grayson, Calver & Styles, 2002), we recommend that dog and cat ownership should be completely banned in

families living in Navy posts, as they are located within wild insular environments (some in National Parks) where pet interactions with wildlife can be particularly severe (Medina et al., 2011) and pets can become feral. (2) The evidence of a feral population of dogs on Navarino Island (where native predators **lack**) urgently needs biological methods of confirmation such as GPS monitoring (Young et al., 2011) and an assessment of its impacts. Feral dogs are still poorly investigated (Boitani et al., 2017) **management** implications for feral dogs are challenging (i.e., they may include the removal of dogs) due to their avoidance of human contact (Boitani & Ciucci, 1995) and restriction to natural habitats. (4) Increasing dog and cat confinement is beneficial to prevent not only pet-wildlife interactions, but also many socio-economic problems. In-house/yard confinement largely depends upon cultural settings (Hsu, Severinghaus & Serpell, 2003; Jackman & Rowan, 2007) and is low in many countries in Africa, Asia, and South America (Reece, 2005), including Chile, which has high percentages of free-roaming dogs in rural settings (67%, Acosta-Jamett et al., 2010; 84-91% Silva-Rodríguez & Sieving, 2012; 92% Sepúlveda et al., 2014). A new Chilean law promoting responsible pet ownership was recently enacted (Ley 21.020, 2017) and could be a first step towards regulating confinement by national legislation which was considered far from sufficient in 2014 by Bonacic & Abarca (2014). Meanwhile, bells or electronic alarms are an option for free-roaming cats to reduce depredation rates on wildlife (Gordon, Matthaei & Van Heezik, 2010; Calver & Thomas, 2011). (5) Finally, an instrument **such** as this survey can reveal what **experienced** problems with free-roaming pets exist; thus, **giving** insight into impacts in the absence of biological studies or complementary to them (e.g., dogs interacting with rare and endemic mammals were only detected through interviews in Silva-Rodríguez & Sieving, 2011). Owners' and non-owners' suggestions can also contribute to creating acceptable measures towards responsible pet ownership (Calver et al., 2011). For the CHBR, in addition to the above mentioned technical and legal solutions, social change to improve pet management for wildlife benefits could be stimulated through education about the vulnerability of native fauna to dogs and cats, attractive education material such as puppet videos, documentaries, or animal-borne films for school pupils, and action days such as walking with leashed dogs or breakfasts with dog owners (further examples in Miller, Ritchie & Weston, 2014).

Acknowledgements

This work was only possible due to the willingness and interest of the local community of Puerto Williams, farm owners, and Chilean Navy families to participate in our survey. We are grateful to the Chilean Navy for facilitating maritime transport to the Navy posts on different islands within the CHBR. We would like to thank Eduardo Silva for his advice in the questionnaire design and Mariela Alarcon and her team from the Scientific Ethical Committee of the University of Magallanes for improving an earlier version of the questionnaire. Amy Wynia, Nancyrose Houston, and two anonymous reviewers made valuable comments on an earlier version of the manuscript. We are also grateful to Andrés Mansilla and Ricardo Rozzi for their support in this study.

References

- Acosta-Jamett G. 2009. The role of domestic dogs in diseases of significance to humans and wildlife health in central Chile. D. Phil. Thesis, University of Edinburgh.
- Acosta-Jamett G, Cleaveland S, Cunningham AA, Bronsvoort BM deC. 2010. Demography of domestic dogs in rural and urban areas of the Coquimbo region of Chile and implications for disease transmission. *Preventive Veterinary Medicine* 94:272-281.
- Anderson CB, Rozzi R, Torres-Mura JC, McGehee SM, Sherriffs MF, Schüttler E, Rosemond AD. 2006. Exotic vertebrate fauna in the remote and pristine sub-Antarctic Cape Horn Archipelago, Chile. *Biodiversity and Conservation* 15:3295-3313.
- Arahoru M, Kuroshimaa H, Horia Y, Takagia S, Chijiiwa H, Fujita K. 2017. Owners' view of their pets' emotions, intellect, and mutual relationship: Cats and dogs compared. *Behavioural Processes*. <http://dx.doi.org/10.1016/j.beproc.2017.02.007>.
- Baker PJ, Boitani L, Harris S, Saunders G, White PCL. 2008. Terrestrial carnivores and human food production: Impact and management. *Mammal Review* 38:123-166.
- Bernard HR. 2006. Research methods in Anthropology. Qualitative and quantitative approaches. Oxford: Altamira Press.
- Boitani L, Ciucci P. 1995. Comparative social ecology of feral dogs and wolves. *Ethology Ecology & Evolution* 7: 49-72.
- Boitani L, Francisci F, Ciucci P, Andreoli G. 2017. The ecology and behavior of feral dogs: A case study from central Italy. In: Serpell J, ed. The domestic dog. Its evolution, behavior and interactions with people. Cambridge: Cambridge University Press, 342-368.

- Bonacic C, Abarca K. 2014. Hacia una política y legislación para el control de poblaciones de cánidos y calidad de vida de las personas: un enfoque multidisciplinario hacia una política y legislación para el control de poblaciones de cánidos y calidad de vida de las personas. *Centro de Políticas Públicas UC* 65:1-13.
- Burnham KP, Anderson DR. 2002. Model selection and multimodel inference. A practical information-theoretic approach. New York: Springer.
- Butler JRA, du Toit JT, Bingham J. 2004. Free-ranging domestic dogs *Canis familiaris* as predators and prey in rural Zimbabwe: threats of competition and disease to large wild carnivores. *Biological Conservation* 115:369-378.
- Calver MC, Grayson J, Lilith M, Dickman CR. 2011. Applying the precautionary principle to the issue of impacts by pet cats on urban wildlife. *Biological Conservation* 144:1895-1901.
- Calver MC, Thomas S. 2011. Effectiveness of the Liberator™ in reducing predation on wildlife by domestic cats. *Pacific Conservation Biology* 16:244-250.
- Chehébar C, Martín S. 1989. Guía para el reconocimiento microscópico de los pelos de los mamíferos de la Patagonia. *Doñana Acta Vertebrata* 16:247-291.
- Cunazza C. 1991. El guanaco, una especie de fauna silvestre con futuro. Corporación Nacional Forestal, Report Gerencia Técnica, Santiago, Chile.
- Echegaray J, Vilà C. 2010. Noninvasive monitoring of wolves at the edge of their distribution and the cost of their conservation. *Animal Conservation* 13:157-161.
- Garde E, Pérez PE, Vanderstichel R, Dalla Villa PF, Serpell JA. 2015. Effects of surgical and chemical sterilization on the behavior of free-roaming male dogs in Puerto Natales, Chile. *Preventive Veterinary Medicine* 123:106-120.
- Gompper ME. 2014a. The dog-human-wildlife interface: assessing the scope of the problem. In: Gompper ME, ed. Free-ranging dogs and Wildlife Conservation. Oxford: Oxford University Press, 9-54.
- Gompper ME. 2014b. Introduction: outlining the ecological influences of a subsidized domesticated predator. In: Gompper ME, ed. Free-ranging dogs and Wildlife Conservation. Oxford: Oxford University Press, 1-8.
- González-Esteban J, Villate I, Irizar I. 1996. Differentiating hair samples of the European mink (*Mustela lutreola*), the American mink (*Mustela vison*) and the European polecat (*Mustela putorius*) using light microscopy. *Journal of Mammalogy* 270:458-461.

- González BA. 2005. Informe sobre la situación de conservación de la población de guanacos (*Lama guanicoe*) en la costa Noroeste de isla Navarino, XII Región de Chile. Report, Fundación Biodiversitas, Chile.
- González B, Zapata B, Marín JC. 2002. Situación de conservación y percepción local sobre la población de guanacos más austral del mundo, Isla Navarino, XII Región de Chile. Report, Pontificia Universidad Católica de Chile.
- Gordon JK, Matthaei C, Van Heezik Y. 2010. Belled collars reduce catch of domestic cats in New Zealand by half. *Wildlife Research* 37:372-378.
- Gramza A, Teel T, VandeWoude S, Crooks K. 2016. Understanding public perceptions of risk regarding outdoor pet cats to inform conservation action. *Conservation Biology* 30:276–286.
- Grayson J, Calver MC, Styles I. 2002. Attitudes of suburban Western Australians to proposed cat control legislation. *Australian Veterinary Journal* 80:536-543.
- Hart LA, Yamamoto M. 2017. Dogs as helping partners and companions for humans. In: Serpell J, ed. The domestic dog. Its evolution, behavior and interactions with people. Cambridge: Cambridge University Press, 247-270.
- Hsu Y, Severinghaus LL, Serpell JA. 2003. Dog keeping in Taiwan: its contribution to the problem of free-roaming dogs. *Journal of Applied Animal Welfare Science* 6:1-23.
- Hughes J, Macdonald DW. 2013. A review of the interactions between free-roaming domestic dogs and wildlife. *Biological Conservation* 157:341-351.
- Hughes KA, Pertierra LR, Molina-Montenegro MA, Convey P. 2015. Biological invasions in terrestrial Antarctica: what is the current status and can we respond? *Biodiversity and Conservation* 24:1031-1055.
- Jackman J, Rowan A. 2007. Free-roaming dogs in developing countries: The benefits of capture, neuter, and return programs. In: Salem DJ, Rowan AN, eds. The state of the animals. Washington, DC: Humane Society Press, 55-78.
- Kays RW, DeWan AA. 2004. Ecological impact of inside/outside house cats around a suburban nature preserve. *Animal Conservation* 7:273-283.
- Ley 21.020. 2017. Sobre tenencia responsable de mascotas y animales de compañía. Available at <https://www.leychile.cl/Navegar?idNorma=1106037&r=6> (accessed 15 September 2017).

- Loyd KAT, Hernandez SM, Carroll JP, Abernathy KJ, Marshall GJ. 2013. Quantifying free-roaming domestic cat predation using animal-borne video cameras. *Biological Conservation* 160:183-189.
- Maccoby EE, Maccoby N. 1954. The interview: A tool of social science. In: Lindzey G, ed. *Handbook of Social Psychology*. Cambridge: Addison-Wesley, 449-487.
- Manor R, Saltz D. 2004. The impact of free-roaming dogs on gazelle kid/female ratio in a fragmented area. *Biological Conservation* 119:231-236.
- Matter HC, Daniels TJ. 2000. Dog ecology and population biology. In: Macpherson CNL, Meslin, F, Wandeler AI, eds. *Dogs, zoonoses and public health*. Wallingford: CABI Publishing, 17-50.
- Medina FM, Bonnaud E, Vidal E, Tershy BR, Zavaleta ES, Donlan CJ, Keitt BS, Le Corre M, Horwath SV, Nogales M. 2011. A global review of the impacts of invasive cats on island endangered vertebrates. *Global Change Biology* 17:3503-3510.
- Meek PD. 1999. The movement, roaming behavior and home range of free-roaming domestic dogs, *Canis lupus familiaris*, in coastal New South Wales. *Wildlife Research* 26:847-855.
- Miller KK, Ritchie EG, Weston MA. 2014. The human dimensions of dog-wildlife interactions. In: Gompper ME, ed. *Free-ranging dogs and Wildlife Conservation*. Oxford: Oxford University Press, 286-304.
- Mitchell BD, Banks PB. 2005. Do wild dogs exclude foxes? Evidence for competition from dietary and spatial overlap. *Australian Ecology* 30:581-591.
- Mittermeier RA, Mittermeier CG, Brooks TM, Pilgrim JD, Konstant WR, da Fonseca GAB, Kormos C. 2003. Wilderness and biodiversity conservation. *Proceedings of the National Academy of Sciences of the United States of America* 100:10309-10313.
- Mundry R. 2014. Statistical issues and assumptions of phylogenetic generalized least squares. In: Garamszegi L, ed. *Modern phylogenetic comparative methods and their application in Evolutionary Biology*. Berlin, Heidelberg: Springer, 131-153.
- Neilson JC, Eckstein RA, Hart BL. 1997. Effects of castration on problem behaviors in male dogs with reference to age and duration of behavior. *Journal of the American Veterinary Medical Association* 211:180-182.
- Nogales M, Vidal E, Medina FM, Bonnaud E, Tershy BR, Campbell KJ, Zavaleta ES. 2013. Feral cats and biodiversity conservation. The urgent prioritization of island management. *BioScience* 63:804-810.

- O'Brien SJ, Johnson WE. 2007. The evolution of cats. *Scientific American* 297:68-75.
- Payne E, Bennett PC, McGreevy PD. 2015. Current perspectives on attachment and bonding in the dog-human dyad. *Psychology Research and Behaviour Management* 8:71-79.
- Pisano E. 1977. Fitogeografía de Fuego-Patagonia chilena. I. Comunidades vegetales entre las latitudes 52 y 56 S. *Anales del Instituto de la Patagonia* 8:121-250.
- Plaza PI, Lambertucci SA. 2017. How are garbage dumps impacting vertebrate demography, health, and conservation? *Global Ecology and Conservation* 12:9-20.
- Randi E. 2008. Detecting hybridization between wild species and their domesticated relatives. *Molecular Ecology* 17:285-293.
- R Core Team. 2016. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. Available at <https://www.R-project.org/>.
- Reece JF. 2005. Dogs and dog control in developing countries. In: Salem DJ, Rowan AN, eds. The state of the animals III, Washington, D.C.: Humane Society Press, 55-64.
- Reponen SEM, Brown SK, Barnett BD, Sacks BN. 2014. Genetic and morphometric evidence on a Galápagos Island exposes founder effects and diversification in the first-known (truly) feral western dog population. *Molecular Ecology* 23:269-283.
- Rozzi R, Massardo F, Berghöfer A, Anderson CB, Mansilla A, Mansilla M, Plana J, Berghöfer U, Araya P, Barros E. 2006. Cape Horn Biosphere Reserve: Nomination document for the incorporation of the Cape Horn Archipelago Territory into the World Biosphere Reserve Network. MaB Program – UNESCO. Punta Arenas: Ediciones de la Universidad de Magallanes.
- Rozzi R, Arango X, Massardo F, Anderson C, Heidinger K, Moses K. 2008. Field environmental philosophy and biocultural conservation: The Omora Ethnobotanical Park Educational Program. *Environmental Ethics* 30:325-336.
- Ruiz-Izaguirre E, van Woersem A, Eilers K, van Wieren SE, Bosch G, van der Zijpp AJ, de Boer IJM. 2014. Roaming characteristics and feeding practices of village dogs scavenging sea-turtle nests. *Animal Conservation* 18:146-156.
- Schalamon J, Ainoedhofer H, Singer G, Petnehazy T, Mayr J, Kiss K, Höllwarth ME. 2006. Analysis of dog bites in children who are younger than 17 years. *Pediatrics* 117:e374-e379.

- Schüttler E, Klenke R, McGehee S, Rozzi R, Jax K. 2009. Vulnerability of ground-nesting waterbirds to predation by invasive American mink in the Cape Horn Biosphere Reserve, Chile. *Biological Conservation* 142:1450-1460.
- Schüttler E, Rozzi R, Jax K. 2011. Towards a societal discourse on invasive species management: A case study of public perceptions of mink and beavers in Cape Horn. *Journal for Nature Conservation* 19:175-184.
- Sepúlveda MA, Singer RS, Silva-Rodríguez E, Stowhas P, Pelican K. 2014. Domestic dogs in rural communities around protected areas: conservation problem or conflict solution? *PLoS ONE* 9(1): e86152.
- Sepúlveda MA, Pelican K, Cross P, Eguren A, Singer R. 2015. Fine-scale movements of rural free-ranging dogs in conservation areas in the temperate rainforest of the coastal range of southern Chile. *Mammalian Biology* 80:290-297.
- Sih A, Bolnick DI, Luttbeg B, Orrock JL, Peacor SD, Pintor LM, Preisser E, Rehage JS, Vonesh JR. 2010. Predator-prey naïveté, antipredator behavior, and the ecology of predator invasions. *Oikos* 119:610-621.
- Silva-Rodríguez EA, Ortega-Solís GR, Jiménez JE. 2010. Conservation and ecological implications of the use of space by chilla foxes and free-ranging dogs in a human-dominated landscape in southern Chile. *Austral Ecology* 35:765-777.
- Silva-Rodríguez EA, Sieving KE. 2011. Influence of care of domestic carnivores on their predation on vertebrates. *Conservation Biology* 25:808-815.
- Silva-Rodríguez EA, Sieving KE. 2012. Domestic dogs shape the landscape-scale distribution of a threatened forest ungulate. *Biological Conservation* 150:103-110.
- Slater MR, Di Nardo A, Pediconi O, Villa PD, Candeloro L, Alessandrini B, Del Papa S. 2008. Free-roaming dogs and cats in central Italy: public perceptions of the problem. *Preventive Veterinary Medicine* 84:27-47.
- Spain CV, Scarlett JM, Houpt KA. 2004. Long-term risks and benefits of early-age gonadectomy in dogs. *Journal of the American Veterinary Medical Association* 224:380-387.
- Symonds MRE, Moussalli A. 2011. A brief guide to model selection, multimodel inference and model averaging in behavioural ecology using Akaike's Information Criterion. *Behavioral Ecology and Sociobiology* 65:13-21.

- 689 Vanak AT, Gompper ME. 2009. Dogs *Canis familiaris* as carnivores: their role and function in
690 intraguild competition. *Mammal Review* 39:265-283.
- 691 Vanak AT, Thaker M, Gompper ME. 2009. Experimental examination of behavioural interactions
692 between free-ranging wild and domestic canids. *Behavioral Ecology and Sociobiology* 64:279-
693 287.
- 694 White PCL, Jennings NV, Renwick AR, Barker NHL. 2005. Questionnaires in ecology: a review
695 of past use and recommendations for best practice. *Journal of Applied Ecology* 42:421-430.
- 696 Williams KJH, Weston MA, Henry S, Maguire GS. 2009. Birds and beaches, dogs and leashes:
697 Dog owners' sense of obligation to leash dogs on beaches in Victoria, Australia. *Human*
698 *Dimensions of Wildlife* 14:89-101.
- 699 Woods M, McDonald RA, Harris S. 2003. Predation of wildlife by domestic cats *Felis catus* in
700 Great Britain. *Mammal Review* 33:174-188.
- 701 Young JK, Olson, KA, Reading RP, Amgalanbaatar S, Berger J. 2011. Is wildlife going to the
702 dogs? Impacts of feral and free-roaming dogs on wildlife populations. *BioScience* 61:125-132.
- 703 Zuur AF, Ieno EN, Elphick CS. 2010. A protocol of data exploration to avoid common statistical
704 problems. *Methods in Ecology & Evolution* 1:3-14.

Figure 1

Map of the Cape Horn Biosphere Reserve, southern Chile.

The Alberto de Agostini and Cape Horn National Parks are core areas of the CHBR and Yendegaia is a recently-created national park. The only large settlement (2,800 inhabitants) is Puerto Williams, on Navarino Island. Eleven Chilean Navy posts are located throughout the reserve on Tierra del Fuego Island (n=2), Navarino Island (n=2), and uninhabited islands (n=7); all are only accessible via maritime transport, except the western Navy post on Navarino Island.




Figure 2

Free-roaming dog and cat sightings on Navarino Island, southern Chile.

Approximate sighting locations of unaccompanied adult dogs and cats, dog pups (abandoned or feral), and kittens (abandoned) from n=227 sightings by 143 participants during 2014/15. Dog sightings are shown in different classes of pack size.





Figure 3

Experienced and suspected problems with dogs and cats in southern Chile.


Problematic experiences with dogs during 2009-2015 A) inside (n=143) and B) outside (n=44) of Puerto Williams, suspected dog problems (first problem mentioned) C) inside (n=221) and D) outside of town (n=202), and suspected cat problems (first problem mentioned) E) outside of town (n=77). Conflicts with people included biting, attacking, frightening, disease transmitting, accidents, and stealing. Dog-domestic animal problems referred to killing, attacking, or feeding on free-ranging domestic animals such as cows, horses (*Equus caballus*), sheep, pigs (*Sus scrofa*), and cats, whereas cat-domestic animal problems referred to preying upon chickens (*Gallus gallus domesticus*). Conflicts with wildlife included killing wild animals such as birds and their eggs, North American beavers (*Castor canadensis*), and guanacos, or harming ecosystems. Conflicts with conspecifics were fights among dog packs or between cats, and disease transmission. Dog feces and waste dispersing were considered as hygienic problems. "Other" includes dog and cat overpopulation, bad image for tourists, and barking. Images of animals represent predominant animals involved in dog/cat problems.

Experienced dog problems (%) Suspected dog problems (%) Suspected cat problems (%)

A Inside town




C Inside town




Problems with:

- People
- Domestic animals
- Conspecifics
- Wildlife
- Hygiene
- Other

B Outside town



D Outside town



E Outside town




Figure 4

Preyed and harassed animals by dogs and cats, and suspected feral dog prey in southern Chile.

Prey brought to owners by 31 of 168 dogs (n=40 mentioned items), animals observed to be harassed by 64 dogs (n=79 items), prey brought to owners by 27 of 52 cats (n=33 items), animals observed to be harassed by 19 cats (n=25 items), and suspected prey of feral dogs (n=494 items) by 244 participants. "Other" includes fish (dog prey); fish, foxes, and seals, (harassed by dogs); bats (cat prey); and bird and horse feces, fish bait, foxes, seals, and vegetable material (suspected feral dog prey). "Small livestock" refer to chickens, pigs, and sheep. All bird species mentioned were native, except for the house sparrow, species mentioned among the cat prey.

**Note: Auto Gamma Correction was used for the image. This only affects the reviewing manuscript. See original source image if needed for review.*






Figure 5

Model averaged odds ratios for models predicting dog and cat wildlife interactions in southern Chile.

Plots show the model averaged parameter estimates as odds ratios on a log scale with 95% confidence intervals (CI) for A) dog model and B) cat model, where LOCATION (dogs and cats) and FOOD (only cats) best predicted whether pets brought wildlife prey home. The other variables had confidence intervals that clearly overlapped the dashed line at 1, which implies that there is no direction of the parameter estimate. Estimates with odds ratios <1 indicate a negative association with the response variable, whereas those >1 indicate a positive association.



Odds ratio (log scale) and 95% CI



Odds ratio (log scale) and 95% CI

Table 1(on next page)

Candidate models for predicting dog and cat wildlife interactions in southern Chile.

A detailed description of the response and predictor variables is provided in Table 2.

<i>Candidate model sets</i>	<i>Response variable</i>	<i>Quantitative value of response variable</i>	<i>Predictor variables</i>	<i>n**</i>
Dog model	PREY	19.0% of dogs brought prey home	FOOD+LOCATION+SEX SIZE* STERILIZED*	163
Cat model	PREY	51.9% of cats brought prey home	FOOD+LOCATION+SEX STERILIZED*	52

* Predictor variable  included in final model set due to collinearity

** We deleted 5  from the dog model (3.0%) and 1  from the cat model (1.9%)

Table 2(on next page)

Variable description of candidate models for predicting dog and cat wildlife interactions in southern Chile.

The variable description refers to the questionnaire (see Supplementary Material S1).

<i>Variable</i>	<i>Variable type</i>	<i>Variable description and question number</i>	<i>Categories</i>
PREY	Response	Dog/cat brought wildlife prey home (Q25)	Yes/no
FOOD	Predictor	Feeding of participant's dog/cat with leftovers, commercial food/meat, or both (Q15)	0=leftovers 1=leftovers and commercial food/meat 2=commercial food/meat
LOCATION	Predictor	Dog/cat lives in rural environment (farm /Navy post) or in Puerto Williams	Rural/village
SEX	Predictor	Dog's/cat's sex (Q2)	F=Female, M=Male
SIZE*	Predictor	Dog's size (mean of two independent estimations by ES and LSA, Q42)	1=small, 1.5=small to medium-sized, 2=medium-sized, 2.5=medium-sized to large, 3=large
STERILIZED	Predictor	Dog/cat is sterilized (Q11)	0=not sterilized 1=pets' age<1 year 2=sterilized

* Variable only used in the dog model, not in the cat model

Table 3(on next page)

Demographic dog and cat data, and husbandry results from southern Chile.

We obtained data on owned dog and cat populations via questionnaires from households in Puerto Williams (n=215), accessible farm households on Navarino Island (n=7), and Navy families (n=22, data from two years) living on the 11 Navy posts on different islands within the CHBR.

	<i>Town households</i> (n=215)		<i>Farm households</i> (n=7)		<i>Navy posts</i> (n=22)	
	<i>Dogs</i>	<i>Cats</i>	<i>Dogs</i>	<i>Cats</i>	<i>Dogs</i>	<i>Cats</i>
Demographic data						
Households with pet ownership (%)	85 (39.5)	30 (14.0)	6 (85.7)	6 (85.7)	19 (86.4)	4 (18.2)
Mean pet number per household (SD)	1.4 (0.9)	1.2 (0.6)	5 (3.3)	2.5 (2.5)	1.3 (0.5)	1.0 (0.0)
Total pet number	121	36	30	15	17	2
Male:female ratio	1.3:1	0.7:1	2:1	0.3:1	0.7:1	0:1
Mean pet age (SD)	3.7 (3.8)	4.8 (4.1)	3.8 (4.6)	3.0 (3.0)	3.0 (3.7)	3.3 (1.1)
Number of offspring in previous year	16	0	21	7	9	0
Local origin (CHBR, %)	66.1	66.7	100	100	29.4	100
Reproductive control						
Females spayed;	66.7;	71.4;	10.0;	8.3;	11.1;	0.0;
males neutered (%)	52.2	93.3	20.0	0.0	16.7	0.0
Health						
Vaccinated against rabies (%)	55.4	33.3	0.0	0.0	64.7	0.0
Treated for parasites (%)	60.3	36.1	100.0	40.0	64.7	50.0
Food provisioning						
Commercial food and/or meat (%)	77.7	94.4	86.7	20.0	82.3	50.0
Leftovers (%)	12.4	0.0	13.3	33.3	5.9	0.0
Mix of above (%)	9.9	5.6	0.0	46.7	11.8	50.0
Dog confinement						
Free-roaming during day or night (%)	30.6	-	46.7	-	82.4	-
24-h free-roaming (%)	19.0	-	30.0	-	41.2	-

1

2

Table 4(on next page)

Best-ranked generalized linear models for predicting dog and cat wildlife interactions in southern Chile.

Summary of model selection for models with $\Delta AIC_c < 2$. K indicates number of parameters per model, ΔAIC_c distance from lowest AIC_c , and ω_i model weight.

<i>Model set</i>	<i>Competing models</i>	<i>k</i>	<i>AIC_c</i>	<i>ΔAIC_c</i>	<i>ω_i</i>
Dog model	FOOD+LOCATION	3	150.72	0.00	0.37
	LOCATION	2	151.03	0.32	0.32
	FOOD+LOCATION+SEX	4	152.32	1.60	0.17
Cat model	FOOD+LOCATION	3	62.19	0.00	0.75

1

2