

Microcos magnifica (~~Malvaceae-Grewioideae~~ or Sparrmanniaceae) a new species of cloudforest tree from Cameroon.

Formatted: Left

Martin Cheek¹

¹ Science, Royal Botanic Gardens, Kew, Richmond, Surrey, TW9 3AE, UK.
m.cheek@kew.org

ABSTRACT.

Background: ~~although~~ Although many new species to science have been discovered from thousands of specimens resulting from botanical inventories to support conservation management in Cameroon in recent years, additional species remain to be formally evaluated taxonomically and described. These include species from genera which have been taxonomically neglected for many decades in Africa, such as *Microcos*.

Methods: This study is based mainly on herbarium specimens and field observations made in Cameroon during a series of botanical surveys. Herbarium material was examined with a Leica Wild M8 dissecting binocular microscope fitted with an eyepiece graticule.

Principal findings: *Microcos magnifica* Cheek (~~Malvaceae-Grewioideae~~ or Sparrmanniaceae) is described as an Endangered (EN B2 ab(iii)) new tree species from the submontane forests of Cameroon. It is illustrated and described, and its conservation status and taxonomic affinities are assessed. It is the first new *Microcos* described from Africa in more than 90 years and is unique on the continent in having sculptured fruits.

Discussion: ~~A~~ A systematic revision, with a molecular phylogenetic study, of *Microcos* in Africa is necessary if the affinities of the species, including *M. magnifica*, are to be reliably established.

Key words. Conservation, Endangered, Mt Kupe, submontane, taxonomy.

INTRODUCTION

During identification of specimens resulting from botanical surveys of Mt Kupe and the Bakossi Mts in SW Region, Cameroon, specimens of a remarkable undescribed *Microcos* came to light, which were ~~designated~~ as *Microcos* sp. A (Cheek in Cheek et al. 2004: 414). Subsequently an additional specimen was discovered in the forests of Ebo, Littoral Region. Here these specimens are formally named as *Microcos magnifica* Cheek, the first new species to science to be described in the genus for Africa for 90 years. The species is remarkable for its sculptured fruit surfaces which are verrucate. Sculptured fruit surfaces are not otherwise known in the African species but do occur in some Asian species.

Microcos Burm. ex L. (1753) is a palaeotropical genus of about 77 species (Govaerts et al. continuously updated) based on *M. paniculata* Burm. ex L. (1753) from Sri Lanka. Linnaeus (1767) later synonymised *Microcos* under *Grewia* L. However, the genus was resurrected by Burret (1926). Burret's authoritative revision (1926) of former Tiliaceae sens. lat. presaged its break-up into today's Brownloideae/Brownlowiaceae, Tiliaceae sensu stricto/Tilioideae

and Grewioideae/Sparrmanniaceae (with the largest number of genera and species) including *Microcos* (Bayer et al. 1999, Bayer & Kubitzki 2003, Cheek 2007a-c). In addition, *Schoutenia* included by Burret in Tiliaceae, is now placed in Dombeyoideae/Pentapetaceae (Cheek 2007d). Burret's was the last global treatment of *Microcos* (1926). He recognised 53 species, of which 19 were recorded from Africa and 34 in Asia.

Of the 99 names in *Microcos* listed in IPNI (continuously updated), Govaerts et al. (continuously updated) accept 77 names. ~~77~~ the majority of these 77 are in S.E. Asia, but with 10 in Africa. The genus is absent from the Neotropics and Madagascar.

Formatted: Font: Not Italic

Illogically, while *Microcos* has been maintained as a separate genus from *Grewia* ~~L.~~ in Asia (e.g. Chung 2003, 2006, Chung et al. 2005a, Chung & Soepadmo 2011), the two genera have often been united under *Grewia* in Africa. For example, in one of the most recent Flora accounts of *Grewia* (including *Microcos*) for Africa, Whitehouse (2001) states "...Kirkup followed Burret in recognising *Microcos* as a distinct genus; this concept has also been followed in SE Asia. Although there are clear differences between *Microcos* and the other sections of *Grewia*-, for consistency I am following the practice set by the other African floras, of not recognising...." This practice is maintained widely today, for example by the excellent and essential African Plant Database (continuously updated).

In fact the two genera are readily recognised as expressed in the key below, modified from that in Whitehouse (2001):

-Trees and climbers, rarely shrubs, of evergreen forest; stigmas entire; fruit unlobed; inflorescences terminal, sometimes axillary also, many-flowered.....***Microcos***

Shrubs, rarely trees, of bushland or woodland; stigmas lobed; fruit 4-lobed, rarely entire; inflorescences usually axillary or leaf-opposed, rarely terminal, usually few-flowered.....***Grewia***

According to the molecular analysis of Brunken & Muellner (2012), *Microcos* is not embedded in *Grewia*, neither are they sister groups, and they fall into distinct clades. Additional characters for separating the two genera are found in the pollen, wood anatomy and in the leaf anatomy, particularly the epidermal cells. ~~-(Chattaway 1934,~~ Chung 2002, Chung et al. 2003, 2005b). *Microcos* was maintained in Bayer & Kubitzki (2003).

The genus *Microcos* has been little studied in Africa, as evidenced by the fact that the first new name in African *Microcos* since 1926 was published in 2004 (*Microcos barombiensis* (K. Schum.) Cheek in Cheek et al. 2004: 414). In the course of matching the material described as new in this paper, it became clear that a revision of the genus for Africa is desirable to address specimen misidentifications and additional apparently undescribed species. It is hoped to address ~~these problems~~ in a future paper.

MATERIALS & METHODS

The electronic version of this article in Portable Document Format (PDF) will represent a published work according to the International Code of Nomenclature for algae, fungi, and plants (ICN), and hence the new names contained in the electronic

version are effectively published under that Code from the electronic edition alone. In addition, new names contained in this work which have been issued with identifiers by IPNI will eventually be made available to the Global Names Index. The IPNI LSIDs can be resolved and the associated information viewed through any standard web browser by appending the LSID contained in this publication to the prefix "<http://ipni.org>". The online version of this work is archived and available from the following digital repositories: PeerJ, PubMed Central, and CLOCKSS.

This study is based mainly on herbarium specimens and field observations made in Cameroon during a series of botanical surveys beginning in 1991. [These surveys were](#) mainly led by the author, ~~which~~ so far [they](#) have resulted in 52,450 specimens being studied at K and YA, of which 37,850 were newly collected, the data stored on the Kew Cameroon specimen Access database (Gosline, p. 11 in Cheek et al. 2004). The top set of specimens was initially deposited at SCA, and later YA, duplicates being sent to K. The fieldwork was approved by the Institutional Review Board of the Royal Botanic Gardens, Kew entitled the Overseas Fieldwork ~~Ceommittee~~ (OFC). ~~eThe~~ ~~most; the most~~ recent [OFC approval is being](#) numbered 807. The most recent invitation to effect research on the flora and vegetation of Cameroon [has the reference number 050/IRAD/DG/CRRA-NK/SSRB-HN/09/2016](#). It is issued under the terms of the 5 year Memorandum of Collaboration between [Institute for Research in Agricultural Development \(IRAD\)](#)-Herbier National du Cameroun and Royal Botanic Gardens, Kew signed 5th Sept 2014, ~~has the reference number 050/IRAD/DG/CRRA-NK/SSRB-HN/09/2016 from~~

All specimens cited have been seen by the author unless indicated n.v. Herbarium citations follow Index Herbariorum (Thiers et al. continuously updated) and binomial authorities IPNI (continuously updated). Material of the suspected new species was compared morphologically with material of all other African *Microcos* (or *Grewia* sect. *Microcos* ([L.](#)) [Wight & Arnott](#)) principally at K, but also using material from WAG. [This comprised about 350 specimens](#). The online search address used for retrieving specimen data from labels at P was <https://science.mnhn.fr/institution/mnhn/collection/p/item/p00179355?listIndex=128&listCount=610http://coldb.mnhn.fr/catalognumber/mnhn/p/p00375109>. [Burret's types of *Microcos* at B were destroyed by allied bombing in 1943 so it was not possible to consult them. This has necessitated that subsequent authors select neotypes of his names, e.g. Whitehouse \(2001\).](#)

The conservation assessment was made using the categories and criteria of IUCN (2012). [The](#), ~~and~~ extent of occurrence was calculated with Geocat (Bachman et al. 2011). Herbarium material was examined with a Leica Wild M8 dissecting binocular microscope. [This was](#) fitted with an eyepiece graticule measuring in units of 0.025 mm at maximum magnification. The drawing was made with the same equipment using Leica 308700 camera lucida attachment.

RESULTS

KEY TO THE SPECIES OF MICROCOS TREE SPECIES IN AFRICA WEST OF [DEMOCRATIC REPUBLIC OF CONGO](#) & THE CONGO RIVER

1. Leaves deeply toothed. Maiombe Mts of Cabinda.....***M. gossweileri***

Leaves entire. Nigeria to Congo-Brazzaville, but unknown from Cabinda.....2

2. Leaf base cuneate, leaf surfaces glabrous; fruits glossy, smooth.....**M. coriacea**

Leaf base truncate or cordate, leaf blade lower surface stellate hairy; fruits matt, [wart-verrucate](#).....**M. magnifica**

Microcos magnifica Cheek species novum

Holotype: Cameroon, S.W. Province, Mt Kupe, Kupe village, main trail towards summit, fr. 9 July 1996, *Etuge* 2886 (holo. K; isotypes BR, K, MO, P, SCA, US, WAG, YA) (Fig. 1, Map 1)

Microcos sp. A, Cheek (in Cheek et al. 2004: 414).

Tree 20–35 m tall, 30–70 cm diameter at breast height, crown small, bole straight, base of bole with 4–5 concave slender buttresses reaching 1 to 1.5 m above the ground where sometimes spreading up to 2.5 m from the trunk and branching. Bark dull medium red–brown, fibrous; slash hard fibrous-granular, without scent or exudates, white, oxidising rapidly from white to red.

Leafy stems 3–5 mm diameter below the third node, finely longitudinally ridged, densely minutely grey–brown puberulent, internodes 2.5 cm long. Leaves obovate, obovate-oblong [or](#) elliptic, 12.5–25.5 x 6.6–13.5 cm (those of sterile stems large, to 28 cm long), acumen 0.4–1.8 cm long, base truncate or truncate and abruptly cordate, margin entire, lateral nerves 11–13 on each side of the midrib, the basal pair more conspicuous by virtue of a pair subsidiary nerves, brochidodromous, domatia absent, tertiary nerves strongly scalariform, quaternary nerves inconspicuous: upper surface with midrib varied, convex, densely and minutely grey–brown puberulent, secondary nerves flat but also puberulent: lower surface with midrib and secondary nerves strongly raised, brownish green, the areolae pale green or brown/khaki densely puberulent with minute pale brown 8–20-armed stellate hairs 0.1–0.2 mm diameter, touching each other, more or less completely concealing the epidermis. Presumed shade leaves (larger, from sterile branches – *Elad* 118) with hairs sparse, separated by 1 or 2 hair diameters, smaller, 0.075–0.1 mm diameter, with only 6–8 (–12) arms. Petiole stout, cylindrical, (1.5–)1.8–2 x 0.3 cm. Stipules caducous, not seen, but leaving an arched scar 4 mm long on the stem each side and 1 mm below the insertion of the leaf base.

Inflorescence and flowers unknown. Infructescence terminal, paniculate, 11–16 x 5.5–13 cm, bearing 5–13 (–12) fruits; peduncle 1.5–2.7 cm; bracts not seen; pedicel absent, fruits articulated at junction with stem.

Fruits fleshy, red [when live](#), drying pink–brown, [ring brown](#), obovoid to ellipsoid 2–2.4 x 1.2–1.5 cm, verrucate and finely longitudinally wrinkled with 20–25 verrucae, verrucae 0.5–1 mm long, patent. Mesocarp: outer part thin and fleshy, inner part thick and densely fibrous. Endocarp obovoid, slightly 3-angled, [woody](#), whitish brown, sutures longitudinal, alternating with three lines of hairs; locule 1, probably by abortion from 3, 1-seeded.

Seed narrowly ovoid, glabrous, slightly laterally compressed, hilum subapical; endosperm extensive, embryo flattened.

Phenology: [fruiting](#)~~Fruiting~~: April to July; flowering: unknown.

Distribution and habitat: SW and Littoral Regions of Cameroon; submontane or submontane-lowland forest with *Medusandra mpomiana*, *Santiria trimera*, *Allanblackia gabonensis*, *Coelocaryon preussii* (Mt. Kupe); *Pycnanthus*, *Staudtia*, *Coelocaryon*, *Petersianthus*, *Strombosia* and *Maesobotrya* (Ebo); 750–1000 m alt.

Etymology: ~~meaning~~ Meaning magnificent, for the spectacular and unusual fruit ornamentation.

Affinity: ~~resembling~~ Resembling *Microcos coriacea* Mast., -but fruits verrucate and matt, not smooth and glossy; leaves with base truncate or cordate, not cuneate; lower surface densely white stellate hairy, not glabrous).

Additional specimens: South West Region. Mt Kupe, 15 Km WNW de Tombel, colline 930m de NW de Ngussi, fr. 21 April 1976, *Letouzey* 14669 (P n.v.; YA 3 sheets); Mt Kupe, Nyasoso, trails above village, 4° 49'N; 9° 41'E, st. 6 Feb. 1995, *Elad* 118 (K, YA n.v.); Nyasoso, Max's trail, fr. 3 June 1996, *Cable* 2806 (K, YA); Kupe village, main trail towards summit, fr. 9 July 1996, *Etuge* 2886 (holo. K; iso. BR, K, MO, P, SCA, US, WAG, YA). Littoral Region, Yingui, Ebo proposed National Park, 6 hours walk S. of Iboti village; between the abandoned villages of Bekob and Masseng, 4 21 50 N; 10 25 20 E, st. 16 Feb. 2006, *Cheek* 12980 (K, SCA, YA)

Conservation: *Microcos magnifica* is here assessed as Endangered (EN B2 ab(iii)) using the IUCN 2012 system, since it is known from four threat-based locations with an extent of occurrence of 303 km² calculated using Geocat (Bachman et al. 2011) and an area of occupancy of 16 km² using IUCN preferred 4 km² grid cells. The species is threatened at all its known locations, most immediately the three in the Mt Kupe area of Ngussi and Nyassoso where clearance of forest continues upslope from the volcanic, fertile lowlands of the Chide valley. The clearance is for small-holder agriculture, principally for food crops. The locations concerned are all far outside the Mt Kupe Ecological Reserve and on the edges of towns. It is quite likely that some or all of the trees that provided the specimens and the forest remnants in which they occurred, have been cleared already (Cheek pers. obs.). In order to reduce the threat to the species here, a local conservation poster featuring the species is intended in order to raise awareness of the existence and importance of its protection.

~~However~~ However, at the fourth location, in the proposed Ebo National Park, the species is secure from immediate threat, there being no resident human population.

~~However~~ However, the future of Ebo as a protected area is not certain, and logging, plantation and mining are all threatened as alternative uses for the land. Since there is no indication that more than a single mature individual has ever been recorded at each of the four locations, it is conceivable that *Microcos magnifica* might be better assessed as Critically Endangered under Criterion D of IUCN (less than 50 mature individuals recorded).

DISCUSSION

The affinities of *M. magnifica* may be with the only two other arborescent species of *Microcos* that occur in West-Central Africa (see key to species above). Of these, only *M. coriacea* is sympatric with *M. magnifica*. At Mt Kupe the two species have differing altitudinal ranges, *M. coriacea* with a range of 200–420 m, based on four records, and *M. magnifica* -with 900–1000 m, based on three records (Cheek in Cheek et al. 2004: 414). It can be postulated that *M. magnifica* has arisen as a submontane derivative of *M. coriacea*. However, among the taxa discovered as new at Mt Kupe at

Formatted: No underline

Formatted: Font: Not Italic

similar altitudes to *M. magnifica* was *Kupea martinetei* Cheek (Cheek et al. 2003) which has its sister species in the Eastern Arc Mts of Tanzania (Cheek 2004).

Formatted: Font: Not Italic

Exactly the same geographic range as *Microcos magnifica*, which extends disjunctly from the western slopes of Mt Kupe to the submontane N-S ridge of the Ebo forest, is seen also in *Uvariopsis submontana* Kenfack (Kenfack et al. 2003) and *Costus kupensis* ~~H.Maas & Maas~~ (Maas-van de Kamer et al. 2016). It is remarkable that none of these conspicuous species has been discovered in the submontane ridge of the Ngovayang massif to the east at Bipindi, nor in the Bakossi Mts, immediately West of Mt Kupe, despite significant botanical surveys in these areas by Zenker and by K-YA teams respectively. This suggests that these distributions are real and not the result of undercollecting. ~~However~~ ~~However~~, all of these species are infrequent and only known from three to six specimens in as many locations.

The discovery of such a distinctive new species in the Kupe-Bakossi, Ebo and adjoining areas is not unusual. Among other species discovered here were (in alphabetical order by genus):

Allophylus ujori Cheek (Cheek & Etuge 2009a), *Ancistrocladus grandiflorus* Cheek (Cheek 2000b), *Brachystephanus kupeensis* Champl. (Champluvier & Darbyshire 2009), *Chassalia laikomensis* Cheek (Cheek & Csiba 2000), *Coffea montekupensis* Stoff. (Stoffelen et al. 1997), *Coffea bakossii* Cheek & Bridson (Cheek et al. & Bridson 2002), *Cola metallica* Cheek (Cheek 2002), *Coleochloa domensis* Muasya & D.A. Simpson (Muasya et al. 2010), *Deinbollia oreophila* Cheek (Cheek & Etuge 2009b), *Diospyros kupensis* Gosline (Gosline & Cheek 1998); *Dovyalis cameroonensis* Cheek & Ngolan (Cheek & Ngolan 2007), *Dracaena kupensis* Mwachala, Cheek, Eb. Fisch. et al. (Mwachala et al. 2007), *Impatiens etindensis* Cheek & Eb. Fisch. et al. (Cheek & Fischer 1999), *Impatiens frithii* Cheek (Cheek & Csiba 2002b), *Isoglossa dispersa* I.Darbysh. (Darbyshire et al. 2011), *Kupea martinetei* Cheek & S.-A.-Williams (Cheek et al. 2003), *Ledermanniella onanae* Cheek (Cheek 2003), *Ledermanniella pollardiana* Cheek & Ameka (Cheek & Ameka 2008), *Memecylon kupeanum* R.D.Stone, Ghogue & Cheek (R.D.Stone et al. 2008), *Mussaenda epiphytica* Cheek (Cheek 2009), *Newtonia duncanthomasii* Mackinder & Cheek (Mackinder & Cheek 2003), *Oxyanthus okuensis* Cheek & Sonké (Cheek & Sonké 2000), *Psychotria darwiniana* Cheek (Cheek et al. 2009), *Psychotria geophylax* Cheek & Sonké and *P. bakossiensis* Cheek & Sonké (2005), *Psychotria kupensis* Cheek (Cheek et al. 2008), *Psychotria moseskemei* Cheek (Cheek & Csiba 2002a), *Rhaptopetalum geophylax* Cheek & Gosline (Cheek et al. 2002) and *Ternstroemia cameroonensis* Cheek (Cheek et al. 2017).

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Formatted: French (Cameroon)

Formatted: French (Cameroon)

Formatted: French (Cameroon)

Formatted: French (Cameroon)

Formatted: French (Cameroon)

Formatted: French (Cameroon)

Formatted: French (Cameroon)

Formatted: French (Cameroon)

Formatted: French (Cameroon)

Formatted: French (Cameroon)

Commented [DL1]: Publication seems to be dated 2016 although issued in 2017?

Formatted: French (Cameroon)

Most of these species are threatened with extinction, since they are narrow endemics with small ranges, restricted to mainly submontane (cloud) forest which is steadily being cleared, mainly for small-scale cultivation of food crops. These species feature in the Red Data Book of Cameroon (Onana & Cheek 2011).

CONCLUSIONS

A systematic revision, with a molecular phylogenetic study, of *Microcos* in Africa is necessary if the affinities of the species, including *M. magnifica*, are to be reliably established.

Acknowledgements

The botanical surveys which resulted in the specimens cited being collected were mainly supported by Earthwatch (1993–2005) and by the Darwin Initiative of the UK Government through the Plant Conservation of Western Cameroon, and Red Data Book of Cameroon projects both led by RBG, Kew, working with the IRAD-National Herbarium of Cameroon. Bate Oben is thanked for assisting by making morphological measurements. George Gosline is thanked for producing the map, and Saba Rokni for providing the Geocat analysis. Gaston Achoundong, Jean-Michel Onana, former heads of the National Herbarium of Cameroon (YA) and their successor Marie Florence Ngo Ngwe are thanked for their collaboration and support of the Royal Botanic Gardens, Kew's activities over the years. Laurence J. Dorr and two other, anonymous, reviewers are thanked for their constructive comments on an earlier version of this manuscript.

References

African Plant Database (version 3.4.0), [\(continuously updated\)](#). Conservatoire et Jardin botaniques de la Ville de Genève and South African National Biodiversity Institute, Pretoria, "Retrieved [March 2017]", from <http://www.ville-ge.ch/musinfo/bd/cjb/africa/>.

Formatted: Font: 12 pt

Formatted: Font: 12 pt, Not Bold

Formatted: Font: 12 pt

Bachman S, Moat J, Hill A, de la Torre J, Scott B. (2011.) Supporting Red List threat assessments with GeoCAT: geospatial conservation assessment tool. *ZooKeys* **150**:117–126. <https://doi.org/10.3897/zookeys.150.2109>

Formatted: Font: Bold

Formatted: Font: Bold

Bayer C, Fay MF, De Bruijn AY, Savolainen V, Morton CM, Kubitzki K, Alverson WS, Chase MW. 1999. Support for an expanded family concept of Malvaceae within a recircumscribed order Malvales: a combined analysis of plastid atpB and rbcL DNA sequences. *Botanical Journal of the Linnean Society* **129**:267–303.

Formatted: Font: Bold

Formatted: Font: Bold

Bayer C, Kubitzki K. 2003. Malvaceae. In: Kubitzki K, Bayer C (eds), *The families and genera of vascular plants 5: flowering plants dicotyledons*: 225–311. Springer-Verlag, Berlin.

Formatted: Font: Bold

Brunken U, Muellner AN. 2012. A New Tribal Classification of Grewioideae (Malvaceae) Based on Morphological and Molecular Phylogenetic Evidence. *Systematic Botany* **37**(3):699–711. <http://www.bioone.org/doi/full/10.1600/036364412X648670>

Formatted: Font: Bold

Formatted: Font: Bold

Formatted: Font: Italic

Formatted: Font: Bold

Burret M. 1926. Beiträge zur Kenntnis der Tiliaceae I. *Notizblatt des Königlich-Botanischen Gartens und Museums zu Berlin-Dahlem* **9**:592–880.

Formatted: Font: Bold

Formatted: Font: Bold

Champluvier D, Darbyshire I. 2009. A revision of the genera *Brachystephanus* and *Oreacanthus* (Acanthaceae) in tropical Africa. *Systematics and Geography of Plants* **79**(2):115–192. DOI: 10.2307/25746

Formatted: Font: Bold

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Bold

Chattaway MM. 1934. Anatomical evidence that *Grewia* and *Microcos* are distinct genera. *Tropical Woods* 38:9-11.

Cheek M. 2000. A synoptic revision of *Ancistrocladus* (Ancistrocladaceae) in Africa, with a new species from western Cameroon. *Kew Bulletin* 55:871-882. DOI: 10.2307/4113632

<https://doi.org/10.2307/4113632>

Cheek M. 2002. Three new species of *Cola* (Sterculiaceae) from western Cameroon, Cameroon. *Kew Bulletin* 57:402-415. DOI: 10.2307/4111117

Cheek M. 2003. A new species of *Ledermanniella* (Podostemaceae) from western Cameroon. *Kew Bulletin* 58:733-737. DOI: 10.2307/4111153

Cheek M. 2004. Kupeaeae, a new tribe of Triuridaceae from Africa. *Kew Bulletin* 58:939-949. <https://doi.org/10.2307/4111207>

Cheek M. 2007a. *Brownlowia* sp. nov. (Sparmanniaceae). In: Heywood VH, Brummitt RK, Culham A, Seberg O (eds), *Flowering plant families of the world*: 307-308. Royal Botanic Gardens, Kew

Cheek M. 2007b. *Tiliaceae*. In: Heywood VH, Brummitt RK, Culham A, Seberg O (eds), *Flowering plant families of the world*: 321-322. Royal Botanic Gardens, Kew

Cheek M. 2007c. *Sparmanniaceae*. In: Heywood VH, Brummitt RK, Culham A, Seberg O (eds), *Flowering plant families of the world*: 307-308. Royal Botanic Gardens, Kew

Cheek M. 2007d. *Pentapetaceae*. In: Heywood VH, Brummitt RK, Culham A, Seberg O (eds), *Flowering plant families of the world*: 247-248. Royal Botanic Gardens, Kew

Cheek M. 2009. *Mussaenda epiphytica* sp. nov. (Rubiaceae) an epiphytic shrub from cloud forest of the Bakossi Mts, western Cameroon. *Nordic Journal of Botany* 27(6): 456-459. DOI: 10.1111/j.1756-1051.2009.00576.x <https://doi.org/10.1111/j.1756-1051.2009.00576.x>

Cheek M, Ameka G. 2008. *Ledermanniella pollardiana* sp. nov. (Podostemaceae) from western Cameroon. *Nordic Journal of Botany* 26:214-217. DOI: 10.1111/j.1756-1051.2008.00162.x

Formatted: Font: Bold

Formatted: Font: Bold

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Bold

Formatted: Font: (Default) Times New Roman, 12 pt

Formatted: Font: 12 pt

Formatted: Font: Bold

Formatted: Font: Italic

Formatted: Font: Bold

Formatted: Font: 12 pt, Not Bold

Formatted: Font: Bold

Formatted: Font: Italic

Formatted: Font: Bold

Formatted: Font: 12 pt, Not Bold

Formatted: Font: Italic

Formatted: Font: Bold

Formatted: Font: Bold

Formatted: Font: Not Italic

Formatted: Font: Bold

Formatted: Font: 12 pt

Formatted: Font: 12 pt

Formatted: Font: Bold

Formatted: Font: Bold

Formatted: Font: Bold

Formatted: Font: Bold

Formatted: Font: Bold

Formatted: Font: Italic

Formatted: Font: Bold

Formatted: Font: 12 pt

Formatted: Font: 12 pt

Formatted: Font: Bold

Formatted: Font: Italic

Formatted: Font: Bold

Cheek M, **Csiba L**, Bridson D. 2002. A new species of *Coffea* (Rubiaceae) from western Cameroon. *Kew Bulletin* **57**:675–680. DOI: [10.2307/4110998](https://doi.org/10.2307/4110998)

Cheek M, Corcoran M, & Horwath A. 2009. Four new submontane species of *Psychotria* (Rubiaceae) with bacterial nodules from western Cameroon. *Kew Bulletin* **63**:405–418. <https://doi.org/10.1007/s12225-008-9056-4>

Cheek M, Csiba L. 2000. A new species and new combination in *Chassalia* (Rubiaceae) of western Cameroon. *Kew Bulletin* **55**:883–888. DOI: 10.2307/4113633

Cheek M, Csiba L. 2002a. A revision of the *Psychotria chalconeura* complex (Rubiaceae) in Guineo-Congolian Africa. *Kew Bulletin* **57**:375–387. DOI: 10.2307/4111113

Cheek M, Csiba L. 2002b. A new epiphytic species of *Impatiens* (Balsaminaceae) from western Cameroon. *Kew Bulletin* **57**(3):669–674. DOI: 10.2307/4110997

Cheek M, Etuge M. 2009a. *Allophylus -conraui* (Sapindaceae) -reassessed and *Allophylus ujori* described from western Cameroon. *Kew Bulletin* **64**(3):495–502. doi:10.1007/s12225-009-9139-x

Cheek M, Etuge M. 2009b. A new submontane species of *Deinbollia* (Sapindaceae) from Western Cameroon and adjoining Nigeria. *Kew Bulletin* **64**(3):503–508. doi:10.1007/s12225-009-9132-4

Cheek M, Fischer E. 1999. A tuberous and epiphytic new species of *Impatiens* (Balsaminaceae) from Southwest Cameroon. *Kew Bulletin* **xx**: 471-475. <https://doi.org/10.2307/4115828>

Cheek M, Gosline G, Csiba L. 2002. A new species of *Rhaptopetalum* (Scytopetalaceae) from western Cameroon. *Kew Bulletin* **57**(3):661–667. DOI: 10.2307/4110996

Cheek M, Horwath A, Haynes D. 2008. *Psychotria kupensis* (Rubiaceae) a new dwarf, litter gathering species from western Cameroon. *Kew Bulletin* **63**:243–246. <https://doi.org/10.1007/s12225-008-9018-x>

Cheek M, Ngolan R. 2007. A reassessment of the *Dovyalis spinosissima* Gilg (Flacourtiaceae) complex in Africa, with a new species from Cameroon. *Kew Bulletin* **61**:595-600. <http://www.jstor.org/stable/20443304>

Cheek M, Pollard BJ, Darbyshire I, Onana J-M, Wild C. 2004. *The Plants of Kupe, Mwanenguba and the Bakossi Mountains, Cameroon. A Conservation Checklist*. Royal Botanic Gardens, Kew. London.

Formatted: Font: Bold

Formatted: Font: Italic

Formatted: Font: Bold

Formatted: Font: 12 pt

Formatted: Font: Bold

Formatted: Font: Not Italic

Formatted: Font: Bold

Formatted: Font: 12 pt

Formatted: Font: Bold

Formatted: Font: Italic

Formatted: Font: Bold

Formatted: Font: Bold

Formatted: Font: Italic

Formatted: Font: Bold

Formatted: Font: Bold

Formatted: Font: Italic

Formatted: Font: Bold

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Bold

Formatted: Font: Bold

Formatted: Font: Italic

Formatted: Font: Bold

Formatted: Font: Bold

Formatted: Font: Bold

Formatted: Font: Bold

Formatted: Font: Bold

Commented [DL2]: Volume number is not given.

Formatted: Font: Italic, Highlight

Formatted: Highlight

Formatted: Font: Not Bold, Highlight

Formatted: Highlight

Formatted: Font: 12 pt, Highlight

Formatted: Highlight

Formatted: Font: 12 pt

Formatted: Font: Bold

Formatted: Font: Italic

Formatted: Font: Bold

Formatted: Font: Bold

Formatted: Font: Bold

Formatted: Font: Italic

Formatted: Font: Bold

Formatted: Font: 12 pt

Formatted: Font: Bold

Formatted: Font: Italic

Formatted: Font: Bold

Formatted: Font: (Default) Times New Roman, 12 pt

Formatted: Font: Bold

Cheek M, Sonké B. 2000. A new species of *Oxyanthus* (Rubiaceae-Gardeniinae) from western Cameroon. *Kew Bulletin* **55**:889–893. DOI: 10.2307/4113634

Formatted: Font: Bold

Formatted: Font: Italic

Cheek M, Sonké B. 2005. Two further new species of *Psychotria* (Rubiaceae) from western Cameroon. *Kew Bulletin* **60**:–293–300. Stable URL:

Formatted: Font: Bold

Formatted: Font: Italic

<http://www.jstor.org/stable/4110940>

Formatted: Font: Bold

Cheek M, Tchiengue B, Tacham WN. 2016. *Ternstroemia cameroonensis* (Ternstroemiaceae), a new medicinally important species of montane tree, nearly extinct in the Highlands of Cameroon. *Blumea* **62**, 2017:–53 – 57.

Formatted: Font: Bold

Formatted: Font: Bold

Formatted: Font: Italic

DOI: <https://doi.org/10.3767/000651917X695362>.

Formatted: Font: Italic

Cheek M, Williams S, Etuge M. 2003. *Kupea martinetugei*, a new genus and species of Triuridaceae from western Cameroon. *Kew Bulletin* **58**:–225–228. DOI: 10.2307/4119366

Formatted: Font: Bold

Formatted: Font: Italic

Formatted: Font: Bold

Chung RCK. 2002. Leaf epidermal micromorphology of *Grewia* L. and *Microcos* L. (Tiliaceae) in Peninsular Malaysia and Borneo. *Gardens' Bulletin Singapore* **54**:–263–286.

Formatted: Font: Bold

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Bold

Chung RCK. 2003. New taxa and new combinations of *Microcos* (Tiliaceae) from Peninsular Malaysia and Borneo. *Kew Bulletin* **58**:–329–349.

Formatted: Font: Bold

Formatted: Font: Italic

Formatted: Font: Not Bold, Italic

Chung RCK. 2006. A revision of *Grewia* (Malvaceae-Grewioideae) in Peninsular Malaysia and Borneo. *Edinburgh Journal of Botany* **62**:–1–27.

Formatted: Font: Bold

Formatted: Font: Bold

Formatted: Font: Italic

Chung RCK, Lim SC, Lim AL, Soepadmo E. 2005ba. Wood anatomy of *Grewia* and *Microcos* from Peninsular Malaysia and Borneo. *Journal of Tropical Forest Science* **17**:–175–196.

Formatted: Font: Bold

Formatted: Font: Bold

Formatted: Font: Italic

Chung RCK, Soepadmo E, Lim AL. 2003. The significance of pollen morphology in the taxonomy of *Grewia* and *Microcos* (Tiliaceae) in Peninsular Malaysia and Borneo. *Gardens' Bulletin Singapore* **55**:–239–256.

Formatted: Font: Italic

Formatted: Font: Bold

Formatted: Font: Bold

Formatted: Font: Italic

Chung RCK, Soepadmo E, Lim AL. 2005a. A synopsis of the Bornean species of *Microcos* L. (Tiliaceae). *Gardens' Bulletin Singapore* **57**:–103–132.

Formatted: Font: Italic

Formatted: Font: Bold

Formatted: Font: Bold

Formatted: Font: Italic

Chung RCK, Soepadmo E. 2011. Taxonomic revision of the genus *Microcos* (Malvaceae-Grewioideae) in Peninsular Malaysia and Singapore. *Blumea* **56**:–273–299. <http://dx.doi.org/10.3767/000651911X619704>

Formatted: Font: Bold

Formatted: Font: Not Bold, Highlight

Formatted: Font: Italic

Formatted: Font: Bold

Darbyshire I, Pearce L, Banks H. 2011. The genus *Isoglossa* (Acanthaceae) in west West Africa. *Kew Bulletin* **66**(3):–425–439. DOI: 10.1007/s12225-011-9292-x

Formatted: Font: Bold

Formatted: Font: Italic

Formatted: Font: Bold

Gosline G, Cheek M. 1998. A new species of *Diospyros* (Ebenaceae) from Southwest Cameroon. *Kew Bulletin* **53**(2):–461–465 DOI: 10.2307/4114512

Formatted: Font: Bold

Formatted: Font: Italic

Formatted: Font: Bold

Govaerts, RHA, Belyaeva I, Hartley H, & Lindon H. (continuously updated).

Microcos Burm. ex L. in: *Plants of the World Online*. Kew Science.

powo.science.kew.org/taxon/urn:lsid:ipni.org:names:328157-2

Formatted: Font: Not Bold

IPNI (continuously updated). The International Plant Names Index. <http://ipni.org/>.

Formatted: Font color: Auto

Formatted: Normal, Pattern: Clear

IUCN. 2012. *IUCN Red List Categories and Criteria*: Version 3.1. Second edition.

Formatted: Font: Not Bold

Gland, Switzerland and Cambridge, UK: IUCN. Available from:

<http://www.iucnredlist.org/> (accessed: 01/2017).

Kenfack D, Gosline G, Gereau, RE, Schatz G. 2003. The genus *Uvariopsis* in Tropical Africa, with a recombination and one new species from Cameroon. *Novon* 13:443-449

Formatted: Font: Italic

Formatted: Font: Bold

Linnaeus C. 1753. *Species plantarum*. Salvii, Stockholm.

Formatted: Font: Bold

Linnaeus C. 1767. *Systema naturae* ed. 12, 1, 2. Salvii, Stockholm.

Formatted: Font: Italic

Mackinder B, Cheek M. 2003. A new species of *Newtonia* (Leguminosae-Mimosoideae) from Cameroon. *Kew Bulletin* 58:447-452. DOI: 10.2307/4120627

Formatted: Font: Bold

Formatted: Font: Italic

Formatted: Font: Bold

Formatted: Font: Italic

Formatted: Font: Bold

Maas-van de Kamer H, Maas PJM, Wieringa JJ, Specht CD. 2016. Monograph of African *Costus*. *Blumea - Biodiversity, Evolution and Biogeography of Plants* 61(3): 280-318. DOI: <https://doi.org/10.3767/000651916X694445>

Formatted: Font: Italic

Formatted: Font: Italic

Muasya A, Harvey YH, Cheek M, Tah K, Simpson DA. 2010. A new species of epiphytic *Coleochloa* (Cyperaceae) from Cameroon. *Kew Bulletin* 65:1-3. DOI: 10.1007/s12225-010-9194-3

Formatted: Font: Bold

Formatted: Font: Bold

Formatted: Font: Italic

Formatted: Font: Italic

Mwachala G, Cheek M, Fischer E, and A.M. Muasya AM. 2007. A new species of *Dracaena* L. (Ruscaceae) from Cameroon. *Kew Bulletin* 64:613-616.

Formatted: Font: Bold

Formatted: Font: Bold

<http://www.jstor.org/stable/20443393>

Formatted: Font: Bold

Onana J-M, Cheek M. 2011. *The Red Data Book, Plants, of Cameroon*. Royal Botanic Gardens, Kew.

Formatted: Font: (Default) Times New Roman, 12 pt

Formatted: Font: Bold

Stoffelen P, Cheek M, Bridson D, Robbrecht E. 1997. A new species of *Coffea* (Rubiaceae) and notes on Mt Kupe (Cameroon). *Kew Bulletin* 52(4):989-994. DOI: 10.2307/3668527

Formatted: Font: Bold

Formatted: Font: Italic

Formatted: Font: Italic

Stone RD, Ghogue J-P & Cheek M. 2008. Revised treatment of *Memecylon* sect. *Afzeliana* (Melastomataceae-Olisbeoideae), including three new species from Cameroon. *Kew Bulletin* 63:227-241. <https://doi.org/10.1007/s12225-008-9033-y>

Formatted: Font: Bold

Formatted: Font: Bold

Formatted: Font: Italic

Formatted: Font: Italic

Thiers B (continuously updated). *Index Herbariorum: A global directory of public herbaria and associated staff*. New York Botanical Garden's Virtual Herbarium.

Formatted: Font: Bold

Formatted: Font: 12 pt

~~[continuously updated]~~. Available from: <http://sweetgum.nybg.org/ih/> (accessed: Jan. 2017).

Formatted: Font: Italic

Whitehouse C. 2001. *Grewia*, pp. 7–63 in Whitehouse et al. Tiliaceae. *Flora of Tropical East Africa*. Royal Botanic Gardens, Kew.

Formatted: Font: Bold

Formatted: Font: Italic

CAPTION FOR FIGURE.

Fig. 1. *Microcos magnifica* **A** habit, fruiting branch; **B** habit sketch; **C** leaf variation: *Etuge* 2686: hairs on upper surface (left); hairs on lower surface (right). **D** leaf variation: *Elad* 118, with detail of hairs on lower surface; **E** leaf variation: *Cable* 2806, with detail of hairs on lower surface; **F** fruit, side view; **G** fruit, left with pericarp removed exposing mesocarp fibres; left while, right longitudinal section (endocarp stippled, endosperm densely stippled), **H** endocarp, left, side view; right, distal end view; **I** endocarp with seed, transverse section. **A**, **C**, **F-K** from *Etuge* 2686; **B** from field observations of *Cheek* 12980; **D** *Elad* 128; **E** *Cable* -2806. All drawn by ANDREW BROWN.

Map 1. Global distribution of *Microcos magnifica*.