

# The importance of local settings: within-year variability in seawater temperature at South Bay, Western Antarctic Peninsula

César A Cárdenas <sup>Corresp., 1</sup>, Marcelo González-Aravena <sup>1</sup>, Pamela A. Santibañez <sup>1</sup>

<sup>1</sup> Departamento Científico, Instituto Antártico Chileno, Punta Arenas, Chile

Corresponding Author: César A Cárdenas  
Email address: ccardenas@inach.cl

The Western Antarctic Peninsula (WAP) has undergone significant changes in air and seawater temperatures during the last 50 years. Although highly stenotherm Antarctic organisms are expected to be severely affected by the increase of seawater temperature, high-resolution datasets of seawater temperature within coastal areas of the WAP (where diverse marine communities have been reported) are not commonly available. Here we report on within-year (2016-2017) variation in seawater temperature at three sites on Doumer Island, Palmer Archipelago, WAP. Within a year, Antarctic organisms in South Bay were exposed to water temperatures in excess of 2° C for more than 25 days and 2.5° C for more than 10 days. We recorded a temperature range between -1.7° to 3.0° C. Warming of seawater temperature was 3.75 times faster after October 2016 than it was before October. Results from this study indicate that organisms at South Bay are already exposed to temperatures that are being used in experimental studies to evaluate physiological responses to thermal stress in WAP organisms. Continuous measurements of short to long-term variability in seawater temperature provides important information for parametrizing meaningful experimental treatments that aim to assess the local effects of environmental variation on Antarctic organisms under future climate scenarios.

**The importance of local settings: within-year variability in seawater temperature at South Bay, Western Antarctic Peninsula**

César A. Cárdenas, Marcelo González-Aravena, Pamela A. Santibáñez

<sup>1</sup>Departamento Científico, Instituto Antártico Chileno (INACH), Plaza Muñoz Gamero 1055, Punta Arenas 6200965, Chile

Corresponding Author:

César A. Cárdenas

Email address: [ccardenas@inach.cl](mailto:ccardenas@inach.cl)

# **Abstract**

The Western Antarctic Peninsula (WAP) has undergone significant changes in air and seawater temperatures during the last 50 years. Although highly stenothermal Antarctic organisms are expected to be severely affected by the increase of seawater temperature, high-resolution datasets of seawater temperature within coastal areas of the WAP (where diverse marine communities have been reported) are not commonly available. Here we report on within-year (2016-2017) variation in seawater temperature at three sites on Doumer Island, Palmer Archipelago, WAP. Within a year, Antarctic organisms in South Bay were exposed to water temperatures in excess of 2° C for more than 25 days and 2.5° C for more than 10 days. We recorded a temperature range between -1.7° to 3.0° C. Warming of seawater temperature was 3.75 times faster after October 2016 than it was before October. Results from this study indicate that organisms at South Bay are already exposed to temperatures that are being used in experimental studies to evaluate physiological responses to thermal stress in WAP organisms. Continuous measurements of short to long-term variability in seawater temperature provides important information for parametrizing meaningful experimental treatments that aim to assess the local effects of environmental variation on Antarctic organisms under future climate scenarios.

# **Introduction**

Although there is an absence of continent-scale warming over Antarctica over the last 100 years (Stenni et al., 2017), during the last 50 years the Western Antarctic Peninsula (WAP) has undergone significant warming in air and seawater temperatures, with the later increasing by ~ 2° C (Cook et al., 2016; Meredith & King, 2005; Stenni et al., 2017). Warming has contributed to the range expansion of various marine flora (e.g., Convey & Smith, 2006; Smith, 1994; Hill et al., 2011), the loss of glacier mass, the retreat of ice shelves, and summer melting, with the latter

being unprecedented over the past 1,000 years (Abram et al., 2013; Paolo, Fricker & Padman, 2015). Increased snowfall on the Antarctic Peninsula has been linked to warming sea surface temperatures in the western Pacific (Thomas et al., 2015, 2017) and the warming of WAP mid-ocean depths (rather than the warming of air temperatures) has been proposed as the primary driver of glacier retreat (e.g., Cook et al., 2016). These changes highlight the complexity of environmental change occurring at the WAP and the need for high-resolution information on regional- to local-scale patterns of climate change, to address its implications for marine polar ecosystems (e.g., Clarke et al., 2007; Meredith, Stefels & van Leeuwe, 2017).

Many Antarctic species are not well adapted to sudden temperature change because they have evolved in one of the most stable environments on Earth (e.g., Peck, 2005; Morley et al., 2016; Peck, Webb & Bailey, 2014). Some Antarctic species are sensitive to even small changes in temperature (2 - 3°C above the annual average), losing the ability to perform essential functions (Ingels et al., 2012; Peck, Morley & Clark, 2010; Peck, Webb & Bailey, 2004; Peck, 2005; Pörtner, Peck & Somero, 2007). For example, the Antarctic scallop *Adamussium colbecki* loses its ability to perform essential activities (Peck, 2005) with increases in seawater temperature of 2-3 °C, and attempts to acclimate individuals to temperatures above 2°C have failed for this species (Bailey, Johnston & Peck, 2005). Experiments using the Antarctic clam *Laternula elliptica*, have shown that half of the individual died in 60 days at 3°C, and for the brittle star *Ophionotus victoriae* half of the individuals died in less than 30 days at 3°C (Turner et al., 2009). In contrast others organisms, such as the sea urchin *Sterechinus neumayeri*, are comparatively resistant to increases in seawater temperature (Ericson et al., 2012; Suckling et al., 2015). Recent studies have highlighted the dramatic effects of warming waters on Antarctic organisms at the community level (Ashton et al., 2017), where small increases in temperature (1°

C) altered community composition by reducing overall species diversity and evenness. The study of resistance, adaptation and tolerance to environmental variation has become a key focus of evaluating how Antarctic ecosystems respond to climate change (Gutt et al., 2012; Ingels et al., 2012; Peck, 2005). Research to date has focused on identifying how Antarctic organisms respond to the expected seawater temperature increase at community (Ashton et al., 2017), assemblage (Peck et al., 2009) and species (physiological and molecular) levels (Cascella et al., 2015; Ericson et al., 2012; Foo et al., 2016; González et al., 2016; Kapsenberg & Hofmann, 2014; Peck et al., 2009, 2014; Schram et al., 2015; Suckling et al., 2015), with the latter being the most studied. To inform the design of studies that are relevant to future climate scenarios high-resolution records of local to regional patterns of water temperature are required.

Due to the physical constraints that the Antarctic continent imposes on equipment and sampling, obtaining long-term high-resolution water temperature data is often logistically challenging. The urgent need to increase the spatio-temporal coverage of surface seawater temperature data, with a focus on coastal areas, has been highlighted by several authors (Gutt et al., 2017; Stenni et al., 2017). Although a number of moorings have been deployed along channels and open waters in the WAP (See [www.soos.aq/activities/soos-at-sea/moorings](http://www.soos.aq/activities/soos-at-sea/moorings)), there remains a lack of long-term, high-resolution temperature data from shallow waters in coastal zones dominated by diverse communities of sessile organisms with limited behavioral thermoregulatory ability (Ashton et al., 2017). Fine-scale temporal (daily) data can be crucial for inferring climatic impacts on survival, growth and reproduction (Kearney, Matzelle & Helmuth, 2011), while monthly data is better suited to studying relationships between population dynamics and climatic conditions (Gutt et al., 2017; Jan et al., 2017; Kennicutt et al., 2016). Observational gaps in surface seawater temperature leave fractionated information that does not provide a

coherent and robust picture of the current and future conditions occurring in Antarctic ecosystems. In order to fill some of these gaps, high-resolution temporal observations are required to advance ecosystem modelling and projections (Gutt et al., 2017).

Here we report for the first time extreme changes in shallow seawater temperature around Doumer Island, Palmer Archipelago (WAP). We present shallow seawater temperatures recorded over one year at high temporal resolution (hours), which can be used to parameterize future physiological experiments.

## Material and Methods

This study was conducted from the Chilean Yelcho Scientific Station in South Bay, Doumer Island, Palmer Archipelago, WAP (Fig. 1), where diverse benthic communities have been described (Cárdenas et al., 2016; Zamorano, 1983) (Fig. 2). Three HOBO pendant temperature/light data loggers (Part #UA-002-XX; Onset Computer Corp; accuracy:  $\pm 0.5^\circ \text{C}$  at  $0^\circ \text{C}$ , resolution:  $0.1^\circ \text{C}$  at  $0^\circ \text{C}$ ) were deployed in January 2016 and recovered in February 2017 at three sites in South Bay (Fig. 1). Sites were separated by  $\sim 1.5 \text{ km}$  and are part of a research project assessing the responses of Antarctic sponge assemblages to climate change. Data loggers from two of the three sites (PY1;  $64^\circ 52' 28.1'' \text{S}$ ;  $63^\circ 34' 36.0'' \text{W}$  and PY4;  $64^\circ 51' 58.6'' \text{S}$ ;  $63^\circ 37' 46.7'' \text{W}$ ) were at 10 m depth, whereas the third data logger (PY2;  $64^\circ 52' 15.3'' \text{S}$ ;  $63^\circ 33' 52.2'' \text{W}$ ) was at a depth of 20 m. Data was recorded every 2 hours at sites PY2 and PY4, and every 1 hour at PY1. Sites PY1 and PY2 were located inside the bay and were characterized by dense beds of *Himantothallus grandifolius* with a sub-canopy of crustose calcareous algae and sponges on a boulder/gravel substrate (Fig. 2). PY4 was located at Cape Kemp, which is regularly exposed to swell and is dominated by a mix of *H. grandifolius*, *Desmarestia* spp. and *Ascoseira mirabilis* on a bedrock substrate. Benthic coverage of sponges at PY4 was higher than

in sites inside the bay (Bravo, 2017). The data loggers at PY1, PY2 and PY4 recorded data for 404, 374 and 312 days, respectively. Data from PY4 stopped by late November (2016), presumably after being damaged by an iceberg. The study was conducted under the permit 806/2015 granted by the Chilean Antarctic Institute (INACH).

In order to describe non-linear, site-specific variability in the within-year patterns of seawater temperature, we fit separate Generalized Additive Models (GAMs) to each of the three sites, as they do not prescribe any particular form for the trend (Ferguson et al., 2008; Hastie & Tibshirani 1990; Orr et al., 2015). We chose GAMs instead of other techniques (e.g. linear regression) because: (1) GAMs do not assume linearity and allow specification of the mean value of the response and the systematic component using a link-function (Wood, 2006; Zuur, Ieno & Smith, 2007); (2) GAMs allow the application of an autoregressive structure to account for dependency of serial correlations in time series data; and (3) GAMs allow the use of thin-plate shrinkage smoothers to model temporal trends (recommended in Wood, 2006). The estimated smooth trends are described by: (i) the shape of the estimated curve, (ii) a hypothesis test for no variation over time, and (iii) *edf*, which describes effect complexity and corresponds to the amount of information used by each smoother. An *edf* of 0 indicates no variation, an *edf* of 1 indicates a linear relationship between the response and the covariate, and higher values correspond to more complex non-linear relationships; and (4) GAMs allow the use of finite differences (Curtis & Simpson, 2014; Monteith et al., 2014) to estimate periods where the rate of change (slope) is statistically different from zero (i.e., periods where the temperature is either increasing or decreasing in a significant manner). This was achieved by computing the first derivatives of the fitted trend. Uncertainty estimates were also calculated for the derivatives to form approximate 95% confidence intervals.

All models were fit using the R-programming language (version 3.3.3 by R Foundation for Statistical Computing, 2017) and the package mgcv (Wood, 2004; Wood, 2006). The R code and scripts are available in the supplemental material.

## Results

All seawater temperature records showed significant seasonality ( $p\text{-value} < 0.01$ ; Fig. 3), with clear minima in winter (July – September) and maxima in summer (January-March). Although data loggers were located at different sites and depths (Fig. 1) they showed a similar seasonal pattern throughout the year (Fig. 4), implying that the three sites are representative of broader shallow waters temperatures in South Bay. In addition, sea surface temperatures within the same time frame from Port Arthur (Palmer Station, ~25 km from Yelcho Station) show similar temperature ranges and patterns suggesting that the temperature variability reported could be representative of the wider Palmer Archipelago. Mean daily temperatures ranged from  $-1.7^{\circ}$  to  $3.0^{\circ}$  C (range =  $4.7^{\circ}$  C, SD = 1.1) at PY1 (the longest record, 404 days), from  $-1.6^{\circ}$  to  $1.00^{\circ}$  C (range =  $2.6^{\circ}$  C, SD = 0.8) at PY4 (the shortest record, 312 days), and from  $-1.8^{\circ}$  to  $1.8^{\circ}$  C (range =  $3.6^{\circ}$  C, SD = 0.8) at PY2. Seawater temperature differed between the 2016 (max  $0.8^{\circ}$  C) and 2017 (max  $3.0^{\circ}$  C) summers at PY1 ( $p\text{-value} < 0.01$ ; Figs. 5 and S1).

To identify periods of statistically significant change in the time series of seawater temperature at PY1 we calculated the first derivative method from a second GAM model ( $k = 58$ ,  $edf = 55.7$ ,  $p\text{-value} < 0.001$ , Figs. 5 and S2), which allowed a closer examination of the temperature variability at PY1 site. Figure 5 shows there were 26 periods of statistically significant change throughout the year, 12 increases and 13 decreases. From all periods of significant increasing temperatures, the rate of temperature increase ( $^{\circ}\text{C day}^{-1}$ ) to which marine organisms were subjected varied between 0.11 to  $0.15^{\circ}\text{C day}^{-1}$  (mean 0.13, SD = 0.01) after



October 2016. In contrast, the rate of increase before October 2016 ranged between 0.04 and 0.07 °C day<sup>-1</sup> (mean = 0.05, SD = 0.02; Table 1).

## Discussion

In recent years, several studies have focused on assessing how Antarctic organisms will adapt to future environmental conditions (e.g., Cascella et al., 2015; Ericson et al., 2012; Peck, 2011, 2016; Peck et al., 2014; Schram et al., 2015). Existing studies have used different species, methodologies and warming rates, which sometimes make comparing species-specific responses and treatment effects difficult (Clark et al., 2017). In addition, most experimental studies have used standard scenarios of climate change that are not necessarily relevant to nearshore areas that may be affected by a higher degree of temperature variability than other marine environments (e.g., Vargas et al., 2017). It is therefore important to incorporate realistic scenarios of temperature change into evaluations of ecosystem responses to climate change across multiple spatial and temporal scales (Gutt et al., 2017).

The present work provides the first yearlong record of shallow seawater temperatures around Doumer Island and one of the few on the Palmer Archipelago, WAP. Results show short-term variability and significant warming events. This constitutes important baseline information that could be used to assess long-term trends in local seawater temperature, and inform future physiological and molecular experimental studies in the area. Although HOBO data loggers have an accuracy of  $\pm 0.5^{\circ}\text{C}$  at  $0^{\circ}\text{C}$  which could increase the temperature range reported in our study, our results are in accordance with those obtained from 24-hour CTD measurements carried out in parallel by other researchers during certain days of the field season. Results demonstrate that marine organisms, at least in some coastal areas of Antarctica, are already exposed seawater temperatures that have been widely used in physiological tolerance

experiments to evaluate the effects of future climate scenarios. For example, observed temperatures in this study of more than 2.0 °C for more than 25 days or 2.5 °C for more than ten days within a year, have been used experimentally to test responses to acute heat stress (e.g. Peck et al., 2009). Further, Waller et al. (2017) reported for the first time the presence of the bivalve *Laternula elliptica* (has been used as a model to test the responses of Antarctic organisms to warming sea temperatures) in intertidal habitats where sediment and air temperature reached more than 7 to 10 °C, respectively. That study (Waller et al., 2017) and the continuous records of water temperature reported here suggest that we may need to rethink the temperature ranges used in experimental setups to better reflect local conditions, as organisms occurring on different regions of Antarctica might respond differently to environmental variation.

Short-term variability in water temperature is relatively low in Antarctic coastal environments, with a mean annual temperature range of ~3.1° C at some places such as Anvers Island (Barnes et al., 2006); hence, it is expected to have very little impact on ecology (e.g., Clarke, 2001). However, recent work highlights the importance of environmental variability at different scales, with acute thermal stress events and even timing of seasonal events being suggested as critical factors driving physiological, ecological and evolutionary processes (See Fordham et al., 2017 and references therein). Although our time-series of continuous, directly-measured sea temperatures is limited to a year, it shows the complete annual pattern and existence of dynamic variations within South Bay, Doumer Island. Our results are in accordance with sea surface temperature data from Port Arthur, Palmer Station in Anvers Island (<http://pal.lternet.edu>), which is approximately 25 km west of Doumer Island, suggesting this pattern occurs at least throughout the Palmer Archipelago (Fig. 4). Additionally, there are some

long-term data sets at WAP for a few areas (e.g. Palmer LTER, Rothera Station-BAS Database) that allowed comparison.

Our results show that marine organisms at South Bay were exposed to a broad annual temperature range from  $-1.7^{\circ}$  to  $3.0^{\circ}$  C with a range of  $4.7^{\circ}$  C. Antarctic marine organisms are recognized as the most stenothermal on Earth (i.e., organisms only capable of surviving over a narrow range of temperature). For example, some species such as the brittlestar *Ophionotus victoriae* cannot tolerate acclimation temperatures of more than  $2 - 3^{\circ}$  C (Peck et al., 2009). Based on an experimental approach using  $3.5^{\circ}$  C seawater temperatures (representative of current summer transient peaks of sea surface recorded at Port Arthur, Anvers Island), Schram et al. (2015) suggested that an increase in the frequency and peaks of warming events may affect feeding preferences of amphipod species. Considering the observed within-year trends in water temperature recorded at South Bay, it is important to assess and monitor the potential effects on physiological and ecological processes due to rising water temperatures. The effects of increased seawater in sessile organisms, such as sponges, which form diverse assemblages in the shallows (Cárdenas et al., 2016), have not been well studied in Antarctica. Evidence from other latitudes suggests some sponge species might be highly vulnerable to small increases in seawater temperature (Lopez-Legentil et al., 2011), however this remains to be tested in Antarctic species.

Previous work has shown that even small variations in environmental condition can have significant effects on the physiology of some organisms (Morley et al. 2010). This is highly relevant considering thermal limits cannot only vary spatially but also between years and even seasons (Morley et al., 2012, 2016). Laboratory studies have used rapid warming rates, which are 100 -10,000 times faster than those observed here and predicted for seawater temperatures (Peck et al., 2009). Our HOBO data loggers recorded a significant change in the warming rates before

and after October 2016 (3.75 times higher after October 2016, Table 1, Fig. 4). Warming rates are positively correlated with Critical Thermal Maximum (CTMax), which is used to describe the effects of heat tolerance on different organisms (Bilyk & DeVries, 2011; Cascella et al., 2015; Mora & Maya, 2006; Peck et al., 2009; Peck et al., 2014; Terblanche et al., 2007). Lower CTMax results from slower warming rates and consequently poorer thermal tolerance. Our results suggest that caution must be applied when interpreting CTMax derived from laboratory experiments because some of those values were obtained by using warming rates that are far faster than that experienced by organisms in their natural habitats (in this study between 0.03 and 0.15° C per day). In addition, differences in warming rates before and after October 2016 (0.05 [SD: 0.01] vs. 0.13 [SD: 0.02] ° C day) can have a significant impact on the mechanisms dictating the survival limits of benthic organisms (see Peck et al., 2009).

Recent studies have shown the importance of including natural local variability (with several treatment levels) in experimental designs to avoid exposing organisms to unrealistic scenarios (Vargas et al., 2017). Local conditions should be taken into account since organisms occurring in areas where high variability occurs, might show different responses to stressors compared to those from areas where environmental variability is lower. For instance, it is possible that organisms inhabiting zones that have more stable temperatures (e.g., East Antarctica) might show different responses to warming than those inhabiting areas exposed to higher temperature variability such as the WAP. The use of continuous measurements of temperature and other environmental data constitute a valuable tool to identify long- and short-term variability in coastal waters. This type of information can provide more realistic scenarios based on local conditions in future studies assessing the effect of environmental variation on Antarctic organisms.

## Conclusions

The use of continuous measurements of environmental data such as seawater temperature, can provide key information to identify current patterns of long- and short-term variability in Antarctic coastal waters. Such information is important in informing realistic scenarios for experimental work assessing the local effect of environmental variation on Antarctic organisms. Our results show that benthic organisms at South Bay were exposed to high local variability in seawater temperature (range:  $-1.7^{\circ}$  to  $3.0^{\circ}$ ), rates of warming ( $0.04$  to  $0.15^{\circ}\text{C day}^{-1}$ ), and duration of elevated temperatures (more than 25 days over  $2^{\circ}\text{C}$ ). These results demonstrate that seawater temperatures at depths of 10 to 20 m during summer in South Bay already reach those predicted temperatures for 2100. Although our data was collected in a relatively short period (1 year), due to scarcity of sea temperature data at WAP environments, especially in coastal areas, it represents an important source of baseline information, that along with future high-resolution temporal observations on local temperature, will help us to better understand and forecast climate impacts on Antarctic marine organisms and ecosystems.

## Acknowledgements

The authors would like to thank F. Beaujot, L. Novoa, D. Bravo, K. Attard, L. Rovelli, E. DaForno, F. Rattia and the INACH personnel at Yelcho Antarctic Research Station for help during field activities. We also thank Shane W. Geange for his assistance leading to an improved version of the manuscript. This paper contributes to the SCAR AnT-ERA programme.

## References

- Abram, NJ, Mulvaney, R, Wolff, EW, Triest, J, Kipfstuhl, S, Trusel, LD, Vimeux F, Fleet L, Arrowsmith, C. 2013. Acceleration of snow melt in an Antarctic Peninsula ice core during the twentieth century. *Nature Geoscience*, 6: 404-411.

- Ashton GV, Morley SA, Barnes DKA, Clark MS, Peck LS. 2017. Warming by 1°C drives species and assemblage level responses in Antarctica's marine shallows. *Current Biology* 27:1-8.
- Bailey DM, Johnston IA, Peck LS. 2005. Invertebrate muscle performance at high latitude: swimming activity in the Antarctic scallop, *Adamussium colbecki*. *Polar Biology*, 28: 464-469.
- Barnes DKA, Fuentes V, Clarke A, Schloss IR, Wallace MI. 2006. Spatial temporal variation in shallow seawater temperatures around Antarctica. *Deep Sea Research II* 53:853-865.
- Bravo D. 2017. Análisis de patrones de biodiversidad, distribución espacial y abundancia del ensamble de esponjas en Isla Doumer (Archipiélago de Palmer: Península Antártica) Licenciado en Biología Marina. Universidad de Valparaíso, Chile.
- Bilyk KT, DeVries AL. 2011. Heat tolerance and its plasticity in Antarctic fishes. *Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology* 158.
- Cárdenas CA, Newcombe EM, Hajdu E, González-Aravena M, Geange SW, Bell JJ. 2016. Sponge richness on algae-dominated rocky reefs in the Western Antarctic Peninsula and the Magellan Strait. *Polar Research* 35:30532.
- Cascella K, Jollivet D, Papot C, Léger N, Corre E, Ravaux J, Clark MS, Toullec J-Y. 2015. Diversification, evolution and sub-functionalization of 70 kDa heat-shock proteins in two sister species of Antarctic Krill: differences in thermal habitats, responses and implications under climate change. *PloS One* 10:e0121642.
- Clark MS, Peck LS. 2009. HSP70 heat shock proteins and environmental stress in Antarctic marine organisms: A mini-review. *Marine Genomics* 2:11-18.
- Clark MS, Sommer U, Sihra JK, Thorne MAS, Morley SA, King M, Viant MR, Peck LS. 2017. Biodiversity in marine invertebrate responses to acute warming revealed by a comparative multi-omics approach. *Global Change Biology* 23: 318-330.
- Clarke A. 2001. Benthic organisms and environmental variability in Antarctica: responses to seasonal, decadal and long-term change. *Ocean and Polar Research* 23:433-440.
- Clarke A, Murphy EJ, Meredith MP, King JC, Peck LS, Barnes DKA, Smith RC. 2007. Climate change and the marine ecosystem of the western Antarctic Peninsula. *Philosophical Transactions of the Royal Society of London B Biological Sciences* 362:149-166. 10.1098/rstb.2006.1958
- Convey P, Smith RIL. 2006. Responses of Antarctic ecosystems to climate change. *Plant Ecology* 182:1-10
- Cook AJ, Holland PR, Meredith MP, Murray T, Luckman A, Vaughan DG. 2016. Ocean forcing of glacier retreat in the western Antarctic Peninsula. *Science* 353:283-286.
- Curtis CJ, Simpson GL. 2014. Trends in bulk deposition of acidity in the UK, 1988-2007, assessed using additive models. *Ecological Indicators* 37:274-286.
- Ericson JA, Ho MA, Miskelly A, King CK, Virtue P, Tilbrook B, Byrne M. 2012. Combined effects of two ocean change stressors, warming and acidification, on fertilization and early development of the Antarctic echinoid *Sterechinus neumayeri*. *Polar Biology* 35:1027-1034.
- Ferguson CA, Carvalho L, Scott EM, Bowman AW, Kirika A. 2008. Assessing ecological responses to environmental change using statistical models. *Journal of Applied Ecology* 45:193-203.

- 318 Fordham DA, Saltr   F, Brown SC, Mellin C, Wigley TML. 2017. Why decadal to century  
319 timescale paleoclimate data are needed to explain present-day patterns of biological  
320 diversity and change. *Global Change Biology* (Online version):1-11.
- 321 Foo SA, Sparks KM, Uthicke S, Karelitz K, Barker B, Byrne M, Lamare M. 2016. Contributions  
322 of genetic and environmental variance in early development of the Antarctic sea urchin  
323 *Sterechinus neumayeri* in response to increased ocean temperature and acidification.  
324 *Marine Biology* 163:130.
- 325 Gonz  lez K, Gait  n-Espitia J, Font A, C  rdenas CA, Gonz  lez-Aravena M. 2016. Expression  
326 pattern of heat shock proteins during acute thermal stress in the Antarctic sea urchin,  
327 *Sterechinus neumayeri*. *Revista Chilena de Historia Natural* 89:2. 10.1186/s40693-016-  
328 0052-z
- 329 Gutt J, Isla E, Bertler AN, Bodeker GE, Bracegirdle TJ, Cavanagh RD, Comiso JC, Convey P,  
330 Cummings V, De Conto R, De Master D, di Prisco G, d'Ovidio F, Griffiths HJ, Khan AL,  
331 L  pez-Mart  nez J, Murray AE, Nielsen UN, Ott S, Post A, Ropert-Coudert Y, Sauc  de T,  
332 Scherer R, Schiaparelli S, Schloss IR, Smith CR, Stefels J, Stevens C, Strugnelli JM,  
333 Trimborn S, Verde C, Verleyen E, Wall DH, Wilson NG, Xavier JC. 2017. Cross-  
334 disciplinarity in the advance of Antarctic ecosystem research. *Marine Genomics* (In  
335 press).
- 336 Gutt J, Adams B, Bracegirdle T, Cowan D, Cummings V, di Prisco G, Gradinger R, Isla E,  
337 McIntyre T, Murphy E, Peck L, Schloss I, Smith C, Suckling C, Takahashi A, Verde C,  
338 Wall D, Xavier J. 2012. Antarctic Thresholds - Ecosystem Resilience and Adaptation: a  
339 new SCAR-Biology Programme. *Polarforschung* 82:147-150.
- 340 Hastie TJ, Tibshirani RJ. 1990. *Generalized additive models*. New York: Chapman & Hall.
- 341 Hill PW, Farrar J, Roberts P, Farrell M, Grant H, Newsham KK, Hopkins DW, Bardgett R, Jones  
342 DL. 2011. Vascular plant success in a warming Antarctic may be due to efficient nitrogen  
343 acquisition. *Nature Climate Change* 1:50-53.
- 344 Ingels J, Vanreusel A, Brandt A, Catarino AI, David B, De Ridder C, Dubois P, Gooday AJ,  
345 Martin P, Pasotti F, Robert H. 2012. Possible effects of global environmental changes on  
346 Antarctic benthos: a synthesis across five major taxa. *Ecology and Evolution* 2:453-485.
- 347 Jan PL, Farcy O, Boireau J, Le Texier E, Baudoin A, Le Gouar P, Puechnaille SJ, Petit EJ. 2017.  
348 Which temporal resolution to consider when investigating the impact of climatic data on  
349 population dynamics? The case of the lesser horseshoe bat (*Rhinolophus hipposideros*).  
350 *Oecologia* 184: 749-761.
- 351 Kapsenberg L, Hofmann GE. 2014. Signals of resilience to ocean change: high thermal tolerance  
352 of early stage Antarctic sea urchins (*Sterechinus neumayeri*) reared under present-day  
353 and future pCO<sub>2</sub> and temperature. *Polar Biology*:967-980.
- 354 Kearney MR, Matzelle A, Helmuth B. 2012. Biomechanics meets the ecological niche: the  
355 importance of temporal data resolution. *Journal of Experimental Biology* 215: 922-933.
- 356 Kennicutt MC, Kim YD, Finnemore-Rogan M, Anandakrishnan S, Chown SL, Colwell S,  
357 Cowan D, Escutia C, Frenot Y, Hall J, Liggett D, McDonald A, Nixdorf U, Siegert MJ,  
358 Storey J, W  hlin A, Weathewax A, Wilson GA, Wilson T, Wooding R, Ackley S,  
359 Biebow N, Blankenship D, Bo S, Baesemann J, C  rdenas CA, Cassano J, Danhong C,  
360 Da  nobeitia JJ, Francis J, Guldahl J, Hashida G, Jimenz Corbalan L, Klepikov A, Lee J,  
361 Leppe M, Lijun F, L  pez-Mart  nez J, Memolli M, Motoyoshi Y, Mousalle Bueno R,  
362 Negrete J, Ojeda C  rdenas MA, Proa  o Silva M, Ramos-Garcia S, Sala H, Sheppard P,  
363 Shin H, Shijie X, Shiraishi K, Stockings T, Trotter S, Vaughan DG, Viera Da Unha De

- Menezes J, Vlasich V, Weijia Q, Winthers J-G, Miller H, Rintoul S, Yang H. 2016. Delivering 21st century Antarctic and Southern Ocean science. *Antarctic Science* 28: 407-423.
- Lautenbach S, Seppelt R, Liebscher J, Dormann CF. 2012. Spatial and temporal trends of global pollination benefit. *PloS One* 7:e35954.
- Lopez-Legentil S, Song B, McMurray SE, Pawlik JR. 2011. Bleaching and stress in coral reef ecosystems: hsp70 expression by the giant barrel sponge *Xestospongia muta*. *Molecular Ecology* 17:1840-1849.
- Meredith MP, King JC. 2005. Rapid climate change in the ocean west of the Antarctic Peninsula during the second half of the 20th century. *Geophysical Research Letters* 32:L19604–L19609.
- Meredith MP, Stefels J, van Leeuwe M. 2017. Marine studies at the western Antarctic Peninsula: Priorities, progress and prognosis. *Deep Sea Research Part II: Tropical Studies in Oceanography* 139:1-8.
- Monteith DT, Evans CD, Henrys PA, Simpson GL, Malcolm IA. 2014. Trends in the hydrochemistry of acid-sensitive surface waters in the UK 1988-2008. *Ecological Indicators* 37:287–303.
- Mora C, Maya MF. 2006. Effect of the rate of temperature increase of the dynamic method on the heat tolerance of fishes. *Journal of Thermal Biology* 31:337-341.
- Morley SA, Griffiths HJ, Barnes DKA, Peck LS. 2010. South Georgia: a key location for linking physiological capacity to distributional changes in response to climate change. *Antarctic Science* 22:774–781.
- Morley SA, Martin SM, Bates AE, Clark MS, Ericson J, Lamare M, Peck LS. 2012. Spatial and temporal variation in the heat tolerance limits of two abundant Southern Ocean invertebrates. *Marine Ecology Progress Series* 450:81-92.
- Morley SA, Bates AE, Lamare M, Richard J, Nguyen KD, Brown J, Peck LS. 2016. Rates of warming and the global sensitivity of shallow water marine invertebrates to elevated temperature. *Journal of the Marine Biological Association of the United Kingdom* 96:159-165.
- Orr HG, Simpson GL, des Clers S, Watts G, Hughes M, Hannaford J, M.J. D, Laizé CLR, Wilby RL, Battarbee RW, Evans R. 2015. Detecting changing river temperatures in England and Wales. *Hydrological Processes* 29:752-766.
- Paolo FS, Fricker HA, Padman L. 2015. Volume loss from Antarctic ice shelves is accelerating. *Science*, 348(6232), 327-331.
- Peck L, Morley S, Clark M. 2010. Poor acclimation capacities in Antarctic marine ectotherms. *Marine Biology* 157:2051-2059.
- Peck LS, Webb K, Bailey, D. 2004. Extreme sensitivity of biological function to temperature in Antarctic marine species. *Functional Ecology* 18: 625-630.
- Peck LS. 2005. Prospects for survival in the Southern Ocean: vulnerability of benthic species to temperature change. *Antarctic Science* 17:497-507.
- Peck LS. 2011. Organisms and responses to environmental change. *Marine Genomics* 4:237–243.
- Peck LS. 2016. A cold limit to adaptation in the sea. *Trends in Ecology and Evolution* 31:13-26.
- Peck LS, Clark MS, Morley SA, Massey A, Rossett H. 2009. Animal temperature limits and ecological relevance: effects of size, activity and rates of change. *Functional Ecology* 23:248-256.




- Peck LS, Morley SA, Richard J, Clark MS. 2014. Acclimation and thermal tolerance in Antarctic marine ectotherms. *The Journal of Experimental Biology* 217:16-22.
- Pörtner HO, Peck L, Somero G. 2007. Thermal limits and adaptation in marine Antarctic ectotherms: an integrative view. *Philosophical Transactions of the Royal Society of London B Biological Sciences* 362:2233–2258.
- Schram JB, McClintock JB, Amsler CD, Baker BJ. 2015. Impacts of acute elevator seawater temperature on the feeding preferences of an Antarctic amphipod toward chemically deterred macroalgae. *Marine Biology* 162:425-433.
- Smith RL. 1994. Vascular plants as bioindicators of regional warming in Antarctica. *Oecologia*, 99(3): 322-328.
- Stenni B, Curran MAJ, Abram NJ, Orsi A, Goursaud S, Masson-Delmotte V, Neukom R, Goosse H, Divine D, van Ommen T, Steig EJ, Dixon DA, Thomas ER, Bertler NAN, Isaksson E, Ekaykin A, Werner M, Frezzotti M. 2017. Antarctic climate variability on regional and continental scales over the last 2000 years. *Climate of the Past* 13:1609-1634.
- Suckling CC, Clark MS, Richard JS, Morley SA, Thorne MS, Harper EM, Peck LS. 2015. Adult acclimation to combined temperature and pH stressors significantly enhances reproductive outcomes compared to short-term exposures. *Journal of Animal Ecology* 84:773-784.
- Terblanche JS, Deere JA, Clusella-Trullas S, Janion C, Chown SL. 2007. Critical thermal limits depend on methodological context. *Proceedings of the Royal Society B: Biological Sciences* 1628:2935–2943.
- Thomas ER, Hosking JS, Tuckwell RR, Warren RA, Ludlow EC. 2015. Twentieth century increase in snowfall in coastal West Antarctica. *Geophysical Research Letters* 42:9387–9393.
- Thomas ER, van Wessem JM, Roberts J, Isaksson E, Schlosser E, Fudge TJ, Vallelonga P, Medley B, Lenaerts J, Bertler N, van den Broeke MR, Dixon DA, Frezzotti M, Stenni B, Curran M, Ekaykin AA. 2017. Regional Antarctic snow accumulation over the past 1000 years. *Climate of the Past* 13: 1491-1513.
- Turner, J, Bindshadler, R, Convey, P, Di Prisco, G, Fahrbach, E, Gutt, J, Hodgson DA, Mayewski PA, Summerhayes, C. 2009. *Antarctic climate change and the environment*. Cambridge, Scientific Committee on Antarctic Research.
- Vargas CA, Lagos NA, Lardies MA, Duarte C, Manriquez PH, Aguilera VM, Broitman B, Widdicombe S, Dupont S. 2017. Species specific responses to ocean acidification should account for local adaptation and adaptative plasticity. *Nature Ecology & Evolution* 1:0084.
- Waller CL, Overall A, Fitzcharles EM, Griffiths H. 2017. First report of *Laternula elliptica* in the Antarctic intertidal zone. *Polar Biology* 40:227–230.
- Wood SN. 2004. Stable and efficient multiple smoothing parameter estimation for generalized additive models. *Journal of the American Statistical Association* 99:673–686.
- Wood SN. 2006. *Generalized Additive Models: an introduction with R*: Chapman & Hall/CRC.
- Zamorano JH. 1983. Zonación y biomasa de la macrofauna bentónica en Bahía South, Archipiélago de Palmer, Antártica. *INACH Serie Científica* 30:27-38.
- Zuur AF, Ieno EN, Smith GM. 2007. *Analysing Ecological Data*, 1st ed. Springer Science+Bussines Media, LLC, New York, USA.

# Figure 1

## Study site


Locations of HOBO temperature/light data loggers deployments for studying sponge assemblages at Doumer Island (Palmer, Archipelago), WAP.



# Figure 2

Benthic communities at Doumer Island (Palmer Archipelago), WAP


A) algae-dominated zones (10 m depth), where B) diverse sponge assemblages have been identified underneath the algal canopy. The area showed in B is approximately 80 cm x 50 cm.



# Figure 3

## Daily mean seawater temperature

Within-year GAM (solid line) of daily mean seawater temperature ( $^{\circ}\text{C}$ ) in (A) PY1-10m (edf = 6.7, p-value < 0.001, 404 days), (B) PY4-10 m (edf = 5.7, P<0.001, 312 days) and (C) PY2-20 m (edf= 7.1, P<0.001, 374 days), respectively, at South Bay, Doumer Island, WAP. Black solid curves are the fitted seasonal model from the GAM. Colored areas correspond to 95% confidence bands and the colored thin lines are the observational data (temperature daily means).






# Figure 4

## Shallow sea water temperatures recorded at Doumer Island


Full records of shallow sea water temperatures recorded every 2 hours at PY2- 20 m (red dots) and PY4-10 m (green dots), and every 1 hour at PY1-10 m (violet dots). Grey diamonds represent sea surface temperature average for the day, taken from Port Arthur (Palmer Station) in Anvers Island (<http://pal.lternet.edu>).



# Figure 5

First derivative of seawater temperature trend from PY1 (10 m)

Fitted GAM trend ( $k = 58$ ) showing estimate time intervals with significant rates of change in seawater temperature. Periods of significant change in the fitted trend, determined by derivative analysis, are highlighted by the thickened colored sections of the trend (red=decrease; blue=increase). Grey areas around the black line (fitted trend) indicate 95% confidence limits, and the dots are the daily mean seawater temperature recorded at PY1 from January 11th, 2016 to February 17th, 2017 in South Bay, Doumer Island, WAP.



**Table 1**(on next page)

Periods of significant temperature increase

Calculated warming rates ( $^{\circ}\text{C day}^{-1}$ ) of periods of significant temperature increase from PY1 fitted GAM ( $k = 58$ ,  $\text{edf} = 55.7$ ,  $p\text{-value} < 0.001$ , Fig. 2).



Periods of significant temperature increase	Rates ( $^{\circ}\text{C day}^{-1}$ )
15 - 19 Jan 2016	0.04
7 - 13 Feb 2016	0.04
7 - 13 May 2016	0.05
8 - 12 Jul 2016	0.05
11 - 19 Aug 2016	0.07
18 - 19 Sep 2016	0.03
25 - 30 Oct 2016	0.06
10 - 17 Nov 2016	0.12
26 Nov – 2 Dec 2016	0.11
12 -21 Dec 2016	0.15
25 Dec 2016 – 3 Jan 2017	0.13
14 - 31 Jan 2017	0.12