

How male sound pressure level influences phonotaxis in virgin female Jamaican field crickets (*Gryllus assimilis*)

Understanding female mate preference is important for determining the strength and direction of sexual trait evolution. The sound pressure level (SPL) acoustic signalers use is often an important predictor of mating success because higher sound pressure levels are detectable at greater distances. If females are more attracted to signals produced at higher sound pressure levels, then the potential fitness impacts of signalling at higher sound pressure levels should be elevated beyond what would be expected from detection distance alone. Here we manipulated the sound pressure level of cricket mate attraction signals to determine how female phonotaxis was influenced. We examined female phonotaxis using two common experimental methods: spherical treadmills and open arenas. Both methods showed similar results, with females exhibiting greatest phonotaxis towards loud sound pressure levels relative to the standard signal (69 vs. 60 dB SPL) but showing reduced phonotaxis towards very loud sound pressure level signals relative to the standard (77 vs. 60 dB SPL). Reduced female phonotaxis towards supernormal stimuli may signify an acoustic startle response, an absence of other required sensory cues, or perceived increases in predation risk.

3

4

5

6

7

8

9

10

11 Abstract

12 Understanding female mate preference is important for determining the strength and
13 direction of sexual trait evolution. The sound pressure level (SPL) acoustic signalers use is often
14 an important predictor of mating success because higher sound pressure levels are detectable at
15 greater distances. If females are more attracted to signals produced at higher sound pressure
16 levels, then the potential fitness impacts of signalling at higher sound pressure levels should be
17 elevated beyond what would be expected from detection distance alone. Here we manipulated the
18 sound pressure level of cricket mate attraction signals to determine how female phonotaxis was
19 influenced. We examined female phonotaxis using two common experimental methods: spherical
20 treadmills and open arenas. Both methods showed similar results, with females exhibiting
21 greatest phonotaxis towards loud sound pressure levels relative to the standard signal (69 vs. 60
22 dB SPL) but showing reduced phonotaxis towards very loud sound pressure level signals relative
23 to the standard (77 vs. 60 dB SPL). Reduced female phonotaxis towards supernormal stimuli may
24 signify an acoustic startle response, an absence of other required sensory cues, or perceived
25 increases in predation risk.

26 Introduction

27 Acoustic communication in many anurans and insects is an essential part of courtship and
 28 reproductive behaviour (Alexander, 1975; Wells, 1977; Gerhardt, 1991; Gerhardt & Huber,
 29 2002). Acoustic sexual signalling serves to attract receptive females from a distance (Walker,
 30 1957; Alexander, 1962; Gerhardt, 1991; Ryan & Keddy-Hector, 1992). Behavioural studies have
 31 identified several specific parameters of male acoustic signals used for species recognition and to
 32 confer attractiveness (Rand & Ryan, 1981; Gerhardt, 1982; Simmons & Zuk, 1992; Wilczynski,
 33 Rand & Ryan, 1995), with a broad assessment of the literature revealing that females generally
 34 prefer more energetic signallers (reviewed by Ryan & Keddy-Hector, 1992).

35 Sound pressure level (SPL) can be an important predictor of mate choice (Stout &
 36 McGhee, 1988; Castellano et al., 2000; Klappert & Reinhold, 2003; Hedwig & Poulet, 2005)
 37 with females tending to prefer more intense signals when given the choice (Whitney & Krebs,
 38 1975; Sullivan, 1983; Wells & Schwartz, 1984; Ryan, 1985; Gerhardt, 1991). For example, Arak
 39 (1988) revealed that female natterjack toads (*Bufo calamita*) can perceive small differences in
 40 male sound pressure levels and typically prefer males that signal at higher sound levels. Female
 41 wax moths (*Achroia grisella*) also prefer males that produce ultrasound signals that contain
 42 higher acoustic energy and power (Jang & Greenfield, 1996).

43 Males that produce signals at a higher sound pressure level should be detectable at greater
 44 distances; males that signal with high sound pressure level should therefore receive a selective
 45 advantage over males that signal with low sound pressure level and subsequently have a shorter
 46 detection distance (Forrest & Green, 1991; Forrest & Raspet, 1994). However, if females also
 47 prefer higher sound pressure level males over lower sound pressure level males then the potential
 48 fitness impacts of higher sound pressure level signalling should be elevated beyond what would
 49 be expected from detection distance alone. It is therefore important to quantify female preference

functions for sound pressure level to gain insights into the evolutionary consequences and patterns of selection on this trait.

Female preference functions for sound pressure level have been understudied in crickets despite extensive studies on female preference for other male signal components. Adult male field crickets produce long distance acoustic mate attraction signals (also known as calls) by rubbing their modified forewings together (Alexander, 1962). When a male closes his wings, the scraper of one wing hits the teeth of the file on the other wing causing the harp to resonate and produce a single pulse of sound (Bennet-Clark, 2003). Males concatenate these pulses into chirps (Alexander, 1962; Bennet-Clark, 2003) and females use these long distance acoustic signals to both orient towards signalling males (phonotaxis) and choose between potential mates (Alexander, 1962). To date, only a handful of the experiments quantifying female cricket preference for sound pressure level have explored a broad range of sound levels (Table 1). Even fewer have explored the extremes of the natural range of sound pressure levels available to females. Further, most studies examining female preference for sound pressure level often simultaneously manipulate other signalling parameters, making it difficult to tease apart the influence that sound pressure level alone has on female preference. For example, Stout and McGhee (1988) examined the relative importance of pulse + interpulse duration (syllable period), chirp rate and signal sound pressure level on female mate preference. Unfortunately, Stout and McGhee (1988) made no reference to the natural range of sound pressure levels found in nature. Instead, they quantified female phonotactic response at a 2 dB, 5 dB and 10 dB SPL increase above their standard male signal of 65 dB SPL. By presenting female European house crickets (*Acheta domesticus*) with a pair of male signals differing in one or more of these parameters, Stout and McGhee (1988) concluded that sound pressure level was more important than chirp rate, but that syllable period was more important than sound pressure level in influencing female mate choice. While there is emerging interest in determining how multiple signal parameters

interact to influence patterns of selection and the evolutionary consequences of signalling (Brooks et al., 2005), it is first worth examining how sound pressure level alone shapes female preference functions.

Here we quantify female Jamaican field crickets (*Gryllus assimilis*) phonotactic response to variation in male long distance acoustic mate attraction signal sound pressure level using two standard research methods, a spherical treadmill and open arena. These two methods fall within two broader categories of testing: open-loop and closed-loop methods, respectively. Open-loop methods (Kramer spherical treadmill also called a trackball or locomotion compensator) tether a female in one place, so that she does not experience changes in sound pressure level as she walks ‘toward’ the mate attraction signal (Kramer, 1976; Weber, Thorson & Huber, 1981; Doherty, 1985; Doherty & Pires, 1987; trackball: Hedwig & Poulet, 2005; Hedwig, 2006). Closed-loop methods (open arena) allow the female to move within the enclosed space, thereby allowing her to experience natural changes in sound pressure level as she approaches the mate attraction signal. Quantifying female preference for the sound pressure level of the acoustic mate attraction signal provides a powerful comparison of whether one technique quantifies female phonotaxis more effectively than the other because sound pressure level is one of the primary components that differ between the two techniques. The focus of our study was therefore two-fold: 1) to examine female phonotaxis toward long distance acoustic mate attraction signals that vary in sound pressure level across the natural range of this species, and 2) to compare female phonotaxis on the spherical treadmill to the open arena.

Methods

Cricket Rearing

Our founding population of *Gryllus assimilis* was originally collected in Bastrop County, Texas, United States (latitude ~ 30° 17’ N, longitude ~ 97° 46’ W, elevation ~145 m) from 15 to 24 September, 2008. We did not require specific permits for collecting invertebrates because

these crickets are neither endangered nor protected. We imported adult crickets and eggs to the greenhouse laboratories at Carleton University, Ottawa, Canada (Canadian Food Inspection Agency permit # 2007-03130). Our greenhouse facilities are Plant Pest Containment Level 1 certified (Canadian Food Inspection Agency permit # P-2012-03836). Our study was conducted in accordance with the guidelines of the Canadian Council on Animal Care.

The crickets were reared in communal plastic bins (L x W x H = 64 cm x 40 cm x 42 cm) with a 14:10 h L:D illumination period (lights on at 0600 h, off at 2000 h) at 28 ± 2 °C. They were fed *ad libitum* food (Harlan Teklad Rodent diet 8604M, Harlan Laboratories, Indianapolis, IN, USA: 24.3% protein, 40.2% carbohydrate, 4.7% lipid, 16.4% fiber, 7.4% ash). Water was provided in plastic containers filled with marbles to provide a surface to perch on to minimize drowning. In late 2012 (12-16 generations after field collection) we haphazardly collected 30 4th instar females from the communal cricket bins (no wing bud development; ovipositor had just become visible). These juveniles were housed together in a separate communal bin (same conditions as described above) and monitored daily for imaginal eclosion. Within 24 hours of imaginal eclosion we transferred the adult females to individual 520mL (D x H = 11 cm x 7 cm) clear plastic circular containers with screened lids (4 cm x 4 cm). Each female was provided with a small piece of cardboard egg container for shelter, *ad libitum* food, and 4-5 water gels every 2-3 days (created by soaking 2-4 mm sized water polymer crystals in distilled water for 4-6 hours; Feeder Source, Cleveland, GA, USA). The light cycles and temperatures were identical to the communal rearing environment.

Standard and Focal Mate Attraction Signals

We created our standard and focal mate attraction signals using Adobe Audition CS5.5 software (Adobe Systems Incorporated, San Jose, California, USA). We fashioned our signals after results published in Whattam and Bertram (2011). Because Whattam and Bertram's (2011) study revealed that long distance acoustic mate attraction signaling was influenced by diet, we

selected our signals' parameters to reflect the mean from a population of males reared on high quality food (recordings made at 26°C). The signaling parameters we used for both standard and focal signals were: carrier frequency = 3719 Hz, pulse duration = 10.14 ms, interpulse duration = 15.21 ms, pulses per chirp = 8, and interchirp duration = 1055 ms (signal available at Figshare doi: 10.6084/m9.figshare.1037378). Whattam and Bertram (2011) revealed that well-fed male crickets signal at 60.60 ± 8.45 dB SPL ($\bar{X} \pm 1$ SD), ranging 34 - 72 dB SPL. We therefore always set our standard signal to broadcast at 60.6 dB SPL (re: 20 μ Pa RMS). We used four focal sound pressure levels which we labelled very quiet, quiet, loud, and very loud, broadcast at 43 dB $\cong \bar{X} - 2$ SD, 52 dB $\cong \bar{X} - 1$ SD, 69 dB $\cong \bar{X} + 1$ SD, and 77 dB $\cong \bar{X} + 2$ SD SPL, respectively. Our focal sound pressure levels thus spanned the natural range and slightly beyond that which females are accustomed to hearing. We measured sound pressure level using an EXTECH Digital Sound Level Meter (Model # 407768; FLIR Systems, Waltham, MA, USA).

Preference Trials

Each female's phonotaxis was tested using two-choice trials where the standard signal was simultaneously presented against one of the four focal signals. Each female was tested across all four focal signals in both the arena and the spherical treadmill, for a total of 8 tests. These comparisons occurred across four consecutive days, 10-13 days post imaginal moult (hereafter referred to as 10-13 days old) because this is the age range when female *G. assimilis* are most phonotactically responsive (Pacheco et al., 2013). On a single test day, an individual female was tested once on the spherical treadmill and once in the arena, each to randomly assigned focal signals. Females were always given at least 1 hour of rest prior to switching methods. The order of the method the female was tested on first was randomized.

In both spherical treadmill and open arena trials females were given 60 seconds to acclimatize in silence (detailed below). Each signal was then broadcast on its own for 30 seconds

(order and speaker side randomized) to ensure the female heard both the focal and standard signals prior to starting the trial. Both signals were then broadcast simultaneously from the speakers with the signal chirps interleaved (alternating), so that a female had the potential to identify which signal was coming from which speaker. Once both focal and standard signals were being broadcast, the trial began and female phonotaxis was recorded (detailed below). Every trial ran a total of 5 minutes. All trials were run in the evening (between 5:00 PM and 8:00 PM) under a 75 wattage red light.

Spherical Treadmill Preference Trials

We ran all spherical treadmill trials in a chamber (L x W x H = 86 x 87 x 57 cm) lined with sound-attenuating foam. The spherical treadmill was located in the middle of the chamber with two speakers on either side of it, each 17 cm from the center of the sphere and directly across from each other. Prior to mounting the female onto the spherical treadmill we ensured the standard and focal signal would broadcast at the correct sound pressure level by pointing the sound level meter probe directly at the active speaker, directly above the spherical treadmill, 17 cm from the active speaker, and setting the volume of the standard or focal signal accordingly.

To mount the female on the spherical treadmill we attached a coil- (micro-compression) spring (diameter: 3mm, length: 8mm; spring constant: 210.15 N/m) to each female's pronotum using low melting point wax. Coil springs were attached on day 9 post final moult (one day prior to the first trial, *sensu* Pacheco et al., 2013). We mounted each focal female on the treadmill by attaching her spring to a magnetic rod above the treadmill. This mounting ensured that while the female cricket was held firmly in place on the top of the polystyrene ball, she could freely turn 360 degrees, as well as walk or run using natural motions (photos in Pacheco et al., 2013). We adjusted the air pressure flowing to the polystyrene ball such that, with the cricket in position, the ball was able to rotate freely in all directions in response to the cricket's walking, running, or

turning movements. We oriented all females in the same neutral position at the start of their trial: tethered on the spherical treadmill facing straight ahead between the two speakers.

Once the female was mounted on the spherical treadmill, but prior to the start of data collection, the female experienced 60 seconds of silence, then 30 seconds of the focal sound and 30 seconds of the standard sound (order and speaker side randomized). The female was free to walk during this acclimatization period. We then broadcast both the focal and standard signals, with the signal chirps interleaved. Data collection on female phonotaxis began as soon as both the focal and the standard signals were being broadcast. Female phonotaxis was recorded relative to the focal speaker. Both standard and focal signals were presented continuously throughout the 5 min trial, and the female's locomotor behavior was recorded from the treadmill at a sample rate of 20 samples per second. Each 5 minute trial yielded a total of 6000 samples of cricket X, Y positions. Temperature was held at 22-23°C in the trial room and was monitored with a Fisher Scientific Traceable Digital Thermometer (Model # 15-077-20; Fisher Scientific, Toronto, Ontario, Canada).

We calculated instantaneous displacement (cm) and velocity (cm/s) vectors from the positional data (X, Y coordinates). Total path length was calculated as the sum of all vector lengths over the 6000 samples. Female preference was quantified using net vector score (after Huber et al., 1984). Net vector score is the female's net movement toward or away from the focal signal during the 5 min trial and takes into account the female's direction (vector angle) and the vector length of every recorded leg movement:

$$Net\ Vector\ Score = \sum_{t=1}^{6000} [\cos(\text{vector angle}(t)) \times \text{vector length}(t)]$$

We defined the angle of the focal speaker as 0°. Females moving directly toward the focal speaker (0°) had positive vector scores [$\cos(0^\circ) = 1$], females moving directly away from the focal speaker (180°) had negative vector scores [$\cos(180^\circ) = -1$], and females moving

perpendicular to the focal speaker (90° or 270°) had vector scores of 0 [$\cos(90^\circ \text{ or } 270^\circ) = 0$]. By multiplying this value by each vector length, and summing over the trial duration, we quantified the female's relative attraction to the focal signal. A large positive net vector score indicated that the female moved quickly toward the focal speaker, a small positive score indicated the female moved slowly toward the focal speaker, a large negative score indicated the female moved quickly away from the focal speaker, and a small negative score indicated the female moved slowly away from the focal speaker (Huber et al., 1984).

Open Arena Preference Trials

We conducted the open arena trials in the same room and temperature conditions as spherical treadmill trials. We ran all trials in a chamber (L x W x H = 111.7 cm x 50 cm x 25 cm) with the walls lined with sound-attenuating foam. Speakers were located at opposite ends of the arena (lengthwise). We ensured the standard and focal signal would broadcast at the correct sound pressure level by pointing the sound level meter probe directly at the active speaker from the center of the arena, 55 cm from the active speaker, and setting the volume of the standard or focal signal accordingly. We demarcated a semicircle "choice zone" (radius = 28 cm) in front of each speaker. We then placed the female in the center of the arena (exactly 55 cm from each speaker) under an opaque plastic container (L x W x H = 10 cm x 8.5 cm x 11 cm). Prior to the start of data collection, the female experienced 60 seconds of silence, then 30 seconds of the focal sound and 30 seconds of the standard sound (order and speaker side randomized). We then broadcast both the focal and standard signals simultaneously, with the signal chirps interleaved. As soon as both signals were being broadcast, we carefully and silently raised the opaque plastic container by pulling on its attached string. Data collection on female phonotaxis began as soon as the opaque container was removed.

Each trial ran for a total of 5 minutes. The following female responses were recorded: the amount of time spent stationary in the middle of the arena without moving after the opaque

plastic container was removed, time spent in the arena outside of the choice zones (no-choice zone) after moving from the acclimatization location, time spent in focal choice zone, time spent in standard choice zone, and number of switches made between focal and standard choice zones. The base of the arena was vacuumed and wiped down after each trial.

Because of the varying amount of time spent in focal and standard choice zone, females were classified as 'preferring' a particular focal signal treatment if, over the course of the trial, she spent relatively more time in the zone with the focal signal than in the zone with the standard signal (i.e. time outside choice zones were not included to determine female choice) *sensu* Bischoff, Gould and Rubenstein (1985) and Dugatkin and Godin (1992). To quantify female preference we calculated the proportion of time spent in the focal choice zone relative to the proportion of time spent in both choice zones combined. A number close to one indicates a strong phonotaxis response from the female with much more time spent in the focal zone than the standard zone; conversely, a number close to zero indicates much more time was spent in the standard zone than in the focal zone.

Statistical Analysis

All data were analyzed using JMP 10.0.0 statistical software (SAS Institute Inc., 100 SAS Campus Drive, Cary, North Carolina, USA). Female phonotaxis data (net vector scores from the spherical treadmill and proportion of time spent in focal zone from the open arena) were analyzed using repeated measures general linear mixed models (GLMM) that included sound pressure level, age, and the interaction between sound pressure level and age as fixed effects; body size (maximum head width) was included as a covariate; individual was treated as a random effect. To compare experimental methods, we converted all phonotaxis data into z scores. Z scores were then analyzed using a repeated measures GLMM. We included experimental method, sound pressure level, age, experimental method x sound pressure level, sound pressure level x age, experimental method x age, and experimental method x sound pressure level x age as fixed

effects; body size was included as a covariate; we treated individual as a random effect. Because none of the interactions were significant in any of the models we re-ran all models without interactions per Engqvist (2005). We only present the results of the partial models (excluding interactions) because partial F-tests revealed that the full models did not explain significantly more variation in female phonotaxis. We used one sample t-tests to determine which sound pressure levels invoked significantly different female phonotaxis levels relative to the standard. We used Tukey's post-hoc HSD tests to compare phonotaxis across different focal sound pressure levels.

Results

Females in the spherical treadmill trials exhibited highly variable phonotaxis in response to the different focal sound pressure levels. Some females walked substantially farther than others, with distance covered ranging from 60 cm to 4900 cm. Females had over a 60-fold difference in their velocity magnitudes, with the slowest moving at only 0.12 cm/s and the fastest moving at 8.22 cm/s. Females also moved in all directions during trials relative to the focal speaker. There was a 4-fold difference in net vector scores, indicating large variation in the strength of female phonotaxis on the spherical treadmill. Sound pressure level significantly influenced net vector scores, with females exhibiting highest phonotaxis towards the loud 69 dB $\cong \bar{X}+1$ SD SPL treatment relative to all other focal sound pressure level treatments (Figure 1; Table 2). Age and size did not significantly influence female phonotaxis. Further, although individuals were included as a random effect in the GLMM, they did not explain any of the variation in phonotaxis. Post hoc one-way comparisons revealed that phonotaxis towards the loud 69 dB $\cong \bar{X}+1$ SD SPL was significantly higher than towards the standard 60.6 dB SPL ($t=3.66$,

p=0.0010). Phonotaxis did not significantly differ relative to the standard sound pressure level in the other three focal treatments (all $p>0.25$).

Females in the open arena trials also exhibited highly variable phonotaxis in response to the different focal sound pressure levels. In 17/120 of the open arena trials (14%) females failed to move out of the no-choice zone; these trials were excluded from analysis. Females usually spent time in both standard and focal choice zones in the remaining open arena trials. Sound pressure level significantly influenced female phonotaxis (relative time spent in focal choice zone). Females were most phonotactic towards the loud 69 dB $\cong \bar{X}+1$ SD SPL, relative to all other focal sound pressure level treatments (Figure 2; Table 2). Age and size did not significantly influence proportion of time spent in focal choice zone. Further, individuals being included in the GLMM as a random effect did not explain any of the variation in phonotaxis. Post hoc one-way comparisons of the open arena data revealed that females spent significantly more time in the loud 69 dB $\cong \bar{X}+1$ SD SPL choice zone than the standard 60.6 dB SPL choice zone ($t=2.47$, $p=0.0197$). Conversely, when the sound pressure level was very loud 77 dB $\cong \bar{X}+2$ SD SPL, females avoided the focal choice zone, instead spending significantly more time in the standard 60.6 dB SPL choice zone ($t=-4.20$, $p=0.0003$). Female time in the focal choice zone did not differ significantly relative to time in the standard sound pressure level choice zone in the quiet and very quiet treatments (all $p>0.70$).

We used z scores to compare phonotaxis across the two methods (spherical treadmill and open arena). Experimental method, age and size did not significantly influence female phonotaxis. Further, individuals being included in the GLMM as a random effect did not explain any of the variation in phonotaxis. Female phonotaxis was only influenced by sound pressure level, with females exhibiting highest phonotaxis towards the loud 69 dB $\cong \bar{X}+1$ SD SPL treatment relative to all other focal sound pressure level treatments (Figure 3; Table 2).

Discussion

The loud focal signal ($69 \text{ dB} \cong \bar{X}+1 \text{ SD SPL}$) invoked greater female phonotaxis than any other sound pressure level investigated (Figures 1-3; Table 2). Overall, our findings suggest that males that signal at high sound pressure levels should experience a fitness advantage over males that signal at lower sound pressure levels, provided the sound pressure level is within the natural range of the species. This caveat is important, given females showed reduced phonotaxis towards the very loud focal signal ($77 \text{ dB} \cong \bar{X}+2 \text{ SD SPL}$), suggesting female preference for elevated sound pressure level is not open ended.

The very loud focal signal ($77 \text{ dB} \cong \bar{X}+2 \text{ SD SPL}$) may represent a novel supernormal stimulus as it falls beyond *G. assimilis*' natural range of 34-71 dB SPL (Whattam & Bertram, 2011). Many of the females exhibited negative phonotaxis when exposed to this supernormal stimulus (Figures 1-3). Weber, Thorson and Huber (1981) showed a similar response in *Gryllus campestris* L, with females avoiding mate attraction signals above 70 dB SPL. This negative response to a supernormal stimuli has been observed previously, such as when female crickets exhibit negative phonotaxis to the number of syllables (pulses) in a chirp (Hedrick & Weber, 1998; Ryan & Keddy-Hector, 1992). Female negative phonotaxis to very loud signals may be an acoustic startle response (Hoy, Nolen & Brodfuehrer, 1989), or the females may have a perceived increases in predation risk (Moiseff, Pollack & Hoy, 1978; Zuk & Kolluru, 1998). It is also possible that absence of other sensory cues (Crapon de Caprona & Ryan, 1990) invoke this negative phonotaxis. In some species, such as *Xiphophorus nigrensis* and *X. pygmaeus* (Poeciliidae) swordtails, mate selection is dependent on the dual effect of both visual and olfactory cues.

Surprisingly, females did not exhibit significant phonotaxis towards the standard signal when the focal signal's sound pressure level was less than that of the standard signal (quiet 52 dB $\cong \bar{X}$ -1 SD SPL or very quiet 43 dB $\cong \bar{X}$ -2 SD SPL vs. 60.6 SPL; Figures 1-3). These findings suggest females may not discriminate between sound levels unless one of the signals presents a higher than average sound pressure level. Regardless, given low sound levels may go undetected in nature due to elevated background noise levels, males should be selected to signal with high sound pressure levels whenever possible. It is unknown whether females can discern low sound pressure levels from the background noise level in the testing chamber (42 \pm 2 dB SPL); future tests should determine whether females exhibit phonotaxis towards very quiet (43 dB $\cong \bar{X}$ -2 SD SPL) signals when the alternative is silence (Brown & Handford 1996).

Our study compared open arena and spherical treadmill methods. Quantifying female preference for signal sound pressure level provides a powerful comparison of whether one experimental method quantifies female phonotaxis more effectively than the other because sound pressure level is one of the primary components that differ between the two methods. We found that the two methods produced virtually identical female phonotaxis results for almost all of the sound pressure levels examined. The only sound pressure level that evoked a slightly different phonotactic response between the two methods was the very loud treatment (77 dB $\cong \bar{X}$ +2 SD SPL). Females in the open arena avoided the very loud focal sound pressure level, spending significantly more time in choice zone of the standard signal. While females in the spherical treadmill also exhibited somewhat reduced phonotaxis towards the very loud focal signal, phonotaxis did not significantly differ from the standard signal. Overall, our findings are consistent with the handful of other studies that compare these experimental methods. Walikonis et al. (1991), Stout, Atkins and Zacharias (1991) and Pires and Hoy (1992) revealed female phonotaxis was the same in the open arena and on the spherical treadmill. Similar to our study,

Stout, Atkins and Zacharias (1991) investigated female *Acheta domesticus* response to differing signal intensities. Conversely, Walikonis et al. (1991) investigated female *A. domesitcus* response to differing syllable periods while Pires and Hoy (1992) investigated female *G. firmus* response to natural calls recorded songs at different temperatures. Given sound pressure level, syllable period, and temperature are important predictors of mating preference, our joint findings suggest that open arena and spherical treadmill methods may be used interchangeably to quantify phonotaxis to acoustic signals.

Caution is warranted, however, when signals are broadcasted at unnaturally high sound pressure levels because of fundamental methodological differences between the open arena and the spherical treadmill. The open arena lets the female to move within the enclosed space, thereby enabling her to experience natural changes in sound pressure level. When individuals in the open arena run away from very loud signals, they experience a ~6 dB drop in amplitude with every doubling of their distance from the very loud signal (assuming a spherical, radiating sound source without obstructions). However, when individuals on the spherical treadmill run away they do not experience a drop in amplitude regardless of the distance they run because they are tethered in place on the spherical treadmill. Future research should explore how important sound pressure level is relative to other signalling parameters.

355 Acknowledgements

356 We thank three anonymous reviewers, along with Jeff Dawson, Jean-Guy J Godin, Sarah J
357 Harrison, Genevieve L Ferguson, and Michelle Loranger for helpful comments on an earlier
358 version of this manuscript. We also thank Ryan Chlebak for designing and digitally rendering the
359 spherical treadmill cowling, Michael Jutting for designing and rendering the electronics of the
360 spherical treadmill, and Andrew Mikhail who completed preliminary tests of the device. Jeff
361 Dawson was intimately involved with all aspects of designing and building the spherical
362 treadmill and its accompanying software; this research could not have been conducted without
363 him. We would like to thank Sarah J Harrison, Genevieve L Ferguson, Ian R Thomson, and
364 Kathryn Dufour for their help with cricket care. We would also like to thank Steven Gibson and
365 the Stengl Lost Pines Biological Station at the University of Texas for hosting our laboratory
366 during the cricket-collecting trip that resulted in the establishment of our *G. assimilis* laboratory
367 population.

368 References

- 369 Alexander RD. 1962. Evolutionary change in cricket acoustical communication. *Evolution*
370 **16**:443-467.
- 371 Alexander RD. 1975. Natural selection and specialized chorusing behavior in acoustical insects.
372 In: Pimentel D, ed. *Insects, Science and Society*. New York: Academic Press
- 373 Arak A. 1988. Female mate selection in the natterjack toad: active choice or passive attraction?
374 *Behavioural Ecology and Sociobiology* **22**:317-327
- 375 Bailey NW. 2008. Love will tear you apart: different components of female choice exert
376 contrasting selection pressures on male field crickets. *Behavioral Ecology* **19**:960-966
- 377 Bennet-Clark HC. 2003. Wing resonances in the Australian field cricket *Teleogryllus*
378 *oceanicus*. *Journal of Experimental Biology* **206**:1479-1496
- 379 Bischoff VRJ, Gould JL, Rubenstein DI. 1985. Tail size and female choice in the guppy (*Poecilia*
380 *reticulata*). *Behavioural Ecology and Sociobiology* **17**:253-255
- 381 Brooks R, Hunt J, Blows MW, Smith MJ, Bussiere LF, Jennions MD. 2005. Experimental
382 evidence for multivariate stabilizing sexual selection. *Evolution* **59**:871-880
- 383 Brown TJ, Handford P. 1996. Acoustic signal amplitude patterns: a computer simulation
384 investigation of the acoustic adaptation hypothesis. *The Cooper Ornithological Society* **98**:608-
385 623
- 386 Castellano S, Ross A, Laoretti F, Doglio S, Giacomini C. 2000. Call intensity and female
387 preferences in the European green toad. *Ethology* **106**:1129-1141
- 388 Cragon de Caprona MD, Ryan MJ. 1990. Conspecific mate recognition in swordtails, *Xiphophorus*
389 *nigrensis* and *X. pygmaeus* (Poeciliidae) – olfactory and visual cues. *Animal Behaviour* **39**:290-
390 296
- 391 Dugatkin LA, Godin J-GJ. 1992. Reversal of female mate choice by copying in the guppy
392 (*Poecilia reticulata*). *Proceedings of Royal Society B* **249**:179-184
- 393 Doherty JA. 1985. Trade-off phenomena in calling song recognition and phonotaxis in the
394 cricket, *Gryllus bimaculatus* (Orthoptera, Gryllidae). *Journal of Comparative Physiology*
395 *A* **156**:787-801
- 396 Doherty JA, Pires A. 1987. A new microcomputer based method for measuring walking
397 phonotaxis in field crickets (Gryllidae). *Journal of Experimental Biology* **130**:425-432
- 398 Engqvist L. 2005. The mistreatment of covariate interaction terms in linear model analyses of
399 behavioural and evolutionary ecology studies. *Animal Behaviour* **70**: 967-971.
- 400 Forrest TG, Green DM. 1991. Sexual selection and female choice in mole crickets (*Scapteriscus*:
401 *Gryllotalpidae*): modelling the effects of intensity and male spacing. *Bioacoustics* **3**:93-109
- 402 Forrest TG, Raspet R. 1994. Models of female choice in acoustic communication. *Behavioural*
403 *Ecology* **5**:293-303
- 404 Gerhardt HC. 1982. Sound pattern recognition in some North American treefrogs (Anura:
405 *Hylidae*): implications for mate choice. *American Zoologist* **22**:581-595
- 406 Gerhardt HC. 1991. Female mate choice in treefrogs: Static and dynamic acoustic criteria.
407 *Animal Behaviour* **42**:615-635

- 408 Gerhardt HC, Huber F. 2002. *Acoustic Communication in Insects and Anurans: Common*
409 *Problems and Diverse Solutions*. Chicago: University of Chicago Press
- 410 Gray DA, Cade WH. 2000. Sexual selection and speciation in field crickets. *Proceedings of the*
411 *National Academy of Science* **97**:14449-14454
- 412 Hedrick AV, Weber, T. 1998. Variance in female responses to the fine structure of male song in
413 the field cricket, *Gryllus integer*. *Behavioural Ecology* **9**:582-591
- 414 Hedrick AV, Hisada M, Mulloney B. 2007. Tama-kugel: hardware and software for measuring
415 direction, distance, and velocity of locomotion by insects. *Journal of Neuroscience Methods*
416 **164**:86-92
- 417 Hedwig B, Poulet JF. 2005. Mechanisms underlying phonotactic steering in the cricket *Gryllus*
418 *bimaculatus* revealed with a fast trackball system. *The Journal of Experimental Biology* **208**:915-
419 927
- 420 Hedwig B. 2006. Pulses, patterns and paths: neurobiology of acoustic behaviour in crickets.
421 *Journal of Comparative Physiology A* **192**:677-689
- 422 Huber F, Kleindienst HU, Weber T, Thorson J. 1984. Auditory behaviour of the cricket: 3.
423 Tracking of male calling song by surgically and developmentally one-eared females, and the
424 curious role of the anterior tympanum. *Journal of Comparative Physiology A* **155**:725-738
- 425 Hoy R, Nolen T, Brodfuehrer P. 1989. The Neuroethology of Acoustic Startle and Escape in
426 Flying Insects. *Journal of Experimental Biology* **146**:287-306
- 427 Jang Y, Greenfield MD. 1996. Ultrasonic communication and sexual selection in wax moths:
428 female choice based on energy and asynchrony of male signals. *Animal Behaviour* **51**:1095-1106
- 429 Klappert K, Reinhold K. 2003. Acoustic preference functions and sexual selection on the male
430 calling song in the grasshopper *Chorthippus biguttulus*. *Animal Behaviour* **65**:225-233
- 431 Kramer E. 1976. The orientation of walking honeybees in odour fields with small concentration
432 gradients. *Physiological Entomology* **1**:27-37
- 433 Lickman K, Murray AM, Cade WH. 1998. Effect of mating on female phonotactic response in
434 *Gryllus integer* (Orthoptera: Gryllidae). *Canadian Journal of Zoology* **76**:1263-1268
- 435 Magnus D. 1958. Experimentelle Untersuchungen zur Bionomie und Ethologie des
436 Kaisermantels. *Zeitschrift fur Tierpsychologie* **15**:397-426
- 437 Moiseff A, Pollack GS, Hoy RR. 1978. Steering responses of flying crickets to sound and
438 ultrasound: Mate attraction and predator avoidance. *Proceedings of the National Academy of*
439 *Sciences* **75**:4052-4056
- 440 Olvido AE, Wagner WE. 2004. Signal components, acoustic preference functions, and sexual
441 selection in a cricket. *Biological Journal of the Linnean Society* **83**:461-472
- 442 Pacheco K, Dawson JW, Jutting M, Bertram SM. 2013. How age influences phonotaxis in virgin
443 female Jamaican field crickets (*Gryllus assimilis*) *PeerJ* **1**:130
- 444 Pires A, Hoy RR. 1992. Temperature coupling in cricket acoustic communication. *Journal of*
445 *Comparative Physiology A*. **171**:79-92
- 446 Prosser MR, Murray AM, Cade WH. 1997. The influence of female age on phonotaxis during
447 single and multiple song presentations in the field cricket, *Gryllus integer* (Orthoptera:
448 Gryllidae). *Journal of Insect Behavior* **10**:437-449

- 449 Ritchie MG. 1992. Variation in male song and female preferences within a population of
450 *Ephippiger ephippiger* (Orthoptera: Tettigoniidae). *Animal Behaviour* **43**:845-855
- 451 Ryan MJ. 1985. *The tungara frog, A study in sexual selection and communication*. Chicago:
452 University Chicago Press.
- 453 Rand AS, Ryan MJ. 1981. The adaptive significance of a complex vocal repertoire in a
454 Neotropical frog. *Zeitschrift fur Tierpsychologie* **57**:209-214
- 455 Ryan MJ, Keddy-Hector A. 1992. Directional patterns of female mate choice and the role of
456 sensory biases. *American Naturalist* **139**:S4-S35
- 457 Scheuber H, Jacot A, Brinkhof MWG. 2004. Female preference for multiple condition-dependent
458 components of a sexually selected signal. *Proceeding of the Royal Society B* **271**:2453-2457
- 459 Simmons LW, Zuk M. 1992: Variability in call structure and pairing success of male field
460 crickets, *Gryllus bimaculatus* – the effects of age, size and parasite load. *Animal Behaviour*
461 **44**:1145-1152
- 462 Sullivan BK. 1983. Sexual Sexual selection in Woodhouse's toad (*Bufo woodhousei*). II. Female
463 choice. *Animal Behaviour* **31**:1011-1017
- 464 Stout JF, McGhee R. 1988. Attractiveness of the male *Acheta domestica* calling song to females.
465 II. The relative importance of syllable period, intensity, and chirp rate. *Journal of Comparative*
466 *Physiology A* **164**: 277-287
- 467 Stout J, Atkins G, Zacharias D. 1991. Regulation of cricket phonotaxis through hormonal control
468 of the threshold of an identified auditory neuron. *Journal of Comparative Physiology A* **169**:765-
469 772
- 470 Stout JF, McGhee R. 1988. Attractiveness of the male *Acheta domestica* calling song to females.
471 II. The relative importance of syllable period, intensity, and chirp rate. *Journal of Comparative*
472 *Physiology A* **164**:277-287
- 473 Wagner WE, Murray AM, Cade WH. 1995. Phenotypic variation in the mating preferences of
474 female field crickets, *Gryllus integer*. *Animal Behaviour* **49**:1269-1281
- 475 Walikonis R, Schoun D, Zacharias D, Henley J, Coburn P, Stout J. 1991. Attractiveness of the
476 male *Acheta domesticus* calling song to females III. The relation of age-correlated changes in
477 syllable period recognition and phonotactic threshold to juvenile hormone III biosynthesis.
478 *Journal of Comparative Physiology A* **169**:751-764
- 479 Walker R. 1957. Specificity in the response of female tree crickets (Orthoptera, Gryllidae,
480 Oecanthinae) to calling songs of males. *Annals of the Entomological Society of America* **50**:626-
481 636
- 482 Weber T, Thorson J, Huber F. 1981. Auditory Behaviour of the Cricket I . Dynamics of
483 Compensated Walking and Discrimination Paradigms on the Kramer Treadmill. *Journal of*
484 *Comparative Physiology* **141**:215-232
- 485 Wells KD. 1977. The social behaviour of anuran amphibians. *Animal Behaviour* **25**:666-693
- 486 Wells KD, Schwartz JJ. 1984. Vocal communication in a neotropical treefrog, *Hyla ebraccata*:
487 Advertisement calls. *Animal Behaviour* **32**:405-420
- 488 Whattam EM, Bertram SM. 2011. Effects of juvenile and adult condition on long-distance call
489 components in the Jamaican field cricket, *Gryllus assimilis*. *Animal Behaviour* **81**:135-144

- 490 Whitney CL, Krebs JR. 1975. Mate selection in Pacific tree frogs. *Nature* **255**:325-326
- 491 Wilczynski W, Rand AS, Ryan MJ. 1995. The processing of spectral cues by the call analysis of
- 492 the tungara frog, *Physalaemus pustulosus*. *Animal Behaviour* **49**:911-929
- 493 [Wytttenbach RA](#), [Hoy RR](#). 1993. Demonstration of the precedence effect in an insect. *The*
- 494 *Journal of the Acoustical Society of America* **94**:777-784
- 495 [Wytttenbach RA](#), [Hoy RR](#). 1997. Spatial acuity of ultrasound hearing in flying crickets. *The*
- 496 *Journal of Experimental Biology* **200**:1999-2006
- 497 Zuk M, Kolluru GR. 1998. Exploitation of Sexual Signals by Predators and Parasitoids. *The*
- 498 *Quarterly Review of Biology* **73**:415-438

Figure 1

Female phonotaxis under different sound pressure levels on the spherical treadmill

Phonotaxis was quantified using net vector scores (net movement toward or away from the focal signal; direct movement towards the focal speaker resulted in positive vector scores while direct movement away from the focal speaker resulted in negative vector scores). Letters represent significant differences across sound pressure levels (Tukey's post-hoc HSD analysis). Asterisks represent significant differences between the focal and standard signal (above median = phonotaxis toward focal signal; below median = phonotaxis toward standard signal). Box plots: horizontal line within each box = median phonotaxis measure; top of each box = 3rd (75%) quartile phonotaxis measure; bottom of each box = 1st (25%) quartile phonotaxis measure; whiskers represent the outermost data point that falls within the distances 1st quartile – 1.5*interquartile range or 3rd quartile + 1.5*interquartile range; whiskers are determined by the upper and lower data point values (excluding outliers) when the data points do not reach the above computed ranges; light grey dots represent outliers, when present.




Table 1 (on next page)

Variation in the sound pressure levels used in female field cricket preference studies.

Study	Species	SPL (dB)	Speaker Distance (cm)
Scheuber et al. 2004	<i>Gryllus campestris</i>	80 *	10.0
Lickman et al. 1998	<i>Gryllus texensis</i>	82-85 *	35.4
Bailey 2008	<i>Teleogryllus oceanicus</i>	70	
Prosser et al. 1997	<i>Gryllus texensis</i>	83 *	35.4
Wagner et al. 1995	<i>Gryllus texensis</i>	84.6	35.5
Hedrick et al. 2007	<i>Gryllus integer</i>	72	
Olvido & Wagner 2004	<i>Allonemobius socius</i>	66±2	5.0
Gray & Cade 2000	<i>Gryllus texensis</i>	84	
Gray & Cade 2000	<i>Gryllus rubens</i>	84	

* reference is made to the natural range of sound pressure levels

Table 2_(on next page)

Repeated measures GLMM results for factors affecting female phonotaxis.

The spherical treadmill model used net vector scores to quantify phonotaxis and had an $R^2_{adj} = 0.0495$. The open model used proportion of time spent in focal choice zone relative to time spent in both focal and standard choice zones to quantify phonotaxis and had an $R^2_{adj} = 0.0356$. The method comparison model used z scores to quantify phonotaxis and had an $R^2_{adj} = 0.0626$. Female phonotaxis was significantly influenced by sound pressure level but was not influenced by female age, female body size, or experimental method. Individual was identified as a random effect in all models, but did not explain any of the phonotaxis variation.

Method	Source	DF	F	P
Spherical Treadmill	Age	3,84	0.8947	0.4475
	Sound Pressure Level	3,84	5.7035	0.0013
	Body Size	1,28	0.1983	0.6595
Open Arena	Age	3,79	0.7662	0.5164
	Sound Pressure Level	3,79	5.3900	0.0020
	Body Size	1,29	0.3029	0.5864
Both Methods	Age	3,192	0.7747	0.5094
	Sound Pressure Level	3,192	9.5844	<0.0001
	Technique	1,193	0.0392	0.8433
	Body Size	1,29	0.4584	0.5035

Figure 2

Female phonotaxis under different sound pressure levels in the open arena.

Phonotaxis was quantified using proportion of time the female spent in the focal choice zone relative to the total amount of time spent in both the focal and standard choice zones. Box plot, asterisk, and letter descriptions provided in Figure 1.




Figure 3

Female phonotaxis under different sound pressure levels in both the spherical treadmill (white bars) and open arena (grey bars).

Phonotaxis was compared across the two experimental methods by converting all phonotaxis data into z scores. Box plot, asterisk, and letter descriptions provided in Figure 1.

