

From water striders to water bugs: The molecular diversity of aquatic Heteroptera (Gerromorpha, Nepomorpha) of Germany based on DNA barcodes

Nadine Havemann^{1,2}, Martin M. Gossner³, Lars Hendrich⁴, Jérôme Morinière⁵, Rolf Niedringhaus⁶, Peter Schäfer⁷, Michael J. Raupach^{Corresp. 1, 2}

¹ Fakultät V, Institut für Biologie und Umweltwissenschaften (IBU), Carl von Ossietzky Universität Oldenburg, Oldenburg, Lower Saxony, Germany

² German Centre of Marine Biodiversity, Senckenberg Nature Research Society, Wilhelmshaven, Lower Saxony, Germany

³ Forest Entomology, Swiss Federal Institute for Forest, Snow and Landscape Research, Birmensdorf, Switzerland

⁴ Sektion Insecta varia, Bavarian State Collection of Zoology, Munich, Bavaria, Germany

⁵ Taxonomic coordinator - German Barcode of Life (GBOL), Bavarian State Collection of Zoology, Munich, Bavaria, Germany

⁶ Department of Biology, Earth and Environmental Sciences, Carl von Ossietzky Universität Oldenburg, Oldenburg, Lower Saxony, Germany

⁷ B.U.G.S. (Biologische Umwelt-Gutachten Schäfer), Telgte, North-Rhine Westphalia, Germany

Corresponding Author: Michael J. Raupach

Email address: michael.raupach@rub.de

With about 5,000 species worldwide, the Heteroptera or true bugs are the most diverse taxon among the hemimetabolous insects in aquatic and semi-aquatic ecosystems. Species may be found in almost every freshwater environment and have very specific habitat requirements, making them excellent bioindicator organisms for water quality. However, a correct determination by morphology is challenging in many species groups due to high morphological variability and polymorphisms within, but low variability between species. Furthermore, it is very difficult or even impossible to identify the immature life stages or females of some species, e.g. of the corixid genus *Sigara*. In this study we tested the effectiveness of a DNA barcode library to discriminate species of the Gerromorpha and Nepomorpha of Germany. We analyzed about 700 specimens of 67 species, with 63 species sampled in Germany, covering more than 90% of all recorded species. Our library included various morphological similar taxa, e.g. species within the genera *Sigara* and *Notonecta* as well as water striders of the genus *Gerris*. Fifty-five species (82%) were unambiguously assigned to a single Barcode Index Number (BIN) by their barcode sequences, whereas BIN sharing was observed for 10 species. Furthermore, we found monophyletic lineages for 52 analyzed species. Our data revealed interspecific K2P distances with below 2.2% for 18 species. Intraspecific distances above 2.2% were shown for 11 species. We found evidence for hybridization between various corixid species (*Sigara*, *Callicorixa*), but our molecular data also revealed exceptionally high intraspecific distances as a consequence of distinct mitochondrial lineages for *Cymatia coleoptrata* and for the pygmy backswimmer *Plea minutissima*. Our study clearly demonstrates the

usefulness of DNA barcodes for the identification of the aquatic Heteroptera of Germany and adjacent regions. In this context, our data set represents an essential baseline for a reference library for bioassessment studies of freshwater habitats using modern high-throughput technologies in the near future. The existing data also opens new questions regarding the causes of observed low inter- and high intraspecific genetic variation and furthermore highlight the necessity of taxonomic revisions for various taxa, combining both molecular and morphological data.

**From water striders to water bugs: The molecular diversity of aquatic Heteroptera
(Gerromorpha, Nepomorpha) of Germany based on DNA barcodes**

Nadine Havemann^{1,2}, Martin M. Gossner³, Lars Hendrich⁴, Jérôme Morinière⁵, Rolf
Niedringhaus⁶, Peter Schäfer⁷, Michael J. Raupach^{1,2}

¹ Institute for Biology and Environmental Sciences, Carl von Ossietzky University Oldenburg,
26111 Oldenburg, Germany.

² German Centre of Marine Biodiversity (DZMB), Senckenberg am Meer, 26382
Wilhelmshaven, Germany.

³ Forest Entomology, Swiss Federal Research Institute WSL, Zürcherstrasse 111, 8903
Birmensdorf, Switzerland.

⁴ Sektion Insecta varia, Bavarian State Collection of Zoology (SNSB – ZSM), 81247 München,
Germany.

⁵ Taxonomic coordinator – German Barcode of Life (GBOL), Bavarian State Collection of
Zoology (SNSB – ZSM), 81247 München, Germany.

⁶ Department of Biology, Earth and Environmental Sciences, Carl von Ossietzky University
Oldenburg, PO Box 2503, 26111 Oldenburg, Germany.

⁷ B.U.G.S. (Biologische Umwelt-Gutachten Schäfer), 48291 Telgte, Germany.

Corresponding Author:

Michael J. Raupach^{1,2}

¹ Carl-von-Ossietzky-Straße 9-11, Oldenburg, 26111, Germany

24 ² Südstrand 44, Wilhelmshaven, 26382, Germany

25 Email address: michael.raupach@rub.de

26

27 Abstract

28 With about 5,000 species worldwide, the Heteroptera or true bugs are the most diverse
 29 taxon among the hemimetabolous insects in aquatic and semi-aquatic ecosystems. Species may
 30 be found in almost every freshwater environment and have very specific habitat requirements,
 31 making them excellent bioindicator organisms for water quality. However, a correct
 32 determination by morphology is challenging in many species groups due to high morphological
 33 variability and polymorphisms within, but low variability between species. Furthermore, it is
 34 very difficult or even impossible to identify the immature life stages or females of some species,
 35 e.g. of the corixid genus *Sigara*. In this study we tested the effectiveness of a DNA barcode
 36 library to discriminate species of the Gerromorpha and Nepomorpha of Germany. We analyzed
 37 about 700 specimens of 67 species, with 63 species sampled in Germany, covering more than
 38 90% of all recorded species. Our library included various morphological similar taxa, e.g.
 39 species within the genera *Sigara* and *Notonecta* as well as water striders of the genus *Gerris*.
 40 Fifty-five species (82%) were unambiguously assigned to a single Barcode Index Number (BIN)
 41 by their barcode sequences, whereas BIN sharing was observed for 10 species. Furthermore, we
 42 found monophyletic lineages for 52 analyzed species. Our data revealed interspecific K2P
 43 distances with below 2.2% for 18 species. Intraspecific distances above 2.2% were shown for 11
 44 species. We found evidence for hybridization between various corixid species (*Sigara*,
 45 *Callicorixa*), but our molecular data also revealed exceptionally high intraspecific distances as a
 46 consequence of distinct mitochondrial lineages for *Cymatia coleoptrata* and for the pygmy
 47 backswimmer *Plea minutissima*. Our study clearly demonstrates the usefulness of DNA barcodes

for the identification of the aquatic Heteroptera of Germany and adjacent regions. In this context, our data set represents an essential baseline for a reference library for bioassessment studies of freshwater habitats using modern high-throughput technologies in the near future. The existing data also opens new questions regarding the causes of observed low inter- and high intraspecific genetic variation and furthermore highlight the necessity of taxonomic revisions for various taxa, combining both molecular and morphological data.

Introduction

Aquatic insects are the dominant invertebrate fauna element in most freshwater ecosystems and are enormously variable in morphology, development, physiology, and ecology (Lancaster & Downes, 2013; Dijkstra, Monaghan & Pauls, 2014). Among the hemimetabolous insects, the Heteroptera or true bugs comprise a significant and diverse component of the world's aquatic insect biota (Polhemus & Polhemus, 2007). They are unique as a group because they comprise both aquatic and terrestrial species, whereas other taxa include only species that are aquatic during some life stage, for example mayflies, stoneflies, or dragonflies (Wesenberg-Lund, 1943; Lancaster & Downes, 2013; Gullan & Cranston, 2014). Two infraorders, the Gerromorpha and Nepomorpha, are considered as primarily aquatic (Polhemus & Polhemus, 2007; Lancaster & Downes, 2013; Gullan & Cranston, 2014; Henry, 2017). With more than 4,400 described species worldwide (Henry, 2017), aquatic Heteroptera are well-known for utilizing an exceptionally broad range of habitats, ranging from the marine and intertidal to the arctic and high alpine (Polhemus & Polhemus, 2007). They may be found in almost every freshwater biotope. Approximately 120 species of the Gerromorpha and 230 species of the Nepomorpha are known from the Palearctic region (Polhemus & Polhemus, 2007). For

Germany, 47 species of the Nepomorpha and 22 species belonging to the Gerromorpha have been recorded so far (Wachmann, Melber & Deckert, 2006; Strauss & Niedringhaus, 2014).

Species of the Nepo- and Gerromorpha exhibit numerous morphological and ecological adaptations to their aquatic environment. For instance, nepomorphan true bugs have a streamlined body, natatorial legs and short antennae, whereas gerromorphan species are well-known for their long slender legs which operate as motive (middle leg) and rudder (hind legs), allowing them to operate on the water surface (e.g. Wesenberg-Lund, 1943; Andersen, 1982; Lancaster & Downes, 2013; Gullan & Cranston, 2014) (Fig. 1). Furthermore, a reduction, loss, and/or polymorphism of wings can be observed in many taxa, which is controlled by environmental conditions and genetic factors (e.g. Zera, Innes & Saks, 1983; Muraji, Miura & Nakasuji, 1989; Spence & Andersen, 1994). With the exception of the omnivorous Corixidae, all aquatic true bugs are predators, feeding on any organism that can be subdued by the injection of a venom cocktail consisting of various toxins and proteolytic enzymes (Polhemus & Polhemus, 2007). On the other hand they serve as important prey for numerous fish and other organisms at higher trophic levels (e.g. McCafferty, 1981; Zimmermann & Spence, 1989; Hutchinson, 1993; Klecka 2014; Boda et al., 2015).

Due to their general high abundance in many freshwater systems, their great value as bioindicators of water quality and their unique morphological and ecological specializations for exploiting specialized microhabitats, these groups have been in the focus of entomological and ecological research for a long time (e.g. Hufnagel, Bakonyi & Vásárhelyi, 1999; Polhemus & Polhemus, 2007; Whiteman & Sites, 2008; Skern, Zweimüller & Schiemer, 2010). Nevertheless, as a result of their highly similar morphology, the determination of various species is quite difficult and requires the help of experienced taxonomists. Furthermore, it is very difficult or

even impossible to identify nymphal stages or females of some species, e.g. some species of the genus *Sigara* Fabricius, 1775. In term of males of the Corixidae, typical diagnostic traits include the shape and size of the tarsus of the first leg (pala), the arrangement of pala pegs, and the morphology of the genitalia (e.g. Jansson, 1986; Savage, 1989).

Because aquatic Heteroptera are of high importance for ecological and conservational studies, however, the correct species identification is essential (e.g. Hufnagel, Bakonyi & Vásárhelyi, 1999; Whiteman & Sites, 2008; Skern, Zweimüller & Schiemer, 2010). This is especially true for juveniles and females which can, depending on the life history of a species, dominate within a population over a given period of a year (e.g. Barahona, Millan & Velasco, 2005; Pfenning & Poethke, 2006; Wachmann, Melber & Deckert, 2006).

In the last few years, new molecular and genomic approaches have become more and more popular to overcome possible drawbacks of this traditional and morphology-based way of species assessment. Given the recent technological advancement of DNA-based methods, in particular in the field of modern high-throughput technologies (e.g. Heather & Chain, 2016), it is expected that such techniques will gradually replace traditional field and lab procedures in bioassessment studies over the coming ten to fifteen years (Leese et al., 2016). For example, the EU COST Action CA15219 on “Developing new genetic tools for bioassessment of aquatic ecosystems in Europe” – or DNAqua-Net (<http://dnaqua.net/>) – aims to accelerate the use of DNA-based approaches for the monitoring and assessment of aquatic habitats (Leese et al., 2016). Following these considerations, the analysis of single specimens, bulk samples or environmental DNA will be performed routinely as part of modern species diversity assessment studies (e.g. Yu et al., 2012; Scheffers et al., 2012; Cristescu, 2014; Kress et al., 2015; Creer et al., 2016). However, the effectiveness of all these approaches relies highly on comprehensive

sequence libraries that act as valid references (e.g. Brandon-Mong et al., 2015; Creer et al., 2016). In this context, DNA barcoding represents undoubtedly the most prominent and popular approach using sequence data for valid species identification (e.g. Hajibabaei et al., 2007; Miller et al., 2016). The concept of DNA barcoding relies on the postulate that the interspecific genetic variation is higher than the intraspecific variation of the selected marker (Hebert, Ratnasingham & deWaard, 2003; Hebert et al., 2003). As a consequence, every species is characterized by a unique DNA barcode cluster. For animals, an approximately 650 base-pair fragment of the mitochondrial cytochrome *c* oxidase subunit I (COI) gene was proposed as the global standard for the identification of unknown specimens in terms of a given classification (*sensu* Hebert, Ratnasingham & deWaard, 2003; Hebert et al., 2003). However, it should be noted that various problems may affect the use of mitochondrial DNA, e.g. recent speciation events (e.g. Balvín et al., 2012; Raupach et al., 2014), heteroplasmy (e.g. Boyce, Zwick & Aquadro, 1989; Kavar et al., 2006; Kmiec, Woloszyńska & Janska, 2006), incomplete lineage sorting (e.g. Petit & Excoffier, 2009), (introgressive) hybridization (e.g. Jansson, 1979a; 1979b; Calabrese, 1982; Spence & Wilcox, 1986; Wilcox & Spence, 1986; Savage & Parkin, 1998; Raupach et al., 2014), the presence of alpha-proteobacteria as *Wolbachia* within terrestrial arthropods (Werren, Zhang & Guo, 1995; Xiao et al., 2011; Werren, Baldo & Clark, 2008), and the existence of mitochondrial pseudogenes (e.g. Leite, 2012; Song, Moulton & Whiting, 2014). Nevertheless, a vast number of studies across a broad range of different animals demonstrate the efficiency of DNA barcoding (e.g. Spelda et al., 2011; Hausmann et al., 2013; Hendrich et al., 2015; Lin, Stur & Ekrem, 2015; Raupach et al., 2015; Barco et al., 2016; Coddington et al., 2016; Morinière et al., 2017).

Despite the fact that more than 45,000 species of true bugs have been described worldwide until now (Henry, 2017), the number of studies analyzing the usefulness of DNA

barcodes to discriminate species of this highly diverse insect taxon is still low. Some studies focus on selected species (Rebijith et al., 2012; Zhou et al., 2012; Lis, Lis & Ziaja, 2013), other on specific families (Grebennikov & Heiss, 2014; Kaur & Sharma, 2017), whereas four publications provide a greater representation of various families (Park et al., 2011; Jung, Duwal & Lee, 2011; Raupach et al., 2014; Tembe, Shouche & Ghate, 2014). However, all these studies focused primarily on terrestrial species, analyzing just small number of species belonging to the Gerromorpha and/or Nepomorpha (Park et al., 2011; Jung, Duwal & Lee, 2011; Raupach et al., 2014). To our knowledge, only two publications analyzed aquatic true bugs specifically until now: Castanhole et al. (2013) investigated the variability of 17 barcode sequences of a few species from Brazil, whereas Ebong et al. (2016) successfully tested the usefulness of DNA barcodes to discriminate various species from Cameroon.

The aim of this study was to build-up a baseline for a comprehensive library of DNA barcodes for aquatic Heteroptera (Gerromorpha, Nepomorpha) of Central Europe with a focus on the German fauna and to test the efficiency of DNA barcodes to discriminate the analyzed species. Moreover, our study provides the first thorough molecular study of the aquatic Heteroptera of Germany. In doing so, we analyzed more than 700 DNA barcodes representing more than 60 species. Our library included various morphological similar taxa, e.g. species of the genera *Sigara* Fabricius, 1775 and *Notonecta* Linnaeus, 1758 as well as water striders of the genus *Gerris* Fabricius, 1794 from different localities in Germany. In addition to this we added various specimens from other European countries for comparison, e.g. specimens of the expansive small-bodied backswimmer *Anisops sardeus* Herrich-Schaeffer, 1849 (Berchi, 2011; Klementová & Svitok, 2014).

Material & Methods

Species collection and identification

All analyzed Gerromorpha and Nepomorpha were collected between the years 2003 and 2017. Most of them were adults ($n = 584$; 96.8%). Specimens were stored in ethanol (96%) immediately after collection and identified by some of the authors (NH, MMG, MJR, PS, RN) using various keys (Nieser, 1982; Jansson, 1986; Savage, 1989; Stoffelen et al., 2013; Strauss & Niedringhaus, 2014) based on the most recent taxonomic classification (Aukema & Rieger, 1995). All specimens were carefully checked multiple times by some of the authors in order to prevent a misidentification. For our analysis we also included 109 DNA barcodes of aquatic bugs that were part of a previous barcoding study of true bugs of Central Europe and in which species identification was verified by the authors for comparison (Raupach et al., 2014). Most of the analyzed bug specimens were collected in Germany ($n = 616$; 86.5%), but various individuals were sampled in Austria (37; 5.2%), Greece (20; 2.8%), Spain (16; 2.3%), Switzerland (8; 1.1%), Italy (7; 1.0%), Poland (6; 0.8%), and Portugal (2; 0.3%) for comparison (Fig. 2). In this context we also included specimens from four species that are not recorded for Germany: I. *Anisops sardeus* Herrich-Schaeffer, 1849, ($n = 5$) from Greece, II. *Mesovelia vittigera* Horváth, 1895 ($n = 4$) from Greece, III. *Sigara dorsalis* (Leach, 1817) ($n = 1$) from Switzerland, and IV. *Velia currens* (Fabricius, 1794) ($n = 3$) from Switzerland. The total data set consisted of 712 DNA barcodes with 63 species that are documented for Germany. Furthermore, the number of analyzed specimens per species ranged from one (8 species) to a maximum of 41 for *Notonecta glauca* Linnaeus, 1758.

DNA barcode amplification, sequencing and data depository

The DNA barcode amplification was either performed at the German Centre of Biodiversity Research (Senckenberg am Meer) in Wilhelmshaven, the Carl von Ossietzky University of Oldenburg, or the Bavarian State Collection of Zoology in Munich (SNSB-ZSM). Following the guidelines of DNA barcoding studies (Ratnasingham & Hebert, 2007), all species were documented by photographs before molecular work started. In the majority of the studied animals, all legs of one side of the body were dissected and used for DNA extraction. In case of larger specimens of the genera *Notonecta* Linnaeus, 1758, *Ilyocoris* Stål, 1861, *Ranatra* Fabricius, 1790, *Nepa* Linnaeus, 1758, and *Aphelocheirus* Westwood, 1833, however, only one leg was used. For some very small specimens with a body length <3 mm, e.g. species of the genus *Microvelia* Westwood, 1834, complete specimens were used for DNA extraction. All voucher specimens as well as DNA extracts are stored in a local collection at the Carl von Ossietzky University of Oldenburg.

The DNA extraction was performed using the NucleoSpin Tissue Kit by Macherey and Nagel (Düren, Germany), following the extraction protocol. Polymerase chain reaction (PCR) has been used for amplifying the COI barcode fragment by using the established primer pairs LCO1480/HCO2198 (Folmer et al., 1994), LCO1480/NANCY (Simon et al., 1994), jgLCO1490/jgHCO2198 (Geller et al., 2013), or LepF1/LepR1 (Hebert et al., 2004) for most specimens. For various specimens of the Gerromorpha, however, a new specific forward primer HETF1 (5'-ATG AAT TAT TCG AAT TGA AAT AGG-3') was designed and used in combination with HCO2198 for amplification, resulting in a somewhat smaller fragment with a length of 579 base pairs (bp) of the barcode region. All primers were modified with M13 forward and reverse tails to provide defined base sequences for sequencing (see Ivanova et al., 2007; Khalaji-Pirbalouty & Raupach, 2014).

Barcode amplicons were amplified using illustra™ puReTaq Ready-To-Go PCR Beads (GE Healthcare, Buckinghamshire, UK) in a total volume of 20 µl, containing 17.5 µl sterile molecular grade H₂O, 2 µl DNA template with an DNA amount between 2 to 150 ng/µl and 0.25 µl of each primer (20 pmol/µl). The PCR thermal conditions included an initial denaturation at 94 °C (5 min), followed by 38 cycles at 94 °C (denaturation, 45 s), 48 °C (annealing, 45 s), 72 °C (extension, 80 s), and a final extension step at 72 °C (7 min). All PCR amplification reactions were conducted using an Eppendorf Mastercycler Pro system (Eppendorf, Hamburg, Germany). Negative and positive controls were included with each round of reactions. Two µl of the amplified products were verified for size conformity by electrophoresis in a 1% agarose gel with GelRed or SYBR Green using commercial DNA size standards, whereas the remaining PCR product was purified with the NucleoSpin Gel and PCR Clean-up Kit (Macherey-Nagel, Düren, Germany). Purified PCR products were cycle-sequenced and sequenced in both directions at a contract sequencing facility (GATC, Konstanz, Germany) using the given M13 tail sequences. Double stranded sequences became assembled and checked for mitochondrial pseudogenes (numts) analyzing the presence of stop codons, frameshifts as well as double peaks in chromatograms with the Geneious program package version 7.0.4 (Biomatters, Auckland, New Zealand) (Kearse et al., 2012). Ambiguous parts at the 5'-end or 3'-end of the sequences were removed. For verification, BLAST (nBLAST, search set: others, program selection: megablast) and/or BOLD (identification engine; species level barcode records) searches were performed to confirm the identity of all new sequences as bug sequences based on already published sequences.

Detailed voucher information, taxonomic classifications, photos, DNA barcode sequences, used primer pairs and trace files (including their quality) are publicly accessible

through the public data set “DS-BAHCE Barcoding Aquatic Heteroptera of Central Europe” (Dataset ID: dx.doi.org/10.5883/DS-BAHCE) on the Barcode of Life Data Systems workbench (BOLD; www.boldsystems.org) (Ratnasingham & Hebert, 2007). All new barcode data were also deposited in GenBank (MG665389-MG665993).

DNA barcode analysis

We analyzed intra- and interspecific distances of the studied aquatic Heteroptera using the provided analytical tools of the BOLD workbench (align sequences: BOLD aligner; ambiguous base/gap handling: pairwise deletion) based on the Kimura 2-parameter model of sequence evolution (K2P; Kimura, 1980). Furthermore, all analyzed COI sequences became subject to the Barcode Index Number (BIN) system implemented in BOLD which clusters DNA barcodes in order to generate operational taxonomic units that closely correspond to species (Ratnasingham & Hebert, 2013). We used a recommended threshold of 2.2% for a rough differentiation of intraspecific as well as interspecific K2P distances (Ratnasingham & Hebert, 2013).

A Neighbor Joining cluster analysis (NJ; Saitou & Nei, 1987) was performed for all studied species for a graphical representation of the genetic differences between sequences and clusters of sequences using MEGA v7.0.18 (Kumar, Stecher & Tamura, 2016). The K2P model was chosen as the model for sequence evolution for comparison purposes with previous studies. For validation, non-parametric bootstrap support values were obtained by resampling and analyzing 1,000 replicates (Felsenstein, 1985). All analysis were based on an alignment that was generated using MUSCLE (Edgar, 2004) implemented in MEGA v7.0.18 for all studied barcode sequences. Additionally, statistical maximum parsimony networks were constructed exemplarily

for species with interspecific distances ranging from zero to 1% (see Table 1) by using TCS networks (Clement, Posada & Crandall, 2002) as part of the software package of PopArt v.1.7 (Leigh & Bryant, 2015). Such networks allow the identification of haplotype sharing between species as a consequence of recent speciation and/or on-going hybridization processes (e.g. Raupach et al., 2010; Raupach et al., 2014).

Results

Our analyzed DNA barcode library comprised 63 species that are documented for Germany, representing 91% of the known aquatic bug species diversity of this country (Nepomorpha: $n = 43$ (92%); Gerromorpha: $n = 20$ (91%)), and additional four species that were collected in other countries and not recorded for Germany. In total, we generated 603 new barcodes of 64 species. The complete alignment of all analyzed sequences ($n = 712$) had a length of 658 bp, with fragments lengths ranging from a minimum of 366 bp to the full barcode fragment size of 658 bp. For some studied specimens of *Cymatia coleoptrata* (Fabricius, 1777) ($n = 22$), our analysis revealed two characteristic deletions of 39 (alignment position: 110 – 148) and nine nucleotides (629 – 637) for all studied specimens (Fig. S1). Average base frequencies were A = 32%, C = 17%, G = 16%, and T = 35%. For eight species only one barcode sequence was generated (Table 1). Intraspecific distances ranged from zero to maximum values of 8.3% (*Plea minutissima* Leach, 1817) and 9.44% (*Cymatia coleoptrata*) (Table 1). Maximum intraspecific pairwise distances with values $>2.2\%$ were found for 11 species (Table 1). In terms of interspecific divergence, values ranged from zero to 18.58%, with 18 species pairs having values $<2.2\%$ (Table 1). We found interspecific distances below 1% for 9 species. Unique BINs were recorded for 55 species, whereas two BINs were identified for 10 species (Table 1). For

two species that were represented only by one specimen, namely *Arctocorisa germari* (Fieber, 1848) and *Corixa dentipes* Thompson, 1869, our sequences did not have the required fragment length of at least 400 bp to fulfill the criteria for BIN assignment. As consequence, no BINs were available for these two species.

Our NJ analysis based on K2P distances revealed two large and distinct clusters, separating all analyzed Gerromorpha and all Nepomorpha specimens from each other (Fig. S2). For a better presentation, the topology has been split on this basis and shown in two figures (Gerromorpha: Fig. 3, Nepomorpha: Fig. 4). We found non-overlapping clusters with bootstrap values >90% for 57 species (85%) (Fig. 3; 4). Of the analyzed 59 species with more than one specimen, 52 (88%) were monophyletic, 3 (5%) paraphyletic, and 4 (7%) polyphyletic (Table 1, Fig S2).

The statistical maximum parsimony network analysis of species with interspecific distances below 1% revealed a close relationship between *Gerris asper* (Fieber, 1860) ($n = 1$) and *Gerris lateralis* Schummel, 1832 ($n = 2$) (Fig. 5). We found three haplotypes with a frequency of 1 (singletons) that were separated by only one or two mutational steps, with haplotype h1 (*Gerris asper*) connected with h2 (*Gerris lateralis*), which was in turn connected with haplotype h3 (*Gerris lateralis*). A similar situation was observed for *Sigara limitata* (Fieber, 1848) ($n = 2$) and *Sigara semistriata* (Fieber, 1848) ($n = 5$) (Fig. 5). Three different haplotypes were identified, with h1 representing all studied specimens of *Sigara semistriata*. Both unique haplotypes of *Sigara limitata* (h2, h3) were directly connected to this haplotype by two or three mutational steps. In the case of *Callicorixa praeusta* (Fieber, 1848) ($n = 23$) and *Callicorixa producta* (Reuter, 1880) ($n = 1$) we found five different haplotypes (Fig. 5), with h1 representing the dominant haplotype which includes 19 specimens of *Callicorixa praeusta* and

the only specimen of *Callicorixa producta*. All other four haplotypes (h2-h5) were only scored in one specimen and connected with h1 by one or two mutational steps. A much more complex network was revealed for *Sigara distincta* (Fieber, 1848) ($n = 7$), *Sigara falleni* (Fieber, 1848) ($n = 12$), and *Sigara iactans* Jansson, 1983 ($n = 12$) (Fig. 6). We identified 16 different haplotypes in total, with six haplotypes (h1-h6) shared by more than one specimen. Three of these haplotypes (h2, h3, h5) were shared by specimens of *Sigara falleni* and *Sigara iactans*. Furthermore, haplotypes of both previously mentioned species were randomly distributed within the network. In many cases, haplotypes of *Sigara falleni* were separated merely by two mutational steps from haplotypes of *Sigara iactans* (e.g. h6 and h13) and vice versa. We found four singletons for *Sigara falleni* and five for *Sigara iactans*. In contrast to these two species, we identified only two haplotypes (h1, h8) for the seven analyzed specimens of *Sigara distincta*. Moreover, most specimens ($n = 6$) were identical (h1) and located at the periphery of the network. The other haplotype (h8), a singleton collected among others at Apen (Lower Saxony), was separated by more than 25 mutational steps from the network and represents the most isolated haplotype in this network by far. Therefore, *Sigara distincta* shared no haplotypes with other species.

Discussion

Our comprehensive DNA barcode library represents an important step for the molecular characterization of the freshwater fauna in Central Europe and adjacent regions. As COI sequences are used routinely in phylogeographic, phylogenetic and evolutionary studies as well, our data can be also implemented in projects analyzing the genetic variation of species in relation to historical, geographical and ecological factors (Galactos, Cognato & Sperling, 2002;

Damgaard, 2005; Damgaard, 2008b; Gagnon & Turgeon, 2010; Ye et al., 2016). Unique BINs were found for 55 species, allowing a valid identification of 82% of the analyzed 67 species. Distinct and monophyletic lineages were revealed for 52 species (78%). Our study also indicates the need of further detailed taxonomic revisions, using state-of-the-art methods for a fine-scaled characterization (Raupach et al., 2016). This is especially true for the species-rich family Corixidae. In the following we will discuss noticeable species with high intraspecific and/or low interspecific distances more in detail.

Interspecific K2P distances with values below 2.2%

The efficiency of DNA barcoding highly depends on distinct mitochondrial lineages, ideally coupled with moderate to high genetic interspecific distances. If sister species, however, have low interspecific distances and haplotype sharing as a result of a recent ancestry and/or ongoing gene flow, DNA barcoding will fail (e.g. Tautz et al., 2003; Frezal & Leblois, 2008; Raupach & Radulovici, 2015). For the analyzed species of the Gerromorpha and Nepomorpha, minimum interspecific K2P distances with values below 2.2% were found for 18 species (Table 1). Distance values ranged from 0% (four species: *Callicorixa praeusta* (Fieber, 1848), *Callicorixa producta* (Reuter, 1880), *Sigara falleni* (Fieber, 1848), *Sigara iactans* Jansson, 1983) to 2.11% (*Sigara venusta* (Douglas & Scott, 1869)). Distinct monophyletic clusters were revealed for *Notonecta obliqua* Thunberg, 1787 and *Notonecta glauca* Linnaeus, 1758 (1.08%), *Notonecta lutea* Müller, 1776 and *Notonecta reuteri* Hungerford, 1928 (1.24%), *Sigara dorsalis* (Leach, 1817) and *Sigara striata* (Linnaeus, 1758) (1.71%) (but see Savage and Parkin, 1998), and *Sigara venusta* (Douglas & Scott, 1869) and *Sigara limitata* (Fieber, 1848)/*Sigara semistriata* (Fieber, 1848) (2.11%), indicating a close relationship of these species pairs with

distinct lineages (Table 1). Furthermore, the analyzed specimen of *Arctocorisa germari* (Fieber, 1848) was nested in the paraphyletic cluster of *Arctocorisa carinata* (C. R. Sahlberg, 1819) (1.03%) (Fig. S1). In this context it should be noted that experimental crosses gave viable hybrids between both *Arctocorisa* species with intermediate characters (Jansson, 1979). These examples show that recent speciation events as well as hybridization may represent important processes in these groups. Future studies including more specimens and other genetic markers should be conducted to resolve the eco-evolutionary events leading to the low interspecific variation. Species pairs with interspecific K2P distances <1% will be discussed more in detail below.

Species pairs with interspecific distances below 1%

I. *Gerris asper* (Fieber, 1860) and *Gerris lateralis* Schummel, 1832

From a morphological perspective, both species are very similar (e.g. Wagner & Zimmermann, 1955; Wachmann, Melber & Deckert, 2006). Not surprisingly, *Gerris asper* is suggested as a south-eastern vicariant of its boreo-montane sister species *Gerris lateralis* (Jeziorski et al., 2012). Whereas *Gerris lateralis* has a distribution ranging from Europe to the Far East of Russia, *Gerris asper* is found in Southern and Central Europe, extending to Afghanistan (Jeziorski et al., 2012). In spite of the fact that our sample sizes were very small (*Gerris asper*: $n = 1$, *Gerris lateralis*: $n = 2$), our molecular data set clearly support the proposed close relationship of both water striders species (Fig. 5, Table 1). Future studies including more specimens covering a larger geographic range are needed to test whether both taxa represent distinct lineages or hybridization still takes place as it is known from other species of this genus (e.g. Calabrese, 1982).

370

371 **II. *Sigara limitata* (Fieber, 1848) and *Sigara semistriata* (Fieber, 1848)**

372 Both species belong to the subgenus *Retrocorixa* Walton, 1940 and have a similar
 373 distribution, ranging from Europe eastwards to Siberia (Jansson, 1986; Wachmann, Melber &
 374 Deckert, 2006; Coulianos, Økland & Økland, 2008). A close relationship as it has been indicated
 375 by our data has not been proposed yet. In contrast to our results, morphological characters
 376 suggest *Sigara venusta* (Douglas & Scott, 1869) as sister species of *Sigara semistriata* (see
 377 Jansson, 1986). As part of our study, *Sigara venusta* represents the sister species of *Sigara*
 378 *limitata* and *Sigara semistriata* with a distance of 2.11% (Fig. 5, Table 1). Due to the fact that
 379 neither *Sigara limitata* nor *Sigara semistriata* were monophyletic and the observed interspecific
 380 distances were very low (0.15%) (Table 1), we suggest a recent ancestry of both species. Hybrids
 381 are currently not known. Future studies are needed to verify this hypothesis.

382

383 **III. *Callicorixa praeusta* (Fieber, 1848) and *Callicorixa producta* (Reuter, 1880)**

384 The genus *Callicorixa* White, 1873 includes five medium sized species (6 – 8 mm) that
 385 are recorded for Europe, with two species documented in Central Europe. Specimens of
 386 *Callicorixa praeusta* can be found throughout most Europe except the Mediterranean region
 387 reaching to the Far East of Russia, whereas the distribution of *Callicorixa producta* ranges from
 388 the Northern parts of Central Europe to Fennoscandia, Northern Russia, Kazakhstan, Mongolia,
 389 and Siberia (Jansson, 1986; Wachmann, Melber & Deckert, 2006; Coulianos, Økland & Økland,
 390 2008). Most identification keys for this genus rely largely on the shape and intensity of dark
 391 areas of the hind tarsus 1 (Jansson, 1986; Savage 1989; Strauss & Niedringhaus, 2014). While
 392 this morphological trait is fairly good for the determination of most typical specimens, existing

variation is rather wide, making it unreliable in many cases (Jansson, 1986). Similar to other species, our DNA barcode data give evidence for a recent ancestry or ongoing gene flow between *Callicorixa praeusta* and *Callicorixa producta* (Fig. 5). However, only one (female) specimen of *Callicorixa producta* was available, demonstrating the need for more detailed studies to clarify the underlying processes.

IV. *Sigara distincta* (Fieber, 1848), *Sigara falleni* (Fieber, 1848), and *Sigara iactans* Jansson, 1983

Some decades ago, a comprehensive revision revealed that the well-known species *Sigara falleni* of the subgenus *Subsigara* Stichel, 1935 was actually a mixture of four closely related and highly similar species, including *Sigara iactans* (see Jansson, 1983). Whereas the identification of females is not always reliable, males of both species can be recognized by the shape of their pala: specimens of *Sigara falleni* are characterized by triangular pala, whereas trapezoidal pala are found for *Sigara iactans* (Jansson, 1983; 1986). Intermediate specimens, however, have been also documented and indicate on-going hybridization between both species (Jansson, 1983; 1986). Water bugs of *Sigara distincta* are found from the British Isles through North and Central Europe to Asia as far as East Siberia and Mongolia (Jansson, 1986; Savage, 1989; Coulianos, Økland & Økland, 2008). A similar distribution is known for *Sigara falleni*, ranging throughout most of Europe eastwards to Siberia and China (Jansson, 1986; Savage, 1989; Coulianos, Økland & Økland, 2008). In contrast to both previous species, *Sigara iactans* is found in two disjunct areas, one in North and Central Europe, and the other in Southeastern Europe (Jansson, 1986; Wachmann, Melber & Deckert, 2006). Our DNA barcode data revealed multiple haplotype sharing between *Sigara falleni* and *Sigara iactans*, supporting the close relationship and on-

going hybridization between both species (Fig. 6). Beside this, our results revealed a close relationship of *Sigara (Subsigara) distincta* with *Sigara falleni* and *Sigara iactans*, as it has been discussed in the past also (Jansson, 1986). However, we found no shared haplotypes yet. Additional studies involving more specimens of a larger geographic region are needed to validate the species status within this subgenus.

Intraspecific K2P distances with values >2.2%

Various phenomena can generate distinct lineages within DNA barcode data, e.g. phylogeographic processes (e.g. Andersen et al., 2000; Damgaard 2005; 2008b; Ye et al., 2016), the presence of maternally inherited endosymbionts as *Wolbachia* (e.g. Lis, Maryńska-Nadachowska & Kajtoch, 2015), or the existence of cryptic species (e.g. Paterson et al., 2016; Jiu et al., 2017). In this context we found 11 species with intraspecific K2P distances >2.2%, ranging from 2.32% (*Mesovelgia vittigera* Horváth, 1895) to a maximum of 9.44 (*Cymatia coleoptrata* (Fabricius, 1777)). For most species, excluding *Sigara iactans* (2.67%), *Sigara falleni* (3.37%), and *Sigara distincta* (5.77%) (see discussion above), we are currently unable to clarify the background of the observed high nucleotide distances and distinct lineages based on the given data set. However, exceptionally high intraspecific distances with values >8% were found within the pygmy backswimmer *Plea minutissima* Leach, 1817 (8.3%) and *Cymatia coleoptrata* (Fabricius, 1777) (9.44%) (Table 1). Both will be discussed more in detail.

Small and cryptic: two highly distinct DNA barcode clusters within *Plea minutissima* Leach, 1817

Pygmy backswimmers are small bugs, usually less than 3.5 mm in length and confine themselves to the vegetation in which they hide and where they prey on mosquito larvae and other small arthropods (Schuh & Slater, 1995). For Europe, only one species of the Pleidae is recorded, namely *Plea minutissima*. As part of our study we found two distinct lineages within the sixteen analyzed specimens with high distances ranging from 8.1 to 8.3%. Both lineages were supported by high bootstrap values (99%) (Fig. 7). Most specimens of lineage A ($n = 8$) were found in Brandenburg and Bavaria, but also two specimens were collected in Lower Saxony (Jaderberg). In contrast to this, all specimens of lineage B ($n = 8$) were collected in Lower Saxony (Jaderberg, Lingen, Norderney). Whether this surprisingly high molecular diversity is a result of effects as incomplete lineage sorting (e.g. Damgaard, 2008) or whether we found evidence for the existence of two sibling species (e.g. Damgaard, 2005), is not within the scope of this study but clearly needs further investigation.

A currently unknown species of the genus *Cymatia* Flor, 1860?

For the genus *Cymatia*, three European species are documented so far: *Cymatia coleoptrata* (Fabricius, 1777), *Cymatia bonsdorffii* (C. R. Sahlberg, 1819), and *Cymatia rogenhoferi* (Fieber, 1864). In terms of a morphological identification, all species can be identified according to their size and hemelytral patterns without doubt (Janssen 1986, Stoffelen et al., 2013). Our study revealed two distinct lineages within the analyzed specimens of *Cymatia coleoptrata* (lineage A and B), with a K2P distances ranging from 9.13 – 9.42% and bootstrap support values of 99% (Fig. 8). Whereas lineage A includes 22 specimens from Lower Saxony ($n = 21$, Lingen) and Baden-Württemberg ($n = 1$, Wolperstwende), lineage B contains two specimens that were collected in Brandenburg (Voßberg). Both specimens of lineage B were

small adult males with a body size between 4.3 and 4.5 mm and were identified using morphological traits as *Cymatia coleoptrata* at first sight. Interestingly, their barcode sequences did not have the characteristic nucleotide deletions of this species (Fig. S1). Furthermore, we found no other similar sequences using the BOLD identification engine (Best ID: *Cymatia coleoptrata*) (date of request: 2017-11-20). Unfortunately, both *Cymatia* vouchers were lost, preventing a closer reanalysis of the specimens. Nevertheless, our results should motivate heteropterologists to study more specimens of this genus, in particular from the Eastern parts of Germany, in order to verify the presence of this putative new species.

Conclusion

In our study we lay the foundations for a comprehensive DNA barcode data set for the aquatic Heteroptera in Central Europe and adjacent regions, which will act as useful reference library for freshwater bioassessment studies using modern high-throughput sequencing technologies. Unique BINs were revealed for 55 species, representing 82% of the analyzed 67 species. Furthermore, monophyletic lineages were found for 52 species (78%). Nevertheless, our molecular data highlights discordance between the generally accepted but exclusively morphologically-based taxonomy and observed molecular diversity within some species of the Gerromorpha and Nepomorpha. The analysis of additional specimens from other localities and of other molecular markers, for example microsatellites or SNPs, will give us more insights into the taxonomic status of these species as well as in the eco-evolutionary processes underlying the observed genetic patterns. However, it should be kept in mind that the traditional aims of taxonomy are unchanged and include various aspects, e.g. detailed high-quality descriptions and delimitation of species, a classification that reflects evolution, a dynamic nomenclature, and fast

and reliable identification tools. Therefore, our DNA barcode library may be considered as a promoter for such studies.

Acknowledgements

We would like to thank Jana Deppermann and Babett Günther (both DZMB, Wilhelmshaven) for their laboratory assistance. Three anonymous reviewers provided helpful comments on the manuscript. We also thank Torsten Berger (Potsdam) and Reinhard Müller (Berlin) for providing further specimens. Furthermore we are very grateful to Gerhard Strauss for giving his permission to use the excellent photos of aquatic true bugs from www.corisa.de. We are also grateful to the team of Paul Hebert in Guelph (Ontario, Canada) for their great support and help and in particular to Sujeevan Ratnasingham for developing the BOLD database infrastructure and the BIN management tools.

References

- Andersen NM. 1982. *The Semiaquatic Bugs (Hemiptera, Gerromorpha). Phylogeny, Adaptations, Biogeography and Classification*. Klampenborg: Scandinavian Science Press Ltd.
- Andersen NM, Cheng L, Damgaard J, Sperling FAH. 2000. Mitochondrial DNA sequence variation and phylogeography of oceanic insects (Hemiptera: Gerridae: *Halobates* spp.). *Marine Biology* 136:421-430. DOI: <https://doi.org/10.1007/s002270050701>.

505 Aukema B, Rieger C. 1995. *Catalogue of the Heteroptera of the Palaearctic Region Vol. 1:*
 506 *Enicocephalomorpha, Dipsocoromorpha, Nepomorpha, Gerromorpha and Leptopodomorpha.*
 507 Amsterdam: Netherlands Entomological Society.
 508

509 Balvín O, Munclinger P, Kratochvíl L, Vilímová J. 2012. Mitochondrial DNA and morphology
 510 show independent evolutionary histories of bedbug *Cimex lectularius* (Heteroptera: Cimicidae)
 511 on bats and humans. *Parasitology Research* 111:457-469. DOI: 10.1007/s00436-012-2862-5.
 512

513 Barahona J, Millan A, Velasco. 2005. Population dynamics, growth and production of *Sigara*
 514 *selecta* (Fieber, 1848) (Hemiptera, Corixidae) in a Mediterranean hypersaline stream.
 515 *Freshwater Biology* 50:2101-2113. DOI: 10.1111/j.1365-2427.2005.01463.x.
 516

517 Barco A, Raupach MJ, Laakmann S, Neumann H, Kneblsberger T. 2016. Identification of
 518 North Sea molluscs with DNA barcoding. *Molecular Ecology Resources* 16:288-297. DOI:
 519 10.1111/1755-0998.12440.
 520

521 Berchi GM. 2011. First record of *Anisops sardeus* (Hemiptera: Heteroptera: Notonectidae) in
 522 Romania. *North-Western Journal of Zoology* 7:339-341.
 523

524 Boda P, Bozóki T, Vásárhelyi T, Bakonyi G, Várbiro G. 2015. Revised and annotated checklist
 525 of aquatic and semi-aquatic Heteroptera of Hungary with comments on biodiversity patterns.
 526 *ZooKeys* 501:89-108. DOI: 10.3897/zookeys.501.8964.
 527

528 Boyce TM, Zwick ME, Aquadro CF. 1989. Mitochondrial DNA in the bark weevils size,
529 structure and heteroplasmy. *Genetics* 123:825-836.
530

531 Brandon-Mong GJ, Gan HM, Sing KW, Lee PS, Lim PE, Wilson JJ. 2015. DNA metabarcoding
532 of insects and allies: an evaluation of primers and pipelines. *Bulletin of Entomological Research*
533 105:717-727. DOI: 0.1017/S0007485315000681.
534

535 Calabrese DM. 1982. Hybridization between *Gerris alacris* Hussey and *Gerris comatus* Drake
536 and Hottes (Hemiptera: Heteroptera: Gerridae) in nature. *Proceedings of the Entomological*
537 *Society of Washington* 84:209-210.
538

539 Castanhole MMU, Marchesin SRC, Pereira LLV, Moreira FFF, Barbosa JF, Valério JR, Itoyama
540 MM. 2013. The first assess of the haplotypes from COI gene sequences in species of spittlebugs
541 (Cicadomorpha: Hemiptera) and aquatic true bugs (Gerromorpha and Nepomorpha: Hemiptera)
542 in Brazil. *Genetics and Molecular Research* 12:5372-5381. DOI: 10.4238/2013.November.7.12.
543

544 Clement M, Posada D, Crandall KA. 2000. TCS: estimating gene genealogies. *Molecular*
545 *Ecology* 9:1657-1659. DOI:10.1046/j.1365-294x.2000.01020.x.
546

547 Coddington JA, Aganrsson I, Cheng R-C, Candek K, Driskell A, Frick H, Gregorič M,
548 Kostanjšek R, Kropf C, Kweskin M, Lokovšek T, Pipan M, Vidergar N, Kunter M. 2016. DNA
549 barcode data accurately assign higher spider taxa. *PeerJ* 4:e2201. DOI: 10.7717/peerj.2201.
550

551 Coulianos C-C, Økland J, Økland KA. 2008. Norwegian water bugs. Distribution and ecology
 552 (Hemiptera-Heteroptera: Gerromorpha and Nepomorpha). *Norwegian Journal of Entomology*
 553 55:179-222.

554

555 Creer S, Deiner K, Frey S, Porazinska D, Taberlet P, Thomas WK, Potter C, Bik HM. 2016. The
 556 ecologist's field guide to sequence-based identification of biodiversity. *Methods in Ecology and*
 557 *Evolution* 7:1008-1018. DOI: 10.1111/2041-210X.12574.

558

559 Cristescu ME. 2014. From barcoding single individuals to metabarcoding biological
 560 communities: towards an integrative approach to the study of global biodiversity. *Trends in*
 561 *Ecology and Evolution* 29:566-571. DOI: <http://dx.doi.org/10.1016/j.tree.2014.08.001>.

562

563 Damgaard J. 2005. Genetic diversity, taxonomy, and phylogeography of the western Palaearctic
 564 water strider *Aquarius najas* (DeGeer) (Heteroptera: Gerridae). *Insect Systematics & Evolution*
 565 36:395-406. DOI: 10.1163/187631205788838537.

566

567 Damgaard J. 2008. MtDNA diversity and species phylogeny of western Palaearctic members of
 568 the *Gerris lacustris* group (Hemiptera-Heteroptera: Gerridae) with implications for "DNA
 569 barcoding" of water striders. *Insect Systematics & Evolution* 39:107-120. DOI:
 570 10.1163/187631208788784156.

571

572 Damgaard J. 2008b. MtDNA diversity and phylogeography of five Palaearctic water striders
573 (Hemiptera-Heteroptera: Gerridae). In: Grozewa S, Simov N, eds. *Advances in Heteroptera*
574 *Research*. Sofia-Moscow: Pensoft Publishers, 17-30.

575

576 Dijkstra K-DB, Monagha MT, Pauls SU. 2014. Freshwater biodiversity and aquatic insect
577 diversification. *Annual Review of Entomology* 59:153-163. DOI:
578 <https://doi.org/10.1146/annurev-ento-011613-161958>.

579

580 Ebong SMA, Petit E, Le Gall P, Chen P-P, Nieser N, Guilbert E, Njiokou F, Marsollier L,
581 Guégan J-F, Pluot-Sigwalt D, Eyangoh S, Harry M. 2016. Molecular species delimitation and
582 morphology of aquatic and sub-aquatic bugs (Heteroptera) in Cameroon. *PLoS ONE* 11:
583 e0154905. DOI: 10.1371/journal.pone.0154905.

584

585 Edgar RC. 2004. MUSCLE: a multiple sequence alignment method with reduced time and space
586 complexity. *BMC Bioinformatics* 5:113. DOI: 10.1186/1471-2105-5-113.

587

588 Eyre MD, Foster GN. 1989. A comparison of aquatic Heteroptera and Coleoptera communities
589 as a basis for environmental and conservation assessments in static water sides. *Journal of*
590 *Applied Entomology* 108:355-362. DOI: 10.1111/j.1439-0418.1989.tb00468.x.

591

592 Felsenstein J. 1985. Confidence limits on phylogenies: an approach using the bootstrap.
593 *Evolution* 39:783-791. DOI: 10.2307/2408678.

594

595 Folmer O, Black M, Hoeh W, Lutz R, Vrijenhoek R. 1994. DNA primers for amplification of
596 mitochondrial cytochrome c oxidase subunit I from diverse metazoan invertebrates. *Molecular*
597 *Marine Biology and Biotechnology* 3:294-299.

598

599 Frezal L, Leblois R. 2008. Four years of DNA barcoding: Current advances and prospects.
600 *Infection, Genetics and Evolution* 8:727-736. DOI: 10.1016/j.meegid.2008.05.005.

601

602 Gagnon M-C, Turgeon. 2010. Disjunct distributions in *Gerris* species (Insecta: Hemiptera:
603 Gerridae): an analysis based on spatial and taxonomic patterns of genetic diversity. *Journal of*
604 *Biogeography* 37:170-178. DOI: 10.1111/j.1365-2699.2009.02195.x.

605

606 Galactos K, Cognato AI, Sperling FAH. 2002. Population genetic structure of two water strider
607 species in the Ecuadorian Amazon. *Freshwater Biology* 47:391-399. DOI: 10.1046/j.1365-
608 2427.2002.00811.x.

609

610 Geller J, Meyer C, Parker M, Hawk H. 2013. Redesign of PCR primers for mitochondrial
611 cytochrome c oxidase subunit I for marine invertebrates and application in all-taxa biotic
612 surveys. *Molecular Ecology Resources* 13:851-61. DOI: 10.1111/1755-0998.12138.

613

614 Grebennikov VV, Heiss E. 2014. DNA barcoding of flat bugs (Hemiptera: Aradidae) with
615 phylogenetic implications. *Arthropod Systematics & Phylogeny* 72:213-219.

616

617 Gullan PA, Cranston PS. 2014. *The Insects - An outline of Entomology. Fifth Edition*. Oxford:
618 Wiley-Blackwell.

619

620 Hajibabaei M, Singer GAC, Hebert PDN, Hickey DA. 2007. DNA barcoding: how it
621 complements taxonomy, molecular phylogenetics and population genetics. *Trends in Genetics*
622 23:167-172. DOI: <https://doi.org/10.1016/j.tig.2007.02.001>.

623

624 Hausmann A, Godfray HCJ, Huemer P, Mutanen M, Rougerie R, van Nieukerken EJ,
625 Ratnasingham S, Hebert PDN. 2013. Genetic patterns in European Geometrid moths revealed by
626 the Barcode Index Number (BIN) System. *PLoS ONE* 8:e84518. DOI:
627 <https://doi.org/10.1371/journal.pone.0084518>.

628

629 Hebert PDN, Cywinska A, Ball SL, deWaard JR. 2003a. Biological identifications through DNA
630 barcodes. *Proceedings of the Royal Society Biological Sciences* 270:313-21. DOI:
631 10.1098/rspb.2002.2218.

632

633 Hebert PDN, Ratnasingham S, deWaard JR. 2003. Barcoding animal life: cytochrome *c* oxidase
634 subunit 1 divergences among closely related species. *Proceedings of the Royal Society*
635 *Biological Sciences* 270:96-99. DOI: 10.1098/rsbl.2003.0025.

636

637 Hebert PDN, Penton EH, Burns JM, Janzen DH, Hallwachs W. 2004. Ten species in one: DNA
638 barcoding reveals cryptic species in the neotropical skipper butterfly *Astraptes fulgerator*.

639 *Proceedings of the National Academy of Sciences of the United States of America* 101:14812-
640 14817. DOI: 10.1073/pnas.0406166101.

641

642 Hendrich L, Morinière J, Haszprunar G, Hebert PD, Hausmann A, Köhler F, Balke M. 2015. A
643 comprehensive DNA barcode database for Central European beetles with a focus on Germany:
644 adding more than 3500 identified species to BOLD. *Molecular Ecology Resources* 15:795-818.
645 DOI: 10.1111/1755-0998.12354.

646

647 Henry TJ. 2017. Biodiversity of Heteroptera. In: Foottit RG, Adler PH, eds. *Insect biodiversity:*
648 *Science and Society*. Hoboken: Wiley, 279-335.

649

650 Hufnagel L, Bakonyi G, Vásárhelyi T. 1999. New approach for habitat characterization based on
651 species lists of aquatic and semiaquatic bugs. *Environmental Monitoring and Assessment* 58:305-
652 316. DOI: <https://doi.org/10.1023/A:1006047130545>.

653

654 Hurst GD, Jiggins FM. 2005. Problems with mitochondrial DNA as a marker in population,
655 phylogeographic and phylogenetic studies: the effects of inherited symbionts. *Proceedings of the*
656 *Royal Society Biological Sciences* 272:1525-1534. DOI: 10.1098/rspb.2005.3056.

657

658 Hutchinson G. 1993. *A treatise on limnology, Vol. 4, The Zoobenthos*. New York: Wiley & Sons.

659

660 Ivanova NV, Zemlak TS, Hanner RH, Hebert PDN. 2007. Universal primer cocktails for fish
661 DNA barcoding. *Molecular Ecology Notes* 7:544-548. DOI: 10.1111/j.1471-8286.2007.01748.x.

662

663 Jansson A. 1979a. Reproductive isolation and experimental hybridization between *Arctocorisa*
664 *carinata* and *A. germari* (Heteroptera, Corixidae). *Annales Zoologici Fennici* 16: 89-104. DOI:
665 <http://www.jstor.org/stable/23734416>.

666

667 Jansson A. 1979b. Experimental hybridization of *Sigara striata* and *S. dorsalis* (Heteroptera,
668 Corixidae). *Annales Zoologici Fennici* 16:105-114. DOI: <http://www.jstor.org/stable/23734417>.

669

670 Jansson A. 1983. Three new palaearctic species of *Sigara* (*Subsigara*) (Heteroptera, Corixidae).
671 *Annales Entomologici Fennici* 49:65-70.

672

673 Jansson A. 1986. The Corixidae (Heteroptera) of Europe and some adjacent regions. *Acta*
674 *Entomologica Fennica* 47:1-94.

675

676 Jeziorski P, Kment P, Ditrich T, Straka M, Sychra J, Dvořák L. 2012. Distribution of *Gerris*
677 *asper* and *G. lateralis* (Hemiptera: Heteroptera: Gerridae) in the Czech Republic. *Klapalekiana*
678 48:191-202.

679

680 Jung S, Duwal RK, Lee S. 2011. COI barcoding of true bugs (Insecta, Heteroptera). *Molecular*
681 *Ecology Resources* 11:266-270. DOI: 10.1111/j.1755-0998.2010.02945.x.

682

683 Kavar T, Pavlovčič P, Sušnik S, Meglič V, Virant-Doberlet M. 2006. Genetic differentiation of
684 geographically separated populations of the southern green stink bug *Nezara viridula*

685 (Hemiptera: Pentatomidae). *Bulletin of Entomological Research* 96:117-128. DOI:
 686 10.1079/BER2005406.

687

688 Kaur H, Sharma K. 2017. COI-based DNA barcoding of some species of Pentatomidae from
 689 North India (Hemiptera: Heteroptera). *Mitochondrial DNA Part A* 28:756-761. DOI:
 690 <http://dx.doi.org/10.1080/24701394.2016.1180513>.

691

692 Kearse M, Moir R, Wilson A, Stones-Havas S, Cheung M, Sturrock S, Buxton S, Cooper A,
 693 Markowitz S, Thierer CDT, Ashton B, Meintjes P, Drummond A. 2012. Geneious Basic: an
 694 integrated and extendable desktop software platform for the organization and analysis of
 695 sequence data. *Bioinformatics* 15:1647-1649. DOI:
 696 <https://doi.org/10.1093/bioinformatics/bts199>.

697

698 Kress WJ, García-Robledo C, Uriarte M, Erickson DL. 2015. DNA barcodes for ecology,
 699 evolution, and conservation. *Trends in Ecology & Evolution* 30:25-35. DOI:
 700 <http://dx.doi.org/10.1016/j.tree.2014.10.008>

701

702 Khalaji-Pirbalouty V, Raupach MJ (2014) A new species of *Cymodoce* Leach, 1814 (Crustacea:
 703 Isopoda: Sphaeromatidae) from the Persian Gulf based on morphological and molecular
 704 characteristics, with a redescription of *Cymodoce tribullis* from Queensland. *Zootaxa* 3826: 230-
 705 254. DOI: 10.11646/zootaxa.3826.1.7.

706

707 Kimura M. 1980. A simple method for estimating evolutionary rate of base substitutions through
708 comparative studies of nucleotide sequences. *Journal of Molecular Evolution* 16:111-120. DOI:
709 <https://doi.org/10.1007/BF01731581>.

710

711 Klecka J. 2014. The role of a water bug, *Sigara striata*, in freshwater food webs. *PeerJ* 2:e389.
712 DOI: 10.7717/peerj.389.

713

714 Klementová BR, Svitok M. 2014. *Anisops sardeus* (Heteroptera): A new expansive species in
715 Central Europe. *Biologia* 69:676-680. DOI: 10.2478/s11756-014-0354-z

716

717 Kmiec B, Woloszynska M, Janska H. 2006. Heteroplasmy as a common state of mitochondrial
718 genetic information in plants and animals. *Current Genetics* 50:149-159. DOI:
719 <https://doi.org/10.1007/s00294-006-0082-1>.

720

721 Kumar S, Stecher G, Tamura K. 2016. MEGA7: Molecular Evolutionary Genetics Analysis
722 version 7.0 for bigger datasets. *Molecular Biology and Evolution* 33:1870-1874. DOI:
723 10.1093/molbev/msw054.

724

725 Lancaster J, Downes BJ. 2013. *Aquatic Entomology. First Edition*. Oxford: Oxford University
726 Press.

727

728 Lejzerowicz F, Esling P, Pillet L, Wildling TA, Black KD, Pawlowski J. 2015. High-throughput
 729 sequencing and morphology perform equally well for benthic monitoring of marine ecosystems.
 730 *Scientific Reports* 5:13932. DOI: 10.1038/srep13932
 731
 732 Leese F, Altermatt F, Bouchez A, Ekrem T, Hering D, Meissner K, Mergen P, Pawlowski J,
 733 Piggott J, Rimet F, Steinke D, Taberlet P, Weigand A, Abarenkov K, Beja P, Bervoets L,
 734 Björnsdóttir S, Boets P, Boggero A, Bones A, Borja Á, Bruce K, Bursić V, Carlsson J, Čiampor
 735 F, Čiamporová-Zatovičová Z, Coissac E, Costa F, Costache M, Creer S, Csabai Z, Deiner K,
 736 DelValls Á, Drakare S, Duarte S, Eleršek T, Fazi S, Fišer C, Flot J, Fonseca V, Fontaneto D,
 737 Grabowski M, Graf W, Guðbrandsson J, Hellström M, HersHKovitz Y, Hollingsworth P,
 738 Japoshvili B, Jones J, Kahlert M, Kalamujic Stroil B, Kasapidis P, Kelly M, Kelly-Quinn M,
 739 Keskin E, Kõljalg U, Ljubešić Z, Maček I, Mächler E, Mahon A, Marečková M, Mejdandzic M,
 740 Mircheva G, Montagna M, Moritz C, Mulk V, Naumoski A, Navodaru I, Padisák J, Pálsson S,
 741 Panksep K, Penev L, Petrusek A, Pfannkuchen M, Primmer C, Rinkevich B, Rotter A, Schmidt-
 742 Kloiber A, Segurado P, Speksnijder A, Stoev P, Strand M, Šulčius S, Sundberg P, Traugott M,
 743 Tsigenopoulos C, Turon X, Valentini A, van der Hoorn B, Várбірó G, Vasquez Hadjilyra M,
 744 Viguri J, Vitonytė I, Vogler A, Vrålstad T, Wägele W, Wenne R, Winding A, Woodward G,
 745 Zegura B, Zimmermann J. 2016. DNAqua-Net: Developing new genetic tools for bioassessment
 746 and monitoring of aquatic ecosystems in Europe. *Research Ideas & Outcomes* 2:e11321. DOI:
 747 10.3897/rio.2.e11321.
 748
 749 Leigh JW, Bryant D. 2015. POPART: Fullfeature software for haplotype network construction.
 750 *Methods in Ecology and Evolution* 6: 1110-1116. DOI: 10.1111/2041-210X.12410.

751

752 Leite LAR. 2012. Mitochondrial pseudogenes in insect DNA barcoding: differing points of view

753 on the same issue. *Biota Neotropica* 12:3. DOI:

754 <http://www.biotaneotropica.org.br/v12n3/en/abstract?thematic-review+bn02412032012>.

755

756 Lin X, Stur E, Ekrem T. 2015. Exploring genetic divergence in a species-rich insect genus using

757 2790 DNA barcodes. *PLoS ONE* 10:e0138993. DOI:

758 <https://doi.org/10.1371/journal.pone.0138993>.

759

760 Lis A, Maryńska-Nadachowska A, Kajtoch Ł. 2015. Relations of *Wolbachia* infection with

761 phylogeography of *Philaenus spumarius* (Hemiptera: Aphrophoridae) populations within and

762 beyond the Carpathian contact zone. *Microbial Ecology* 70:509-521. DOI: 10.1007/s00248-015-

763 0570-2.

764

765 Lis B, Lis JA, Ziaja DJ. 2013. Identification of the nymphal stages of two European seed bugs,

766 *L. equestris* and *L. simulans* (Hemiptera: Heteroptera: Lygaeidae), using DNA barcodes. *Zootaxa*

767 3608:147-150. DOI: <http://dx.doi.org/10.11646/zootaxa.3608.2.5>.

768

769 Lobo J, Shokralla S, Costa MH, Hajibabaei M, Costa FO. 2017. DNA metabarcoding for high-
770 throughput monitoring of estuarine microbenthic communities. *Scientific Reports* 7:15618. DOI:
771 10.1038/s41598-017-15823-6.

772

773 McCafferty W. 1981. *Aquatic Entomology*. Boston: Science Books International.

774

775 Miller SE, Hausmann A, Hallwachs W, Janzen DH. 2016. Advancing taxonomy and
776 bioinventories with DNA barcodes. *Philosophical Transactions of the Royal Society B:*
777 *Biological Sciences* 371:20150339. DOI: <http://dx.doi.org/10.1098/rstb.2015.0339>.

778

779 Mohrbeck I, Raupach MJ, Martínez Arbizu P, Knebelsberger T, Laakmann S. 2015. High-
780 throughput sequencing – the key to rapid biodiversity assessment of marine Metazoa? *PLoS*
781 *ONE* 10:e0140342. DOI: <https://doi.org/10.1371/journal.pone.0140342>

782

783 Morinière J, Hendrich L, Balke M, Beermann AJ, König T, Hess M, Koch S, Müller R, Leese F,
784 Hebert PDN, Hausmann A, Schubart CD, Haszprunar G. 2017. A DNA barcode library for
785 Germany`s mayflies, stoneflies and caddisflies (Ephemeroptera, Plecoptera and Trichoptera).
786 *Molecular Ecology Resources* 17:1293-1307.

787

788 Morinière J, Cancian de Araujo B, Lam AW, Hausmann A, Balke M, Schmidt S, Hendrich L,
789 Doczkal D, Fartmann B, Arvidsson S, Haszprunar G. 2016. Species identification in Malaise trap
790 samples by DNA barcoding based on NGS technologies and a scoring matrix. *PLoS ONE*
791 11:e0155497. DOI: 10.1371/journal.pone.0155497.

792

793 Muraji M, Miura T, Nakasuji F. 1989. Phenological studies on the wing dimorphism of a semi-
794 aquatic bug, *Microvelia douglasi* (Heteroptera: Veliidae). *Researches on Population Ecology*
795 31:129-138. DOI: <https://doi.org/10.1007/BF02515811>.

796

797 Nieser N. 1982. De Nederlandse water- en oppervlakte wantsen (Heteroptera: Nepomorpha en
798 Gerromorpha). *Wetenschappelijke mededelingen van de koninklijke Nederlandse*
799 *Natuurhistorische Vereniging* 155:1-103. (in Dutch)

800

801 Park DS, Foottit R, Maw E, Hebert PDN. 2011. Barcoding bugs: DNA-based identification of
802 the true bugs (Insecta: Hemiptera: Heteroptera). *PLoS ONE* 6:e18749. DOI:
803 <https://doi.org/10.1371/journal.pone.0018749>.

804

805 Paterson ID, Mangan R, Downie DA, Coetzee JA, Hill MP, Burke AM, Downey PO, Henry TJ,
806 Compton SG. 2016. Two in one: cryptic species discovered in biological control agent
807 populations using molecular data and crossbreeding experiments. *Ecology and Evolution* 6:6139-
808 6150. DOI: [10.1002/ece3.2297](https://doi.org/10.1002/ece3.2297).

809

810 Peckarsky BL. 1982. Aquatic insect predator-prey relations. *BioScience* 32:261-266. DOI:
811 <https://doi.org/10.2307/1308532>.

812

813 Petit RJ, Excoffier L. 2009. Gene flow and species delimitation. *Trends in Ecology and*
814 *Evolution* 24:386-393. DOI: [10.1016/j.tree.2009.02.011](https://doi.org/10.1016/j.tree.2009.02.011).

815

816 Pfenning B, Poethke HJ. 2006. Variability in the life history of the water strider *Gerris lacustris*
817 (Heteroptera: Gerridae) across small spatial scales. *Ecological Entomology* 31:123-130. DOI:
818 10.1111/j.0307-6946.2006.00763.x.

819

820 Polhemus JT, Polhemus DA. 2008. Global diversity of true bugs (Heteroptera; Insecta) in
821 freshwater. *Hydrobiologia* 595:379-391. DOI: <https://doi.org/10.1007/s10750-007-9033-1>.

822

823 Ratnasingham S, Hebert PD. 2007. BOLD: The Barcode of Life Data System
824 (www.barcodinglife.org). *Molecular Ecology Notes* 7:355-364. DOI: 10.1111/j.1471-
825 8286.2007.01678.x.

826

827 Ratnasingham S, Hebert PD. 2013. A DNA-based registry for all animal species: The Barcode
828 Index Number (BIN) system. *PLoS ONE* 8:e66213. DOI:
829 <https://doi.org/10.1371/journal.pone.0066213>.

830

831 Raupach MJ, Astrin JJ, Hannig K, Peters MK, Stoeckle MY, Wägele J-W. 2010. Molecular
832 species identification of Central European ground beetles (Coleoptera: Carabidae) using nuclear
833 rDNA expansion segments and DNA barcodes. *Frontiers in Zoology* 7:26. DOI:
834 <https://doi.org/10.1186/1742-9994-7-26>.

835

836 Raupach MJ, Hendrich L, Kuchler SM, Deister F, Morinière J, Gossner MM. 2014. Building-up
837 of a DNA barcode library for true bugs (Insecta: Hemiptera: Heteroptera) of Germany reveals

838 taxonomic uncertainties and surprises. *PLoS ONE* 9:e106940. DOI:
839 <https://doi.org/10.1371/journal.pone.0106940>.
840
841 Raupach MJ, Radulovici AE. 2015. Looking back on a decade of barcoding crustaceans.
842 *ZooKeys* 539:53-81. DOI: 10.3897/zookeys.539.6530.
843
844 Raupach MJ, Barco A, Steinke D, Beermann J, Laakmann S, Mohrbeck I, Neumann H, Kihara
845 TC, Pointner K, Radulovici A, Segelken-Voigt A, Wesse C, Knebelsberger T. 2015. The
846 application of DNA barcodes for the identification of marine crustaceans from the North Sea and
847 adjacent regions. *PLoS ONE* 10:e0139421. DOI: <https://doi.org/10.1371/journal.pone.0139421>.
848
849 Raupach MJ, Amann R, Wheeler QD, Roos C. 2016. The application of “-omics” technologies
850 for the classification and identification of animals. *Organisms, Diversity & Evolution* 16:1-12.
851 DOI: 10.1007/s13127-015-0234-6.
852
853 Rebijith KB, Asokan R, Kumar NK, Srikumar KK, Ramamurthy VV, Bhat PS. 2012. DNA
854 barcoding and development of species-specific markers for the identification of tea mosquito
855 bugs (Miridae: Heteroptera) in India. *Environmental Entomology* 41:1239-1245. DOI:
856 10.1603/EN12096.
857
858 Saitou N, Nei M. 1987. The Neighbor-joining method: A new method for reconstructing
859 phylogenetic trees. *Molecular Biology and Evolution* 4:406-425. DOI:
860 <https://doi.org/10.1093/oxfordjournals.molbev.a040454>.

861

862 Savage AA. 1989. Adults of British aquatic Hemiptera Heteroptera: A key with ecological notes.

863 *Freshwater Biological Association Scientific Publications* 50:1-173.

864

865 Savage A, Parkin E. 1998. The diagnostic features of British *Sigara striata*, *S. dorsalis* and

866 intermediate specimens (Corixidae), with a new key for identification of adult males. *Freshwater*

867 *Forum* 10:35-48. DOI: <http://aquaticcommons.org/id/eprint/4589>.

868

869 Scheffers BR, Joppa LN, Pimm SL, Laurance WF. 2012. What we know and don't know about

870 Earth's missing biodiversity. *Trends in Ecology & Evolution* 27:501-510. DOI:

871 10.1016/j.tree.2012.05.008.

872

873 Schuh RT, Slater JA. 1995. *True Bugs of the World (Hemiptera: Heteroptera): Classification*

874 *and Natural History*. Ithaca: Cornell University Press.

875

876 Schuster G. 1983. Zur Trennung von *Gerris lateralis* Schumm. und *Gerris asper* Fieb. *Bericht*

877 *der Naturforschenden Gesellschaft Augsburg* 38:3-6. (in German)

878

879 Shackleton M, Rees GN. 2015. DNA barcoding Australian macroinvertebrates for monitoring

880 programs: benefits and current short comings. *Marine and Freshwater Research* 67:380-390.

881 DOI: <https://doi.org/10.1071/MF14331>.

882

883 Simon C, Frati F, Beckenbach A, Crespi B, Liu H, Flook P. 1994. Evolution, weighting, and
884 phylogenetic utility of mitochondrial gene sequences and a compilation of conserved polymerase
885 chain reaction primers. *Annals of the Entomological Society of America* 87:651-701. DOI:
886 <https://doi.org/10.1093/aesa/87.6.651>.
887
888 Skern M, Zweimüller I, Schiemer F. 2010. Aquatic Heteroptera as indicators for terrestrialisation
889 of floodplain habitats. *Limnologica - Ecology and Management of Inland Waters* 40:241-250.
890 DOI: <https://doi.org/10.1016/j.limno.2009.09.002>.
891
892 Song H, Moulton MJ, Whiting MF. 2014. Rampant nuclear insertion of mtDNA across diverse
893 lineages within Orthoptera (Insecta). *PLoS ONE* 9:e110508. DOI:
894 <https://doi.org/10.1371/journal.pone.0110508>.
895
896 Soós N, Petri A, Nagy-László Z, Csabai. 2010. *Anisops sardeus* Herrich-Schaeffer, 1849: first
897 records from Hungary (Heteroptera: Notonectidae). *Folia Entomologica Hungarica* 71:15-18.
898
899 Spelda J, Reip HS, Oliveira-Biener U, Melzer RR. 2011. Barcoding Fauna Bavarica: Myriapoda
900 - a contribution to DNA sequence-based identifications of centipedes and millipedes (Chilopoda,
901 Diplopoda). *ZooKeys* 156:123-139. DOI: <http://doi.org/10.3897/zookeys.156.2176>.
902
903 Spence JR, Wilcox RS. 1986. The mating system of two hybridizing species of water striders
904 (Gerridae). II. Alternative tactics of males and females. *Behavioral Ecology and Sociobiology*
905 19:87-95. DOI: <https://doi.org/10.1007/BF00299943>.

906

907 Spence JR, Andersen NM. 1994. Biology of water striders – interactions between systematics

908 and ecology. *Annual Review of Entomology* 39:101-128. DOI:

909 <https://doi.org/10.1146/annurev.en.39.010194.000533>.

910

911 Stoffelen E, Henderickx H, Vercauteren T, Lock K, Bosmans R. 2013. *De water- en*

912 *oppervlaktewantsen van België*. Brussel: Koninklich Belgisch Instituut voor

913 Natuurwetenschappen. (in Dutch)

914

915 Strauss G, Niedringhaus R. 2014. *Die Wasserwanzen Deutschlands - Bestimmungsschlüssel für*

916 *alle Nepo- und Gerromorpha*. Scheeßel: WABV. (in German)

917

918 Tautz D, Arctander P, Minelli A, Thomas RH, Vogler AP. 2003. A plea for DNA taxonomy.

919 *Trends in Ecology and Evolution* 18:70-74. DOI: [https://doi.org/10.1016/S0169-5347\(02\)00041-](https://doi.org/10.1016/S0169-5347(02)00041-1)

920 1.

921

922 Tembe S, Shouche Y, Ghate HV. DNA barcoding of Pentatomorpha bugs (Hemiptera:

923 Heteroptera) from Western Ghats of India. *Meta Gene* 2:737-745. DOI:

924 [10.1016/j.mgene.2014.09.006](https://doi.org/10.1016/j.mgene.2014.09.006).

925

926 Wachmann E, Melber A, Deckert J. 2006. *Die Tierwelt Deutschlands 77. Teil. Wanzen Band 1.*

927 *Dipsocoromorpha, Nepomorpha, Gerromorpha, Leptopodomorpha, Cimicomorpha (Teil 1).*

928 Keltern: Goecke & Evers. (in German)

929

930 Wagner E, Zimmermann S. 1955. Beitrag zur Systematik der Gattung *Gerris* F. (Hemiptera-
931 Heteroptera, Gerridae). *Zoologischer Anzeiger* 155:177-190. (in German)

932

933 Werren JH, Zhang W, Guo LR. 1995. Evolution and phylogeny of *Wolbachia*: reproductive
934 parasites of arthropods. *Proceedings of the Royal Society B: Biological Sciences* 261:55-71 DOI:
935 10.1098/rspb.1995.0117.

936

937 Werren JH, Baldo L, Clark ME. 2008. *Wolbachia*: master manipulators of invertebrate biology.
938 *Nature Reviews* 6:741-751. DOI: 10.1038/nrmicro1969.

939

940 Wesenberg-Lund C. 1943. *Biologie der Süßwasserinsekten* (Reprint). Luxemburg: Springer. (in
941 German)

942

943 Whiteman NK, Sites RW. 2008. Aquatic insects as umbrella species for ecosystem protection in
944 Death Valley National Park. *Journal of Insect Conservation* 12:499-509. DOI: 10.1007/s10841-
945 007-9090-9.

946

947 Wilcox RS, Spence JR. 1986. The mating system of two hybridizing species of water striders
948 (Gerridae). I. Ripple signal functions. *Behavioral Ecology and Sociobiology* 19:79-85. DOI:
949 <https://doi.org/10.1007/BF00299942>.

950

951 Xiao J-H, Wang N-X, Murphy RW, Cook J, Jia L-Y, Huang D-W. 2011. *Wolbachia* infection
 952 and dramatic intraspecific mitochondrial DNA divergence in a fig wasp. *Evolution* 66:1907-
 953 1916. DOI: 10.1111/j.1558-5646.2011.01561.x.

954

955 Ye Z, Zhu G, Damgaard J, Chen X. Chen P, Bu W. 2016. Phylogeography of a semi-aquatic bug,
 956 *Microvelia horvathi* (Hemiptera: Veliidae): an evaluation of historical, geographical and
 957 ecological factors. *Scientific Reports* 6:21932. DOI: 10.1038/srep21932.

958

959 Zera AJ, Innes DJ, Saks ME. 1983. Genetic and environmental determinants of wing
 960 polymorphism in the waterstrider *Limnoporus canaliculatus*. *Evolution* 31:513-522. DOI:
 961 10.2307/2408264.

962

963 Zhou C, Kandemir I, Walsh DB, Zalom FG, Lavine LC. 2012. Identification of *Lygus hesperus*
 964 by DNA barcoding reveals insignificant levels of genetic structure among distant and habitat
 965 diverse populations. *PLoS ONE* 7:e34528. DOI: <https://doi.org/10.1371/journal.pone.0034528>.

966

967 Zimmermann , Spence JR. 1989. Prey use of the fishing spider *Dolomedes triton* (Pisauridae,
 968 Araneae): an important predator of the neuston community. *Oecologia* 80:187-194. DOI:
 969 10.1007/BF00380149.


Figure 1(on next page)

Representative images of analyzed aquatic bug species.

A: *Nepa cinerea* Linnaeus, 1758 (Nepidae), **B:** *Corixa affinis* Leach, 1817 (Corixidae), **C:** *Sigara (Subsigara) scotti* (Douglas & Scott, 1868) (Corixidae), **D:** *Ilyocoris cimicoides* (Linnaeus, 1758) (Naucoridae), **E:** *Aphelocheirus aestivalis* (Fabricius, 1794) (Aphelocheiridae), **F:** *Notonecta viridis* Delcourt, 1909 (Notonectidae), **G:** *Plea minutissima* Leach, 1817 (Pleidae), **H:** *Mesovelvia furcata* Mulsant & Rey, 1852 (Mesovelidae), **I:** *Hydrometra gracilenta* Horváth, 1899 (Hydrometridae), **J:** *Hebrus ruficeps* Thompson, 1871 (Hebridae), **K:** *Velia caprai* Tamanini, 1947 (Velidae), **L:** *Gerris costae* (Herrich-Schaeffer, 1850) (Gerridae). Scale bars = 1 mm. All images were obtained from www.corisa.de.



A



B




C




D




E




F



G



H



I



J



K



L

Figure 2

Sampling sites of the studied aquatic true bugs (Gerromorpha, Nepomorpha) across Europe.




Figure 3(on next page)

Neighbor Joining (NJ) topology of the analyzed species of the Gerromorpha based on Kimura 2-parameter distance.

Triangles indicate the relative number of individual's sampled (height) and sequence divergence (width). Blue triangles indicate species with intraspecific maximum pairwise distances $>2.2\%$, red triangles species pairs with interspecific distances $<2.2\%$. Numbers next to nodes represent non-parametric bootstrap values $>80\%$ (1,000 replicates). Asterisks indicate species not recorded in Germany. All images were obtained from www.corisa.de.




Figure 4(on next page)

Neighbor Joining (NJ) topology of the analyzed species of the Nepomorpha based on Kimura 2-parameter distance.

Triangles indicate the relative number of individual's sampled (height) and sequence divergence (width). Blue triangles indicate species with intraspecific maximum pairwise distances $>2.2\%$, red triangles species with interspecific distances $<2.2\%$. Numbers next to nodes represent non-parametric bootstrap values $\geq 80\%$ (1,000 replicates). Asterisks indicate species not recorded in Germany. All images were obtained from www.corisa.de.




Figure 5(on next page)

Maximum statistical parsimony network of various species of the Gerromorpha and Nepomorpha with interspecific K2P-based distances of COI sequences <1%.

A *Gerris asper* (Fieber, 1860) ($n = 1$) and *Gerris lateralis* Schummel, 1832 ($n = 2$); B *Sigara limitata* (Fieber, 1848) ($n = 2$) and *Sigara semistriata* (Fieber, 1848) ($n = 5$); C *Callicorixa praeusta* (Fieber, 1848) ($n = 23$) and *Callicorixa producta* (Reuter, 1880) ($n = 1$). Used settings included default settings for connection steps whereas gaps were treated as fifth state. Each line represents a single mutational change whereas small black dots and small black lines indicate missing haplotypes. The diameter of the circles is proportional to the number of haplotypes sampled (see open half circles with numbers). Color codes were given for each species. Scale bars = 1 mm. Aquatic bug images were obtained from www.corisa.de.








Figure 6(on next page)


Maximum statistical parsimony network of three *Sigara* species with interspecific K2P-based distances of COI sequences <1%.


Used settings included default settings for connection steps whereas gaps were treated as fifth state. Each line represents a single mutational change whereas small black dots and small black lines indicate missing haplotypes. The diameter of the circles is proportional to the number of haplotypes sampled (see open half circles with numbers). Color codes were given for each species. Scale bars = 1 mm. Aquatic bug images were obtained from www.corisa.de.



 *Sigara distincta* (Fieber, 1848)
(*n* = 7)



 *Sigara falleni* (Fieber, 1848)
(*n* = 12)




 *Sigara iactans* Jansson, 1983
(*n* = 12)

Figure 7 (on next page)

Subtree of the Neighbour Joining topology of the analyzed specimens of *Plea minutissima* Leach, 1817.

Branches with specimen ID-Number from BOLD and species names. Numbers next to internal branches are non-parametric bootstrap values (in %). Scale bar = 1 mm. Image obtained from www.corisa.de.




Figure 8(on next page)

Subtree of the Neighbour Joining topology of the analyzed species of the genus *Cymatia* Flor, 1860.

Numbers next to internal branches are non-parametric bootstrap values (in %). Images obtained from www.corisa.de.




Table 1(on next page)

BOLD distance analysis of the studied Gerromorpha and Nepomorpha.

With the number of analyzed specimens (*n*), phylogenetic categories (PC), barcode index number (BIN), maximum intraspecific pairwise K2P distances (MID), minimum interspecific pairwise K2P distances to the nearest neighbor species (DNN), and the nearest neighbor species (NNS). Maximum intraspecific distances >2.2% and minimum interspecific distances <2.2% are marked in bold. At least one specimen of the compared species showed a distance value above or below this threshold in terms of a pairwise comparison. Asterisks (*) indicate species not recorded for Germany

Family	Species	n	PC	BIN	MID	DNN	NNS
Aphelocheiridae	<i>Aphelocheirus aestivalis</i>	2	mono	ABX0398	0	11.86	<i>Notonecta maculata</i>
Corixidae	<i>Arctocorisa carinata</i>	5	para	AAJ7903, ACY1261	2.36	1.03	<i>Arctocorisa germari</i>
	<i>Arctocorisa germari</i>	1	n. a.	-	0	1.03	<i>Arctocorisa carinata</i>
	<i>Callicorixa praeusta</i>	23	para	AAK1938	0.31	0	<i>Callicorixa producta</i>
	<i>Callicorixa producta</i>	1	n. a.	AAK1938	0	0	<i>Callicorixa praeusta</i>
	<i>Corixa affinis</i>	13	mono	ACY0615	1.92	5.92	<i>Corixa panzeri</i>
	<i>Corixa dentipes</i>	1	n. a.	-	0	6.08	<i>Corixa punctata</i>
	<i>Corixa panzeri</i>	2	mono	ACX9506	0	5.92	<i>Corixa affinis</i>
	<i>Corixa punctata</i>	21	mono	ACB1799	0.77	6.08	<i>Corixa dentipes</i>
	<i>Cymatia bondsdorffii</i>	4	mono	ABX0396	0.62	12.34	<i>Cymatia coleoprata</i>
	<i>Cymatia coleoprata</i>	24	mono	ACB1796, ADD1561	9.44	12.4	<i>Cymatia bondsdorffii</i>
	<i>Cymatia rogenhoferi</i>	1	n. a.	ACB2132	0	12.7	<i>Cymatia coleoprata</i>
	<i>Glaenocorisa propinqua</i>	8	mono	ABX4248	1.55	9.96	<i>Sigara semistriata</i>
	<i>Hesperocorixa castanea</i>	14	mono	ABX0447	0.32	13.22	<i>Paracorixa concinna</i>
	<i>Hesperocorixa linnaei</i>	8	mono	ABX0448	0	11.89	<i>Sigara venusta</i>
	<i>Hesperocorixa sahlbergi</i>	39	mono	AAN0795	1.7	11.34	<i>Corixa panzeri</i>
	<i>Micronecta griseola</i>	2	mono	AAK6480	0	10.63	<i>Micronecta poweri</i>
	<i>Micronecta poweri</i>	2	mono	ACB1970	2.39	10.63	<i>Micronecta griseola</i>
	<i>Micronecta scholtzi</i>	6	mono	AAK6479	0.16	18.58	<i>Sigara semistriata</i>
	<i>Paracorixa concinna</i>	11	mono	ABV3365, ADG5371	1.71	7.03	<i>Sigara semistriata</i>
	<i>Sigara distincta</i>	7	poly	ABY7152, ABV4484	5.77	0.37	<i>Sigara falleni</i>
	<i>Sigara dorsalis*</i>	1	n. a.	AAJ6688	0	1.71	<i>Sigara striata</i>
	<i>Sigara falleni</i>	12	poly	AAH9524, ABY7152	3.37	0	<i>Sigara iactans</i>
	<i>Sigara fossarum</i>	3	mono	AAJ6707, ADD1512	2.82	2.72	<i>Sigara scotti</i>
	<i>Sigara hellensii</i>	2	mono	ADH9592, ACT7694	4.41	9.09	<i>Sigara distincta</i>
	<i>Sigara iactans</i>	12	poly	ABY7152, AAH9524	2.67	0	<i>Sigara falleni</i>
	<i>Sigara lateralis</i>	14	mono	AAJ6697	0.81	9.84	<i>Sigara striata</i>
	<i>Sigara limitata</i>	2	para	ACM1221	0.48	0.15	<i>Sigara semistriata</i>
	<i>Sigara nigrolineata</i>	16	mono	ACB1978	0.46	10.12	<i>Sigara semistriata</i>
	<i>Sigara scotti</i>	12	mono	ACY0807	1.08	2.72	<i>Sigara fossarum</i>
	<i>Sigara semistriata</i>	5	poly	ACM1221	0	0.15	<i>Sigara limitata</i>
	<i>Sigara stagnalis</i>	6	mono	ACY0713	0.55	11.45	<i>Paracorixa concinna</i>
	<i>Sigara striata</i>	10	mono	AAJ6688	0.93	1.71	<i>Sigara dorsalis</i>
	<i>Sigara venusta</i>	2	mono	ABA5309	0	2.11	<i>Sigara semistriata</i>
Naucoridae	<i>Ilyocoris cimicoides</i>	17	mono	AAF2590	1.03	15.06	<i>Hesperocorixa sahlbergi</i>
Nepidae	<i>Nepa cinerea</i>	10	mono	AAK8359	0.34	17.06	<i>Notonecta maculata</i>
	<i>Ranatra linearis</i>	16	mono	AAL1328	0.84	15.03	<i>Notonecta lutea</i>
Notonectidae	<i>Anisops sardeus*</i>	5	mono	ABV0079	1.24	12.84	<i>Notonecta maculata</i>
	<i>Notonecta glauca</i>	41	mono	AAK4442	1.71	1.08	<i>Notonecta obliqua</i>
	<i>Notonecta lutea</i>	19	mono	AAN1701	0.68	1.24	<i>Notonecta reuteri</i>
	<i>Notonecta maculata</i>	10	mono	AAN1703	2.43	6.56	<i>Notonecta glauca</i>
	<i>Notonecta obliqua</i>	9	mono	AAK4442	0.64	1.08	<i>Notonecta glauca</i>
	<i>Notonecta reuteri</i>	5	mono	ACE8526	0.46	1.24	<i>Notonecta lutea</i>
	<i>Notonecta viridis</i>	10	mono	ABV0133	1.18	5.03	<i>Notonecta glauca</i>
Pleidae	<i>Plea minutissima</i>	17	mono	ACY0868, AAF3832	8.3	10.92	<i>Notonecta lutea</i>
Gerridae	<i>Aquarius najas</i>	7	mono	AAN1521	2.14	11.75	<i>Gerris thoracicus</i>
	<i>Aquarius paludum</i>	19	mono	AAI7450	1.24	12.61	<i>Gerris argentatus</i>
	<i>Gerris argentatus</i>	32	mono	ADD1846	0.72	6.55	<i>Gerris odontogaster</i>
	<i>Gerris asper</i>	1	n. a.	ABA3327	0	0.34	<i>Gerris lateralis</i>
	<i>Gerris costae</i>	11	mono	ACI6181	0	7.48	<i>Gerris thoracicus</i>

	<i>Gerris gibbifer</i>	11	mono	ACB1756	0.88	8.91	<i>Gerris lacustris</i>
	<i>Gerris lacustris</i>	38	mono	ACT3584	1.05	8.91	<i>Gerris gibbifer</i>
	<i>Gerris lateralis</i>	2	mono	ABA3327	0.17	0.34	<i>Gerris asper</i>
	<i>Gerris odontogaster</i>	19	mono	ABU6679, ADD1838	1.59	6.55	<i>Gerris argentatus</i>
	<i>Gerris thoracicus</i>	6	mono	ACB1745	0.35	7.48	<i>Gerris costae</i>
	<i>Limnopus rufoscutellatus</i>	3	mono	AAV0261	0.88	11.86	<i>Gerris asper</i>
Hebridae	<i>Hebrus pusillus</i>	2	mono	AAN0981	0.15	14.32	<i>Hebrus ruficeps</i>
	<i>Hebrus ruficeps</i>	7	mono	AAI6967	0.15	14.32	<i>Hebrus pusillus</i>
Hydrometridae	<i>Hydrometra gracilenta</i>	9	mono	AAN0857	0.46	13.06	<i>Hydrometra stagnorum</i>
	<i>Hydrometra stagnorum</i>	21	mono	AAK5632	0.62	13.06	<i>Hydrometra gracilenta</i>
Mesoveliidae	<i>Mesovelia furcata</i>	17	mono	AAN2451	1.39	16.24	<i>Mesovelia vittigera</i>
	<i>Mesovelia vittigera*</i>	4	mono	ACD4048	2.32	16.24	<i>Mesovelia furcata</i>
Veliidae	<i>Microvelia buenoi</i>	1	n. a.	ACY1789	0	15.06	<i>Gerris costae</i>
	<i>Microvelia reticulata</i>	27	mono	AAG4341	0.77	15.04	<i>Gerris asper</i>
	<i>Velia caprai</i>	20	mono	AAN0455	1.1	4.94	<i>Velia saulii</i>
	<i>Velia currens*</i>	3	mono	ADI1962	0	2.82	<i>Velia saulii</i>
	<i>Velia saulii</i>	1	n. a.	ABX0836	0	2.82	<i>Velia currens</i>