

BAREFOOT RUNNING AND REACTION TIME.

1 **Barefoot running does not affect simple reaction time: an exploratory study.**

2 Nicholas J. Snow¹, Jason F. L. Blair², Graham Z. MacDonald³, Jeannette M. Byrne^{2, †}, Fabien A.
3 Basset^{2, †}

4 ¹ Faculty of Medicine, Memorial University of Newfoundland, St. John's, NL, Canada.

5 ² School of Human Kinetics and Recreation, Memorial University of Newfoundland, St. John's,
6 NL, Canada

7 ³ Human Performance Laboratory, Faculty of Kinesiology, University of Calgary, Calgary, AB,
8 Canada

9

10 † Co-principal Investigator.

11

12 Corresponding Author:

13 Fabien Basset²

14 Physical Education Building, Memorial University of Newfoundland, St. John's, NL, A1C 5S7

15 Email address: fbasset@mun.ca

BAREFOOT RUNNING AND REACTION TIME.

Abstract

Background. Converging evidence comparing barefoot (BF) and shod (SH) running highlights differences in foot-strike patterns and running economy, as well as somatosensory feedback. Anecdotal evidence from SH runners attempting BF running suggests a greater attentional demand may be experienced during BF running. However, little work to date has examined whether there is an attentional cost of BF versus SH running. **Objective.** This exploratory study aimed to examine whether an acute bout of BF running would impact simple reaction time (SRT) compared to SH running, in a sample of runners naïve to BF running. **Methods.** Eight male distance runners completed SRT testing during 10 minutes of BF or SH treadmill running at 70% maximal aerobic speed ($17.9 \pm 1.4 \text{ km hr}^{-1}$). To test SRT, participants were required to press a hand-held button in response to the flash of a light bulb placed in the center of their visual field. SRT was tested at 1-minute intervals during running. . BF and SH conditions were completed in a pseudo-randomized and counterbalanced crossover fashion. SRT was defined as the time elapsed between the light bulb flash and the button press. SRT errors were also recorded and were defined as the number of trials in which a button press was not recorded in response to the light bulb flash. **Results.** Overall, SRT later in the exercise bouts showed a statistically significant increase compared to earlier. Statistically significant decrements in SRT were present at 7 minutes versus 5 minutes ($0.29 \pm 0.02 \text{ s}$ vs. $0.27 \pm 0.02 \text{ s}$, $p < 0.05$) and at 9 minutes versus 2 minutes ($0.29 \pm 0.03 \text{ s}$ vs. $0.27 \pm 0.03 \text{ s}$, $p < 0.05$); however, BF running did not influence this increase in SRT ($p > 0.05$) or the number of SRT errors (17.6 ± 6.6 trials vs. 17.0 ± 13.0 trials, $p > 0.05$). **Discussion.** In a sample of distance runners naïve to BF running, there was no statistically significant difference in SRT or SRT errors during acute bouts of BF and SH running. We interpret these results to mean that BF running does not have a greater attentional

Commented [md1]: Remove space

BAREFOOT RUNNING AND REACTION TIME.

39 cost compared to SH running during a SRT task throughout treadmill running. Literature
40 suggests that stride-to-stride gait modulation during running may occur predominately via
41 mechanisms that preclude conscious perception, thus potentially attenuating effects of increased
42 somatosensory feedback experienced during BF running. Additionally, the completion of
43 treadmill running, as opposed to over-ground, may have masked possible attentional differences
44 across running conditions. Future research should explore the present experimental paradigm in a
45 larger sample using over-ground running trials, as well as employing different or more complex
46 tests of attention.

Introduction

Despite a considerable amount of research focusing on footwear's role in injury prevention (Gallant & Pierrynowski, 2014), injury rates have remained constant over the past 40 years (Lieberman, 2012). Approximately 85% of runners experience running-related musculoskeletal injuries throughout their running career, and 30-70% of runners are treated for these injuries annually (Nielsen, Buist, Sorensen, Lind, & Rasmussen, 2012). This high prevalence of running-related injuries has led to investigations into the mechanisms contributing to their etiology (Hreljac, 2005), and to alternative solutions beyond the classic recommendation of a change in footwear characteristics. In this context, barefoot (BF) running has been proposed as an alternative solution (Murphy, Curry, & Matzkin, 2013), and has gained substantial traction in the public (Hryvniak, Dicharry, & Wilder, 2014). Indeed, recent literature describes several distinct differences between shod (SH) and BF running. Of interest are changes in foot-strike patterns (Divert, Baur, Mornieux, Mayer, & Belli, 2005; Lieberman et al., 2010; Squadron & Gallozzi, 2009), movement kinematics (Squadron & Gallozzi, 2009), and muscle activation (Snow, Basset, & Byrne, 2016; von Tscharner, Goepfert, & Nigg, 2003).

BF running has been promoted as a method to increase foot plantar sensation (Robbins, Gouw, & Hanna, 1989; Robbins, Gouw, McClaran, & Waked, 1993; Robbins & Hanna, 1987), feedback that is believed to be masked during SH running (Robbins et al., 1993; Robbins, Waked, & McClaran, 1995). Improved plantar sensory feedback associated with BF running has been suggested to modify kinematic variables (e.g., stride length, stride frequency) in a way that ultimately alters foot-strike patterns (Daoud et al., 2012; Lieberman, 2012) to reduce plantar pain and the risk of injury due to repetitive impact (e.g., during SH running with a heel-to-toe foot-strike; Lieberman et al., 2010). Given that running involves chronic repetitive movement, it is no

BAREFOOT RUNNING AND REACTION TIME.

surprise that kinematic differences in the running gait can greatly impact one's risk of injury (Daoud et al., 2012; Hreljac, 2005). Furthermore, running is associated with a high risk of acute injury due to trips, falls, and sprains (Hsu, 2012; Knobloch, Yoon, & Vogt, 2008); thus, the prevalence of both acute and chronic injuries in running demonstrate that efficient cognitive processing and rapid reaction to perturbations or obstacles is imperative during a bout of running.

Prior work has suggested that the speed of basic information processing is a valid indicator of higher cognitive function (Kail & Salthouse, 1994; Woods, Wyma, Yund, Herron, & Reed, 2015), and that decrements in higher-order cognitive operations can be reflected by diminished performance in simple tasks evaluating the speed of processing (Kail & Salthouse, 1994; Woods et al., 2015). Simple reaction time (SRT) is a task frequently used to measure speed of processing (Woods et al., 2015). Tests of SRT often involve making a physical response (e.g., pressing a button) to the presentation of a visual stimulus (e.g., light bulb flash). SRT is thus defined as minimum amount of time needed to respond to the stimulus (Woods et al., 2015). Furthermore, when an SRT task is simultaneously combined with another demanding situation (e.g., exercise), it causes a dual-task situation (Watanabe & Funahashi, 2017). In the dual-task, poorer performance often results in one or both tasks, relative to when they are preformed alone (Watanabe & Funahashi, 2017). This dual-task interference effect has been established as an important indicator of humans' limited capacity for information processing (Watanabe & Funahashi, 2017). Past evidence has demonstrated that concurrent acute exercise can result in decrements in performance on SRT tasks (Brisswalter, Arcelin, Audiffren, & Delignières, 1997; McMorris & Keen, 1994). In particular, a dual-task interference effect was observed when a non-preferred running stride frequency was adopted (Brisswalter, Durand, Delignieres, & Legros,

BAREFOOT RUNNING AND REACTION TIME.

1995; Collardeau, Brisswalter, & Audiffren, 2001). This evidence suggests that maintaining a non-preferred gait pattern required considerable attention, sufficient to cause participants' SRT performance to deteriorate. If a non-preferred running stride frequency can alter SRT, then SRT could also be compromised under BF running conditions that produce greater sensory feedback and cause participants to pay more attention to their foot-strike (Abernethy, Hanna, & Plooy, 2002; Brisswalter et al., 1995; Hanson, Whitaker, & Heron, 2009). In line with this idea, a recent study suggested that BF running requires a greater level of attention than SH, in particular due to a greater need to focus on placing footfalls on the ground (Alloway, Alloway, Magyari, & Floyd, 2016).

The literature also indicates that individuals with worse reaction time performance are prone to increased ankle instability (Konradsen & Ravn, 1990, 1991), and may be at a greater risk of falls (Richardson, Exkner, Allet, Kim, & Ashton-Miller, 2017) and acute lower-limb injuries such as ankle sprains (Beynnon, Renström, Alosa, Baumhauer, & Vacek, 2001; D. F. Murphy & Connolly, 2003; Willems et al., 2005). Thus, given the greater sensory feedback associated with BF running (Robbins et al., 1989; Robbins & Hanna, 1987), it is possible that BF running requires a greater attentional demand relative to SH running, potentially leading to decrements in SRT and increased acute injury risk due to sprains, collisions, stumbles, or falls.

Therefore, the purpose of the present study was to determine if an acute bout of BF running influenced SRT compared to SH running, in a sample of competitive distance runners naïve to BF running. We hypothesized that SRT would be increased during BF running, relative to SH running at a similar exercise intensity.

Materials & Methods

Commented [md2]: Insert space

Commented [md3]: Excellent!

Commented [md4]: Remove space

BAREFOOT RUNNING AND REACTION TIME.

This study was approved by the Interdisciplinary Committee on Ethics in Human Research at the Memorial University of Newfoundland (ethics approval number: 20130246-HK), with informed consent being gathered in accordance with the principles outlined by the Declaration of Helsinki.

Research participants were recruited using posters distributed throughout the Memorial University of Newfoundland campus athletic facilities and Physical Education Building, as well as local running retail outlets in the community of St. John's, NL, Canada. A standard recruitment email was also distributed to local running clubs and the Memorial University of Newfoundland's varsity running team.

Participants

Inclusion criteria consisted of the following: i) male participants; ii) aged between 19 and 30 years; iii) experienced runners, operationally defined as a function of running experience (≥ 2 years of active running training), frequency (≥ 4 days of running per week), and duration (≥ 30 minutes per day spent running); iv) competitive racing experience (≤ 18 minutes to complete 5 km; ≤ 40 minutes to complete 10 km); v) experience training at a high intensity (3-4 min km⁻¹ running pace, ≥ 1 day per week); vi) free of any chronic illnesses including cardiometabolic, neurological, or psychiatric diagnoses, or neuromuscular and musculoskeletal injuries, for at least 3 months; vii) active use of a minimalist running shoe (≤ 4 mm heel-toe drop, measured as the difference between the sole height of the heel and toe of the shoe), with greater than 3 months of experience and at least 3 days of use per week, for at least 30 minutes per session; and viii) naïve to BF running.

Twenty-three participants responded to recruitment materials. Of these participants, nine were eligible to participate. However, one participant withdrew from the study after initial

BAREFOOT RUNNING AND REACTION TIME.

138 enrollment, due to personal time constraints. Thus, eight adult male distance runners were
139 enrolled in the present study. No *a priori* power calculations were performed to arrive at this
140 sample size; rather, this convenience-sample of eight participants was examined as an
141 exploratory study, to generate data intended to inform further research on the present topic.

Commented [md5]: Excellent!

142 Participants were all competitive runners in the general preparatory phase of their training.
143 Participants were experienced with treadmill running. Participants completed both a short- and
144 long-form Physical Activity Readiness Questionnaire (PAR-Q) (Canadian Society for Exercise
145 Physiology (CSEP), 2002) to ensure that they were injury-free for at least 3 months prior to
146 enrolling in the study, and to screen for any injuries or health conditions that would preclude
147 their inclusion in the study. Participants were instructed to refrain from strenuous exercise 36
148 hours prior to testing (Dannecker & Koltyn, 2014); to avoid caffeine, alcohol, drugs, or
149 supplements 24 hours prior to testing; and were required to obtain at least 6 hours of sleep the
150 night prior to each testing session.

151 **Experimental Conditions**

152 Each participant was subjected to both BF and SH experimental conditions in a crossover
153 fashion (Figure 1). For the BF running condition, rubber-gripped toe-socks (Gaiam No Slip Yoga
154 Socks, Gaiam Inc., Boulder, CO, USA) were used. All participants were naïve to BF running and
155 running in the socks provided. As participants had likely not developed the appropriate responses
156 to minimize discomforts associated with true barefoot running (Lieberman, 2012; Robbins et al.,
157 1993), the toe-socks were used to help minimize potential abrasions or friction burns associated
158 with true barefoot running on a treadmill (Snow et al., 2016). For the SH running condition,
159 participants were required to bring in a pair of their own running shoes (Snow et al., 2016), to
160 mitigate any negative influences footwear discomfort may have on SRT (Mündermann, Nigg,

BAREFOOT RUNNING AND REACTION TIME.

Humble, & Stefanyshyn, 2003). In accordance with previous studies, these running shoes were defined as shoes which were > 225 g in mass, had a > 5 mm heel-toe drop, and were with or without medial arch support or impact attenuation features (Esculier, Dubois, Dionne, Leblond, & Roy, 2015; Rixe, Gallo, & Silvis, 2012). Participants did not wear minimalist footwear to complete the experiment.

Experimental Set-up and Protocol

The experimental protocol was administered over two testing sessions separated by ≥ 48 hours, as depicted in Figure 1. All testing sessions were conducted in the morning.

Testing session one. During this baseline session, participants' informed consent, anthropometrics (i.e., body mass, height), and demographic information were collected first. Participants were then familiarized with the experimental conditions and set-up, before completing an incremental treadmill exercise test. All running trials were conducted on a Cybex 750T motorized treadmill (Cybex International, Inc., Medway, MA, USA) set at a constant 1% grade to account for air-resistance experienced when running outdoors (Jones & Doust, 1996).

For familiarization, participants were instructed to run at a self-selected treadmill speed in both their standard running footwear and the toe-socks provided (BF running, 10.7 ± 0.8 km h⁻¹; SH running, 11.0 ± 0.5 km h⁻¹), for 2.5 minutes each, in random order. Heart rate (HR) was not samples during these running trials. Furthermore, these running trials were not intended to habituate participants to the footwear conditions in advance of SRT testing, but rather to provide them with an expectation of how each running condition felt, to aid in their decision to either continue in the subsequent study session or withdraw. Immediately following this familiarization session, exercise testing was conducted, and consisted of an incremental treadmill test to exhaustion, to determine maximal O₂ uptake ($\dot{V}O_{2max}$) and maximal aerobic speed (MAS),

Commented [md6]: Sampled?

BAREFOOT RUNNING AND REACTION TIME.

defined as the participants' running speed at $\dot{V}O_{2max}$ (Basset, Chouinard, & Boulay, 2003). Participants were instructed to wear their preferred footwear for exercise testing, given prior evidence that running economy is greater when participants wear shoes with a higher comfort rating (Luo, Stergiou, Worobets, Nigg, & Stefanyshyn, 2009). The incremental test started at a treadmill speed of 7.0 km hr⁻¹ and was increased by 1.0 km hr⁻¹ every 2 minutes until participants reached volitional exhaustion (Leger & Boucher, 1980). To ensure participants reached $\dot{V}O_{2max}$ upon volitional exhaustion (as opposed to peak $\dot{V}O_2$), they recovered for 5 minutes at walking speed prior to the treadmill being increased to 105 % of MAS (Rossiter, Kowalchuk, & Whipp, 2006). Participants were instructed to maintain 105 % of MAS until they reached their limit of tolerance (Rossiter et al., 2006).

Exercise metabolic rate of the incremental test was recorded with an indirect calorimetry system (AEI Technologies, Inc., Pittsburgh, PA, USA). Oxygen uptake ($\dot{V}O_2$), carbon dioxide ($\dot{V}CO_2$), breathing frequency (Bf), and tidal volume (V_T) were continuously collected with an automated open-circuit gas analysis system using O₂ and CO₂ analyzers (Model S-3A and Anarad AR-400, Ametek, Pittsburgh, PA), and a pneumo-tachometer (Model S-430, Vacumetrics/Vacumed Ltd., Ventura, CA) with a 4.2 L mixing chamber. Respiratory exchange ratio (RER) and minute ventilation ($\dot{V}_E$) were calculated as the quotient of $\dot{V}CO_2$ on $\dot{V}O_2$ and as the product of Bf by V_T , respectively. Online HR data were wirelessly transmitted to the AEI indirect calorimetric system with a Polar HR monitor (Polar Electro, Oy, Finland), via telemetry. Prior to testing, volume and gas analyzers were calibrated with a 3 L calibration syringe and medically certified O₂ and CO₂ calibration gases that were 16% O₂ and 4% CO₂, respectfully. The data were online digitalized from an A/D card to a computer for monitoring the metabolic rate (AEI Metabolic System Software, AEI Technologies, Inc., Pittsburgh, PA, USA). Results of the

BAREFOOT RUNNING AND REACTION TIME.

$\dot{V}O_{2\max}$ testing were used to determine running speeds implemented during later experimental trials.

Testing session two. On day two of testing participants completed two, 10-minute running trials, one BF and the other SH. During each of the 10-minute trials participants' SRT was tested 10 times every minute. Participants' HR was not measured during these running trials. Both BF and SH running conditions were completed at 70% of the participants' MAS, which was believed to coincide with a level of physiological arousal that optimizes SRT (Brisswalter et al., 2002; Collardeau et al., 2001). Indeed, past work suggests that exercise at such an intensity and duration can facilitate cognitive processes during exercise (Lambourne & Tomporowski, 2010; Tomporowski, 2003). This exercise intensity was selected in order to emphasize the potential effect that footwear (or lack thereof) would have on SRT, without the contaminating effect of fatigue or inappropriate exercise intensity. Conditions were pseudo-randomized and counterbalanced across the study sample, such that the order of BF and SH running was reversed for every other participant, to prevent an order effect of running conditions on SRT performance.

SRT was defined as the time required to press a hand-held button in response to the flash of a 40 W soft-white light bulb placed in the center of the participants' visual field. The SRT device used presently was developed at the Memorial University of Newfoundland, and used in previous research examining exercise effects on SRT (Behm, Bambury, Cahill, & Power, 2004). The between- and within-session reliability of SRT measurements was shown by intra-class correlation coefficients of 0.60 and 0.79 (moderate to good reliability), respectively, with no statistically significant ($p < 0.05$) differences between test and re-test values (Behm et al., 2004). Participants held the button apparatus in their dominant hand during all SRT procedures. The apparatus was also affixed to participants' wrist with a fabric-lined Velcro™ strap, to prevent

BAREFOOT RUNNING AND REACTION TIME.

dropping. To eliminate any auditory distractions, participants wore ear-plugs along with a noise-cancelling headset. To eliminate any visual distractions, barricades restricted participant's peripheral field of view. Both triggering of the light bulb (SRT stimulus) and the button-press (SRT response) were recorded at 2000 Hz, sampled using a BIOPAC MP100 biological amplifier, and displayed using AcqKnowledge 3.9.1 software (BIOPAC Systems, Inc., Goleta, CA, USA). All SRT data were stored offline on a computer and later pre-processed using AcqKnowledge and Microsoft Excel software.

Prior to commencing each running condition, participants completed three, 20-stimulus, SRT familiarization trials while standing on the treadmill at rest (Brisswalter et al., 1997, 1995). Prior to both SRT familiarization and exercise periods, participants were instructed to focus on a small target just below the light bulb and to respond quickly and vigilantly to the presented stimuli. Following SRT familiarization, participants warmed up for 5 minutes at their self-selected treadmill speed, completed the required condition (i.e. BF or SH), and then rested for 3 minutes prior to repeating the protocol, completing the second condition. In combination with the subsequent familiarization SRT trials, participants were inactive (i.e., not running) for a total of 6 minutes. This time allowed for recovery of HR between running conditions (Saltin et al., 1968), with the intention of not influencing cognitive performance under the subsequent condition (Lambourne & Tomporowski, 2010). During each 10-minute condition (i.e., BF and SH), SRT testing was administered in blocks of 10 SRT stimuli, delivered over the last 50 seconds of each minute, with each of the stimuli separated by a random interval to prevent anticipation of subsequent trials (Schmidt & Lee, 2005). The above setup was intended to produce a dual-task effect, in order to examine which running condition would have a greater

BAREFOOT RUNNING AND REACTION TIME.

influence on participants' SRT. To complement the objective SRT information provided we also asked participants to briefly comment on their experience of BF versus SH running.

Video footage was gathered on the right lower-limb, for both SH and BF running conditions to assess stride frequency. A Sony HDR-CX430VB 30 Hz video camera (Sony Computer Entertainment America, San Mateo, CA, USA) was positioned perpendicular to the treadmill at a distance of 1.5 metres and a height of 0.75 metres. Video footage was collected at a 30 Hz frame rate, in accordance with previous literature (Macpherson, Taylor, McBain, Weston, & Spears, 2016; Nikodelis, Moscha, Metaxiotis, & Kollias, 2011), with a total of 30 running strides per participant being collected for each running condition (i.e., BF and SH). Raw video data were converted to MPEG-4 using Sony PMB software (Sony Computer Entertainment America, San Mateo, CA, USA) for further analysis.

Data Analyses

$\dot{V}O_{2max}$. Participants' $\dot{V}O_{2max}$ was considered the peak value in O_2 uptake using a 30-second moving window average technique. MAS was the corresponding treadmill speed at $\dot{V}O_{2max}$ (American College of Sports Medicine (ACSM), 2013). HR_{max} was defined as the peak HR value obtained during the $\dot{V}O_{2max}$ test (ACSM, 2013).

Stride frequency. The Kinovea (Version 2.0) high-resolution video analysis software platform (<http://www.kinovea.org/>) was used to determine the frame of foot-contact and toe-off during each 10-second window, for each minute, during each 10-minute trial, for both the BF and SH running conditions (Damsted, Nielsen, & Larsen, 2015; Padulo et al., 2015). To minimize error in video interpretation, blind cross-checks were performed by two researchers (NJS, JMB). Foot-contact and toe-off were used to determine stride frequency (strides s^{-1}) by

BAREFOOT RUNNING AND REACTION TIME.

counting the number of complete strides per 10 seconds of video data. This number was then multiplied by 6 to provide the final stride frequency estimate (strides min⁻¹).

SRT. Participants' SRT was considered the time difference (in seconds) between the initiation of the SRT stimulus (light bulb) and the completion of the SRT response (button press) (Magill, 2011). As such, SRT encompassed participants' overall response time, which is comprised of both reaction time (i.e., time between stimulus presentation and initiation of response) and movement time (i.e., time between response initiation and response completion) (Magill, 2011). Therefore, our SRT measure contained a global measure of both stimulus detection (reaction time) and response execution (movement time), but did not distinguish between the two. Any SRT trial < 0.160 seconds was considered an anticipated response and was to be omitted from the data set (Brisswalter et al., 1997, 1995; Collardeau et al., 2001). However, zero SRT responses met this criterion, and thus no SRT trials were omitted from the data set on this basis. We also analyzed SRT errors, which consisted of instances where participants did not respond to the stimulus (light bulb), or when SRT trials > 1.0 second were recorded (Woods et al., 2015). No SRT trials exceeded 1.0 second, and so SRT errors were considered only those trials that un-recorded. The number of un-recorded SRT responses were counted by the AcqKnowledge software during each minute and averaged across each total running trial (BF, SH). Remaining SRT trials were averaged over each minute for their respective running conditions (i.e., BF and SH running). Increasing SRT (in seconds) was indicative of an increase in perceptual latency and attentional load.

Statistical Analyses

All data were examined for normality using the Shapiro-Wilk test and visual examination of histogram plots. Because of sensitivity to sample size variations in statistics-based normality

BAREFOOT RUNNING AND REACTION TIME.

tests, as well as the robustness of within-subjects designs to normality violations, we used a stringent significance level of $p < 0.001$ in objective examinations of the data distributions (i.e., Shapiro-Wilk test) (Gamst, Meyers, & Guarino, 2008).

The effect of BF versus SH running on SRT was tested using a two-way (2×10) repeated-measures analysis of variance (rmANOVA) with the factors Condition (BF running, SH running) and Time (1-10 min). To determine if SRT during BF running, SH running, or running in general was different from rest, the average SRT from the familiarization trials, the BF and SH running conditions, and both exercise conditions combined was compared using a one-way (4 levels) rmANOVA with the factor Condition (Familiarization, BF running, SH running, Combined running). *Post hoc* pairwise comparisons were conducted when necessary using the Bonferroni correction. Mean SRT errors (number of trials) were compared across BF and SH running conditions using a paired samples *t*-test. Finally, average stride frequency (strides min^{-1}) was compared across conditions (BF running, SH running) using a paired samples *t*-test.

All results are presented as means $\pm$ one standard deviation (SD). Statistical significance was set at $p < 0.05$. Effect sizes (Cohen's *d*) and 95% confidence intervals (95%CI) of effect sizes were calculated for our primary outcome (SRT in seconds) using Microsoft Excel, and interpreted as "trivial" < 0.20 ; "small" 0.20-0.49; "medium" 0.50-0.79; "large" > 0.80 (Cohen, 1988). All statistical analyses were conducted in SPSS (Version 20, IBM Corporation, Armonk, NY, USA).

Results

Data Inspection

All data were deemed normally distributed on the basis of Shapiro-Wilk statistics (SRT $W_{(8)} = 0.774-0.990$, $p = 0.015-0.996$; stride frequency $W_{(8)} = 0.876-0.912$, $p = 0.171-0.368$) and

BAREFOOT RUNNING AND REACTION TIME.

histogram plot inspection. For SRT data, Mauchly's Test of Sphericity was not statistically significant ($p > 0.05$); thus, sphericity was assumed for interpreting the results of the rmANOVA on SRT values.

Participants

Baseline characteristics. Participants were on average 25.1 ± 3.7 years of age, with a body mass and height of 78.4 ± 8.9 kg and 180.7 ± 7.8 cm, respectively. Their training experience ranged from 1 to 8 years (3.1 ± 2.1 yr), and all participants were experienced in using minimalist footwear (11.5 ± 11.4 mo). Participants completed an average of 8.1 ± 3.5 training sessions per week, including 1.90 ± 0.8 sessions of interval training at $> 75\% \dot{V}O_{2\max}$. Total training volume was 90.0 ± 44.7 km per week, and $65.3 \pm 44.9\%$ of training volume was completed using minimalist footwear (72.3 ± 62.0 km wk^{-1}). Average 10 km personal best race time (mm:ss) was $37:26 \pm 2:50$.

$\dot{V}O_{2\max}$ testing. Participants' $\dot{V}O_{2\max}$ was on average 61.4 ± 6.7 mL $\text{min}^{-1} \text{kg}^{-1}$, corresponding to "excellent" fitness (ACSM, 2013). In fact, all participants achieved $\dot{V}O_{2\max}$ scores in the 95th to 99th percentile based on age and sex norms (ACSM, 2013). Average $\text{HR}_{\max}$ was 191 ± 4 bpm. Mean maximal aerobic speed (MAS) was 17.9 ± 1.3 km hr^{-1} , while 70% MAS (for BF and SH running SRT trials) was 12.5 ± 0.9 km hr^{-1} .

SRT

SRT results from BF running, SH running, and combined across conditions are shown in Figure 2. A statistically significant main effect of Time ($F_{(9, 63)} = 3.097$, $p = 0.004$) was present when assessing SRT. Statistically significant increases in SRT were present at 7 minutes relative to 5 minutes (0.29 ± 0.02 s vs. 0.27 ± 0.02 s, $p < 0.05$, $d = -0.99$, 95%CI = -1.98 to +0.09, large effect), and at 9 minutes relative to 2 minutes (0.29 ± 0.03 s vs. 0.27 ± 0.03 s, $p < 0.05$, $d = -0.67$,

BAREFOOT RUNNING AND REACTION TIME.

95%CI = -1.63 to +0.37, moderate effect). There was neither a statistically significant main effect of Condition ($F_{(1, 7)} = 1.002, p = 0.350$) nor a statistically significant Condition $\times$ Time interaction effect ($F_{(9, 63)} = 1.233, p = 0.292$). Examination of effect sizes between conditions indicated that overall, BF running had a small negative effect on SRT ($d = -0.32, 95\%CI = -3.29$ to $+2.65$). The largest negative effect BF running had on SRT was at 8 minutes ($d = -0.80, 95\%CI = -1.65$ to $+0.06$, large effect; BF SRT = 0.30 ± 0.03 s, SH SRT = 0.27 ± 0.03 s) when compared to SH running. However, this effect was not statistically significant ($p > 0.05$).

Figure 3 illustrates average SRT values for the familiarization trials and each experimental condition, with an increase in SRT indicating a decrement in SRT performance. When comparing the average SRT for the familiarization trial (0.25 ± 0.03 s), SH running (0.27 ± 0.02), BF running (0.28 ± 0.03 s), and combined trials across both conditions (0.28 ± 0.02 s), there was no statistically significant main effect of Condition ($F_{(3, 21)} = 2.944, p = 0.057$).

Figure 4 demonstrates SRT errors during BF and SH running trials. There were no SRT errors during familiarization periods. However, due to limitations in the experimental set-up, we were not able to distinguish participants' non-responses (participant error) from intended button-presses that were not registered by the experimental setup (experimental error). Absent SRT responses (i.e., SRT errors) represented 17.6% and 17.0% of total SRT trials under the BF and SH running conditions, respectively (17.6 ± 6.6 trials vs. 17.0 ± 13.0 trials). There was no statistically significant difference in SRT errors across conditions ($t_{(7)} = 1.07, p = 0.918$).

Finally, seven participants reported feeling an increase in attentional demands during BF relative to SH running, while one participant noted no difference. In general, participants (7/8) highlighted: (i) a need to focus more on their footfalls to prevent uncomfortable landings; (ii) a

BAREFOOT RUNNING AND REACTION TIME.

perceived change in foot-strike patterns; and (iii) pain or burning on the plantar surface of the foot.

Stride Frequency

Stride frequency showed a statistically significant increase during BF running, relative to SH running (88.26 ± 5.58 strides min^{-1} vs. 86.09 ± 5.70 strides min^{-1} , $p < 0.05$, $d = 0.38$, small effect).

Discussion

The purpose of the present study was to determine whether there was a difference in SRT during acute bouts of BF and SH running, in competitive distance runners naïve to BF running. Despite a statistically significant increase in SRT during later time-points of the exercise bouts compared to earlier, we did not observe a statistically significant difference in SRT across footwear conditions. However, we found a statistically significant increase in stride frequency during BF running; and participants anecdotally reported having perceived an increase in attentional demands during BF relative to SH running. Nevertheless, we observed no objective alteration of the sensorimotor processing and attentional load during BF compared to SH running, as measured using SRT.

SRT and Attentional Demands

Reaction time is an ecologically-relevant measure of perceptual-motor cost of running (Schmidt & Lee, 2005), due to the high prevalence of acute injuries sustained during running (Hsu, 2012; Knobloch et al., 2008), combined with the common nature of situations requiring reactions to extrinsic stimuli (Magill, 2011). In some instances, individuals must react quickly and suddenly to an unexpected stimulus to avoid injury, making it imperative to avoid any threat to reaction time performance (Magill, 2011; Schmidt & Lee, 2005). On this basis, increased SRT

BAREFOOT RUNNING AND REACTION TIME.

has been linked to a possible increased injury risk during running, due to falls and sprains (Beynnon et al., 2001; Konradsen & Ravn, 1990, 1991; Murphy & Connolly, 2003; Richardson et al., 2017; Willems et al., 2005). Past work highlights that runners need to pay more attention to their foot-strikes during BF running (Alloway et al., 2016), and to alter their running kinematics to avoid noxious plantar stimuli (Lieberman et al., 2010). Consequently, we hypothesized that BF running would produce a detrimental effect on SRT performance, with reference to SH running; yet we observed no statistically significant effect of BF running on SRT or SRT errors.

In the present study, participants anecdotally reported an increase in attentional demands during BF relative to SH running, noting: (i) a need to focus more on their footfalls to prevent uncomfortable landings; (ii) a perceived change in foot-strike patterns, particularly during the latter minutes of BF running; and (iii) pain or burning on the plantar surface of the foot. A recent study directly comparing BF and SH running trials on an indoor track showed that runners had to pay greater attention to their foot-strikes during the BF condition, as evidenced by greater working memory when stepping on targets during running (Alloway et al., 2016). This observation is supported by work that has indicated that SRT performance is decreased in the presence of externally applied cutaneous stimulation (Hanson et al., 2009). When considering the concept of dual-task interference, which emphasizes participants' limited attentional capacity (Watanabe & Funahashi, 2017), it could be expected that cognitive task performance would suffer in the presence of an attentionally demanding procedure such as BF running. Therefore, at the same relative intensity, it is possible BF running does not have any additional attentional demand compared to SH running (Brisswalter et al., 1997). It is also plausible that by providing participants with rubber-gripped toe socks for the BF running condition, plantar sensory

Commented [md7]: Remove extra space

BAREFOOT RUNNING AND REACTION TIME.

feedback was masked relative to a true BF running condition, introducing a confound to our experiment.

Stride Frequency

The statistically significant increase in stride frequency with BF running may reflect kinematic differences between BF and SH running (Divert et al., 2005; Ekizos, Santuz, & Arampatzis, 2017), which could be the result of increased somatosensory feedback present during BF running, and intended to avoid painful foot-strikes (Hsu, 2012; Lieberman, 2012). Indeed, increased sensory feedback during BF running can alter foot-strike patterns, for instance by modulating ankle coordination prior to foot-strike (Kurz & Stergiou, 2004), reducing ground reaction forces (Lieberman et al., 2010), and increasing stride frequency along with decreased contact time, respectively (Ekizos et al., 2017; Kurz & Stergiou, 2004). During SH running, this “protective” feedback is believed to be impaired (Robbins et al., 1995). We anticipated that such differences in somatosensory feedback would have a limiting effect on attentional capacity, reflected in a reduction in SRT performance. Yet, in spite of our observation that stride frequency increase to a statistically significant degree during BF running, SRT did not change.

Possible Explanations

Despite anecdotal reports and past research suggesting a potential attentional difference between BF and SH footwear conditions, we did not observe such an effect. There is evidence in support of our finding, that SRT was not different during BF versus SH running. For example, Klint et al. (2008) showed that proprioceptive and cutaneous feedback from the foot and leg can modulate stepping patterns and increase variation in kinematics during BF locomotion, independent of higher processing. They concluded that higher centers are likely reserved for

BAREFOOT RUNNING AND REACTION TIME.

more complex movements and processes outside of gait. Similarly, others have intimated that afferent feedback during locomotion is processed in subcortical regions, or via transcortical reflex pathways, without higher order processing (Nielsen, 2003). This is likely why neural interactions governing gait modulation are rapid, allowing stride-to-stride variation in stepping patterns (Nigg, 2001).

Greater variation of stride kinematics during BF running might therefore be associated with injury prevention and pain reduction, due to a reduction in repeated impact (Lieberman et al., 2010). However, if the increased afferent feedback experienced during BF running does not undergo higher processing (Klint et al., 2008), then runners would be at not greater risk of acute injuries due to stumbles or falls (Beynnon et al., 2001; Konradsen & Ravn, 1990, 1991; D. F. Murphy & Connolly, 2003; Richardson et al., 2017; Willems et al., 2005). This is supported by our finding that BF running did not have a statistically significant effect on SRT. Although speculative in nature, this lack of conscious processing of somatosensory and proprioceptive information would serve a protective role for persons engaging in BF running.

Methodological Considerations and Future Directions

There are a few noteworthy methodological considerations in the current work, which could have influenced our present findings. Throughout the course of prolonged exercise fatigue can negatively influence corticospinal and neuromuscular output (Meardon, Hamill, & Derrick, 2011; Ross, Middleton, Shave, George, & Nowicky, 2007), and consequently reduce perceptual-motor performance (Brisswalter et al., 2002). The present results support existing evidence in that SRT performance tended to increase towards the end of the exercise bout, a result that has been previously reported in the literature (Brisswalter et al., 1995; Brisswalter et al., 2002; Collardeau et al., 2001; McMorris & Keen, 1994). However, this was a short and moderately-

Commented [md8]: Did these authors suggest that this subcortical activity could occur without perceptual detection?

Commented [md9]: Could then these changes in gait be occurring at rates well beyond the measuring capacity of SRT in the study (SRT sampling rate)?

Commented [md10]: Remove extra space

BAREFOOT RUNNING AND REACTION TIME.

intense bout of steady-state exercise (ACSM, 2013), with an appropriately-timed rest period (Saltin et al., 1968); so fatigue was not likely a major contributor to decreased performance (Lambourne & Tomporowski, 2010; Tomporowski, 2003). Indeed, the decrease in performance observed may have been a result of the mode of exercise (i.e., treadmill exercise), as compared to exercise intensity or fatigue (Lambourne & Tomporowski, 2010). In other words, given the dual-task nature of the present experiment, SRT performance may simply have suffered in response to participants' avoiding falling off the treadmill. Nevertheless, there was no statistically significant difference in SRT during resting familiarization trials compared to exercise. In addition, without the use of treadmill running, it would not have been possible for us to employ the present SRT task.

Additionally, our measure of SRT was unable to decompose participants' overall response time, into its constituent components of reaction time (i.e., time between stimulus presentation and initiation of response), which measures stimulus detection; and movement time (i.e., time between response initiation and response completion), which measures response execution (Magill, 2011). Past work has shown that acute exercise preferentially influences movement time over reaction time in SRT tasks (Beyer, Sage, Staines, Middleton, & McIlroy, 2017; Davranche, Burle, Audiffren, & Hasbroucq, 2005, 2006), indicating that exercise-induced changes in response time are related more to faster movement execution than changes in cognitive function (Beyer et al., 2017). Thus, to elucidate the cognitive influence of BF running it would be prudent for further work to examine a greater number of dimensions of task performance, including separating reaction and movement times. Similarly, examining more complex cognitive tasks (e.g., discrimination RT) may better discern the cognitive impacts of BF running, as opposed to SRT which simply examines speed of information processing (Alloway et

Commented [md11]: As I review figure 3, It would appear that may be there is a sig difference between SRT during resting/ familiarization when compared to exercise. However, the data for SRT during resting/ familiarization is not apparent in the data set attachment.

BAREFOOT RUNNING AND REACTION TIME.

al., 2016; Beyer et al., 2017). Finally, it is conceivable that the small sample size in this exploratory study may have threatened the validity of the observed results. Consequently, future work could benefit from examining a larger sample of runners.

Commented [md12]: May add, longer trial periods, and recreational runners who have not had experience running in minimalist running shoes.

Conclusions

In the present exploratory study, an acute bout of BF versus SH running did not impact SRT. It is possible that increased afferent feedback during BF running (Kurz & Stergiou, 2004; Robbins et al., 1993) is responded to in subcortical regions or transcortical reflex pathways (Nielsen, 2003), without affecting the attentional requirements of the task. Additionally, this may be the case only for simple tasks such as SRT. Alternatively, it is possible that our small sample size did not have sufficient power to reveal a significance across BF and SH running conditions. Nevertheless, the present results suggest that although differences in running kinematics across BF and SH running may lead to differences in musculoskeletal injuries (Daoud et al., 2012; Hreljac, 2005), it is not likely that BF running will impact runners' risk of attention-related acute injuries such as trips or falls (Hsu, 2012; Knobloch, Yoon, & Vogt, 2008). Future work should examine whether more complex perceptual-motor tasks and more sensitive outcomes will be affected by BF versus SH running. Further efforts should also examine whether the present observations will emerge in larger sample of runners. Finally, it is prudent to examine whether changes in SRT will manifest when runners are performing over-ground on a stable running surface, as opposed to during treadmill running.

Acknowledgements

The current study received no external funding support, and the authors have no conflicts of interest to declare. We wish to sincerely thank Mr. Blaise Dubois for his tremendous support

BAREFOOT RUNNING AND REACTION TIME.

503 during experimental planning, Dr. Normand Teasdale for his generous assistance in reviewing
504 the original manuscript before its initial submission, and Dr. Thamir Alkanani for his technical
505 contributions.

References

- Abernethy, B., Hanna, A., & Plooy, A. (2002). The attentional demands of preferred and non-preferred gait patterns. *Gait and Posture*, 15(3), 256–265. [https://doi.org/10.1016/S0966-6362\(01\)00195-3](https://doi.org/10.1016/S0966-6362(01)00195-3)
- Alloway, R. G., Alloway, T. P., Magyari, P. M., & Floyd, S. (2016). An Exploratory Study Investigating the Effects of Barefoot Running on Working Memory. *Perceptual and Motor Skills*, 122(2), 432–443. <https://doi.org/10.1177/0031512516640391>
- American College of Sports Medicine (ACSM). (2013). *ACSM's Guidelines for Exercise Testing and Prescription*. (9th ed.). Baltimore, MD: Lippincott Williams and Wilkins.
- Basset, F. A., Chouinard, R., & Boulay, M. R. (2003). Training profile counts for time-to-exhaustion performance. *Canadian Journal of Applied Physiology = Revue Canadienne de Physiologie Appliquee*, 28(4), 654–666.
- Behm, D. G., Bambury, A., Cahill, F., & Power, K. (2004). Effect of acute static stretching on force, balance, reaction time, and movement time. *Medicine and Science in Sports and Exercise*, 36(8), 1397–1402. <https://doi.org/10.1249/01.MSS.0000135788.23012.5F>
- Beyer, K. B., Sage, M. D., Staines, W. R., Middleton, L. E., & McIlroy, W. E. (2017). A single aerobic exercise session accelerates movement execution but not central processing. *Neuroscience*, 346, 149–159. <https://doi.org/10.1016/j.neuroscience.2017.01.020>
- Beynnon, B. D., Renström, P. a, Alosa, D. M., Baumhauer, J. F., & Vacek, P. M. (2001). Ankle ligament injury risk factors: a prospective study of college athletes. *Journal of Orthopaedic Research : Official Publication of the Orthopaedic Research Society*, 19(2), 213–220. [https://doi.org/10.1016/S0736-0266\(00\)90004-4](https://doi.org/10.1016/S0736-0266(00)90004-4)
- Brisswalter, J., Arcelin, R., Audiffren, M., & Delignières, D. (1997). Influence of physical

BAREFOOT RUNNING AND REACTION TIME.

- 529 exercise on simple reaction time: effect of physical fitness. *Perceptual and Motor Skills*,
- 530 85(3 Pt 1), 1019–27. <https://doi.org/10.2466/pms.1997.85.3.1019>
- 531 Brisswalter, J., Collardeau, M., & René, A. (2002). Effects of Acute Physical Exercise
- 532 Characteristics on cognitive performance. *Sports Medicine (Auckland, N.Z.)*, 32(9), 555–
- 533 566.
- 534 Brisswalter, J., Durand, M., Delignieres, D., & Legros, P. (1995). Optimal and non-optimal
- 535 demand in a dual task of pedalling and simple reaction time: effects on energy expenditure
- 536 and cognitive performance. *Journal of Human Movement Studies*, 29, 15–34.
- 537 Canadian Society for Exercise Physiology (CSEP). (2002). Physical activity readiness
- 538 questionnaire - PAR-Q. Retrieved from [http://www.csep.ca/cmfiles/publications/parq/par-](http://www.csep.ca/cmfiles/publications/parq/par-q.pdf)
- 539 [q.pdf](http://www.csep.ca/cmfiles/publications/parq/par-q.pdf)
- 540 Cohen, J. (1988). *Statistical power analysis for the behavioral sciences*. (2nd ed.). Hillsdale, NJ:
- 541 Lawrence Earlbaum Associates.
- 542 Collardeau, M., Brisswalter, J., & Audiffren, M. (2001). Effects of a prolonged run on simple
- 543 reaction time of well trained runners. *Perceptual and Motor Skills*, 93(3), 679–689.
- 544 <https://doi.org/10.2466/PMS.93.7.679-689>
- 545 Damsted, C., Nielsen, R. O., & Larsen, L. H. (2015). Reliability of video-based quantification of
- 546 the knee- and hip ankle at foot strike during running. *Int J Sports Phys Ther*, 10(2), 147–54.
- 547 Dannecker, E. a., & Koltyn, K. F. (2014). Pain during and within hours after exercise in healthy
- 548 adults. *Sports Medicine*, 44(7), 921–942. <https://doi.org/10.1007/s40279-014-0172-z>
- 549 Daoud, A. I., Geissler, G. J., Wang, F., Saretsky, J., Daoud, Y. A., & Lieberman, D. E. (2012).
- 550 Foot strike and injury rates in endurance runners: a retrospective study. *Medicine and*
- 551 *Science in Sports and Exercise*, 44(7), 1325–1334.

BAREFOOT RUNNING AND REACTION TIME.

- 552 Davranche, K., Burle, B., Audiffren, M., & Hasbroucq, T. (2005). Information processing during
553 physical exercise: A chronometric and electromyographic study. *Experimental Brain*
554 *Research*, 165(4), 532–540. <https://doi.org/10.1007/s00221-005-2331-9>
- 555 Davranche, K., Burle, B., Audiffren, M., & Hasbroucq, T. (2006). Physical exercise facilitates
556 motor processes in simple reaction time performance: An electromyographic analysis.
557 *Neuroscience Letters*, 396(1), 54–56. <https://doi.org/10.1016/j.neulet.2005.11.008>
- 558 Divert, C., Baur, H., Mornieux, G., Mayer, F., & Belli, A. (2005). Stiffness adaptations in shod
559 running. *Journal of Applied Biomechanics*, 21(4), 311–321.
- 560 Ekizos, A., Santuz, A., & Arampatzis, A. (2017). Transition from shod to barefoot alters
561 dynamic stability during running. *Gait & Posture*, 56(April), 31–36.
562 <https://doi.org/https://doi.org/10.1016/j.gaitpost.2017.04.035>
- 563 Esculier, J.-F., Dubois, B., Dionne, C. E., Leblond, J., & Roy, J.-S. (2015). A consensus
564 definition and rating scale for minimalist shoes. *Journal of Foot and Ankle Research*, 8(1),
565 42. <https://doi.org/10.1186/s13047-015-0094-5>
- 566 Gallant, J. L., & Pierrynowski, M. R. (2014). A theoretical perspective on running-related
567 injuries. *Journal of the American Podiatric Medical Association*, 104(2), 211–220.
- 568 Gamst, G., Meyers, L. S., & Guarino, A. J. (2008). ANOVA Assumptions. In *Analysis of*
569 *Variance Designs* (pp. 49–84). New York, NY: Cambridge University Press.
- 570 Hanson, J. V. M., Whitaker, D., & Heron, J. (2009). Preferential processing of tactile events
571 under conditions of divided attention. *Neuroreport*, 20(15), 1392–1396.
572 <https://doi.org/10.1097/WNR.0b013e3283319e25>
- 573 Hreljac, A. (2005). Etiology, prevention, and early intervention of overuse injuries in runners: a
574 biomechanical perspective. *Physical Medicine and Rehabilitation Clinics of North America*,

BAREFOOT RUNNING AND REACTION TIME.

- 575 16(3), 651–67, vi.
- 576 Hryvniak, D., Dicharry, J., & Wilder, R. (2014). Barefoot running survey: Evidence from the
- 577 field. *Journal of Sport and Health Science*, 3(2), 131–136.
- 578 <https://doi.org/10.1016/j.jshs.2014.03.008>
- 579 Hsu, A. R. (2012). Topical Review: Barefoot Running. *Foot & Ankle International*, 33(9), 787–
- 580 794. <https://doi.org/10.3113/FAI.2012.0787>
- 581 Jones, A. M., & Doust, J. H. (1996). A 1% treadmill grade most accurately reflects the energetic
- 582 cost of outdoor running. *Journal of Sports Sciences*, 14(4), 321–327.
- 583 Kail, R., & Salthouse, T. A. (1994). Processing speed as a mental capacity. *Acta Psychologica*
- 584 (Amst), 86, 199–225.
- 585 Klint, R. A. F., Nielsen, J. B., Cole, J., Sinkjaer, T., & Grey, M. J. (2008). Within-step
- 586 modulation of leg muscle activity by afferent feedback in human walking. *The Journal of*
- 587 *Physiology*, 586(Pt 19), 4643–4648. <https://doi.org/10.1113/jphysiol.2008.155002>
- 588 Knobloch, K., Yoon, U., & Vogt, P. M. (2008). Acute and overuse injuries correlated to hours of
- 589 training in master running athletes. *Foot & Ankle International / American Orthopaedic*
- 590 *Foot and Ankle Society [and] Swiss Foot and Ankle Society*, 29(7), 671–676.
- 591 <https://doi.org/10.3113/FAI.2008.0671>
- 592 Konradsen, L., & Ravn, J. B. (1990). Ankle instability caused by prolonged peroneal reaction
- 593 time. *Acta Orthopaedica Scandinavica*, 61(15), 388–390.
- 594 Konradsen, L., & Ravn, J. B. (1991). Prolonged peroneal reaction time in ankle instability.
- 595 *International Journal of Sports Medicine*, 12(3), 290–292. [https://doi.org/10.1055/s-2007-](https://doi.org/10.1055/s-2007-1024683)
- 596 1024683
- 597 Kurz, M. J., & Stergiou, N. (2004). Does footwear affect ankle coordination strategies? *Journal*

BAREFOOT RUNNING AND REACTION TIME.

- 598 of the American Podiatric Medical Association, 94(1), 53–58.
- 599 Lambourne, K., & Tomporowski, P. (2010). The effect of exercise-induced arousal on cognitive
- 600 task performance: a meta-regression analysis. *Brain Research*, 1341, 12–24.
- 601 <https://doi.org/10.1016/j.brainres.2010.03.091>
- 602 Leger, L., & Boucher, R. (1980). An indirect continuous running multistage field test: the
- 603 Universite de Montreal track test. *Canadian Journal of Applied Sport sciences. Journal*
- 604 *Canadien Des Sciences Appliquees Au Sport*, 5(2), 77–84.
- 605 Lieberman, D. E. (2012). What we can learn about running from barefoot running: an
- 606 evolutionary medical perspective. *Exercise and Sport Sciences Reviews*, 40(2), 63–72.
- 607 Lieberman, D. E., Venkadesan, M., Werbel, W. A., Daoud, A. I., D’Andrea, S., Davis, I. S., ...
- 608 Pitsiladis, Y. (2010). Foot strike patterns and collision forces in habitually barefoot versus
- 609 shod runners. *Nature*, 463(7280), 531–535.
- 610 Luo, G., Stergiou, P., Worobets, J., Nigg, B., & Stefanyshyn, D. (2009). Improved footwear
- 611 comfort reduces oxygen consumption during running. *Footwear Science*, 1, 25–29.
- 612 Macpherson, T. W., Taylor, J., McBain, T., Weston, M., & Spears, I. R. (2016). Real-time
- 613 measurement of pelvis and trunk kinematics during treadmill locomotion using a low-cost
- 614 depth-sensing camera: A concurrent validity study. *Journal of Biomechanics*, 49(3), 474–
- 615 478. <https://doi.org/10.1016/j.jbiomech.2015.12.008>
- 616 Magill, R. A. (2011). *Motor Learning and Control: Concepts and Applications* (9th ed.). New
- 617 York, NY: McGraw-Hill.
- 618 McMorris, T., & Keen, P. (1994). Effect of exercise on simple reaction times of recreational
- 619 athletes. *Perceptual and Motor Skills*, 78, 123–130.
- 620 Meardon, S. A., Hamill, J., & Derrick, T. R. (2011). Running injury and stride time variability

BAREFOOT RUNNING AND REACTION TIME.

- 621 over a prolonged run. *Gait & Posture*, 33(1), 36–40.
- 622 Mündermann, A., Nigg, B. M., Humble, R. N., & Stefanyshyn, D. J. (2003). Orthotic comfort is
- 623 related to kinematics, kinetics, and EMG in recreational runners. *Medicine and Science in*
- 624 *Sports and Exercise*, 35(24), 1710–1719.
- 625 <https://doi.org/10.1249/01.MSS.0000089352.47259.CA>
- 626 Murphy, D. F., & Connolly, A. J. (2003). Risk factors for lower extremity injury: a review of the
- 627 literature. *Br J Sports Med*, 37, 13–29. <https://doi.org/10.1136/bjsm.37.1.13>
- 628 Murphy, K., Curry, E. J., & Matzkin, E. G. (2013). Barefoot running: does it prevent injuries?
- 629 *Sports Medicine (Auckland, N.Z.)*, 43(11), 1131–1138.
- 630 Nielsen, J. B. (2003). How we walk: central control of muscle activity during human walking.
- 631 *The Neuroscientist : A Review Journal Bringing Neurobiology, Neurology and*
- 632 *psychiatryNeuroscientist*, 9(3), 195–204.
- 633 Nielsen, R. O., Buist, I., Sorensen, H., Lind, M., & Rasmussen, S. (2012). Training errors and
- 634 running related injuries: a systematic review. *International Journal of Sports Physical*
- 635 *Therapy*, 7(1), 58–75.
- 636 Nikodelis, T., Moscha, D., Metaxiotis, D., & Kollias, I. (2011). Commercial video frame rates
- 637 can produce reliable results for both normal and CP spastic gait's spatiotemporal, angular,
- 638 and linear displacement variables. *Journal of Applied Biomechanics*, 27(3), 266–271.
- 639 Padulo, J., Vando, S., Chamari, K., Chaouachi, A., Bagnò, D., & Pizzolato, F. (2015). Validity of
- 640 the MarkWiiR for kinematic analysis during walking and running gaits. *Biology of Sport*,
- 641 32(1), 53–58. <https://doi.org/10.5604/20831862.1127282>
- 642 Richardson, J. K., Exkner, J. T., Allet, L., Kim, H., & Ashton-Miller, J. (2017). Complex and
- 643 Simple Clinical Reaction Times Are Associated with Gait, Balance, and Major Fall Injury

BAREFOOT RUNNING AND REACTION TIME.

in Older Subjects with Diabetic Peripheral Neuropathy. *American Journal of Physical*

Medicine & Rehabilitation, 96(1), 8–16.

<https://doi.org/doi:10.1097/PHM.0000000000000604>.

Rixe, J. A., Gallo, R. A., & Silvis, M. L. (2012). The barefoot debate: can minimalist shoes reduce running-related injuries? *Current Sports Medicine Reports*, 11(3), 160–165.

Robbins, S. E., Gouw, G. J., & Hanna, A. M. (1989). Running-related injury prevention through innate impact-moderating behavior. *Medicine and Science in Sports and Exercise*, 21(2), 130–139.

Robbins, S. E., Gouw, G. J., McClaran, J., & Waked, E. (1993). Protective sensation of the plantar aspect of the foot. *Foot and Ankle*, 14(6), 347–352.

Robbins, S. E., & Hanna, A. M. (1987). Running-related injury prevention through barefoot adaptations. *Medicine and Science in Sports and Exercise*, 19(2), 148–156.

Robbins, S., Waked, E., & McClaran, J. (1995). Proprioception and stability: foot position awareness as a function of age and footwear. *Age and Ageing*, 24, 67–72.

<https://doi.org/10.1093/ageing/24.1.67>

Ross, E. Z., Middleton, N., Shave, R., George, K., & Nowicky, A. (2007). Corticomotor excitability contributes to neuromuscular fatigue following marathon running in man. *Experimental Physiology*, 92(2), 417–26. <https://doi.org/10.1113/expphysiol.2006.035972>

Rossiter, H. B., Kowalchuk, J. M., & Whipp, B. J. (2006). A test to establish maximum O₂ uptake despite no plateau in the O₂ uptake response to ramp incremental exercise. *Journal of Applied Physiology (Bethesda, Md.: 1985)*, 100(3), 764–770.

Saltin, B., Blomqvist, G., Mitchell, J. H., Johnson, R. L. J., Wildenthal, K., & Chapman, C. B. (1968). Response to exercise after bed rest and after training. *Circulation*, 38(5 Suppl),

Commented [md13]: Small font in many places

BAREFOOT RUNNING AND REACTION TIME.

667 VII1-78.

668 Schmidt, R. A., & Lee, T. D. (2005). *Motor Control and Learning: A Behavioral Emphasis*.
669 Champaign, IL, USA: Human Kinetics.

670 Snow, N. J., Basset, F. A., & Byrne, J. (2016). *An Acute Bout of Barefoot Running Alters*
671 *Lower-limb Muscle Activation for Minimalist Shoe Users*. *International Journal of Sports*
672 *Medicine*, 37(5), 382–387. <https://doi.org/10.1055/s-0035-1565140>.

673 Squadrone, R., & Gallozzi, C. (2009). Biomechanical and physiological comparison of barefoot
674 and two shod conditions in experienced barefoot runners. *The Journal of Sports Medicine*
675 *and Physical Fitness*, 49(1), 6–13.

676 Tomporowski, P. D. (2003). Effects of acute bouts of exercise on cognition. *Acta Psychologica*,
677 112(3), 297–324.

678 von Tscharn, V., Goepfert, B., & Nigg, B. M. (2003). Changes in EMG signals for the muscle
679 tibialis anterior while running barefoot or with shoes resolved by non-linearly scaled
680 wavelets. *Journal of Biomechanics*, 36(8), 1169–1176.

681 Watanabe, K., & Funahashi, S. (2017). Toward an understanding of the neural mechanisms
682 underlying dual-task performance: Contribution of comparative approaches using animal
683 models. *Neuroscience & Biobehavioral Reviews*, 84(February 2017), 12–28.
684 <https://doi.org/10.1016/j.neubiorev.2017.08.008>

685 Willems, T. M., Witvrouw, W., Delbaere, K., Mahieu, N., De Bourdeaudhuij, I., & De Clercq,
686 D. (2005). Intrinsic risk factors for inversion ankle sprains in male subjects: a prospective
687 study. *American Journal of Sports Medicine*, 33(3), 415–423.
688 <https://doi.org/10.1177/0363546504268137>

689 Woods, D. L., Wyma, J. M., Yund, E. W., Herron, T. J., & Reed, B. (2015). Factors influencing

Commented [md14]: Small font in many cases.

BAREFOOT RUNNING AND REACTION TIME.

690 the latency of simple reaction time. *Frontiers in Human Neuroscience*, 9(March), 1–12.
691 <https://doi.org/10.3389/fnhum.2015.00131>