

Uncovering the hidden players in Lepidoptera biology: the heritable microbial endosymbionts

Anne Duplouy¹ and Emily A. Hornett²

¹ Organismal and Evolutionary Biology Research Program, University of Helsinki, Helsinki, Finland

² Department of Zoology, University of Cambridge, Cambridge, UK

ABSTRACT

The Lepidoptera is one of the most widespread and recognisable insect orders. Due to their remarkable diversity, economic and ecological importance, moths and butterflies have been studied extensively over the last 200 years. More recently, the relationship between Lepidoptera and their heritable microbial endosymbionts has received increasing attention. Heritable endosymbionts reside within the host's body and are often, but not exclusively, inherited through the female line. Advancements in molecular genetics have revealed that host-associated microbes are both extremely prevalent among arthropods and highly diverse. Furthermore, heritable endosymbionts have been repeatedly demonstrated to play an integral role in many aspects of host biology, particularly host reproduction. Here, we review the major findings of research of heritable microbial endosymbionts of butterflies and moths. We promote the Lepidoptera as important models in the study of reproductive manipulations employed by heritable endosymbionts, with the mechanisms underlying male-killing and feminisation currently being elucidated in moths and butterflies. We also reveal that the vast majority of research undertaken of Lepidopteran endosymbionts concerns *Wolbachia*. While this highly prevalent bacterium is undoubtedly important, studies should move towards investigating the presence of other, and interacting endosymbionts, and we discuss the merits of examining the microbiome of Lepidoptera to this end. We finally consider the importance of understanding the influence of endosymbionts under global environmental change and when planning conservation management of endangered Lepidoptera species.

Submitted 22 January 2018

Accepted 27 March 2018

Published 8 May 2018

Corresponding author

Emily A. Hornett, eh481@cam.ac.uk

Academic editor

Joseph Gillespie

Additional Information and
Declarations can be found on
page 28

DOI 10.7717/peerj.4629

© Copyright

2018 Duplouy and Hornett

Distributed under

Creative Commons CC-BY 4.0

OPEN ACCESS

Subjects Conservation Biology, Ecology, Entomology, Evolutionary Studies

Keywords Symbiosis, *Wolbachia*, Conservation, Climate change, Butterfly, Moth, Microbes

INTRODUCTION

Symbiosis was originally described as the living together of dissimilar organisms in an intimate association (*de Bary, 1879*). This broad term is commonly used to encompass relationships between two or more organisms that range from parasitic, through commensal (one party gains a benefit, whilst the other is not significantly affected) to mutualistic (both parties benefit). We now know that the nature of an association is often much more complex, and varies greatly depending on factors such as the local environment, the host genetic background or condition, and the longevity of the

relationship. Thus it is perhaps now more pertinent to understand symbiosis as an interaction in which two or more organisms of different species are in a persistent relationship, with no pre-conceived idea of the nature of the interaction.

One of the most intimate associations between species is that between a host organism and a microbial endosymbiont (a symbiont living within the body of its host). This lifestyle substantially affects the relationship between the two parties as survival and reproduction of host and microbe are intrinsically linked. Where the endosymbiont is intracellular—residing within the cytoplasm of host cells—it is predominantly inherited through the female line (although intrasperm paternal transmission has also been described ([Watanabe et al., 2014](#))). Such maternal inheritance produces selection upon the symbiont to favour the cytoplasmic lineage of the host (in essence the females)—a phenomenon that has resulted in the evolution of remarkable manipulations of host reproductive biology, including sex-ratio distortion ([O'Neill, Hoffmann & Werren, 1998](#); [Bandi et al., 2001](#); [Engelstädter & Hurst, 2009](#)). With increasing pace, evidence is gathering that diverse endosymbionts interact with many aspects of arthropod host biology including host reproduction ([Werren, Zhang & Guo, 2004](#)), development ([Fraune & Bosch, 2010](#)), immunity ([Gross et al., 2009](#); [Nyholm & Graf, 2012](#)), behaviour ([Dion et al., 2011](#)), body colour ([Tsuchida et al., 2010](#)), nutritional stress resistance ([Brownlie et al., 2009](#)), pathogen load ([Graham & Wilson, 2012](#)), dispersal ([Goodacre et al., 2009](#)), host plant specialisation ([Leonardo & Muiru, 2003](#)), thermal tolerance ([Dunbar et al., 2007](#)), nutrition ([Douglas, 1998](#)) and metabolism ([McCutcheon, McDonald & Moran, 2009](#)). Furthermore, symbiosis has been purported to be a key factor underlying natural variation, as well as an instigator of novelty and a promoter of speciation ([Margulis & Fester, 1991](#); [Brucker & Bordenstein, 2012](#)).

Since the advent of the diagnostic PCR assay in the mid-1980s, organisms can be routinely screened for known endosymbionts. As a consequence of this development and recent advancements in genomics and bioinformatics (including high-throughput amplicon sequencing of microbial genes and metagenomics), we now recognise that all organisms are infected by a diverse range of microbes, including viruses, fungi and bacteria, and that many arthropods carry heritable endosymbionts. A recent study estimated that 52% of terrestrial arthropod species are infected with the intracellular bacteria *Wolbachia*, with a further 24% and 13% species infected with *Cardinium* and *Rickettsia* bacteria, respectively ([Weinert et al., 2015](#)). How species initially acquire heritable endosymbionts is not yet fully understood. While phylogenetic evidence suggests that horizontal transfer of endosymbionts on an evolutionary scale must be common, many barriers—ecological, geographical and physiological—exist that perturb the spread of endosymbionts between species and prevent the formation of novel symbioses. Successful transfer of an endosymbiont between species depends on the ability of the microbe to first enter and then survive in a novel host environment, followed by successful migration to the host germline to ensure propagation. The symbiont must then be able to invade the host population, or at least be maintained at low frequency. Thus the 'fit' between a host and symbiont can be quite specific, with host

biology playing an important role in the ability of the symbiont to thrive in the novel species. Failure in the formation of persistent associations may also be due to the endosymbiont causing harm to their new hosts ([Hutchence et al., 2011](#)). Where movement of heritable endosymbionts has been observed, it is often via ecological connectors such as shared host food sources ([Huigens et al., 2000](#); [Duron, Wilkes & Hurst, 2010](#); [Caspi-Fluger et al., 2012](#); [Chrostek et al., 2017](#)) or common symbiont-vector parasites or parasitoids ([Heath et al., 1999](#); [Vavre et al., 1999](#); [Huigens et al., 2004](#); [Jaenike et al., 2007](#); [Gehrer & Vorburger, 2012](#)). Horizontal transfer is perhaps more successful between related hosts ([Russell et al., 2009](#)); it has been suggested that within *Acraea* butterflies, *Wolbachia* has moved between species either via a common parasitoid, or through hybridisation and subsequent introgression. It is also possible that the different species inherited the bacteria from a recent common ancestor ([Jiggins et al., 2000b](#)).

The Lepidoptera are remarkably diverse and widely recognisable, encompassing butterflies and moths that are economically and ecologically important. While many aspects of Lepidopteran biology have been well studied, it is only recently that the pervasiveness of host-associated microbes in this group has been appreciated. Heritable endosymbionts have been the subject of several reviews ([Bandi et al., 2001](#); [Moran, McCutcheon & Nakabachi, 2008](#); [Duron & Hurst, 2013](#)), and here we focus upon studies of these influential elements in the Lepidoptera. Butterflies and moths are particularly important in the study of heritable endosymbionts due to the Lepidoptera sex determination system. In contrast to most other arthropod groups, the female is the heterogametic sex (females have one Z and one W sex chromosome, males have two Z chromosomes). The mechanisms and repercussions of reproductive manipulations caused by inherited microbial endosymbionts, which are commonly observed in butterflies and moths, are therefore likely to be very different from that observed in arthropods with alternative sex determination systems. Furthermore, in the Lepidoptera, heritable endosymbiont prevalence is commonly very high, and vertical transmission of the infection is often near perfect. Together with the maternal inheritance of intracellular endosymbionts such as *Wolbachia*, this creates linkage of the infection not only with similarly maternally inherited host mitochondria, but also with the female W chromosome. Formation of this wider co-inherited network may have implications for host genetic diversity and even the sex determination system itself.

In this review, we summarise the main body of research that has been conducted to date in order to form a springboard for future work and to emphasise to researchers from traditionally disparate fields as ecology, genomics and conservation, that in order to fully understand the biology of an organism, one must take into account its endosymbionts. For clarity this review is divided into areas of current research: (1) Manipulation of host reproduction; (2) impact upon host fitness; (3) symbiont-mediated protection; (4) host genetics and (5) behavioural modification. We then highlight outstanding questions and future directions, including consideration of the influence of endosymbionts under global environmental change, and in species of conservation concern.

Survey methodology

The authors have drawn upon knowledge gained from over a decade in butterfly-endosymbiont research. Extensive literature searches were performed using repositories such as NCBI PubMed and Google Scholar, and using keywords including 'endosymbiont,' 'microbe' and 'heritable symbiont,' along with 'butterfly,' 'moth' and 'Lepidoptera.' Social media platforms such as Twitter provided a useful tool to obtain up to date information of relevant publications. Research on heritable endosymbionts of arthropods in general was also gathered with the aim to provide information about areas in heritable endosymbiont-arthropod research that is lacking for Lepidoptera. Particular effort was made to compile a comprehensive list of butterfly and moth species that are published as infected with heritable endosymbionts.

The influence of heritable microbial endosymbionts on Lepidopteran biology

Concordant with general insect surveys, the Lepidoptera are commonly infected with heritable microbial endosymbionts. In an early screen of Panamanian arthropods, *Wolbachia* was detected in 16.3% of the 43 Lepidoptera species tested ([Werren, Windsor & Guo, 1995](#)). Further surveys identified *Wolbachia* in 29% of 24 species of *Acraea* butterflies from Uganda ([Jiggins et al., 2001](#)), 45% of 49 species of butterflies studied in Japan ([Tagami & Miura, 2004](#)), 50% of 56 Indian butterfly species ([Salunke et al., 2012](#)), 58.3% of 120 Lepidoptera species in West Siberia ([Ilinsky & Kosterin, 2017](#)) and 79% of 24 species of African *Bicyclus* butterflies ([Duploux & Brattström, 2017](#)). Additionally, in a broad survey of ants, moths and butterflies (specifically Lycaenidae and Nymphalidae) for five heritable symbionts, *Wolbachia* (39 of 158 species) and *Spiroplasma* (five of 200 species) were found to infect Lepidopteran species ([Russell et al., 2012](#)). In general, these estimates are likely to be highly conservative, due to the presence of undetected low frequency infections, geographical and temporal variation in infection, tissue-specificity and PCR false negatives. Geographic structure in infection incidence and prevalence is a particularly important consideration and especially evident in endosymbiont-Lepidoptera systems, e.g. *Wolbachia*-*Hypolimnna bolina* butterflies ([Charlat et al., 2005](#)). In a recent survey of published records of *Wolbachia* infections in the Lepidoptera, generalised geographic structure in infection frequency was observed, with lower frequencies towards higher latitudes ([Ahmed et al., 2015](#)).

In [Tables S1](#) and [S2](#) we compile a comprehensive list of butterfly and moth species, respectively, reported as carrying heritable endosymbionts from published sources. We find that research of heritable endosymbionts in Lepidoptera is heavily dominated by studies of *Wolbachia* as opposed to that of other infections (*Wolbachia* in 248/253 butterfly species and in 109/115 moth species). While arthropod-infecting endosymbiont diversity is notable, including such divergent taxa as *Rickettsia*, *Spiroplasma*, *Arsenophonus*, *Flavobacteria*, *Cardinium* and the microsporidia, much of the early arthropod endosymbiont literature focused upon the Alphaproteobacteria genus *Wolbachia* ([Hertig & Wolbach, 1924](#)). Due to its presence in many agricultural pests and

disease vectors, and also owing to the range of reproductive manipulations it employs in the host, *Wolbachia* is still widely, but justly, studied.

Tables S1 and S2 reveal that *Wolbachia* is common across Lepidopteran families, being found in all five families of ‘true’ butterflies (the Papilionoidea), and also in the skippers (Hesperiidae). *Wolbachia* strains have been divided into separate genetic lineages termed supergroups. It is clear from the compiled data that the *Wolbachia* strains carried by Lepidoptera are almost exclusively from supergroups A and B, with B group *Wolbachia* predominating over A group *Wolbachia*. Of species where the *Wolbachia* supergroup has been determined (80/109 moths and 208/248 butterflies), 85% of moth species carry B group and 25% A group; while 79% of butterfly species carry B group *Wolbachia* and 26% carry A group. Note that these data include multiple infections (i.e. some species harbour both A and B strains of *Wolbachia*). These findings concur with research analysing 90 *Wolbachia* strains associated with Lepidoptera: 84% of the strains belonged to supergroup B (76/90), with the remainder (14/90) belonging to supergroup A (Ahmed, Breinholt & Kawahara, 2016). A further study identified 22 *Wolbachia*-infected Lepidoptera species in Japan, 19 of which had infections from supergroup B (86%), with the remaining three from supergroup A (Tagami & Miura, 2004). It is unclear why B group *Wolbachia* are particularly prevalent in the Lepidoptera; is there a greater ‘fit’ between Lepidoptera and B group *Wolbachia*, i.e. are B group *Wolbachia* more likely to become established, or are B group *Wolbachia* those ancestrally associated with the Lepidoptera thus seeding this group stochastically? It is also interesting to note that there appears to be one particularly common strain of *Wolbachia* in Lepidoptera. In a study of 53 Lepidoptera species, 11 species across three families are infected with *Wolbachia* ST41, the next most common strain types (ST40 and ST125) were found in three species each (Ahmed, Breinholt & Kawahara, 2016). Whether *Wolbachia* ST41 is especially adept at moving between species, and/or whether it is particularly successful at establishing and maintaining itself with the host remains to be fully investigated.

The second most common heritable endosymbiont recorded in butterflies and moths is *Spiroplasma*—a bacterial genus belonging to the class Mollicutes (Table S1: 5/253 butterfly species and Table S2: 5/115 moth species). Until such endosymbionts receive the same level of attention as *Wolbachia*, or there is a move towards a generalised metagenomic approach to identify symbiotic microbes, little can be said of the extent of their presence or action in Lepidoptera. However, while there is a propensity for Lepidoptera to be specifically screened for *Wolbachia* infections (thereby creating a bias towards detection of *Wolbachia*), discovery of sex-ratio distorter identity is commonly a phenotype forward investigation, i.e. a sex-ratio bias in progeny or in a population is observed, and then the causative factor is identified. Thus, there should be no bias in the responsible infection found in these studies. Despite this, it appears that Lepidoptera are different from many other groups, e.g. ladybirds, in that *Wolbachia* is almost always responsible for the observed sex-ratio bias. In comparison with other arthropod groups such as the Diptera, Hymenoptera and Hemiptera, heritable endosymbiont diversity does appear to be particularly low in the Lepidoptera. A systematic review (Russell et al., 2012)

compiling data of infection screens of arthropods for the heritable endosymbionts *Arsenophonus*, *Cardinium*, *Hamiltonella*, *Spiroplasma* and *Wolbachia*, found that only the latter two genera of bacteria were present in Lepidoptera species (*Spiroplasma*: 5/205, *Wolbachia*: 140/481 species infected). Thus we can say that for *Arsenophonus*, *Cardinium* and *Hamiltonella*, where 263, 183 and 251 Lepidopteran species were assayed respectively, such infections, should they exist at all in Lepidoptera, are remarkably rare. A later compilation of data of arthropods screened for heritable endosymbionts found that *Rickettsia* bacteria were also not commonly found in Lepidoptera, with only one species (an unidentified Noctuidae moth) infected out of 14–32 species (variation in number reported here due to several individuals tested having no taxonomic assignment in the study ([Weinert et al., 2015](#))).

Manipulation of host reproduction

The Lepidoptera are becoming model systems for the study of endosymbiont manipulation of host reproduction. Many species are infected with maternally inherited bacteria that have evolved the ability to alter host reproduction to either increase the proportion of infected females in the population, or increase the reproductive fitness of infected females relative to their uninfected counterparts. In Lepidoptera endosymbionts are currently known to manipulate host reproduction in three ways: through male-killing (MK), feminisation and cytoplasmic incompatibility (CI) ([Fig. 1](#)). While these methods facilitate the maintenance of the symbiont in the host population, there are often severe repercussions for host biology and evolution. We provide a list of butterflies and moths that have been recorded as being infected with endosymbionts that manipulate the reproductive biology of the host ([Table 1](#)).

Male-killing

Male-killing is particularly well-known in the Lepidoptera. Here, male offspring are killed early in development (most usually as an egg, but also as first instar larvae) producing a female-biased sex ratio within an infected female's offspring ([Fig. 1](#)). Should the male killer infect many females, the host population as a whole may become female-biased. Several hypotheses have been proposed to explain why maternally inherited endosymbionts kill male hosts. If infected females gain a fitness benefit from the death of their male siblings over uninfected females (whose male siblings survive), the infection will invade and spread through the host population. Such benefits may include a reduction in the likelihood of detrimental inbreeding (as there are no brothers with which to mate) or a reduction in competition for resources (as there are half as many siblings with which to compete) ([Hurst & Majerus, 1992](#); [Hurst, Hurst & Majerus, 1997](#)). In ladybirds, *Wolbachia*-infected female neonates gain an important first meal by consuming their dead brothers, while uninfected females lack this ready source of nutrients ([Elnagdy, Majerus & Handley, 2011](#)). However, in Lepidopteran systems, the relative fitness benefit for infected females remains elusive as many of the species studied lay their eggs singly, thus making the likelihood of inbreeding, sibling egg cannibalism or competition unlikely (e.g. *Danaus chrysippus* ([Jiggins et al., 2000a](#))).




Figure 1 Endosymbiont-induced manipulation of Lepidoptera reproduction. In the Lepidoptera, endosymbionts are currently known to manipulate host reproduction in three ways in order to increase their transmission to the next generation. Male-killing: female hosts infected with male-killing endosymbionts only give rise to infected female offspring, with male offspring dying early in development. Feminisation: female hosts infected with feminising endosymbionts only give rise to infected female offspring, with male offspring having been feminised so that they are genetically male (ZZ) but phenotypically female. Uninfected males may arise through inefficient transmission of the infection. Cytoplasmic incompatibility (CI): crosses between uninfected females and infected males result in few or no viable offspring, as the result of an incompatibility induced by the endosymbiont in the male. Infected females are able to rescue this incompatibility and hence are able to produce viable (infected) offspring when mated with infected males. For male-killing and feminisation the endosymbiont acts as a sex-ratio distorter, creating a female-bias in the offspring, and potentially in the population if the infection is highly prevalent. [Full-size DOI: 10.7717/peerj.4629/fig-1](https://doi.org/10.7717/peerj.4629/fig-1)

Despite the lack of evidence of any fitness benefit being provided to infected females, MK has been recorded numerous times in the Lepidoptera, possibly due to the readily observable phenotype of all-female broods and the long history of Lepidoptera being collected and reared in captivity. Early work recorded the presence of female-biases in wild-caught collections and captive bred broods in both *Acraea encedon* (Poulton, 1914; Owen, 1965, 1970), and *Hypolimnas bolina* (Poulton, 1923, 1926) butterflies. Later, MK *Wolbachia* was identified as the causative agent in both *A. encedon* (Jiggins, Hurst & Majerus, 1998; Hurst et al., 1999) and *H. bolina* (Dyson, Kamath & Hurst, 2002). We now know that populations of *Acraea* butterflies carry highly prevalent MK *Wolbachia* infections, with more than 80% and 95% of Ugandan *A. encedon* and *A. encedana* females being infected, respectively (Jiggins et al., 2000a; Jiggins, Hurst & Majerus, 2000). The *H. bolina* system has become remarkable due to the extensive spatial and temporal variation in the dynamics of the interaction across the South-east Asian to Eastern Pacific range of the butterfly (Charlat et al., 2005; Hornett et al., 2009). The island of Samoa is particularly notable due to its well-documented history of a highly biased sex ratio of 100 females to every male, caused by 99% of female butterflies

Table 1 Lepidoptera species carrying heritable endosymbionts that manipulate host reproduction.

	Host	Endosymbiont	Phenotype	Source
Butterflies	Lycaenidae			
	<i>Talicauda nyseus</i>	<i>Wolbachia</i>	Sex-ratio distortion	Ankola, Brueckner & Puttaraju (2011)
	<i>Zizina emelina</i>	<i>Wolbachia</i>	MK	Sakamoto et al. (2011)
	Nymphalidae			
	<i>Acraea acerata</i>	<i>Wolbachia</i>	CI	Jiggins et al. (2001)
	<i>Acraea encedana</i>	<i>Wolbachia</i>	MK	Jiggins et al. (2000a)
	<i>Acraea encedon</i>	<i>Wolbachia</i>	MK	Jiggins, Hurst & Majerus (1998); Jiggins et al. (2000a)
	<i>Acraea eponina</i>	<i>Wolbachia</i>	MK	Jiggins et al. (2001)
	<i>Acraea stoikensis</i>	<i>Wolbachia</i>	MK	Hassan & Idris (2013)
	<i>Danaus chrysippus</i>	<i>Spiroplasma ixodetis</i>	MK	Jiggins et al. (2000b)
	<i>Hypolimnas bolina</i>	<i>Wolbachia</i>	MK &/or CI	Dyson, Kamath & Hurst (2002); Charlat et al. (2006); Hornett et al. (2008)
	Pieridae			
	<i>Colias erate poliographus</i>	<i>Wolbachia</i>	CI	Narita, Shimajiri & Nomura (2009)
	<i>Eurema hecabe</i>	<i>Wolbachia</i>	Feminisation, CI	Narita et al. (2011)
	<i>Eurema mandarina</i>	<i>Wolbachia</i>	Feminisation, CI	Hiroki et al. (2002, 2004)
Moths	Crambidae			
	<i>Ostrinia furnacalis</i>	<i>Wolbachia</i>	MK	Kageyama et al. (2002)
	<i>Ostrinia orientalis</i>	<i>Wolbachia</i>	Sex-ratio distortion	Kageyama et al. (2004)
	<i>Ostrinia scapulalis</i>	<i>Wolbachia</i>	MK	Kageyama & Traut (2004)
	<i>Ostrinia zaguliaevi</i>	<i>Wolbachia, Spiroplasma ixodetis</i>	MK	Kageyama et al. (2004); Tabata et al. (2011)
	<i>Ostrinia zealis</i>	Undefined agent	Sex-ratio distortion	Kageyama et al. (2004)
	Erebidae			
	<i>Lymantria dispar</i>	Undefined agent	MK	Higashiura, Ishihara & Schaefer (1999)
	Noctuidae			
	<i>Cerapteryx graminis</i>	<i>Spiroplasma sp.</i>	Sex-ratio distortion	Graham, Hartley & Wilson (2011)
	<i>Spodoptera exempta</i>	<i>Wolbachia</i>	MK	Graham & Wilson (2012)
	<i>Spodoptera littoralis</i>	Undefined agent	MK	Brimacombe (1980)
	Plutellidae			
	<i>Plutella xylostella</i>	<i>Wolbachia</i>	Sex-ratio distortion	Delgado & Cook (2009)
	Pyrallidae			
	<i>Cadra cautella</i>	<i>Wolbachia</i>	CI	Sasaki & Ishikawa (1999)
	<i>Ephestia kuehniella</i>	<i>Wolbachia</i>	CI	Sasaki & Ishikawa (1999)
	Tortricidae			
	<i>Epiphyas postvittana</i>	Undefined agent	MK	Geier, Briese & Lewis (1978)
	<i>Homona magnanima</i>	RNA virus	Late MK	Morimoto et al. (2001); Nakanishi et al. (2008)

Notes:

A list of butterfly and moth species that have been recorded as carrying heritable endosymbionts that manipulate the reproduction of the host. Endosymbiont induced phenotypes are given as MK: Male-killing, Late MK: Male-killing occurring late in development; CI: Cytoplasmic Incompatibility; Feminisation; or Sex-ratio distortion (where further investigation is needed to determine the nature of the sex-ratio bias).

being infected with a MK *Wolbachia* (Dyson & Hurst, 2004). It appears that MKs are often found at a particularly high frequency within butterfly populations, contrasting patterns seen in other taxa studied such as the ladybirds, where generally less than 49% of females carry an infection (Hurst & Jiggins, 2000). In the lycaenid *Zizina emelina*, at least one of the two *Wolbachia* strains described in Japanese populations is a MK that rapidly increased in prevalence from 65% to 86% within a three year period (Sakamoto et al., 2011).

The consequences of a highly distorted sex ratio are likely to be large (for discussions of evolutionary consequences see (Charlat, Hurst & Mercot, 2003; Engelstädter & Hurst, 2007)). As perhaps can be expected, one direct effect is that a large number of females remain unmated. In Makerere, Uganda, 94% of *Wolbachia*-infected *A. encedon* females were virgins (Jiggins, Hurst & Majerus, 2000). Of Samoan *H. bolina* 50% of infected females were unmated, with the females that did mate showing significant fertility deficiencies, implying sperm limitation (Dyson & Hurst, 2004). However, despite the detrimental impacts of MKs upon the reproductive biology of their hosts, natural host populations infected with high prevalence infections can persist: the 100:1 female to male sex ratio of the Samoan *H. bolina* population persisted for over 100 years (Dyson & Hurst, 2004). Only recently did the dynamics of this interaction change, with the host evolving resistance of the MK activity (Hornett et al., 2006; Charlat et al., 2007b).

Wolbachia are not the only endosymbionts that selectively kill male Lepidoptera. In the nymphalid butterfly, *D. chrysippus*, a *Spiroplasma* bacteria, related to a MK strain previously found in ladybirds, underlies the observed MK (Jiggins et al., 2000a). Similarly, while *Ostrinia* corn borer moths are especially well-known to harbour MK *Wolbachia* strains (i.e. the adzuki bean borer *Ostrinia scapularis* (Kageyama & Traut, 2004), and the Asian corn borer *O. furnacalis* (Sakamoto et al., 2007)), a MK *Spiroplasma* related to that found in *D. chrysippus* infects the butterfly borer *O. zaguliaevi* (Tabata et al., 2011). Mirroring the pattern seen in *H. bolina*, spatial variation of the MK *Spiroplasma* infection was observed in *D. chrysippus* (Smith et al., 1998; Herren et al., 2007), with 40% of females infected in Uganda vs. 4% in East Kenya (Jiggins et al., 2000a). Intriguingly, in this system infection appears to be correlated with a colour pattern allele. Although the forces generating this correlation are unknown, it may be the case that particular host genotypes are more susceptible to, or more efficient at transmitting, the infection than others (Herren et al., 2007).

In most study systems the precise mechanisms of MK are unclear, and variation across taxa is expected given that MK occurs in arthropods with widely disparate sex determination systems. Dependent on host context several mechanisms have been proposed including defective male chromatin remodelling (*Wolbachia*-infected *Drosophila* (Riparbelli et al., 2012)); targeting the dosage compensation complex (*Spiroplasma*-infected *Drosophila* (Veneti et al., 2005)); damaging the host's X chromosome to induce embryonic apoptosis (*Spiroplasma*-infected *Drosophila* (Harumoto et al., 2016)), and affecting maternally inherited centrosomes (*Arsenophonus*-infected *Nasonia* wasps: (Ferree et al., 2008)). In a *Wolbachia*-infected moth, *O. scapularis*, MK is unusual in that males (genotype ZZ) selectively die early in development,

whereas females (ZW) die if cured of the *Wolbachia* infection following antibiotic treatment (Fig. 2). Studies of this system suggest that MK *Wolbachia* interferes with the sex-specific splicing pattern of the *Ostrinia* homologue of the sex determination gene *doublesex*, *Osdsx* (Sugimoto et al., 2010), producing a mismatch between the genotypic sex and expression of the phenotypic sex and leading to sex-specific death (Sugimoto & Ishikawa, 2012). Later examination of the levels of dosage compensation (Z-linked gene expression) in male and female embryos destined to die, revealed that misdirection of dosage compensation underlies the observed mortality. Males destined to die (from *Wolbachia*-infected females) have higher levels of expression of Z-linked genes than normal; while females destined to die (from females cured of the *Wolbachia* infection) have lower expression levels of Z-linked genes than normal (Sugimoto et al., 2015). In a related moth, *O. furnacalis*, RNA-Seq data of *Wolbachia*-infected embryos demonstrated that MK *Wolbachia* down-regulated a masculinizing gene, *Masc*, essential in controlling both sex determination and dosage compensation in Lepidoptera, compared to uninfected embryos. The decrease in *Masc* mRNA levels is reported to cause the MK phenotype via a failure of dosage compensation, and injection of in vitro transcribed *Masc* cRNA into *Wolbachia*-infected embryos rescued male progeny (Fukui et al., 2015).

The mechanism of MK in *Ostrinia* moths may be different to that underlying MK in other Lepidoptera. In *H. bolina* butterflies no female specific death is observed following antibiotic treatment to reduce or remove MK *Wolbachia* (Charlat et al., 2007a). It is interesting to note however, that the *doublesex* homologue in *H. bolina* may be involved in MK in this butterfly as it resides within the chromosomal region defined as containing a suppressor of MK action (Hornett et al., 2014). It would therefore be interesting to compare the *Wolbachia* strains and MK mechanisms of *Ostrinia* and *H. bolina*. Likewise, a comparison between the modes of action of the MK strain of *Wolbachia* in *Ostrinia* moths, which kills males as a consequence of feminising them through alteration of expression of *Osdsx*, and of 'true' feminising *Wolbachia* such as that infecting *Eurema* butterflies (see *Feminisation*) may shed light on how one genus of bacteria can induce multiple reproductive manipulations in their hosts and whether there is a functional link.

Finally, the Oriental tea tortrix moth *Homona magnanima* also carries a MK (Morimoto et al., 2001), however male death in this case occurs much later in development (termed 'late MK'), and appears to be associated with two novel RNA sequences (Nakanishi et al., 2008). Late MK was originally only recorded in mosquitoes, with the causative agent being a microsporidian (Andreadis & Hall, 1979), however subsequent studies have now observed similar phenomena in other taxa including *Drosophila* flies (Jaenike, 2007). The extent of this type of manipulation, and the mechanisms underlying it, is still to be determined in insects, including Lepidoptera.

Feminisation

The feminisation of genetic males into functional phenotypic females (Stouthamer, Breeuwer & Hurst, 1999) is another strategy employed by maternally inherited




Figure 2 *Wolbachia*-induced male-killing and interference of sex determination in *Ostrinia scapularis* moths. (A) Uninfected females give rise to a normal 1:1 sex ratio in progeny: female offspring have ZW sex chromosomes and express the female isoform of the *Ostrinia* homologue of a gene in the sex determination cascade, *doublesex* (*dsx*), called Osd_{sx}^F ; male offspring have two Z sex chromosomes and express the male *dsx* isoform Osd_{sx}^M . (B) *Wolbachia* infected females only give rise to infected progeny. Male offspring die early in development due to a mismatch between the genotypic sex (ZZ) and phenotypic sex (Osd_{sx}^F). (C) *Wolbachia*-infected females cured of the infection as larvae by antibiotic treatment only give rise to uninfected males. Female offspring die early in development due to a mismatch between their genotypic sex (ZW) and phenotypic sex (Osd_{sx}^M). (D) *Wolbachia*-infected females cured of the infection as adults prior to oviposition by antibiotic treatment give rise to sexual mosaics which have the male ZZ genotype but both Osd_{sx}^F and Osd_{sx}^M . Note: there are two female isoforms of *dsx* in *Ostrinia scapularis*: Osd_{sx}^{FL} and Osd_{sx}^{FS} ; these are simplified to Osd_{sx}^F in this schematic. White circles: uninfected individual; Red circles: *Wolbachia*-infected individual; Dark grey circle, *Wolbachia*-infected female cured as larva; Light grey circle, *Wolbachia*-infected female cured as adult.

Full-size  DOI: 10.7717/peerj.4629/fig-2

endosymbionts to distort the host sex ratio towards the transmitting sex (females) (Fig. 1). While best known from the work on the association between *Wolbachia* and the terrestrial isopod *Armadillidium vulgare* (Juchault, Rigaud & Mocquard, 1992; Rigaud, Juchault & Mocquard, 1997; Bouchon, Rigaud & Juchault, 1998; Cordaux et al., 2004), feminisation also occurs in other female-heterogametic arthropods such as leafhoppers (XX/X0) (Negri et al., 2006), and Lepidoptera (ZZ/ZW). Other than *Wolbachia*, the Bacteroidetes bacterium *Cardinium* can also feminise males (Chigira & Miura, 2005; Groot & Breeuwer, 2006), however *Cardinium* has not yet been reported in butterflies or moths.

Observation of female-biased lines of pierid *Eurema* butterflies in Japan (Kato, 2000) led to the identification of a feminising *Wolbachia* in *Eurema mandarina* (formerly *E. hecabe* Y type) (Hiroki et al., 2002). *E. hecabe* (formerly *E. hecabe* B type) was later also discovered to carry a feminising *Wolbachia* indistinguishable from that of *E. mandarina*, thus suggesting that the infection transferred between the allopatric butterfly hosts via a shared predator or parasite, or via hybrid introgression between the species (Narita et al., 2011). When *E. mandarina* infected larvae were fed antibiotics to cure them of the infection, many of the adults emerged displaying sexually intermediate traits in their wings, reproductive organs and genitalia. Moreover, age at which antibiotics were administered was found to be important, with the highest level of intermediate sexual traits being exhibited when first instar larvae were treated. This

work demonstrated that endosymbionts might continually influence and interact with their host (Narita *et al.*, 2007) rather than have phenotypes that are effective only at a discrete time point in the lifecycle of the host.

The process of feminising in *E. mandarina* is more complex than originally thought. Against expectation, female butterflies infected with the feminising strain of *Wolbachia*, wFem, had only one, paternally derived, Z chromosome. This was proposed to be due to meiotic drive against the maternal Z, preventing the formation of the expected ZZ feminised males. It was also suggested that wFem lines have lost the W chromosome, and rely on wFem for female development as curing the infection with antibiotics results in all-male offspring (Kern *et al.*, 2015). Later work demonstrated that *Wolbachia* itself was responsible for the disruption of maternal Z chromosome inheritance in wFem infected females, as well as the feminisation of female ZO individuals that have lost the female-determining W chromosome (Kageyama *et al.*, 2017).

Cytoplasmic incompatibility

Perhaps the most commonly observed reproductive manipulation employed by endosymbionts in insects is CI. Unlike for MK or feminisation, the sex ratio of host populations infected by CI-inducing endosymbionts is generally not altered. Instead the symbiont induces an incompatibility upon mating between infected males and females of a different infection status (i.e. uninfected or infected with a different symbiont strain), leading to the death of all or a proportion of the offspring (Yen & Barr, 1971, 1973) (Fig. 1). This incompatibility is proposed to occur due to a modification of the infected male's sperm that can be rescued when the female is similarly infected (mod-res mechanism). This specific rescue function is lacking in uninfected females or females carrying a different infection (Hoffman & Turelli, 1997; Charlat, Calmet & Mercot, 2001; Poinso, Charlat & Mercot, 2003). Infected females therefore have a reproductive benefit of successfully producing a full complement of progeny (relative to uninfected females), when mated to uninfected or similarly infected males in the population. While *Wolbachia* is often described as the causative agent, *Cardinium* also has this ability (Hunter, Perlman & Kelly, 2003; Perlman, Kelly & Hunter, 2008).

Cytoplasmic incompatibility has been observed in a number of Lepidoptera including the Mediterranean flour moth *Ephesia kuehniella* and the almond moth *Cadra cautella* (Sasaki & Ishikawa, 1999). Interestingly, CI *Wolbachia* was discovered to lower the amount of fertile sperm transferred in *C. cautella* during second matings. However, no effect was shown on the amount of apyrene (non-fertile) sperm suggesting that *Wolbachia* may only target fertile sperm production (Lewis *et al.*, 2011). Further work is required to expand our knowledge of CI mechanisms in Lepidoptera. Similarly to MK, CI *Wolbachia* are often observed at high frequency in Lepidoptera populations. In a study of seven Japanese populations of the pierid butterfly *Colias erate poliographus*, CI *Wolbachia* occurred at 85–100% prevalence. The high infection frequency was ascribed to strong CI (i.e. a high proportion of the progeny from an incompatible cross die) and perfect vertical transmission of the bacteria (Tagami & Miura, 2004; Narita, Shimajiri & Nomura, 2009). Where CI reaches very high frequency within a host

population, the incompatibility between infected males and uninfected females is rarely observed as few females remain uninfected. However, selection for beneficial effects of infection and an eventual shift towards a mutualistic relationship between host and symbiont would remain.

Symbioses can be extraordinarily complex; individuals can carry multiple endosymbionts with differing phenotypes. For instance, *E. hecabe* butterflies carry a feminising *Wolbachia* strain, but also a second strain that causes CI (Hiroki *et al.*, 2004). This was the first indication that different strains of *Wolbachia* could infect a single individual and cause different phenotypes. Host context is important in the expression of endosymbiont-induced phenotypes—one symbiont strain can have the ability to cause more than one phenotype, including reproductive manipulations that were originally assumed to be distinct from each other. This has been exemplified in the butterfly *H. bolina*: in populations where *H. bolina* has evolved suppression of the action of MK, surviving infected males are incompatible with uninfected females in the population, i.e. expression of the CI phenotype (Hornett *et al.*, 2008). This finding indicates a potential functional or mechanistic link between the two phenotypes. However phenotypic switching between CI and MK through mutations cannot yet be ruled out. An intriguing possibility is whether feminisation is also mechanistically linked to CI and MK. Some evidence that may suggest this latter link is provided in studies of the moth *O. scapularis*. As mentioned above, male moths that die as a result of infection with MK *Wolbachia*, were found to carry the female isoform of a homologue of the sex-determining gene, *doublesex*, and hence were feminised prior to death (Sugimoto & Ishikawa, 2012).

Artificial transinfection of *Wolbachia* strains have provided further evidence of the relative importance of endosymbiont or host in determining the nature of the phenotype expressed. While in some cases transfer of *Wolbachia* from the natural host into a novel host did not alter the phenotype expressed (e.g. *Wolbachia* causes MK in the natural host *O. scapularis* and in the transinfected host *E. kuehniella* (Fujii *et al.*, 2001)), host context is important in others. Transfer of CI *Wolbachia* wCau-A from *C. cautella* to *E. kuehniella* resulted in the expression of MK in the novel host (Sasaki, Kubo & Ishikawa, 2002). The strength of the phenotype may also alter in the novel host: the level of CI induced by *Wolbachia* in the transinfected host *O. scapularis*, was higher than that in its natural host *E. kuehniella*, indicating that host factors as well as endosymbiont strain are important in determining the phenotype expressed (Sakamoto *et al.*, 2005).

Impact upon host fitness

It is becoming increasingly evident that many heritable endosymbionts do not manipulate host reproduction, and yet are still maintained within the host population. Host-associated microbes are now thought to be commonly beneficial to their host. For an inherited endosymbiont, the trade-off between virulence and transmission can lead to a reduction in its pathogenicity towards the host, and evolution towards mutualism (Weeks *et al.*, 2007). At the extreme end of the spectrum are the obligatory

endosymbionts, which are necessary for host survival or reproduction. The growing number of cases include: *Wolbachia* required for oogenesis in the wasp *Asobara tabida* (Dedeine et al., 2001; Dedeine, Bouletreau & Vavre, 2005); *Wigglesworthia* bacteria acting as an obligate nutritional mutualist in tsetse flies (Aksoy, 1995); and *Buchnera* bacteria providing essential nutrients to aphids (Buchner, 1965).

Many more endosymbionts are facultatively (non-essentially) beneficial, with fitness benefits including increasing host survival (Fry & Rand, 2002) or fecundity (Vavre, Girin & Bouletreau, 1999; Weeks & Stouthamer, 2004). Studies of the beneficial effects of endosymbiont infection in the Lepidoptera provide an unusual example in *Parnassius apollo*. In one isolated population this near threatened butterfly regularly exhibits deformed or reduced wings; however, while 86% of normal winged butterflies are found infected with *Wolbachia*, this percentage drops to 30% in individuals displaying deformed wings, and 0% in individuals with reduced wings. Although this is suggestive of a protective role of *Wolbachia* in the ontogenetic development of the butterfly, further study needs to be carried out to prove causality (Lukasiewicz, Sanak & Węgrzyn, 2016).

Microbial endosymbionts can contribute to insect adaptation by providing complementary or novel metabolic capacities, allowing the insect host to exploit host plant nutritional resources. One such instance has been observed in the phytophagous leaf-mining moth *Phyllonorycter blancardella*. In this system, a bacterial endosymbiont, most likely *Wolbachia*, indirectly affects larval nutrition by manipulating the physiology of the host plant to create photosynthetically active green patches in otherwise senescent yellow leaves. The phenotype, termed 'green-island,' produces areas of leaf viable for host feeding in a nutritionally constrained stage of the lifecycle. Curing the larvae of endosymbionts resulted in the non-production of 'green-islands,' and consequent increased compensatory larval feeding and higher mortality (Kaiser et al., 2010). The mechanism behind green island formation involves increased levels of cytokinins (CKs), plant hormones important in plant senescence and nutrient translocation. *Wolbachia* have been shown to be involved in the release of CKs by the larvae, creating these nutritionally enhanced areas of leaf. Whether the CKs are bacterial-derived or produced by the insect in response to *Wolbachia* infection (or a combination of both) remains to be fully understood (Body et al., 2013; Giron & Glevarec, 2014). Several strains of *Wolbachia* from both A- and B-supergroups have been identified in 13 Gracillariidae leaf-mining moth species, while none were found in ancestral Gracillariidae. Acquisition of the green-island phenotype appears to have occurred several times independently across the Gracillariidae in association with different *Wolbachia* infections (Gutzwiller et al., 2015).

Generally, vertically inherited endosymbionts are unlikely to be maintained in host populations if they are highly costly. However, direct fitness or physiological costs of infection have been observed where the symbiont also manipulates host reproduction. CI *Wolbachia* are maintained in the host population despite reducing male fertility (Snook et al., 2000) or female fecundity (Hoffman, Turelli & Harshman, 1990) in *Drosophila* flies, or detrimentally affecting fecundity, adult survival and locomotor

performance in the parasitoid wasp *Leptopilina heterotoma* (Fleury et al., 2000). Among Lepidoptera examples, presence of MK *Spiroplasma* in *D. chrysippus* in Kenya was negatively correlated with forewing length, suggesting that the bacteria may adversely affect development time or the growth rate of larvae (Herren et al., 2007). Presumably these physiological costs are counter-balanced by the reproductive manipulations employed by heritable endosymbionts, thus enabling the symbiont to persist. Although not covered in detail here, we note that in contrast, a symbiont that is also (or only) horizontally transmitted, can be highly detrimental to the host yet still be maintained in the host population. Indeed, host death may be its source of transmission to a novel host.

The gregarine protozoan infection, *Ophryocystis elektroscirrha*, of the Monarch butterfly (*D. plexippus*) is one of the most studied cases of direct fitness costs of symbionts in Lepidoptera. While it is not heritable in the sense of being intracellular, we include it here as it is passed vertically from mother to offspring via the surface of the egg. An infected female inadvertently coats her eggs with protozoan spores that cover the outside of her abdomen during oviposition. Newly hatched larvae ingest these spores while consuming the eggshell (McLaughlin & Myers, 1970). The parasite, which requires the adult host stage for transmission, rarely kills larvae or pupae under natural conditions, however the degree of virulence and transmission trade-off varies depending on the level of infection at the adult stage. Where individual *D. plexippus* butterflies carry high densities of the protozoa, they have both reduced survival and flight capacity compared to individuals with lower density infections (Altizer & Oberhauser, 1999; de Roode, Yates & Altizer, 2008; de Roode & Altizer, 2010).

Symbiont-mediated protection

Although understudied in Lepidopteran systems, an exciting avenue of research in arthropods revolves around a symbiont's ability to afford the host some level of resistance to its natural enemies, often through interference with pathogen or parasite replication or transmission (reviewed in insects in (Brownlie & Johnson, 2009)). This may be particularly the case for heritable endosymbionts, where symbiont and host fitness is inextricably linked—competing infections may elicit a response by the endosymbiont to protect the host, and thus simultaneously itself (Haine, 2008). Such symbiont-mediated protection has been documented in numerous taxa, particularly the Diptera, including recent studies demonstrating the ability of *Wolbachia* to supply their *Drosophila* host with anti-viral protection (Hedges et al., 2008; Teixeira, Ferreira & Ashburner, 2008; Martinez et al., 2014). Similarly, maternally transmitted *Spiroplasma* were found to protect *D. neotestacea* against the sterilising effects of a parasitic nematode (Jaenike et al., 2010), and enhance the survival of *D. hydei* parasitized by wasps (Xie, Vilchez & Mateos, 2010).

Symbiont-mediated protection appears to be extremely diverse. Aphids are host to a range of inherited symbionts, several of which provide protection against parasitoid wasp attacks (Oliver et al., 2003; Ferrari et al., 2004) or fungal infections (Ferrari et al., 2004; Scarborough, Ferrari & Godfray, 2005). In the European beewolf wasp, *Philanthus triangulum*, *Streptomyces* bacteria are stored in special antennae glands and deposited

together with the egg in the oviposition chamber. The bacteria secrete antibiotics protecting the developing wasp larvae against fungal pathogens ([Kaltenpoth et al., 2005](#)). There is even some evidence that symbionts can protect their host from predators by producing toxic compounds. For example, a bacterial endosymbiont (that is both vertically and horizontally transmitted) closely related to *Pseudomonas aeruginosa* produces the polyketide toxin pederin, which protects *Paederua* beetle larvae from predatory wolf spiders ([Kellner & Dettner, 1996](#); [Kellner, 1999](#); [Piel, Höfer & Hui, 2004](#)). Furthermore, endosymbionts may have the ability to inhibit a range of pathogens by priming the host immune system ([Braquart-Varnier et al., 2008](#); [Moreira et al., 2009](#); [Hughes et al., 2011](#)), suggesting that symbionts can interact with, and alter integral components of, host biology.

Conversely, it is important to acknowledge that endosymbiont infection can also increase pathogen load. While *Wolbachia* confers protection against a variety of pathogens and parasites in a wide range of hosts, pathogen or parasite levels can also be enhanced by the presence of an endosymbiont ([Hughes, Rivero & Rasgon, 2014](#)): in the moth *Spodoptera exempta*, *Wolbachia* triggers a higher rate of virus infection and therefore lowers host fitness ([Graham & Wilson, 2012](#)).

Host genetics

Host population genetics

Sex ratio distorting symbionts are likely to have a severe impact upon host population biology ([Engelstädter & Hurst, 2007](#)). If the prevalence of a sex-ratio distorter is high, the sex ratio of the population can become severely biased. In consequence, the hosts' effective population size (N_e) will be reduced. Where there is little gene flow into the population (i.e. low immigration), a reduction of the effective population size may affect the amount of standing genetic variation and the potential for the host population to respond and adapt to environmental change. In contrast, if gene flow does occur, spatial variation in sex ratio (as seen in the butterfly *H. bolina* ([Charlat et al., 2005](#); [Hornett et al., 2009](#))) may result in asymmetric gene flow between populations. Although both sexes typically contribute equally to the gene pool of the next generation, immigration of an individual into a population in which the sex ratio is skewed against it (e.g. a male into a highly female-biased region) can have a much larger genetic impact (i.e. contribute more) than if that individual immigrated into an unbiased sex-ratio population ([Telschow et al., 2006](#)). MK symbionts are also thought to hinder the spread of beneficial alleles and facilitate the spread of deleterious alleles, due to constrained gene flow from infected to uninfected individuals within the population ([Engelstädter & Hurst, 2007](#)). In a further complication, strains expressing different reproductive manipulations may be incompatible. Although most famous for its MK *Wolbachia* infections, some populations of *H. bolina* also carry a CI-inducing strain of *Wolbachia*. This CI strain is phylogenetically distant from the MK strain, and crosses between MK-infected females and CI-infected males are fully incompatible with no progeny surviving. The incompatibility produced has led to strong competition between the two strains, with the CI-strain being able to not only spread successfully through

uninfected populations, but to also resist invasion by the MK-strain carried by butterflies from neighbouring island populations (Charlat et al., 2006). Extending from this model, a study (Zug & Hammerstein, 2018) recently showed that when direct fitness benefits are taken into account in parallel to reproductive costs, the CI-strain is likely to also be able to spread across MK-infected *H. bolina* populations. Taken together, this suggests that successful establishment of particular butterfly genotypes is affected by the endosymbionts they harbour.

Linkage with host mitochondrial DNA

Maternally inherited symbionts residing within the cytoplasm of cells can alter the diversity and population genetics of the host's mitochondrial genome (mtDNA). Co-inherited symbionts and mitochondria are in linkage disequilibrium, therefore when a cytoplasmic symbiont invades a population, the initially associated mitochondrial haplotype (mitotype) may 'hitch-hike' and correspondingly increase in frequency. Should such a selective sweep have occurred recently, the effective population size and genetic diversity of mtDNA would be reduced to that of the infected individuals (Johnstone & Hurst, 1996), and the geographic structure of mitochondrial variation lost. The latter has been observed in *Acraea* butterflies (Jiggins, 2003) and the comma butterfly *Polygonia c-album* (Kodandaramaiah et al., 2011). The tight association between endosymbiont and mtDNA can therefore seriously confound the results of any study using mtDNA genes as neutral genetic markers (Hurst & Jiggins, 2005). Reconstruction of phylogenetic trees using mitochondrial markers are hence likely to be misleading, particularly within shallower branches, when the study species is infected. In the Diamondback moth, *Plutella xylostella*, the main correlate of mtDNA variation is presence or absence of the *plutWb1* *Wolbachia* infection (Delgado & Cook, 2009), and the lycaenid butterfly *Lampides boeticus* may have experienced accelerated population differentiation due to *Wolbachia* infection (Lohman et al., 2008). Recognition of these processes should lead to an increasing number of Lepidopteran studies interested in using mtDNA markers to systematically screen for maternally inherited symbionts.

Where there is perfect transmission of the maternally inherited symbiont from the host to its offspring, infected individuals all carry the same mitotype, while uninfected individuals remain polymorphic. This pattern has been repeatedly observed in natural populations of insects, including in the Lepidoptera. In the butterfly *H. bolina*, a strong association between one specific *Wolbachia* strain and one particular mitotype supported the hypothesis that the MK infection occurred with very high vertical transmission efficiency and rare horizontal transmission. In *H. bolina*, this strain of *Wolbachia* is thought to have undergone a recent selective sweep and was introduced into this butterfly through introgression, potentially from another *Hypolimnias* species, *H. alimena*. Conversely the infection and associated mitotype may have been introgressed from *H. bolina* to *H. alimena* (Charlat et al., 2009; Duploux et al., 2010; Sahoo et al., 2018). Similarly, in the *Acraea* butterflies, a study of mitochondrial variants demonstrated that a MK *Wolbachia*, together with the associated mitotype, had introgressed from *A. encedana* into *A. encedon* within the last 16,000 years. As female

butterflies are heterogametic (ZW), this event could potentially also lead to the introgression of genes on the female W chromosome (Jiggins, 2003). This scenario appears to have occurred in *D. chrysippus* infected with a MK *Spiroplasma*, as all infected females carry the same W chromosome variant (Smith, Gordon & Allen, 2010). This aside, the nuclear DNA is generally less likely to be in linkage with inherited symbionts. Gompert *et al.* (2008) studying North American *Lycaeides* butterflies reported that the spread of an endosymbiont (and associated mitotype) through the host population produced substantial mito-nuclear discordance. Therefore, the evolutionary history of an individual's nuclear and mitochondrial genomes may be very different from each other. Such discordance may have far-reaching effects on host metabolism and physiology, as coevolution between nuclear and mitochondrial components of essential pathways is broken down.

Speciation by symbiosis

The concept that symbionts can be important promoters of speciation and diversity has been around for a long time (Wallin, 1927; Laven, 1959; Thompson, 1987; Breeuwer & Werren, 1990; Hurst & Schilthuizen, 1998; Bordenstein, 2003), but has recently been rejuvenated with the development of microbiome analyses (Brucker & Bordenstein, 2012). Contemporary evidence of microbe-assisted speciation involves pre-mating reproductive isolation through behavioural barriers such as mate preference, associated with the microbiome of the potential partners (Koukou *et al.*, 2006; Miller, Ehrman & Schneider, 2010; Sharon *et al.*, 2010; Chafee *et al.*, 2011). The underlying mechanisms may involve alteration of the sex pheromones, interference with sensory organs, or effects upon immune-competence and hence mate attractiveness. Ecological isolation may also be heavily influenced by microbial symbionts. Although the genetic basis of niche or habitat specificity is widely accepted, there is also increasing evidence that symbionts may play a role in determining host resource availability (Akman *et al.*, 2002; Hosokawa *et al.*, 2010), and thus may facilitate niche separation.

Additionally, endosymbionts might enable host speciation through post-mating isolation. In particular, strong bi-directional CI may result in reproductive isolation between hosts carrying different CI symbiont strains (Hurst & Schilthuizen, 1998; Werren, 1998; Bordenstein, 2003). In order for speciation to follow CI, a stable infection polymorphism must be maintained across host populations. This has been demonstrated in many systems including the butterfly *H. bolina* (Charlat *et al.*, 2006). Theory predicts that two bi-directional CI-inducing symbionts can be stable for even high migration rates (Telschow, Hammerstein & Werren, 2005). What is more contentious is that for speciation to occur, the CI produced must be very strong (i.e. no offspring surviving from such crosses), and the symbiont must be maintained at a high transmission rate over time, to allow significant nuclear divergence (Engelstädter & Hurst, 2009).

Male-killing has also been linked to speciation in the butterfly *D. chrysippus*. In Kenya two forms exist: *D. c. chrysippus* and *D. c. dorippus*, separated by a hybrid zone. Each subspecies has an individual colour pattern controlled by locus *C*, which is intermediate in

the hybrid (*Cc*). The *C* locus lies on an autosome that has fused with the *W* chromosome within the hybrid zone, physically linking colour pattern with female determination. A locus on this same autosome has also been associated with susceptibility to MK by *Spiroplasma*. The hybrid zone is characterised by female-biased sex ratios, caused by MK *Spiroplasma* that infects *D. c. chrysippus* or hybrid females, but rarely *D. c. dorippus* females. As immigrant males into the hybrid zone are predominantly *D. c. dorippus*, gene flow between the two subspecies is restricted: *D. c. chrysippus*/hybrid female × *D. c. dorippus* male crosses produce female-biased broods (Smith et al., 2016).

Sex determination

The maternal inheritance of intracellular endosymbionts has led to a great degree of interaction of the symbiont with the sex determination pathways of the host (reviewed in (Cordaux, Bouchon & Grève, 2011; Kageyama, Narita & Watanabe, 2012; Ma, Vavre & Beukeboom, 2014) and so not discussed in detail here). Maternally inherited endosymbionts distort the host sex ratio in order to enhance the fitness of the transmitting female sex. The mechanisms underlying these phenotypes often require considerable manipulation of host sex determination. We have seen above that in several cases MK and feminising *Wolbachia* can interfere with central components of the sex determination pathways in Lepidoptera. When a feminising element is highly prevalent in a host population, sex determination may be inextricably linked to the presence or absence of feminising activity (Hiroki et al., 2002), but may also enter into conflict with other genetic elements not under similar maternal inheritance. Furthermore, evolution of host suppressors of feminisation may move the system away from the original ZZ/ZW sex determination system. In *E. mandarina*, *Wolbachia* disrupts the inheritance of maternal Z chromosomes in *Wolbachia*-infected females, and feminises the resulting Z0 individuals that have lost the female-determining W chromosome (Kageyama et al., 2017). The host may then be prompted to evolve a strategy to counteract the feminising effects of the symbiont. It has been speculated that in the pillbug *A. vulgare*, a masculinising factor in the form of a dominant autosomal *M* gene has evolved in the host to counter the effect of the feminising endosymbiont (Rigaud & Juchault, 1993; Caubet et al., 2000).

Evolution of host resistance

Co-evolution between a host and a detrimental symbiont can result in the evolution of host genetic modifiers of symbiont presence or action. Despite this, and considering the wide array of costly effects that endosymbionts can impose on their hosts, it is perhaps surprising that there are relatively few well documented examples of the host having evolved genetic resistance to an endosymbiont. Indeed no suppression of the detrimental phenotype is observed in several studies where it may have been expected (Hurst, Jiggins & Robinson, 2001; Veneti, Toda & Hurst, 2004; Dyer & Jaenike, 2005). However, artificial transinfection experiments have provided an indirect method of discovering whether a host has evolved resistance to an endosymbiont, and have suggested that suppression of reproductive manipulation phenotypes may actually be common.

In the moth *C. cautella*, which is naturally infected with two *Wolbachia* strains (*wCauA* and *wCauB*), artificial transinfection of CI-inducing *wCauA* to a sister host species, *E. kuehniella*, resulted in the transferred bacteria inducing MK instead of CI in the novel host (Sasaki, Kubo & Ishikawa, 2002; Sasaki, Massaki & Kubo, 2005). By interpreting these data in the light of the hidden MK theory (where MK is masked by the presence of a fixed suppressor), this switch in phenotype between species could be interpreted as the ‘unmasking’ of MK when released into a background devoid of host suppression genes. More generally, resistance may also underlie the loss of infections from populations or host species, however this is obviously hard to document in nature.

The selective pressure for host resistance is particularly strong when the sex ratio is severely biased (Düsing, 1884; Fisher, 1930; Hamilton, 1967), and therefore one would expect the evolution of resistance particularly in cases of highly prevalent sex-ratio distorters. As mentioned above, the Samoan population of the butterfly *H. bolina* had an extraordinarily female-biased sex ratio of 100 females per male, caused by 99% of the females being infected with MK *Wolbachia* (Dyson & Hurst, 2004). However between 2001 and 2006 the dynamics of the interaction changed dramatically when *H. bolina* evolved suppression of the MK trait, allowing infected males to survive and rapidly re-establishing a 1:1 sex ratio within approximately 10 generations of the host butterfly (Charlat et al., 2007b). The presence of a zygotically acting dominant suppressor locus had previously been documented in SE Asian *H. bolina* populations (Hornett et al., 2006).

Sex ratio distorting endosymbionts can also have much wider implications upon host genetics. Recent work on the same Samoan population of *H. bolina* investigating the genomic impact of the rapid spread of suppression revealed that a substantial selective sweep had taken place, covering at least 25 cM of the chromosome carrying the suppressor locus. In addition to large changes in the frequency of genetic variants across this broad region, the sweep was associated with the appearance of several novel alleles. This suggests that the suppressor spread following migration of butterflies carrying the locus, potentially from SE Asia, rather than from a de novo mutation occurring within the population. It is also interesting to note that the suppressor of MK has been located to the chromosome containing *doublesex* (Hornett et al., 2014)—a sex determination gene demonstrated to be involved in *Wolbachia*-induced MK in *Ostrinia* moths.

Horizontal transfer of genetic material

While the extent of horizontal (lateral) gene transfer (HGT) between eukaryotes and prokaryotes remains uncertain, technological advances in genomics followed by an accumulation of microbial and host genomic data, have revealed that endosymbionts, particularly those that are vertically inherited, may readily exchange genetic material with their host. HGT from a prokaryote symbiont to its eukaryote host has been reported in many insects including beetles, flies, parasitoid wasps, mosquitoes and butterflies (Hotopp et al., 2007; Nikoh et al., 2008; Klasson et al., 2009; Werren et al., 2010) and has recently been reviewed in detail (Husnik & McCutcheon, 2018). Such movement of genes can afford the receiving organism important benefits. For instance, horizontally transferred

bacterial DNA that is involved in the detoxification of cyanide has been identified in several moths and butterflies, allowing these insects to utilise otherwise noxious plants ([Wybouw et al., 2014](#)). However, the discovery of bacterial DNA within the host's genome does not necessarily imply functionality, and definitive proof of function is difficult to obtain, indeed many transferred *Wolbachia* genes are not expressed at a significant level in the host ([Hotopp et al., 2007](#); [Nikoh et al., 2008](#)). To date, the identification of a 350 bp long *Wolbachia* gene insert in the genome of the butterfly *Melitaea cinxia*, is the only reported example of an HGT from an endosymbiont to a Lepidoptera species ([Ahmed, Breinholt & Kawahara, 2016](#)), its origin and functionality have yet to be demonstrated.

Horizontal gene transfer is also known to occur in the opposite direction, from eukaryote host to symbiont. *Wolbachia* genome projects have indicated that genome fragments have been transferred from host to the bacteria, including in the *H. bolina* system. The MK *Wolbachia* strain sequenced appears to be extremely receptive to exogenous genetic material ([Duploux et al., 2013](#)). In addition to cross-level transfer of genes, bacteria within a host may also exchange genetic material. Bacteria are known to be promiscuous with regard to DNA, with movement of bacteriophages between co-infecting symbiont species providing a convenient method of transfer of genes. Some endosymbiont traits are associated with phage presence ([Oliver et al., 2009](#)) and thus this movement offers the potential for transfer of traits between co-infecting symbiont strains ([Duron & Hurst, 2013](#)). Indeed, extensive HGT involving the bacteriophage WO has been reported between several *Wolbachia* strains infecting diverse hosts including within the Lepidoptera, Diptera and Hymenoptera ([Masui et al., 2000](#); [Bordenstein & Wernegreen, 2004](#)).

Behavioural modification

The transmission of many parasites is facilitated by their ability to manipulate the behaviour of their hosts ([Lefevre et al., 2009](#)). Reported cases are often restricted to viral and fungal pathogens; for instance, some baculoviruses and fungi cause summit disease—a syndrome that induce caterpillars to climb to high vegetation prior to being killed so that any spores released are carried further by the wind ([Maitland, 1994](#); [Yamazaki, Sugiura & Fukasawa, 2004](#)). Behavioural modification of arthropod hosts by heritable endosymbionts is less evident, and where observed are perhaps more attributable to indirect effects of infection. *Rickettsia* bacteria have been associated with limiting long distance dispersal in a spider ([Goodacre et al., 2009](#)), and *Wolbachia* has been demonstrated to reduce wasp locomotor performance ([Fleury et al., 2000](#)). Models of MK endosymbionts in metapopulations have suggested that MKs can increase host dispersal rates ([Bonte, Hovestadt & Poethke, 2008](#)). These patterns may be attributed to the evolution of adaptive modifications by the symbiont to promote its own transmission. However another explanation is that these behavioural changes are merely side effects of physiological alterations without any adaptive causality.

In the butterfly, *D. plexippus*, the protozoan *O. elektroscirrha* has attracted much attention because of its potential involvement in the famous migratory behaviour of its host. This parasite is known to reduce the flight capacity of the host

(Altizer & Oberhauser, 1999; Bradley & Altizer, 2005)—a trait that creates an important trade-off as the butterflies' dispersive behaviour allows the spread of the protozoa across the species range, and thus increases the chance of it infecting naive populations. For the butterfly, migration offers an opportunity of escaping highly infected habitats where they may risk reduced fitness (Altizer, Bartel & Han, 2011). Altizer, Oberhauser & Brower (2000) demonstrated that variation in protozoa prevalence correlates with host movement—non-migratory populations have high infection prevalence whereas populations that migrate long distances show less than 10% prevalence of infection. More recently it was found that where migratory behaviour has been lost, the risk of infection is increased (Satterfield, Maerz & Altizer, 2015). Thus in part the presence of the protozoa may have led to Monarch butterflies forming both resident and migratory populations.

Further indirect behavioural consequences of microbial infection are also possible. In order to escape the fecundity and physiological costs of mating with an incompatible mate, individuals may evolve new adaptive mating strategies, including increased polyandry or mate discrimination (reviewed in (Miller & Schneider, 2012)). *Wolbachia* influences mate-choice in the two-spotted spider mite, where uninfected females preferentially mate with uninfected males (Vala *et al.*, 2004), while in *Drosophila paulistorum*, *Wolbachia* titer and mate discrimination are positively correlated (Miller, Ehrman & Schneider, 2010). In *Acraea* butterfly populations harbouring high frequency MK bacteria (thereby having highly female-biased sex ratios), infected females more often remained unmated than uninfected females (Jiggins, Hurst & Majerus, 2000). While this is suggestive of preferential mating by the male, further work needs to be carried out to test this. However, *Acraea* butterflies afford another example: in butterflies, males are often the competing sex and court the females. When the butterfly population is strongly female-biased due to the presence of a highly prevalent sex ratio distorting endosymbiont, the roles of the sexes may reverse. Such sex-role reversal was observed in *Acraea* butterflies infected with MK *Wolbachia*. Although male 'hill-topping' (swarming at the tops of hills) is common throughout the genus (Jiggins, 2002), in *A. encedon* the lack of males induced females to swarm instead, and to exhibit behaviours soliciting the males' attention (Jiggins, Hurst & Majerus, 2000).

Male-killing endosymbionts may also result in female reproduction becoming sperm limited. In a comparison of *H. bolina* populations varying in MK *Wolbachia* prevalence, the prediction that female mating rates would decline with increasing MK infection prevalence as males became increasingly rare was not borne out. Unexpectedly the opposite occurs—as the population sex-ratio becomes more biased, the female mating rate increased until a point at which the lack of males makes it impossible for females to find a mate. It was suggested that female promiscuity increased in response to increasing male 'fatigue.' Males from more highly female-biased populations produced smaller spermatophores thus necessitating females to become more solicitous (Charlat *et al.*, 2007c).

Outstanding questions and future directions

We here are promoting the Lepidoptera as important models in the study of endosymbiont induced reproductive manipulations, with MK, feminisation and CI all

being evident in butterflies and moths. Current research is uncovering the genetic and functional basis underlying these phenotypes but many outstanding questions remain: Are all three reproductive manipulations found in Lepidoptera functionally linked? How commonly can a single endosymbiont strain confer more than one phenotype? How do different endosymbiont genera confer similar phenotypes in their host (e.g. both *Wolbachia* and *Spiroplasma* cause MK in Lepidoptera), and are the mechanisms related? How does MK, feminisation and CI in Lepidoptera differ from that expressed in taxa with divergent sex determination systems? Also, how do sex-ratio distorting endosymbionts affect the long-term evolution of the host. Given recent advances in genomics this now can include investigations of the genomic impact of a sustained population sex-ratio bias. Sex-linked traits in particular may be expected to be affected.

More questions are provoked when research into heritable endosymbionts associated with other arthropod taxa is considered. Of particular interest is the evidence accruing that symbionts often afford the host some level of protection against pathogens and parasitoids. But how frequent is this phenomenon in butterflies and moths? Also, can we see these effects in combination with reproductive manipulations, producing a trade-off between detrimental and mutualistic effects of infection? Conversely, where we see highly prevalent and persistent endosymbiont infections in host populations that do not induce reproductive manipulations, do these symbionts offer the host protection? While there are clearly many outstanding questions to examine in the Lepidoptera, in this next section we focus upon four further areas of research that will move Lepidoptera-heritable endosymbiont research forward.

Comparative endosymbiont genomics

The genomes of many arthropod heritable endosymbionts have now been assembled, however very few of those sequenced are associated with Lepidoptera hosts. A comparative genomics approach can be used to elucidate endosymbiont evolution and function in its host including identifying candidate genes involved in reproductive manipulations such as CI (as in *Drosophila* ([Sutton et al., 2014](#); [LePage et al., 2017](#))), and parthenogenesis induction (in parasitoid wasps ([Newton et al., 2016](#); [Lindsey et al., 2016](#))). A recent comparison of 16 *Wolbachia* genomes identified a core *Wolbachia* genome of 496 sets of orthologous genes, 14 of which were unique to *Wolbachia* among the Rickettsiales bacteria, of which it is a member ([Lindsey et al., 2016](#)). This study included the MK *Wolbachia* strain *wBol1b* from *H. bolina* butterflies, which was revealed to be closely related to a CI *Wolbachia* infecting *Culex pipiens* mosquitoes, *wPip*. A comparison of the two strains identified a number of genes specific to *wBol1b* that could be potential candidates involved in the induction of MK ([Duploux et al., 2013](#)). An interesting future research direction that may inform on the diversity and genetic basis of MK, would be to expand this line of enquiry by comparing the genome of *wBol1b* with other MK and non-MK *Wolbachia* genomes. Candidate loci could also be investigated in other MK-inducing symbiont genomes such as *Spiroplasma*. While the genomes of many *Spiroplasma* bacteria have been characterised from various arthropods (see ([Bolaños, Servín-Garcidueñas & Martínez-Romero, 2015](#)) for a minireview), including

the MK *Spiroplasma* endosymbiont MSRO found in *D. melanogaster* (Paredes et al., 2015), to our knowledge none have as yet been published that specifically associate with Lepidoptera.

As high-throughput sequencing costs reduce, the genomes of increasing numbers of Lepidoptera are being sequenced (for a review of the current status see (Triant, Cinel & Kawahara, 2018)). A happy indirect consequence of this is that endosymbiont genome sequences can be retrieved as a by-product of host genome sequencing. This is a particularly useful tool when studying intracellular endosymbionts that are not readily culturable, and hence difficult to directly isolate and sequence (such as *Wolbachia*). This approach has been used to reconstruct the genome of *Wolbachia* infecting the moth *Operophtera brumata* (Derks et al., 2015), and that of *Wolbachia*, wAus, associated with the moth *P. australiana* (Ward & Baxter, 2017). Interestingly, and similarly to wBol1b from *H. bolina*, both strains were most closely related to the CI *Wolbachia* wPip from the mosquito *C. pipiens* (Derks et al., 2015; Ward & Baxter, 2017); however, in the case of wAus, two genes previously determined to be involved in CI caused by *Wolbachia* from *D. melanogaster* were not found in the genome of wAus (Ward & Baxter, 2017). Further work needs to be conducted to characterise the nature of the interaction between *Wolbachia* and host before more insight can be gained through genomic comparisons.

What else is in there? Moving towards a metagenomics approach

This review has revealed a marked bias in Lepidopteran heritable endosymbiont research—*Wolbachia* is by far the most studied endosymbiont in butterflies and moths. While the incidence of *Wolbachia* is undoubtedly high in Lepidoptera and its effects upon its hosts important, the development of routine PCR assays and resources specific to this one genus of bacteria may have inflated its significance relative to other endosymbionts. Thus a practical limitation of the current methodology in the study of heritable endosymbionts in Lepidoptera is the lack of an unbiased approach to determine what microbes butterflies and moths carry. This is changing with the development of culture-independent methods of ascertaining what microbes, particularly bacteria, are present within an organism. High-throughput sequencing of the hypervariable bacterial 16S rRNA gene, and metagenomics allow the characterisation of whole bacterial communities of hosts. Particular to heritable endosymbiont research, attention should be given to the tissue from which DNA is sourced, as heritable bacteria are not necessarily found in the commonly sequenced gut tissue or lumen. Amplifying bacterial DNA from whole insects or the reproductive tracts may yield a clearer idea of the vertically inherited symbionts present. We also have to consider what constitutes a heritable endosymbiont; many Lepidopteran gut bacteria are transitory and/or environmentally acquired for example via the food plant as larvae (Mason & Raffa, 2014; Hammer et al., 2017), or nectar as adults and as such may not evolve symbiotically with the host. However, gut bacteria may be transmitted by the female to the progeny via for example the egg coating, which neonates often consume upon hatching. One challenge will be to distinguish which of the microbes present in a community are symbiotic, and further, which are vertically

transferred. Therefore close behind microbiome characterisations of Lepidoptera will be experimental manipulations of the microbiome and the sequencing of progeny to ascertain heritability.

Revealing the microbiome of Lepidoptera will open up a new set of questions such as do gut microbes and heritable endosymbionts interact? Can endosymbionts affect the composition of the microbiome? Do their effects interact? One promising avenue of research is the antimicrobial activity of gut symbionts. The moth *S. littoralis* harbours a gut bacterium *Enterococcus mundtii* that secretes an antimicrobial peptide (mundticin KS) against invading bacteria, but not against other resident gut bacteria. This antimicrobial activity directly inhibits competitors, but also potential pathogens, from the gut of its host. In *S. littoralis*, this extracellular symbiont persists across host developmental stages and is a major constituent of the microbiome across generations, suggesting that it can be vertically inherited, and that it may form a long-term symbiotic association with its host (Shao et al., 2017).

A further avenue for future research is the presence and impact of non-bacterial heritable endosymbionts. In particular there is increasing recognition that viruses may be vertically inherited and can have dynamic interactions with their host (reviewed in insects in (Longdon & Jiggins, 2012)). The moth *Helicoverpa armigera*, a crop pest, is infected with a vertically (and horizontally) inherited densovirus (HaDENV-1) that appears to be mutualistic. In wild larvae a negative interaction exists between the symbiotic densovirus and the presence of a nucleopolyhedrovirus (HaNPV) that is widely used as a pesticide against *H. armigera*. Laboratory work confirmed that larvae carrying HaDENV-1 had significantly higher resistance to the HaNPV pesticide, and also to low doses of *Bacillus thuringiensis* toxin. Additionally, HaDENV-1 infected individuals have a higher developmental rate and higher fecundity than that of their uninfected counterparts (Xu et al., 2014). In contrast, in the moth *H. magnanima* a novel RNA virus appears to be responsible for 'late' MK while being benign to female moths, thus acting as a reproductive manipulator (Nakanishi et al., 2008). Metagenomic sequencing has identified viruses across diverse arthropods (Li et al., 2015), and while often pathogenic a recent study identified a vertically inherited sigma virus in the nymphalid butterfly *Pararge aegeria*, that may have a more symbiotic role. In this species transmission of the virus was predominantly maternal (through eggs), with paternal (through sperm) transmission rates being much lower. Wild populations of *P. aegeria* experience high levels of infection, with a mean viral prevalence of 74%, and marked population structure in the genetic diversity of the virus (PAegRV). The nature of the relationship between *P. aegeria* and PAegRV remains to be determined (Longdon et al., 2017).

Global environmental change: can endosymbionts facilitate or constrain adaptation?

Predicting if or how organisms adapt to environmental change is a critical and timely question. Every organism interacts with a multitude of abiotic and biotic factors, including heritable endosymbionts, and knowledge of how these influence each other is imperative in understanding an organism's adaptive potential. Global environmental

change is likely to alter the level and direction of natural selection in host/symbiont co-evolution (*Wolinska & King, 2009*). In one direction, endosymbionts may increase the host's potential repertoire for responding to environmental changes such as temperature, while we also recognise that the destabilisation of often finely tuned host-symbiont interactions may be severely detrimental for natural populations.

As poikilotherms—organisms that do not maintain internal thermal homeostasis—butterflies and moths are very susceptible to extreme temperatures (*Denlinger & Yocum, 1998*). While they utilise a range of mechanisms, including behavioural and physiological responses, to regulate temperature, every species is defined by thermal limits. Recent work has indicated that microbial symbionts of insects can often facilitate or constrain adaptation to environmental changes, including temperature. For instance, aphids carry symbionts that proffer heat stress protection (*Montllor, Maxmen & Purcell, 2002; Russell & Moran, 2006*), including a point mutation (a change in a single nucleotide), which governs host thermal tolerance (*Dunbar et al., 2007*). The temperature insects are exposed to during development is also important in the maintenance of symbionts (*Anbutsu, Goto & Fukatsu, 2008*), or to the phenotype expressed by the symbiont in the host (*Hurst et al., 2000*). With global environmental change, de-stabilisation of the host-symbiont interaction may become more frequent and have severe consequences for many species. The sudden loss of an obligatory mutualistic symbiont, for example, would almost certainly lead to a host population decline (for further discussion of host-symbiont interactions and temperature see (*Wernegreen, 2012; Corbin et al., 2016; Moran, 2016*)).

Furthermore, the nature of the relationship between host and symbiont may be indirectly affected by the changing climate. A few degrees rise in temperature can alter the geographic range of Lepidoptera (*Parmesan et al., 1999*). For many species, such range shifts and colonisation events should only be possible if the plants they utilise were following a similar expansion, such as in the host-limited butterfly *Gonepteryx rhamni* (*Gutiérrez & Thomas, 2000*). Additionally, range shifts may lead to a switch in host plant species or increased generalisation (*Braschler & Hill, 2007*), bringing subsequent repercussions for Lepidopteran-endosymbiont interactions. For example, in the moth *P. blaucardella*, where endosymbionts nutritionally benefit the host by creating photosynthetically active green patches in otherwise senescent leaves of the host plant (*Kaiser et al., 2010*), a shift in host plant use could make this 'green-island' strategy ineffective in a novel plant with a different chemical makeup. In contrast, novel host plant utilisation may also be facilitated by endosymbionts, including through enhanced provisioning of nutrients, or detoxification (reviewed in (*Hansen & Moran, 2014*)).

Finally, habitat degradation and fragmentation is likely to have several implications for natural host-symbiont dynamics. Habitat destruction has the effect of crowding insect populations into smaller patches, and through fragmentation and subsequent isolation, the amount of gene flow between populations becomes reduced. These factors may increase disease transmission within a population, and alter geographical variance in endosymbiont presence and prevalence.

Screening butterflies and moths of conservation concern for endosymbionts

The Lepidoptera are model organisms in the fields of conservation and climate change research. However, despite the high occurrence of endosymbionts in Lepidoptera, current conservation planning rarely includes data on endosymbiont infections of the species under consideration, a deficit that may profoundly influence the outcome of any management undertaken. For effective conservation, or to understand how species will respond and adapt to environmental and anthropogenic changes, it is important that we try to understand the intricate relationships that microbes have with the hosts in which they reside. Fortunately there is increasing recognition of this importance with several recent studies reporting endosymbiont infections in populations of endangered or near threatened Lepidoptera ([Nice et al., 2009](#); [Sakamoto et al., 2011](#); [Patricelli et al., 2013](#); [McHugh et al., 2013](#); [Lukasiewicz, Sanak & Węgrzyn, 2016](#); [Fenner et al., 2017](#)). One study surveying 22 species of conservation concern (comprising members of the Lycaenidae, Nymphalidae, Hesperidae and Noctuidae) for *Wolbachia* found 19 to be infected ([Hamm et al., 2014](#)). [Nice et al. \(2009\)](#) examined the nature of a *Wolbachia* infection in the North American endangered Karner blue butterfly, *Lycaeides melissa samuelis*. Screening for endosymbionts revealed that across the western edge of this butterfly's range there was a widespread *Wolbachia* infection. They went on to simulate demographic effects of the spread of *Wolbachia* into uninfected populations and suggested that the spread of such an infection might further reduce already small population sizes. The authors show concern that the *Wolbachia* infection was prevalent in many of the largest and least impacted populations of this butterfly. This is significant as these populations are likely candidates from whom captive propagation efforts would draw individuals, and so the chance of inadvertently infecting a naturally uninfected population is high.

Release of wild individuals or of those reared in captivity, either as part of conservation management schemes or for commercial purposes (birthdays or weddings), might have unexpected and undesirable impacts if not monitored correctly. Rearing Lepidoptera, which often occurs at high densities, can allow the accumulation of pathogens. Releasing these individuals back into the field may therefore alter the parasite load and consequent fitness of the receiving population. Movement of individuals between populations may also affect the natural spatial pattern of endosymbiont diversity and prevalence: novel microbes may be introduced, or symbionts that have locally adapted in the donor population may affect the host in dramatically different ways in the novel population or environment. Further consequences with regard to endosymbiont infection are likely to be numerous, for example competition between native and novel infections may result in a shift in the natural equilibrium between the host and its native microbes or the introduction of cytoplasmic endosymbionts may also introduce linked variants such as host mtDNA haplotypes or female-linked nuclear DNA. Furthermore, as we have seen in the butterfly *H. bolina*, movement of individuals could also introduce host resistance loci that irrevocably alter the dynamics of host-symbiont interaction, and may have a

wider impact upon the host genome. In general, if a novel association does form and/or spread, there follows rapid evolution of both host and symbiont, with phenotypic alterations that alter or optimise the new symbiosis (for an example see *(Weeks et al., 2007)*).

CONCLUSION

The Lepidoptera have emerged as important models in the study of the genetic and functional basis of the reproductive manipulations heritable endosymbionts employ, particularly with regard to *Wolbachia* bacteria. The results of this cumulative work is suggestive of the role of endosymbionts in the evolution of host sex determination itself. We have no doubt Lepidopteran endosymbiont research will continue to highlight the omnipresence and importance of *Wolbachia* but we suggest that more attention should now be given to the presence and interaction of other heritable endosymbionts Lepidoptera carry. Metagenomic approaches enable an unbiased view of the microbial community residing within moths and butterflies, while comparative endosymbiont genomics may illuminate the genetic mechanisms underlying the phenotypes endosymbionts induce in their host. Finally, given the importance of Lepidoptera as key indicators of climate change and the growing numbers of species listed as endangered, the study of heritable microbial endosymbiont in the Lepidoptera should transition from being a pure science filled with interesting curiosities, to a necessity that will contribute to the preservation of natural biodiversity and inform conservation management.

ACKNOWLEDGEMENTS

We wish to thank S. Charlat, J. Marden, A. Teacher, M. Riegler, H. Kokko, G. Hurst and an anonymous reviewer for constructive comments on the manuscript.

ADDITIONAL INFORMATION AND DECLARATIONS

Funding

This work was supported by an Academy of Finland postdoctoral grant #266021 to Anne Duploux, and a Marie Curie Actions IO Fellowship no. 330136 to Emily A. Hornett. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Grant Disclosures

The following grant information was disclosed by the authors:

Academy of Finland postdoctoral: 266021.

Marie Curie Actions IO Fellowship: 330136.

Competing Interests

The authors declare that they have no competing interests.

Author Contributions

- Anne Duploux conceived and designed the experiments, analysed the data, contributed reagents/materials/analysis tools, prepared figures and/or tables, authored or reviewed drafts of the paper, approved the final draft.
- Emily A. Hornett conceived and designed the experiments, analysed the data, contributed reagents/materials/analysis tools, prepared figures and/or tables, authored or reviewed drafts of the paper, approved the final draft.

Data Availability

The following information was supplied regarding data availability:

The research in this article did not generate any data or code; this is a literature review.

Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.4629#supplemental-information>.

REFERENCES

- Ahmed MZ, Araujo-Jnr EV, Welch JJ, Kawahara AY. 2015. *Wolbachia* in butterflies and moths: geographic structure in infection frequency. *Frontiers in Zoology* 12(1):16 DOI 10.1186/s12983-015-0107-z.
- Ahmed MZ, Breinholt JW, Kawahara AY. 2016. Evidence for common horizontal transmission of *Wolbachia* among butterflies and moths. *BMC Evolutionary Biology* 16:118 DOI 10.1186/s12862-016-0660-x.
- Akman L, Yamashita A, Watanabe H, Oshima K, Shiba T, Hattori M, Aksoy S. 2002. Genome sequence of the endocellular obligate symbiont of tsetse flies, *Wigglesworthia glossinidia*. *Nature Genetics* 32(3):402–407 DOI 10.1038/ng986.
- Aksoy S. 1995. *Wigglesworthia* gen. nov. and *Wigglesworthia glossinidia* sp. nov., taxa consisting of the mycetocyte-associated, primary endosymbionts of tsetse flies. *International Journal of Systematic Bacteriology* 45(4):848–851 DOI 10.1099/00207713-45-4-848.
- Altizer S, Bartel R, Han BA. 2011. Animal migration and infectious disease risk. *Science* 331(6015):296–302 DOI 10.1126/science.1194694.
- Altizer SM, Oberhauser KS. 1999. Effects of the protozoan parasite *Ophryocystis elektroscirrha* on the fitness of monarch butterflies (*Danaus plexippus*). *Journal of Invertebrate Pathology* 74(1):76–88 DOI 10.1006/jipa.1999.4853.
- Altizer SM, Oberhauser KS, Brower LP. 2000. Associations between host migration and the prevalence of a protozoan parasite in natural populations of adult monarch butterflies. *Ecological Entomology* 25(2):125–139 DOI 10.1046/j.1365-2311.2000.00246.x.
- Anbutsu H, Goto S, Fukatsu T. 2008. High and low temperatures differently affect infection density and vertical transmission of male-killing *Spiroplasma* symbionts in *Drosophila* hosts. *Applied and Environmental Microbiology* 74(19):6053–6059 DOI 10.1128/AEM.01503-08.
- Andreadis TG, Hall DW. 1979. Significance of transovarial infections of *Amblyospora* sp. (Microspora: Thelohaniidae) in relation to parasite maintenance in the mosquito *Culex salinarius*. *Journal of Invertebrate Pathology* 34(2):152–157 DOI 10.1016/0022-2011(79)90095-8.
- Ankola K, Brueckner D, Puttaraju HP. 2011. *Wolbachia* endosymbiont infection in two Indian butterflies and female-biased sex ratio in the Red Pierrot, *Talicauda nyseus*. *Journal of Biosciences* 36(5):845–850 DOI 10.1007/s12038-011-9149-3.

- Bandi C, Dunn AM, Hurst GD, Rigaud T. 2001.** Inherited microorganisms, sex-specific virulence and reproductive parasitism. *Trends in Parasitology* **17**(2):88–94
DOI [10.1016/s1471-4922\(00\)01812-2](https://doi.org/10.1016/s1471-4922(00)01812-2).
- Body M, Kaiser W, Dubreuil G, Casas J, Giron D. 2013.** Leaf-miners co-opt microorganisms to enhance their nutritional environment. *Journal of Chemical Ecology* **39**(7):969–977
DOI [10.1007/s10886-013-0307-y](https://doi.org/10.1007/s10886-013-0307-y).
- Bolaños LM, Servín-Garcidueñas LE, Martínez-Romero E. 2015.** Arthropod–*Spiroplasma* relationship in the genomic era. *FEMS Microbiology Ecology* **91**(2):1–8 DOI [10.1093/femsec/fiu008](https://doi.org/10.1093/femsec/fiu008).
- Bonte D, Hovestadt T, Poethke H-J. 2008.** Male-killing endosymbionts: influence of environmental conditions on persistence of host metapopulation. *BMC Evolutionary Biology* **8**(1):243 DOI [10.1186/1471-2148-8-243](https://doi.org/10.1186/1471-2148-8-243).
- Bordenstein SR. 2003.** Symbiosis and the origin of species. In: Bourtzis K, Miller TA, eds. *Insect Symbiosis*. Boca Raton: CRC Press, 283–304.
- Bordenstein SR, Wernegreen JJ. 2004.** Bacteriophage flux in endosymbionts (*Wolbachia*): infection frequency, lateral transfer, and recombination rates. *Molecular Biology and Evolution* **21**(10):1981–1991 DOI [10.1093/molbev/msh211](https://doi.org/10.1093/molbev/msh211).
- Bouchon D, Rigaud T, Juchault P. 1998.** Evidence for widespread *Wolbachia* infection in isopod crustaceans: molecular identification and host feminization. *Proceedings of the Royal Society B: Biological Sciences* **265**(1401):1081–1090 DOI [10.1098/rspb.1998.0402](https://doi.org/10.1098/rspb.1998.0402).
- Bradley CA, Altizer S. 2005.** Parasites hinder monarch butterfly flight: implications for disease spread in migratory hosts. *Ecology Letters* **8**(3):290–300 DOI [10.1111/j.1461-0248.2005.00722.x](https://doi.org/10.1111/j.1461-0248.2005.00722.x).
- Braquart-Varnier C, Lachat M, Herbinière J, Johnson M, Caubet Y, Bouchon D, Sicard M. 2008.** *Wolbachia* mediate variation of host immunocompetence. *PLOS ONE* **3**(9):e3286
DOI [10.1371/journal.pone.0003286](https://doi.org/10.1371/journal.pone.0003286).
- Braschler B, Hill JK. 2007.** Role of larval host plants in the climate-driven range expansion of the butterfly *Polygonia c-album*. *Journal of Animal Ecology* **76**(3):415–423
DOI [10.1111/j.1365-2656.2007.01217.x](https://doi.org/10.1111/j.1365-2656.2007.01217.x).
- Breeuwer JA, Werren JH. 1990.** Microorganisms associated with chromosome destruction and reproductive isolation between two insect species. *Nature* **346**(6284):558–560
DOI [10.1038/346558a0](https://doi.org/10.1038/346558a0).
- Brimacombe LC. 1980.** All-female broods in field and laboratory populations of the Egyptian cotton leafworm, *Spodoptera littoralis* (Boisduval) (Lepidoptera: Noctuidae). *Bulletin of Entomological Research* **70**(3):475–481 DOI [10.1017/s0007485300007756](https://doi.org/10.1017/s0007485300007756).
- Brownlie JC, Cass BN, Riegler M, Witsenburg JJ, Iturbe-Ormaetxe I, McGraw EA, O’Neill SL. 2009.** Evidence for metabolic provisioning by a common invertebrate endosymbiont, *Wolbachia pipientis*, during periods of nutritional stress. *PLOS Pathogens* **5**(4):e1000368
DOI [10.1371/journal.ppat.1000368](https://doi.org/10.1371/journal.ppat.1000368).
- Brownlie JC, Johnson KN. 2009.** Symbiont-mediated protection in insect hosts. *Trends in Microbiology* **17**(8):348–354 DOI [10.1016/j.tim.2009.05.005](https://doi.org/10.1016/j.tim.2009.05.005).
- Brucker RM, Bordenstein SR. 2012.** Speciation by symbiosis. *Trends in Ecology & Evolution* **27**(8):443–451 DOI [10.1016/j.tree.2012.03.011](https://doi.org/10.1016/j.tree.2012.03.011).
- Buchner P. 1965.** *Endosymbiosis of Animals with Plant Microorganisms*. New York: Interscience Publishers, John Wiley.
- Caspi-Fluger A, Inbar M, Mozes-Daube N, Katzir N, Portnoy V, Belausov E, Hunter MS, Zchori-Fein E. 2012.** Horizontal transmission of the insect symbiont *Rickettsia* is plant-mediated. *Proceedings of the Royal Society B: Biological Sciences* **279**(1734):1791–1796
DOI [10.1098/rspb.2011.2095](https://doi.org/10.1098/rspb.2011.2095).

- Caubet Y, Hatcher MJ, Mocquard JP, Rigaud T. 2000.** Genetic conflict and changes in heterogametic mechanisms of sex determination. *Journal of Evolutionary Biology* **13**(5):766–777 DOI [10.1046/j.1420-9101.2000.00225.x](https://doi.org/10.1046/j.1420-9101.2000.00225.x).
- Chafee ME, Zecher CN, Gourley ML, Schmidt VT, Chen JH, Bordenstein SR, Clark ME, Bordenstein SR. 2011.** Decoupling of host-symbiont-phage coadaptations following transfer between insect species. *Genetics* **187**(1):203–215 DOI [10.1534/genetics.110.120675](https://doi.org/10.1534/genetics.110.120675).
- Charlat S, Calmet C, Mercot H. 2001.** On the mod resc model and the evolution of *Wolbachia* compatibility types. *Genetics* **159**:1415–1422.
- Charlat S, Davies N, Roderick GK, Hurst GDD. 2007a.** Disrupting the timing of *Wolbachia*-induced male-killing. *Biology Letters* **3**(2):154–156 DOI [10.1098/rsbl.2006.0584](https://doi.org/10.1098/rsbl.2006.0584).
- Charlat S, Duploux A, Hornett EA, Dyson EA, Davies N, Roderick GK, Wedell N, Hurst GDD. 2009.** The joint evolutionary histories of *Wolbachia* and mitochondria in *Hypolimnas bolina*. *BMC Evolutionary Biology* **9**(1):64 DOI [10.1186/1471-2148-9-64](https://doi.org/10.1186/1471-2148-9-64).
- Charlat S, Engelstädter J, Dyson EA, Hornett EA, Duploux A, Tortosa P, Davies N, Roderick GK, Wedell N, Hurst GDD. 2006.** Competing selfish genetic elements in the butterfly *Hypolimnas bolina*. *Current Biology* **16**(24):2453–2458 DOI [10.1016/j.cub.2006.10.062](https://doi.org/10.1016/j.cub.2006.10.062).
- Charlat S, Hornett EA, Dyson EA, Ho PPY, Loc NT, Schilthuizen M, Davies N, Roderick GK, Hurst GDD. 2005.** Prevalence and penetrance variation of male-killing *Wolbachia* across Indo-Pacific populations of the butterfly *Hypolimnas bolina*. *Molecular Ecology* **14**(11):3525–3530 DOI [10.1111/j.1365-294X.2005.02678.x](https://doi.org/10.1111/j.1365-294X.2005.02678.x).
- Charlat S, Hornett EA, Fullard JH, Davies N, Roderick GK, Wedell N, Hurst GDD. 2007b.** Extraordinary flux in sex ratio. *Science* **317**(5835):214 DOI [10.1126/science.1143369](https://doi.org/10.1126/science.1143369).
- Charlat S, Hurst G, Mercot H. 2003.** Evolutionary consequences of *Wolbachia* infections. *Trends in Genetics* **19**(4):217–223 DOI [10.1016/S0168-9525\(03\)00024-6](https://doi.org/10.1016/S0168-9525(03)00024-6).
- Charlat S, Reuter M, Dyson EA, Hornett EA, Duploux A, Davies N, Roderick GK, Wedell N, Hurst GDD. 2007c.** Male-killing bacteria trigger a cycle of increasing male fatigue and female promiscuity. *Current Biology* **17**(3):273–277 DOI [10.1016/j.cub.2006.11.068](https://doi.org/10.1016/j.cub.2006.11.068).
- Chigira A, Miura K. 2005.** Detection of “candidate *Cardinium*” bacteria from the haploid host *Brevipalpus californicus* (Acari: Tenuipalpidae) and effect on the host. *Experimental and Applied Acarology* **37**(1–2):107–116 DOI [10.1007/s10493-005-0592-4](https://doi.org/10.1007/s10493-005-0592-4).
- Chrostek E, Pelz-Stelinski K, Hurst GDD, Hughes GL. 2017.** Horizontal transmission of intracellular insect symbionts via plants. *Frontiers in Microbiology* **8**:628 DOI [10.3389/fmicb.2017.02237](https://doi.org/10.3389/fmicb.2017.02237).
- Corbin C, Heyworth ER, Ferrari J, Hurst GDD. 2016.** Heritable symbionts in a world of varying temperature. *Heredity* **118**(1):10–20 DOI [10.1038/hdy.2016.71](https://doi.org/10.1038/hdy.2016.71).
- Cordaux R, Bouchon D, Grève P. 2011.** The impact of endosymbionts on the evolution of host sex-determination mechanisms. *Trends in Genetics* **27**(8):332–341 DOI [10.1016/j.tig.2011.05.002](https://doi.org/10.1016/j.tig.2011.05.002).
- Cordaux R, Michel-Salzat A, Frelon-Raimond M, Rigaud T, Bouchon D. 2004.** Evidence for a new feminizing *Wolbachia* strain in the isopod *Armadillidium vulgare*: evolutionary implications. *Heredity* **93**(1):78–84 DOI [10.1038/sj.hdy.6800482](https://doi.org/10.1038/sj.hdy.6800482).
- de Bary A. 1879.** *Die Erscheinungen der Symbiose*. Strassburg: Verlag von Karl J. Trübner.
- de Roode JC, Altizer S. 2010.** Host-parasite genetic interactions and virulence-transmission relationships in natural populations of monarch butterflies. *Evolution* **64**(2):502–514 DOI [10.1111/j.1558-5646.2009.00845.x](https://doi.org/10.1111/j.1558-5646.2009.00845.x).

- de Roode JC, Yates AJ, Altizer S. 2008.** Virulence-transmission trade-offs and population divergence in virulence in a naturally occurring butterfly parasite. *Proceedings of the National Academy of Sciences of the United States of America* **105**(21):7489–7494 DOI [10.1073/pnas.0710909105](https://doi.org/10.1073/pnas.0710909105).
- Dedeine F, Bouletreau M, Vavre F. 2005.** *Wolbachia* requirement for oogenesis: occurrence within the genus *Asobara* (Hymenoptera, Braconidae) and evidence for intraspecific variation in *A. tabida*. *Heredity* **95**(5):394–400 DOI [10.1038/sj.hdy.6800739](https://doi.org/10.1038/sj.hdy.6800739).
- Dedeine F, Vavre F, Fleury F, Loppin B, Hochberg ME, Bouletreau M. 2001.** Removing symbiotic *Wolbachia* bacteria specifically inhibits oogenesis in a parasitic wasp. *Proceedings of the National Academy of Sciences of the United States of America* **98**(11):6247–6252 DOI [10.1073/pnas.101304298](https://doi.org/10.1073/pnas.101304298).
- Delgado AM, Cook JM. 2009.** Effects of a sex-ratio distorting endosymbiont on mtDNA variation in a global insect pest. *BMC Evolutionary Biology* **9**(1):49 DOI [10.1186/1471-2148-9-49](https://doi.org/10.1186/1471-2148-9-49).
- Denlinger D, Yocum G. 1998.** Physiology of heat sensitivity. In: Hallman G, Denlinger D, eds. *Temperature Sensitivity in Insects and Application in Integrated Pest Management*. Boulder: Westview Press, 7–57.
- Derks MFL, Smit S, Salis L, Schijlen E, Bossers A, Mateman C, Pijl AS, de Ridder D, Groenen MAM, Visser ME, Megens H-J. 2015.** The genome of winter moth (*Operophtera brumata*) provides a genomic perspective on sexual dimorphism and phenology. *Genome Biology and Evolution* **7**(8):2321–2332 DOI [10.1093/gbe/evv145](https://doi.org/10.1093/gbe/evv145).
- Dion E, Polin SE, Simon J-C, Outreman Y. 2011.** Symbiont infection affects aphid defensive behaviours. *Biology Letters* **7**(5):743–746 DOI [10.1098/rsbl.2011.0249](https://doi.org/10.1098/rsbl.2011.0249).
- Douglas AE. 1998.** Nutritional interactions in insect-microbial symbioses: aphids and their symbiotic bacteria *Buchnera*. *Annual Review of Entomology* **43**(1):17–37 DOI [10.1146/annurev.ento.43.1.17](https://doi.org/10.1146/annurev.ento.43.1.17).
- Dunbar HE, Wilson ACC, Ferguson NR, Moran NA. 2007.** Aphid thermal tolerance is governed by a point mutation in bacterial symbionts. *PLOS Biology* **5**(5):e96 DOI [10.1371/journal.pbio.0050096](https://doi.org/10.1371/journal.pbio.0050096).
- Duploux A, Brattström O. 2017.** *Wolbachia* in the genus *Bicyclus*: a forgotten player. *Microbial Ecology* **75**(1):255–263 DOI [10.1007/s00248-017-1024-9](https://doi.org/10.1007/s00248-017-1024-9).
- Duploux A, Hurst GDD, O'Neill SL, Charlat S. 2010.** Rapid spread of male-killing *Wolbachia* in the butterfly *Hypolimnas bolina*. *Journal of Evolutionary Biology* **23**(1):231–235 DOI [10.1111/j.1420-9101.2009.01891.x](https://doi.org/10.1111/j.1420-9101.2009.01891.x).
- Duploux A, Iturbe-Ormaetxe I, Beatson SA, Szubert JM, Brownlie JC, McMeniman CJ, McGraw EA, Hurst GDD, Charlat S, O'Neill SL, Woolfit M. 2013.** Draft genome sequence of the male-killing *Wolbachia* strain wBoll reveals recent horizontal gene transfers from diverse sources. *BMC Genomics* **14**(1):20 DOI [10.1186/1471-2164-14-20](https://doi.org/10.1186/1471-2164-14-20).
- Duron O, Hurst GD. 2013.** Arthropods and inherited bacteria: from counting the symbionts to understanding how symbionts count. *BMC Biology* **11**(1):45 DOI [10.1186/1741-7007-11-45](https://doi.org/10.1186/1741-7007-11-45).
- Duron O, Wilkes TE, Hurst GDD. 2010.** Interspecific transmission of a male-killing bacterium on an ecological timescale. *Ecology Letters* **13**(9):1139–1148 DOI [10.1111/j.1461-0248.2010.01502.x](https://doi.org/10.1111/j.1461-0248.2010.01502.x).
- Düsing C. 1884.** Die Regulierung des Geschlechtsverhältnisses bei der Vermehrung der Menschen, Tiere und Pflanzen. *Jenaische Zeitschrift für Naturwissenschaft* **17**:593–940.
- Dyer KA, Jaenike J. 2005.** Evolutionary dynamics of a spatially structured host-parasite association: *Drosophila innubila* and male-killing *Wolbachia*. *Evolution* **59**(7):1518–1528 DOI [10.1111/j.0014-3820.2005.tb01801.x](https://doi.org/10.1111/j.0014-3820.2005.tb01801.x).

- Dyson EA, Hurst G. 2004. Persistence of an extreme sex-ratio bias in a natural population. *Proceedings of the National Academy of Sciences of the United States of America* 101(17):6520–6523 DOI 10.1073/pnas.0304068101.
- Dyson EA, Kamath MK, Hurst GDD. 2002. *Wolbachia* infection associated with all-female broods in *Hypolimnas bolina* (Lepidoptera: Nymphalidae): evidence for horizontal transmission of a butterfly male killer. *Heredity* 88(3):166–171 DOI 10.1038/sj.hdy.6800021.
- Elnagdy S, Majerus MEN, Handley L-JL. 2011. The value of an egg: resource reallocation in ladybirds (Coleoptera: Coccinellidae) infected with male-killing bacteria. *Journal of Evolutionary Biology* 24(10):2164–2172 DOI 10.1111/j.1420-9101.2011.02346.x.
- Engelstädter J, Hurst GDD. 2007. The impact of male-killing bacteria on host evolutionary processes. *Genetics* 175(1):245–254 DOI 10.1534/genetics.106.060921.
- Engelstädter J, Hurst GDD. 2009. The ecology and evolution of microbes that manipulate host reproduction. *Annual Review of Ecology, Evolution, and Systematics* 40(1):127–149 DOI 10.1146/annurev.ecolsys.110308.120206.
- Fenner J, Seltzer J, Peyton S, Sullivan H, Tolson P, Walsh R, Hill J, Counterman B. 2017. Demographic variation of *Wolbachia* infection in the endangered Mitchell's Satyr butterfly. *Insects* 8(2):50 DOI 10.3390/insects8020050.
- Ferrari J, Darby AC, Daniell TJ, Godfray HCJ, Douglas AE. 2004. Linking the bacterial community in pea aphids with host-plant use and natural enemy resistance. *Ecological Entomology* 29(1):60–65 DOI 10.1111/j.1365-2311.2004.00574.x.
- Ferree PM, Avery A, Azpurua J, Wilkes T, Werren JH. 2008. A bacterium targets maternally inherited centrosomes to kill males in *Nasonia*. *Current Biology* 18(18):1409–1414 DOI 10.1016/j.cub.2008.07.093.
- Fisher RA. 1930. *The Genetical Theory of Natural Selection*. Oxford: Clarendon Press.
- Fleury F, Vavre F, Ris N, Fouillet P, Bouletreau M. 2000. Physiological cost induced by the maternally-transmitted endosymbiont *Wolbachia* in the *Drosophila* parasitoid *Leptopilina heterotoma*. *Parasitology* 121(5):493–500 DOI 10.1017/s0031182099006599.
- Fraune S, Bosch TCG. 2010. Why bacteria matter in animal development and evolution. *BioEssays* 32(7):571–580 DOI 10.1002/bies.200900192.
- Fry AJ, Rand DM. 2002. *Wolbachia* interactions that determine *Drosophila melanogaster* survival. *Evolution* 56(10):1976–1981 DOI 10.1111/j.0014-3820.2002.tb00123.x.
- Fujii Y, Kageyama D, Hoshizaki S, Ishikawa H, Sasaki T. 2001. Transfection of *Wolbachia* in Lepidoptera: the feminizer of the adzuki bean borer *Ostrinia scapularis* causes male killing in the Mediterranean flour moth *Ephesia kuehniella*. *Proceedings of the Royal Society B: Biological Sciences* 268(1469):855–859 DOI 10.1098/rspb.2001.1593.
- Fukui T, Kawamoto M, Shoji K, Kiuchi T, Sugano S, Shimada T, Suzuki Y, Katsuma S. 2015. The endosymbiotic bacterium *Wolbachia* selectively kills male hosts by targeting the masculinizing gene. *PLOS Pathogens* 11(7):e1005048 DOI 10.1371/journal.ppat.1005048.
- Gehrer L, Vorburger C. 2012. Parasitoids as vectors of facultative bacterial endosymbionts in aphids. *Biology Letters* 8(4):613–615 DOI 10.1098/rsbl.2012.0144.
- Geier PW, Briese DT, Lewis T. 1978. The light-brown apple moth, *Epiphyas postvittana* (Walker) 2. Uneven sex ratios and a condition contributing to them in the field. *Australian Journal of Ecology* 3(4):467–488 DOI 10.1111/j.1442-9993.1978.tb01192.x.
- Giron D, Glevarec G. 2014. Cytokinin-induced phenotypes in plant-insect interactions: learning from the bacterial world. *Journal of Chemical Ecology* 40(7):826–835 DOI 10.1007/s10886-014-0466-5.

- Gompert Z, Forister ML, Fordyce JA, Nice CC. 2008.** Widespread mito-nuclear discordance with evidence for introgressive hybridization and selective sweeps in *Lycaeides*. *Molecular Ecology* 17(24):5231–5244 DOI 10.1111/j.1365-294X.2008.03988.x.
- Goodacre SL, Martin OY, Bonte D, Hutchings L, Woolley C, Ibrahim K, Thomas CG, Hewitt GM. 2009.** Microbial modification of host long-distance dispersal capacity. *BMC Biology* 7(1):32 DOI 10.1186/1741-7007-7-32.
- Graham RI, Hartley L, Wilson K. 2011.** Characterisation of a nucleopolyhedrovirus and *Spiroplasma* sp. bacterium associated with outbreaking populations of the Antler moth *Cerapteryx graminis*. *Journal of Invertebrate Pathology* 107(1):90–93 DOI 10.1016/j.jip.2011.01.006.
- Graham RI, Wilson K. 2012.** Male-killing *Wolbachia* and mitochondrial selective sweep in a migratory African insect. *BMC Evolutionary Biology* 12(1):204–211 DOI 10.1186/1471-2148-12-204.
- Groot TVM, Breeuwer JAJ. 2006.** *Cardinium* symbionts induce haploid thelytoky in most clones of three closely related *Brevipalpus* species. *Experimental and Applied Acarology* 39(3–4):257–271 DOI 10.1007/s10493-006-9019-0.
- Gross R, Vavre F, Heddi A, Hurst GDD, Zchori-Fein E, Bourtzis K. 2009.** Immunity and symbiosis. *Molecular Microbiology* 73(5):751–759 DOI 10.1111/j.1365-2958.2009.06820.x.
- Gutiérrez D, Thomas CD. 2000.** Marginal range expansion in a host-limited butterfly species *Gonepteryx rhamni*. *Ecological Entomology* 25(2):165–170 DOI 10.1046/j.1365-2311.2000.00241.x.
- Gutzwiller F, Dedeine F, Kaiser W, Giron D, López-Vaamonde C. 2015.** Correlation between the green-island phenotype and *Wolbachia* infections during the evolutionary diversification of Gracillariidae leaf-mining moths. *Ecology and Evolution* 5(18):4049–4062 DOI 10.1002/ece3.1580.
- Haine ER. 2008.** Symbiont-mediated protection. *Proceedings of the Royal Society B: Biological Sciences* 275:353–361 DOI 10.1098/rspb.2007.1211.
- Hamilton WDW. 1967.** Extraordinary sex ratios. A sex-ratio theory for sex linkage and inbreeding has new implications in cytogenetics and entomology. *Science* 156(3774):477–488 DOI 10.1126/science.156.3774.477.
- Hamm CA, Handley CA, Pike A, Forister ML, Fordyce ML, Fordyce JA, Nice CC. 2014.** *Wolbachia* infection and Lepidoptera of conservation concern. *Journal of Insect Science* 14(1):6 DOI 10.1093/jis/14.1.6.
- Hammer TJ, Janzen DH, Hallwachs W, Jaffe SP, Fierer N. 2017.** Caterpillars lack a resident gut microbiome. *Proceedings of the National Academy of Sciences of the United States of America* 114(36):9641–9646 DOI 10.1073/pnas.1707186114.
- Hansen AK, Moran NA. 2014.** The impact of microbial symbionts on host plant utilization by herbivorous insects. *Molecular Ecology* 23(6):1473–1496 DOI 10.1111/mec.12421.
- Harumoto T, Anbutsu H, Lemaitre B, Fukatsu T. 2016.** Male-killing symbiont damages host's dosage-compensated sex chromosome to induce embryonic apoptosis. *Nature Communications* 7:12781 DOI 10.1038/ncomms12781.
- Hassan SSM, Idris E. 2013.** Male-killing in African butterflies. *Trends in Evolutionary Biology* 5(1):2 DOI 10.4081/eb.2013.e2.
- Heath BD, Butcher RD, Whitfield WG, Hubbard SF. 1999.** Horizontal transfer of *Wolbachia* between phylogenetically distant insect species by a naturally occurring mechanism. *Current Biology* 9(6):313–316 DOI 10.1016/s0960-9822(99)80139-0.

- Hedges LM, Brownlie JC, O'Neill SL, Johnson KN. 2008. *Wolbachia* and virus protection in insects. *Science* 322(5902):702 DOI 10.1126/science.1162418.
- Herren JK, Gordon I, Holland PWH, Smith D. 2007. The butterfly *Danaus chrysippus* (Lepidoptera: Nymphalidae) in Kenya is variably infected with respect to genotype and body size by a maternally transmitted male-killing endosymbiont (*Spiroplasma*). *International Journal of Tropical Insect Science* 27(2):62 DOI 10.1017/S1742758407818327.
- Hertig M, Wolbach SB. 1924. Studies on *Rickettsia*-like micro-organisms in insects. *Journal of Medical Research* 44:329–374.7.
- Higashiura Y, Ishihara M, Schaefer PW. 1999. Sex ratio distortion and severe inbreeding depression in the gypsy moth *Lymantria dispar* L. in Hokkaido, Japan. *Heredity* 83(3):290–297 DOI 10.1038/sj.hdy.6885590.
- Hiroki M, Kato Y, Kamito T, Miura K. 2002. Feminization of genetic males by a symbiotic bacterium in a butterfly, *Eurema hecabe* (Lepidoptera: Pieridae). *Naturwissenschaften* 89(4):167–170 DOI 10.1007/s00114-002-0303-5.
- Hiroki M, Tagami Y, Miura K, Kato Y. 2004. Multiple infection with *Wolbachia* inducing different reproductive manipulations in the butterfly *Eurema hecabe*. *Proceedings of the Royal Society B: Biological Sciences* 271(1549):1751–1755 DOI 10.1098/rspb.2004.2769.
- Hoffman AA, Turelli M. 1997. Cytoplasmic incompatibility in insects. In: O'Neill SL, Hoffman AA, Werren JH, eds. *Influential Passengers: Inherited Microorganisms and Arthropod Reproduction*. Oxford: Oxford University Press, 42–80.
- Hoffman AA, Turelli M, Harshman LG. 1990. Factors affecting the distribution of cytoplasmic incompatibility in *Drosophila simulans*. *Genetics* 126:933–948.
- Hornett EA, Charlat S, Duploux AMR, Davies N, Roderick GK, Wedell N, Hurst GDD. 2006. Evolution of male-killer suppression in a natural population. *PLOS Biology* 4(9):e283 DOI 10.1371/journal.pbio.0040283.
- Hornett EA, Charlat S, Wedell N, Jiggins CD, Hurst GDD. 2009. Rapidly shifting sex ratio across a species range. *Current Biology* 19(19):1628–1631 DOI 10.1016/j.cub.2009.07.071.
- Hornett EA, Duploux AMR, Davies N, Roderick GK, Wedell N, Hurst GDD, Charlat S. 2008. You can't keep a good parasite down: evolution of a male-killer suppressor uncovers cytoplasmic incompatibility. *Evolution* 62(5):1258–1263 DOI 10.1111/j.1558-5646.2008.00353.x.
- Hornett EA, Moran B, Reynolds LA, Charlat S, Tazzyman S, Wedell N, Jiggins CD, Hurst GDD. 2014. The evolution of sex ratio distorter suppression affects a 25 cM genomic region in the butterfly *Hypolimnas bolina*. *PLOS Genetics* 10(12):e1004822 DOI 10.1371/journal.pgen.1004822.
- Hosokawa T, Koga R, Kikuchi Y, Meng X-Y, Fukatsu T. 2010. *Wolbachia* as a bacteriocyte-associated nutritional mutualist. *Proceedings of the National Academy of Sciences of the United States of America* 107(2):769–774 DOI 10.1073/pnas.0911476107.
- Hotopp JCD, Clark ME, Oliveira DCSG, Foster JM, Fischer P, Torres MCM, Giebel JD, Kumar N, Ishmael N, Wang S, Ingram J, Nene RV, Shepard J, Tomkins J, Richards S, Spiro DJ, Ghedin E, Slatko BE, Tettelin H, Werren JH. 2007. Widespread lateral gene transfer from intracellular bacteria to multicellular eukaryotes. *Science* 317(5845):1753–1756 DOI 10.1126/science.1142490.
- Hughes GL, Koga R, Xue P, Fukatsu T, Rasgon JL. 2011. *Wolbachia* infections are virulent and inhibit the human malaria parasite *Plasmodium falciparum* in *Anopheles gambiae*. *PLOS Pathogens* 7(5):e1002043 DOI 10.1371/journal.ppat.1002043.

- Hughes GL, Rivero A, Rasgon JL. 2014. *Wolbachia* can enhance *Plasmodium* infection in mosquitoes: implications for malaria control? *PLOS Pathogens* **10**(9):e1004182 DOI [10.1371/journal.ppat.1004182](https://doi.org/10.1371/journal.ppat.1004182).
- Huigens ME, de Almeida RP, Boons PAH, Luck RF, Stouthamer R. 2004. Natural interspecific and intraspecific horizontal transfer of parthenogenesis-inducing *Wolbachia* in *Trichogramma* wasps. *Proceedings of the Royal Society B: Biological Sciences* **271**(1538):509–515 DOI [10.1098/rspb.2003.2640](https://doi.org/10.1098/rspb.2003.2640).
- Huigens ME, Luck RF, Klaassen RHG, Maas MFPM, Timmermans MJTN, Stouthamer R. 2000. Infectious parthenogenesis. *Nature* **405**(6783):178–179 DOI [10.1038/35012066](https://doi.org/10.1038/35012066).
- Hunter MS, Perlman SJ, Kelly SE. 2003. A bacterial symbiont in the *Bacteroidetes* induces cytoplasmic incompatibility in the parasitoid wasp *Encarsia pergandiella*. *Proceedings of the Royal Society B: Biological Sciences* **270**(1529):2185–2190 DOI [10.1098/rspb.2003.2475](https://doi.org/10.1098/rspb.2003.2475).
- Hurst GDD, Hurst LD, Majerus M. 1997. Cytoplasmic sex-ratio distorters. In: O'Neill SL, Hoffmann AA, Werren JH, eds. *Influential Passengers: Inherited Microorganisms and Arthropod Reproduction*. Oxford: Oxford University Press, 125–154.
- Hurst GDD, Jiggins FM. 2000. Male-killing bacteria in insects: mechanisms, incidence, and implications. *Emerging Infectious Diseases* **6**(4):329–336 DOI [10.3201/eid0604.000402](https://doi.org/10.3201/eid0604.000402).
- Hurst GDD, Jiggins FM. 2005. Problems with mitochondrial DNA as a marker in population, phylogeographic and phylogenetic studies: the effects of inherited symbionts. *Proceedings of the Royal Society B: Biological Sciences* **272**(1572):1525–1534 DOI [10.1098/rspb.2005.3056](https://doi.org/10.1098/rspb.2005.3056).
- Hurst GDD, Jiggins FM, Hinrich Graf von der Schulenburg J, Bertrand D, West SA, Goriacheva II, Zakharov IA, Werren JH, Stouthamer R, Majerus MEN. 1999. Male-killing *Wolbachia* in two species of insect. *Proceedings of the Royal Society B: Biological Sciences* **266**(1420):735–740 DOI [10.1098/rspb.1999.0698](https://doi.org/10.1098/rspb.1999.0698).
- Hurst GD, Jiggins FM, Robinson SJ. 2001. What causes inefficient transmission of male-killing *Wolbachia* in *Drosophila*? *Heredity* **87**(2):220–226 DOI [10.1046/j.1365-2540.2001.00917.x](https://doi.org/10.1046/j.1365-2540.2001.00917.x).
- Hurst GD, Johnson AP, Schulenburg JH, Fuyama Y. 2000. Male-killing *Wolbachia* in *Drosophila*: a temperature-sensitive trait with a threshold bacterial density. *Genetics* **156**:699–709.
- Hurst GDD, Majerus MEN. 1992. Why do maternally inherited microorganisms kill males? *Heredity* **71**(1):81–95 DOI [10.1038/hdy.1993.110](https://doi.org/10.1038/hdy.1993.110).
- Hurst GDD, Schilthuizen M. 1998. Selfish genetic elements and speciation. *Heredity* **80**(1):2–8 DOI [10.1038/sj.hdy.6883370](https://doi.org/10.1038/sj.hdy.6883370).
- Husnik F, McCutcheon JP. 2018. Functional horizontal gene transfer from bacteria to eukaryotes. *Nature Reviews Microbiology* **16**(2):67–79 DOI [10.1038/nrmicro.2017.137](https://doi.org/10.1038/nrmicro.2017.137).
- Hutchence KJ, Fischer B, Paterson S, Hurst GDD. 2011. How do insects react to novel inherited symbionts? A microarray analysis of *Drosophila melanogaster* response to the presence of natural and introduced *Spiroplasma*. *Molecular Ecology* **20**(5):950–958 DOI [10.1111/j.1365-294X.2010.04974.x](https://doi.org/10.1111/j.1365-294X.2010.04974.x).
- Ilinsky Y, Kosterin OE. 2017. Molecular diversity of *Wolbachia* in Lepidoptera: prevalent allelic content and high recombination of MLST genes. *Molecular Phylogenetics and Evolution* **109**:164–179 DOI [10.1016/j.ympev.2016.12.034](https://doi.org/10.1016/j.ympev.2016.12.034).
- Jaenike J. 2007. Spontaneous emergence of a new *Wolbachia* phenotype. *Evolution* **61**(9):2244–2252 DOI [10.1111/j.1558-5646.2007.00180.x](https://doi.org/10.1111/j.1558-5646.2007.00180.x).
- Jaenike J, Polak M, Fiskin A, Helou M, Minhas M. 2007. Interspecific transmission of endosymbiotic *Spiroplasma* by mites. *Biology Letters* **3**(1):23–25 DOI [10.1098/rsbl.2006.0577](https://doi.org/10.1098/rsbl.2006.0577).

- Jaenike J, Unckless R, Cockburn SN, Boelio LM, Perlman SJ. 2010. Adaptation via symbiosis: recent spread of a *Drosophila* defensive symbiont. *Science* **329**(5988):212–215
DOI [10.1126/science.1188235](https://doi.org/10.1126/science.1188235).
- Jiggins FM. 2002. Widespread “hilltopping” in *Acraea* butterflies and the origin of sex-role-reversed swarming in *Acraea encedon* and *A. encedana*. *African Journal of Ecology* **40**(3):228–231
DOI [10.1046/j.1365-2028.2002.00359.x](https://doi.org/10.1046/j.1365-2028.2002.00359.x).
- Jiggins FM. 2003. Male-killing *Wolbachia* and mitochondrial DNA: selective sweeps, hybrid introgression and parasite population dynamics. *Genetics* **164**:5–12.
- Jiggins FM, Bentley JK, Majerus MEN, Hurst GDD. 2001. How many species are infected with *Wolbachia*? Cryptic sex ratio distorters revealed to be common by intensive sampling. *Proceedings of the Royal Society B: Biological Sciences* **268**(1472):1123–1126
DOI [10.1098/rspb.2001.1632](https://doi.org/10.1098/rspb.2001.1632).
- Jiggins FM, Hurst GDD, Dolman CE, Majerus MEN. 2000b. High-prevalence male-killing *Wolbachia* in the butterfly *Acraea encedana*. *Journal of Evolutionary Biology* **13**(3):495–501
DOI [10.1046/j.1420-9101.2000.00180.x](https://doi.org/10.1046/j.1420-9101.2000.00180.x).
- Jiggins FM, Hurst GDD, Jiggins CD, Schulenburg JHGvd, Majerus MEN. 2000a. The butterfly *Danaus chrysippus* is infected by a male-killing *Spiroplasma* bacterium. *Parasitology* **120**(5):439–446
DOI [10.1017/s0031182099005867](https://doi.org/10.1017/s0031182099005867).
- Jiggins FM, Hurst GDD, Majerus MEN. 1998. Sex ratio distortion in *Acraea encedon* (Lepidoptera: Nymphalidae) is caused by a male-killing bacterium. *Heredity* **81**(1):87–91
DOI [10.1046/j.1365-2540.1998.00357.x](https://doi.org/10.1046/j.1365-2540.1998.00357.x).
- Jiggins FM, Hurst GDD, Majerus M. 2000. Sex-ratio-distorting *Wolbachia* causes sex-role reversal in its butterfly host. *Proceedings of the Royal Society B: Biological Sciences* **267**(1438):69–73
DOI [10.1098/rspb.2000.0968](https://doi.org/10.1098/rspb.2000.0968).
- Johnstone RA, Hurst GDD. 1996. Maternally inherited male-killing microorganisms may confound interpretation of mitochondrial DNA variability. *Biological Journal of the Linnean Society* **58**(4):453–470
DOI [10.1006/bjil.1996.0047](https://doi.org/10.1006/bjil.1996.0047).
- Juchault P, Rigaud T, Mocquard JP. 1992. Evolution of sex-determining mechanisms in a wild population of *Armadillidium vulgare* Latr. (Crustacea, Isopoda): competition between two feminizing parasitic sex factors. *Heredity* **69**(4):382–390
DOI [10.1038/hdy.1992.138](https://doi.org/10.1038/hdy.1992.138).
- Kageyama D, Narita S, Watanabe M. 2012. Insect sex determination manipulated by their endosymbionts: incidences, mechanisms and implications. *Insects* **3**(1):161–199
DOI [10.3390/insects3010161](https://doi.org/10.3390/insects3010161).
- Kageyama D, Nishimura G, Hoshizaki S, Ishikawa Y. 2002. Feminizing *Wolbachia* in an insect, *Ostrinia furnacalis* (Lepidoptera: Crambidae). *Heredity* **88**(6):444–449
DOI [10.1038/sj.hdy.6800077](https://doi.org/10.1038/sj.hdy.6800077).
- Kageyama D, Nishimura G, Ohno S, Takanashi T, Hoshizaki S, Ishikawa Y. 2004. *Wolbachia* infection and an all-female trait in *Ostrinia orientalis* and *Ostrinia zaguliaevi*. *Entomologia Experimentalis et Applicata* **111**(1):79–83
DOI [10.1111/j.0013-8703.2004.00153.x](https://doi.org/10.1111/j.0013-8703.2004.00153.x).
- Kageyama D, Traut W. 2004. Opposite sex-specific effects of *Wolbachia* and interference with the sex determination of its host *Ostrinia scapularis*. *Proceedings of the Royal Society B: Biological Sciences* **271**(1536):251–258
DOI [10.1098/rspb.2003.2604](https://doi.org/10.1098/rspb.2003.2604).
- Kageyama D, Ohno M, Sasaki T, Yoshido A, Konagaya T, Jouraku A, Kuwazaki S, Kanamori H, Katayose Y, Narita S, Miyata M, Riegler M, Sahara K. 2017. Feminizing *Wolbachia* endosymbiont disrupts maternal sex chromosome inheritance in a butterfly species. *Evolution Letters* **1**(5):232–244
DOI [10.1002/evl3.28](https://doi.org/10.1002/evl3.28).

- Kaiser W, Huguet E, Casas J, Commin C, Giron D. 2010. Plant green-island phenotype induced by leaf-miners is mediated by bacterial symbionts. *Proceedings of the Royal Society B: Biological Sciences* 277(1692):2311–2319 DOI 10.1098/rspb.2010.0214.
- Kaltenpoth M, Göttler W, Herzner G, Strohm E. 2005. Symbiotic bacteria protect wasp larvae from fungal infestation. *Current Biology* 15(5):475–479 DOI 10.1016/j.cub.2004.12.084.
- Kato Y. 2000. Overlapping distribution of two groups of the butterfly *Eurema hecabe* differing in the expression of seasonal morphs on Okinawa-jima Island. *Zoological Science* 17(4):539–547 DOI 10.2108/0289-0003(2000)17[539:ODOTGO]2.0.CO;2.
- Kellner R. 1999. What is the basis of pederin polymorphism in *Paederus riparius* rove beetles? The endosymbiotic hypothesis. *Entomologia Experimentalis et Applicata* 93(1):41–49 DOI 10.1046/j.1570-7458.1999.00560.x.
- Kellner R, Dettner K. 1996. Differential efficacy of toxic pederin in deterring potential arthropod predators of *Paederus* (Coleoptera: Staphylinidae) offspring. *Oecologia* 107(3):293–300 DOI 10.1007/bf00328445.
- Kern P, Cook JM, Kageyama D, Riegler M. 2015. Double trouble: combined action of meiotic drive and *Wolbachia* feminization in *Eurema* butterflies. *Biology Letters* 11(5):20150095 DOI 10.1098/rsbl.2015.0095.
- Klasson L, Kambris Z, Cook PE, Walker T, Sinkins SP. 2009. Horizontal gene transfer between *Wolbachia* and the mosquito *Aedes aegypti*. *BMC Genomics* 10(1):33 DOI 10.1186/1471-2164-10-33.
- Kodandaramaiah U, Weingartner E, Janz N, Dalén L, Nylin S. 2011. Population structure in relation to host-plant ecology and *Wolbachia* infestation in the comma butterfly. *Journal of Evolutionary Biology* 24(10):2173–2185 DOI 10.1111/j.1420-9101.2011.02352.x.
- Koukou K, Pavlikaki H, Kiliass G, Werren JH, Bourtzis K, Alahiotis SN. 2006. Influence of antibiotic treatment and *Wolbachia* curing on sexual isolation among *Drosophila melanogaster* cage populations. *Evolution* 60:87–96 DOI 10.1111/j.0014-3820.2006.tb01084.x.
- Laven H. 1959. Speciation by cytoplasmic isolation in the *Culex pipiens*-complex. *Cold Spring Harbor Symposia on Quantitative Biology* 24:166–173 DOI 10.1101/sqb.1959.024.01.017.
- Lefèvre T, Adamo SA, Biron DG, Missé D, Hughes D, Thomas F. 2009. Invasion of the body snatchers: the diversity and evolution of manipulative strategies in host–parasite interactions. *Advances in Parasitology* 68:45–83 DOI 10.1016/S0065-308X(08)00603-9.
- Leonardo TE, Muiru GT. 2003. Facultative symbionts are associated with host plant specialization in pea aphid populations. *Proceedings of the Royal Society B: Biological Sciences* 270(Suppl. 2):S209–S212 DOI 10.1098/rsbl.2003.0064.
- LePage DP, Metcalf JA, Bordenstein SR, On J, Perlmutter JI, Shropshire JD, Layton EM, Funkhouser-Jones LJ, Beckmann JF, Bordenstein SR. 2017. Prophage WO genes recapitulate and enhance *Wolbachia*-induced cytoplasmic incompatibility. *Nature* 543(7644):243–247 DOI 10.1038/nature21391.
- Lewis Z, Champion de Crespigny FE, Sait SM, Tregenza T, Wedell N. 2011. *Wolbachia* infection lowers fertile sperm transfer in a moth. *Biology Letters* 7(2):187–189 DOI 10.1098/rsbl.2010.0605.
- Li C-X, Shi M, Tian J-H, Lin X-D, Kang Y-J, Chen L-J, Qin X-C, Xu J, Holmes EC, Zhang Y-Z. 2015. Unprecedented genomic diversity of RNA viruses in arthropods reveals the ancestry of negative-sense RNA viruses. *eLIFE* 4:e05378 DOI 10.7554/eLife.05378.
- Lindsey ARI, Werren JH, Richards S, Stouthamer R. 2016. Comparative genomics of a parthenogenesis-inducing *Wolbachia* symbiont. *G3: Genes|Genomes|Genetics* 6(7):2113–2123 DOI 10.1534/g3.116.028449.

- Lohman DJ, Peggie D, Pierce NE, Meier R. 2008. Phylogeography and genetic diversity of a widespread Old World butterfly, *Lampides boeticus* (Lepidoptera: Lycaenidae). *BMC Evolutionary Biology* 8(1):301 DOI 10.1186/1471-2148-8-301.
- Longdon B, Day JP, Schulz N, Leftwich PT, de Jong MA, Breuker CJ, Gibbs M, Obbard DJ, Wilfert L, Smith SCL, McGonigle JE, Houslay TM, Wright LI, Livraghi L, Evans LC, Friend LA, Chapman T, Vontas J, Kambouraki N, Jiggins FM. 2017. Vertically transmitted rhabdoviruses are found across three insect families and have dynamic interactions with their hosts. *Proceedings of the Royal Society B: Biological Sciences* 284(1847):20162381 DOI 10.1098/rspb.2016.2381.
- Longdon B, Jiggins FM. 2012. Vertically transmitted viral endosymbionts of insects: do sigma viruses walk alone? *Proceedings of the Royal Society B: Biological Sciences* 279(1744):3889–3898 DOI 10.1098/rspb.2012.1208.
- Łukasiewicz K, Sanak M, Węgrzyn G. 2016. A lack of *Wolbachia*-specific DNA in samples from apollo butterfly (*Parnassius apollo*, Lepidoptera: Papilionidae) individuals with deformed or reduced wings. *Journal of Applied Genetics* 57(2):271–274 DOI 10.1007/s13353-015-0318-1.
- Ma W-J, Vavre F, Beukeboom LW. 2014. Manipulation of arthropod sex determination by endosymbionts: diversity and molecular mechanisms. *Sexual Development* 8(1–3):59–73 DOI 10.1159/000357024.
- Maitland DP. 1994. A parasitic fungus infecting yellow dungflies manipulates host perching behavior. *Proceedings of the Royal Society B: Biological Sciences* 258(1352):187–193 DOI 10.1098/rspb.1994.0161.
- Margulis L, Fester R. 1991. *Symbiosis as a Source of Evolutionary Innovation: Speciation and Morphogenesis*. Cambridge: The MIT Press.
- Martinez J, Longdon B, Bauer S, Chan Y-S, Miller WJ, Bourtzis K, Teixeira L, Jiggins FM. 2014. Symbionts commonly provide broad spectrum resistance to viruses in insects: a comparative analysis of *Wolbachia* strains. *PLOS Pathogens* 10(9):e1004369 DOI 10.1371/journal.ppat.1004369.
- Mason CJ, Raffa KF. 2014. Acquisition and structuring of midgut bacterial communities in Gypsy Moth (Lepidoptera: Erebidae) larvae. *Environmental Entomology* 43(3):595–604 DOI 10.1603/en14031.
- Masui S, Kamoda S, Sasaki T, Ishikawa H. 2000. Distribution and evolution of bacteriophage WO in *Wolbachia*, the endosymbiont causing sexual alterations in arthropods. *Journal of Molecular Evolution* 51(5):491–497 DOI 10.1007/s002390010112.
- McCutcheon JP, McDonald BR, Moran NA. 2009. Convergent evolution of metabolic roles in bacterial co-symbionts of insects. *Proceedings of the National Academy of Sciences of the United States of America* 106(36):15394–15399 DOI 10.1073/pnas.0906424106.
- McHugh A, Bierzychudek P, Greever C, Marzulla T, Van Buskirk R, Binford G. 2013. A molecular phylogenetic analysis of *Speyeria* and its implications for the management of the threatened *Speyeria zerene hippolyta*. *Journal of Insect Conservation* 17(6):1237–1253 DOI 10.1007/s10841-013-9605-5.
- McLaughlin RE, Myers J. 1970. *Ophryocystis elektroscirrha* sp. n., a neogregarine pathogen of the Monarch butterfly *Danaus plexippus* (L.) and the Florida Queen butterfly *D. gilippus berenice* Cramer. *Journal of Protozoology* 17(2):300–305 DOI 10.1111/j.1550-7408.1970.tb02375.x.
- Miller WJ, Ehrman L, Schneider D. 2010. Infectious speciation revisited: impact of symbiont-depletion on female fitness and mating behavior of *Drosophila paulistorum*. *PLOS Pathogens* 6(12):e1001214 DOI 10.1371/journal.ppat.1001214.

- Miller WJ, Schneider D. 2012.** Endosymbiotic microbes as adaptive manipulators of arthropod behavior and natural driving sources of host speciation. In: Hughes D, Brodeur J, Thomas F, eds. *Host Manipulation by Parasites*. Oxford: Oxford University Press, 119–137.
- Montllor CB, Maxmen A, Purcell AH. 2002.** Facultative bacterial endosymbionts benefit pea aphids *Acyrtosiphon pisum* under heat stress. *Ecological Entomology* **27**(2):189–195
DOI [10.1046/j.1365-2311.2002.00393.x](https://doi.org/10.1046/j.1365-2311.2002.00393.x).
- Moran NA. 2016.** When obligate partners melt down. *mBio* **7**(6):e01904–e01916
DOI [10.1128/mBio.01904-16](https://doi.org/10.1128/mBio.01904-16).
- Moran NA, McCutcheon JP, Nakabachi A. 2008.** Genomics and evolution of heritable bacterial symbionts. *Annual Review of Genetics* **42**(1):165–190
DOI [10.1146/annurev.genet.41.110306.130119](https://doi.org/10.1146/annurev.genet.41.110306.130119).
- Moreira LA, Iturbe-Ormaetxe I, Jeffery JA, Lu G, Pyke AT, Hedges LM, Rocha BC, Hall-Mendelin S, Day A, Riegler M, Hugo LE, Johnson KN, Kay BH, McGraw EA, van den Hurk AF, Ryan PA, O'Neill SL. 2009.** A *Wolbachia* symbiont in *Aedes aegypti* limits infection with dengue, Chikungunya, and Plasmodium. *Cell* **139**(7):1268–1278
DOI [10.1016/j.cell.2009.11.042](https://doi.org/10.1016/j.cell.2009.11.042).
- Morimoto S, Nakai M, Ono A, Kunimi Y. 2001.** Late male-killing phenomenon found in a Japanese population of the oriental tea tortrix, *Homona magnanima* (Lepidoptera: Tortricidae). *Heredity* **87**(4):435–440 DOI [10.1046/j.1365-2540.2001.00924.x](https://doi.org/10.1046/j.1365-2540.2001.00924.x).
- Nakanishi K, Hoshino M, Nakai M, Kunimi Y. 2008.** Novel RNA sequences associated with late male killing in *Homona magnanima*. *Proceedings of the Royal Society B: Biological Sciences* **275**(1640):1249–1254 DOI [10.1098/rspb.2008.0013](https://doi.org/10.1098/rspb.2008.0013).
- Narita S, Kageyama D, Hiroki M, Sanpei T, Hashimoto S, Kamitoh T, Kato Y. 2011.** *Wolbachia*-induced feminisation newly found in *Eurema hecabe*, a sibling species of *Eurema mandarina* (Lepidoptera: Pieridae). *Ecological Entomology* **36**(3):309–317
DOI [10.1111/j.1365-2311.2011.01274.x](https://doi.org/10.1111/j.1365-2311.2011.01274.x).
- Narita S, Kageyama D, Nomura M, Fukatsu T. 2007.** Unexpected mechanism of symbiont-induced reversal of insect sex: feminizing *Wolbachia* continuously acts on the butterfly *Eurema hecabe* during larval development. *Applied and Environmental Microbiology* **73**(13):4332–4341
DOI [10.1128/AEM.00145-07](https://doi.org/10.1128/AEM.00145-07).
- Narita S, Shimajiri Y, Nomura M. 2009.** Strong cytoplasmic incompatibility and high vertical transmission rate can explain the high frequencies of *Wolbachia* infection in Japanese populations of *Colias erate poliographus* (Lepidoptera: Pieridae). *Bulletin of Entomological Research* **99**(4):385–391 DOI [10.1017/S0007485308006469](https://doi.org/10.1017/S0007485308006469).
- Negri I, Pellicchia M, Mazzoglio PJ, Patetta A, Alma A. 2006.** Feminizing *Wolbachia* in *Zyginidia pullula* (Insecta, Hemiptera), a leafhopper with an XX/X0 sex-determination system. *Proceedings of the Royal Society B: Biological Sciences* **273**(1599):2409–2416
DOI [10.1098/rspb.2006.3592](https://doi.org/10.1098/rspb.2006.3592).
- Newton ILG, Clark ME, Kent BN, Bordenstein SR, Qu J, Richards S, Kelkar YD, Werren JH. 2016.** Comparative genomics of two closely related *Wolbachia* with different reproductive effects on hosts. *Genome Biology and Evolution* **8**(5):1526–1542 DOI [10.1093/gbe/evw096](https://doi.org/10.1093/gbe/evw096).
- Nice CC, Gompert Z, Forister ML, Fordyce JA. 2009.** An unseen foe in arthropod conservation efforts: the case of *Wolbachia* infections in the Karner blue butterfly. *Biological Conservation* **142**:3137–3146 DOI [10.1016/j.biocon.2009.08.020](https://doi.org/10.1016/j.biocon.2009.08.020).
- Nikoh N, Tanaka K, Shibata F, Kondo N, Hizume M, Shimada M, Fukatsu T. 2008.** *Wolbachia* genome integrated in an insect chromosome: evolution and fate of laterally transferred endosymbiont genes. *Genome Research* **18**(2):272–280 DOI [10.1101/gr.7144908](https://doi.org/10.1101/gr.7144908).

- Nyholm SV, Graf J. 2012. Knowing your friends: invertebrate innate immunity fosters beneficial bacterial symbioses. *Nature Reviews Microbiology* 10(12):815–827 DOI 10.1038/nrmicro2894.
- Oliver KM, Degnan PH, Hunter MS, Moran NA. 2009. Bacteriophages encode factors required for protection in a symbiotic mutualism. *Science* 325(5943):992–994 DOI 10.1126/science.1174463.
- Oliver KM, Russell JA, Moran NA, Hunter MS. 2003. Facultative bacterial symbionts in aphids confer resistance to parasitic wasps. *Proceedings of the National Academy of Sciences of the United States of America* 100(4):1803–1807 DOI 10.1073/pnas.0335320100.
- Owen DF. 1965. Change in sex ratio in an African butterfly. *Nature* 206(4985):744 DOI 10.1038/206744a0.
- Owen DF. 1970. Inheritance of sex ratio in the butterfly *Acraea encedon*. *Nature* 225(5233):662–663 DOI 10.1038/225662a0.
- O'Neill SL, Hoffmann AA, Werren JH. 1998. *Influential Passengers: Inherited Microorganisms and Arthropod Reproduction*. Oxford: Oxford University Press.
- Paredes JC, Herren JK, Schüpfer F, Marin R, Claverol S, Kuo C-H, Lemaitre B, Béven L. 2015. Genome sequence of the *Drosophila melanogaster* male-killing *Spiroplasma* strain MSRO endosymbiont. *mBio* 6(2):e02437-14 DOI 10.1128/mBio.02437-14.
- Parnesan C, Ryrholm N, Stefanescu C, Hill JK, Thomas CD, Descimon H, Huntley B, Kaila L, Kullberg J, Tammaru T, Tennent WJ, Thomas JA, Warren M. 1999. Poleward shifts in geographical ranges of butterfly species associated with regional warming. *Nature* 399(6736):579–583 DOI 10.1038/21181.
- Patricelli D, Sielezniew M, Ponikwicka-Tyszko D, Ratkiewicz M, Bonelli S, Barbero F, Witek M, Buś MM, Rutkowski R, Balletto E. 2013. Contrasting genetic structure of rear edge and continuous range populations of a parasitic butterfly infected by *Wolbachia*. *BMC Evolutionary Biology* 13(1):14 DOI 10.1186/1471-2148-13-14.
- Perlman SJ, Kelly SE, Hunter MS. 2008. Population biology of cytoplasmic incompatibility: maintenance and spread of *Cardinium* symbionts in a parasitic wasp. *Genetics* 178(2):1003–1011 DOI 10.1534/genetics.107.083071.
- Piel J, Höfer I, Hui D. 2004. Evidence for a symbiosis island involved in horizontal acquisition of pederin biosynthetic capabilities by the bacterial symbiont of *Paederus fuscipes* beetles. *Journal of Bacteriology* 186(5):1280–1286 DOI 10.1128/jb.186.5.1280-1286.2004.
- Poinsot D, Charlat S, Mercot H. 2003. On the mechanism of *Wolbachia*-induced cytoplasmic incompatibility: confronting the models with the facts. *BioEssays* 25(3):259–265 DOI 10.1002/bies.10234.
- Poulton EB. 1914. W. A. Lamborn's breeding experiments upon *Acraea encedon* (Linn.), in the Lagos District of West Africa, 1910–1912. *Zoological Journal of the Linnean Society* 32(218):391–416 DOI 10.1111/j.1096-3642.1914.tb01463.x.
- Poulton EB. 1923. All female families of *Hypolimnas bolina* L., bred in Fiji by H. W. Simmonds. *Proceedings of the Royal Entomological Society London* 9–12.
- Poulton EB. 1926. Sex-ratio of *Hypolimnas bolina* in Viti Levu, Fiji. *Proceedings of the Royal Entomological Society London* 1:29–32.
- Rigaud T, Juchault P. 1993. Conflict between feminizing sex ratio distorters and an autosomal masculinizing gene in the terrestrial isopod *Armadillidium vulgare* Latr. *Genetics* 133:247–252.
- Rigaud T, Juchault P, Mocquard JP. 1997. The evolution of sex determination in isopod crustaceans. *BioEssays* 19(5):409–416 DOI 10.1002/bies.950190508.

- Riparbelli MG, Giordano R, Ueyama M, Callaini G. 2012. *Wolbachia*-mediated male killing is associated with defective chromatin remodeling. *PLOS ONE* 7(1):e30045 DOI 10.1371/journal.pone.0030045.
- Russell JA, Funaro CF, Giraldo YM, Goldman-Huertas B, Suh D, Kronauer DJC, Moreau CS, Pierce NE. 2012. A veritable menagerie of heritable bacteria from ants, butterflies, and beyond: broad molecular surveys and a systematic review. *PLOS ONE* 7(12):e51027 DOI 10.1371/journal.pone.0051027.
- Russell JA, Goldman-Huertas B, Moreau CS, Baldo L, Stahlhut JK, Werren JH, Pierce NE. 2009. Specialization and geographic isolation among *Wolbachia* symbionts from ants and lycaenid butterflies. *Evolution* 63(3):624–640 DOI 10.1111/j.1558-5646.2008.00579.x.
- Russell JA, Moran NA. 2006. Costs and benefits of symbiont infection in aphids: variation among symbionts and across temperatures. *Proceedings of the Royal Society B: Biological Sciences* 273(1586):603–610 DOI 10.1098/rspb.2005.3348.
- Sahoo RK, Lohman DJ, Wahlberg N, Müller CJ, Brattström O, Collins SC, Peggie D, Aduse-Poku K, Kodandaramaiah U. 2018. Evolution of *Hypolimnys* butterflies (Nymphalidae): Out-of-Africa origin and *Wolbachia*-mediated introgression. *Molecular Phylogenetics and Evolution* 123:50–58 DOI 10.1016/j.ympev.2018.02.001.
- Sakamoto Y, Hirai N, Tanikawa T, Yago M, Ishii M. 2011. Infection by two strains of *Wolbachia* and sex ratio distortion in a population of the endangered butterfly *Zizina emelina* (Lepidoptera: Lycaenidae) in Northern Osaka prefecture, Central Japan. *Annals of the Entomological Society of America* 104(3):483–487 DOI 10.1603/AN09168.
- Sakamoto H, Ishikawa Y, Sasaki T, Kikuyama S, Tatsuki S, Hoshizaki S. 2005. Transinfection reveals the crucial importance of *Wolbachia* genotypes in determining the type of reproductive alteration in the host. *Genetical Research* 85(3):205 DOI 10.1017/S0016672305007573.
- Sakamoto H, Kageyama D, Hoshizaki S, Ishikawa Y. 2007. Sex-specific death in the Asian corn borer moth (*Ostrinia furnacalis*) infected with *Wolbachia* occurs across larval development. *Genome* 50(7):645–652 DOI 10.1139/g07-041.
- Salunke BK, Salunke RC, Dhotre DP, Walujkar SA, Khandagale AB, Chaudhari R, Chandode RK, Ghate HV, Patole MS, Werren JH, Shouche YS. 2012. Determination of *Wolbachia* diversity in butterflies from Western Ghats, India, by a multigene approach. *Applied and Environmental Microbiology* 78(12):4458–4467 DOI 10.1128/AEM.07298-11.
- Sasaki T, Ishikawa H. 1999. *Wolbachia* infections and cytoplasmic incompatibility in the Almond moth and the Mediterranean Flour moth. *Zoological Science* 16(5):739–744 DOI 10.2108/zsj.16.739.
- Sasaki T, Kubo T, Ishikawa H. 2002. Interspecific transfer of *Wolbachia* between two Lepidopteran insects expressing cytoplasmic incompatibility: a *Wolbachia* variant naturally infecting *Cadra cautella* causes male killing in *Ephesia kuehniella*. *Genetics* 162:1313–1319.
- Sasaki T, Massaki N, Kubo T. 2005. *Wolbachia* variant that induces two distinct reproductive phenotypes in different hosts. *Heredity* 95(5):389–393 DOI 10.1038/sj.hdy.6800737.
- Satterfield DA, Maerz JC, Altizer S. 2015. Loss of migratory behaviour increases infection risk for a butterfly host. *Proceedings of the Royal Society B: Biological Sciences* 282(1801):20141734 DOI 10.1098/rspb.2014.1734.
- Scarborough CL, Ferrari J, Godfray HCJ. 2005. Aphid protected from pathogen by endosymbiont. *Science* 310(5755):1781 DOI 10.1126/science.1120180.
- Shao Y, Chen B, Sun C, Ishida K, Hertweck C, Boland W. 2017. Symbiont-derived antimicrobials contribute to the control of the Lepidopteran gut microbiota. *Cell Chemical Biology* 24(1):66–75 DOI 10.1016/j.chembiol.2016.11.015.

- Sharon G, Segal D, Ringo JM, Hefetz A, Zilber-Rosenberg I, Rosenberg E. 2010. Commensal bacteria play a role in mating preference of *Drosophila melanogaster*. *Proceedings of the National Academy of Sciences of the United States of America* **107**(46):20051–20056 DOI [10.1073/pnas.1009906107](https://doi.org/10.1073/pnas.1009906107).
- Smith DAS, Gordon IJ, Allen JA. 2010. Reinforcement in hybrids among once isolated semispecies of *Danaus chrysippus*(L.) and evidence for sex chromosome evolution. *Ecological Entomology* **35**:77–89 DOI [10.1111/j.1365-2311.2009.01143.x](https://doi.org/10.1111/j.1365-2311.2009.01143.x).
- Smith DA, Gordon IJ, Depew LA, Owen DF. 1998. Genetics of the butterfly *Danaus chrysippus* (L.) in a broad hybrid zone, with special reference to sex ratio, polymorphism and intragenomic conflict. *Biological Journal of the Linnean Society* **65**(1):1–40 DOI [10.1006/bijl.1998.0240](https://doi.org/10.1006/bijl.1998.0240).
- Smith DAS, Gordon IJ, Traut W, Herren J, Collins S, Martins DJ, Saitoti K, Ileri P, French-Constant R. 2016. A neo-W chromosome in a tropical butterfly links colour pattern, male-killing, and speciation. *Proceedings of the Royal Society B: Biological Sciences* **283**(1835):20160821 DOI [10.1098/rspb.2016.0821](https://doi.org/10.1098/rspb.2016.0821).
- Snook RR, Cleland SY, Wolfner ME, Karr TL. 2000. Offsetting effects of *Wolbachia* infection and heat shock on sperm production in *Drosophila simulans*: analyses of fecundity, fertility and accessory gland proteins. *Genetics* **155**:167–178.
- Stouthamer R, Breeuwer JAJ, Hurst GDD. 1999. *Wolbachia pipientis*: microbial manipulator of arthropod reproduction. *Annual Review of Microbiology* **53**(1):71–102 DOI [10.1146/annurev.micro.53.1.71](https://doi.org/10.1146/annurev.micro.53.1.71).
- Sugimoto TN, Fujii T, Kayukawa T, Sakamoto H, Ishikawa Y. 2010. Expression of a doublesex homologue is altered in sexual mosaics of *Ostrinia scapularis* moths infected with *Wolbachia*. *Insect Biochemistry and Molecular Biology* **40**(12):847–854 DOI [10.1016/j.ibmb.2010.08.004](https://doi.org/10.1016/j.ibmb.2010.08.004).
- Sugimoto TN, Ishikawa Y. 2012. A male-killing *Wolbachia* carries a feminizing factor and is associated with degradation of the sex-determining system of its host. *Biology Letters* **8**(3):412–415 DOI [10.1098/rsbl.2011.1114](https://doi.org/10.1098/rsbl.2011.1114).
- Sugimoto TN, Kayukawa T, Shinoda T, Ishikawa Y, Tsuchida T. 2015. Misdirection of dosage compensation underlies bidirectional sex-specific death in *Wolbachia*-infected *Ostrinia scapularis*. *Insect Biochemistry and Molecular Biology* **66**:72–76 DOI [10.1016/j.ibmb.2015.10.001](https://doi.org/10.1016/j.ibmb.2015.10.001).
- Sutton ER, Harris SR, Parkhill J, Sinkins SP. 2014. Comparative genome analysis of *Wolbachia* strain wAu. *BMC Genomics* **15**(1):928 DOI [10.1186/1471-2164-15-928](https://doi.org/10.1186/1471-2164-15-928).
- Tabata J, Hattori Y, Sakamoto H, Yukuhiro F, Fujii T, Kugimiya S, Mochizuki A, Ishikawa Y, Kageyama D. 2011. Male killing and incomplete inheritance of a novel *Spiroplasma* in the moth *Ostrinia zaguliaevi*. *Microbial Ecology* **61**(2):254–263 DOI [10.1007/s00248-010-9799-y](https://doi.org/10.1007/s00248-010-9799-y).
- Tagami Y, Miura K. 2004. Distribution and prevalence of *Wolbachia* in Japanese populations of Lepidoptera. *Insect Molecular Biology* **13**(4):359–364 DOI [10.1111/j.0962-1075.2004.00492.x](https://doi.org/10.1111/j.0962-1075.2004.00492.x).
- Teixeira L, Ferreira A, Ashburner M. 2008. The bacterial symbiont *Wolbachia* induces resistance to RNA viral infections in *Drosophila melanogaster*. *PLOS Biology* **6**(12):e1000002 DOI [10.1371/journal.pbio.1000002](https://doi.org/10.1371/journal.pbio.1000002).
- Telschow A, Engelstadter J, Yamamura N, Hammerstein P, Hurst GDD. 2006. Asymmetric gene flow and constraints on adaptation caused by sex ratio distorters. *Journal of Evolutionary Biology* **19**(3):869–878 DOI [10.1111/j.1420-9101.2005.01049.x](https://doi.org/10.1111/j.1420-9101.2005.01049.x).
- Telschow A, Hammerstein P, Werren JH. 2005. The effect of *Wolbachia* versus genetic incompatibilities on reinforcement and speciation. *Evolution* **59**(8):1607–1619 DOI [10.1111/j.0014-3820.2005.tb01812.x](https://doi.org/10.1111/j.0014-3820.2005.tb01812.x).
- Thompson JN. 1987. Symbiont-induced speciation. *Biological Journal of the Linnean Society* **32**(5):385–393 DOI [10.1111/j.1095-8312.1987.tb00439.x](https://doi.org/10.1111/j.1095-8312.1987.tb00439.x).

- Triant DA, Cinel SD, Kawahara AY. 2018.** Lepidoptera genomes: current knowledge, gaps and future directions. *Current Opinion in Insect Science* 25:99–105 DOI 10.1016/j.cois.2017.12.004.
- Tsuchida T, Koga R, Horikawa M, Tsunoda T, Maoka T, Matsumoto S, Simon J-C, Fukatsu T. 2010.** Symbiotic bacterium modifies aphid body color. *Science* 330(6007):1102–1104 DOI 10.1126/science.1195463.
- Vala F, Egas M, Breeuwer JAJ, Sabelis MW. 2004.** *Wolbachia* affects oviposition and mating behaviour of its spider mite host. *Journal of Evolutionary Biology* 17(3):692–700 DOI 10.1046/j.1420-9101.2003.00679.x.
- Vavre F, Fleury F, Lepetit D, Fouillet P, Bouletreau M. 1999.** Phylogenetic evidence for horizontal transmission of *Wolbachia* in host-parasitoid associations. *Molecular Biology and Evolution* 16(12):1711–1723 DOI 10.1093/oxfordjournals.molbev.a026084.
- Vavre FF, Girin CC, Boulétreau MM. 1999.** Phylogenetic status of a fecundity-enhancing *Wolbachia* that does not induce thelytoky in *Trichogramma*. *Insect Molecular Biology* 8(1):67–72 DOI 10.1046/j.1365-2583.1999.810067.x.
- Veneti Z, Bentley JK, Koana T, Braig HR, Hurst GDD. 2005.** A functional dosage compensation complex required for male killing in *Drosophila*. *Science* 307(5714):1461–1463 DOI 10.1126/science.1107182.
- Veneti Z, Toda MJ, Hurst GD. 2004.** Host resistance does not explain variation in incidence of male-killing bacteria in *Drosophila bifasciata*. *BMC Evolutionary Biology* 4:52 DOI 10.1186/1471-2148-4-52.
- Wallin IE. 1927.** *Symbiogenesis and the Origin of Species*. Baltimore: Williams & Wilkins.
- Ward CM, Baxter SW. 2017.** Draft genome assembly of a *Wolbachia* endosymbiont of *Plutella australiana*. *Genome Announcements* 5(43):e01134-17 DOI 10.1128/genomea.01134-17.
- Watanabe K, Yukuhiro F, Matsuura Y, Fukatsu T, Noda H. 2014.** Intrasperm vertical symbiont transmission. *Proceedings of the National Academy of Sciences of the United States of America* 111(20):7433–7437 DOI 10.1073/pnas.1402476111.
- Weeks AR, Stouthamer R. 2004.** Increased fecundity associated with infection by a cytophaga-like intracellular bacterium in the predatory mite, *Metaseiulus occidentalis*. *Proceedings of the Royal Society B: Biological Sciences* 271(Suppl_4):S193–S195 DOI 10.1098/rsbl.2003.0137.
- Weeks AR, Turelli M, Harcombe WR, Reynolds KT, Hoffmann AA. 2007.** From parasite to mutualist: rapid evolution of *Wolbachia* in natural populations of *Drosophila*. *PLOS Biology* 5(5):e114 DOI 10.1371/journal.pbio.0050114.
- Weinert LA, Araujo-Jnr EV, Ahmed MZ, Welch JJ. 2015.** The incidence of bacterial endosymbionts in terrestrial arthropods. *Proceedings of the Royal Society B: Biological Sciences* 282(1807):20150249 DOI 10.1098/rspb.2015.0249.
- Wernegreen JJ. 2012.** Mutualism meltdown in insects: bacteria constrain thermal adaptation. *Current Opinion in Microbiology* 15(3):255–262 DOI 10.1016/j.mib.2012.02.001.
- Werren JH. 1998.** *Wolbachia* and speciation. In: Howard DJ, Berlocher SH, eds. *Endless Forms: Species and Speciation*. New York: Oxford University Press, 245–260.
- Werren JH, Richards S, Desjardins CA, Niehuis O, Gadau J, Colbourne JK, Nasonia Genome Working Group. 2010.** Functional and evolutionary insights from the genomes of three parasitoid *Nasonia* species. *Science* 327(5963):343–348 DOI 10.1126/science.1178028.
- Werren JH, Windsor D, Guo L. 1995.** Distribution of *Wolbachia* among neotropical arthropods. *Proceedings of the Royal Society B: Biological Sciences* 262(1364):197–204 DOI 10.1098/rspb.1995.0196.

- Werren JH, Zhang W, Guo LR. 2004. Evolution and phylogeny of *Wolbachia*: reproductive parasites of arthropods. *Proceedings of the Royal Society B: Biological Sciences* 261(1360): 55–63 DOI 10.1098/rspb.1995.0117.
- Wolinska J, King KC. 2009. Environment can alter selection in host–parasite interactions. *Trends in Parasitology* 25(5):236–244 DOI 10.1016/j.pt.2009.02.004.
- Wybouw N, Dermauw W, Tirry L, Stevens C, Grbić M, Feyereisen R, Van Leeuwen T. 2014. A gene horizontally transferred from bacteria protects arthropods from host plant cyanide poisoning. *eLife* 3:e02365 DOI 10.7554/elife.02365.
- Xie J, Vilchez I, Mateos M. 2010. *Spiroplasma* bacteria enhance survival of *Drosophila hydei* attacked by the parasitic wasp *Leptopilina heterotoma*. *PLOS ONE* 5(8):e12149 DOI 10.1371/journal.pone.0012149.
- Xu P, Liu Y, Graham RI, Wilson K, Wu K. 2014. Densovirus is a mutualistic symbiont of a global crop pest (*Helicoverpa armigera*) and protects against a Baculovirus and Bt biopesticide. *PLOS Pathogens* 10(10):e1004490 DOI 10.1371/journal.ppat.1004490.
- Yamazaki K, Sugiura S, Fukasawa Y. 2004. Epizootics and behavioral alteration in the arctiid caterpillar *Chionarctia nivea* (Lepidoptera: Arctiidae) caused by an entomopathogenic fungus, *Entomophaga aulicae* (Zygomycetes: Entomophthorales). *Entomological Science* 7(3):219–223 DOI 10.1111/j.1479-8298.2004.00066.x.
- Yen JH, Barr AR. 1971. New hypothesis of the cause of cytoplasmic incompatibility in *Culex pipiens* L. *Nature* 232(5313):657–658 DOI 10.1038/232657a0.
- Yen JH, Barr AR. 1973. The etiological agent of cytoplasmic incompatibility in *Culex pipiens*. *Journal of Invertebrate Pathology* 22(2):242–250 DOI 10.1016/0022-2011(73)90141-9.
- Zug R, Hammerstein P. 2018. Evolution of reproductive parasites with direct fitness benefits. *Heredity* 120(3):266–281 DOI 10.1038/s41437-017-0022-5.