

Comparison of cryptobenthic reef fish communities among microhabitats in the Red Sea (#21327)

1

First submission

Editor guidance

Please submit by **15 Dec 2017** for the benefit of the authors (and your \$200 publishing discount).



Structure and Criteria

Please read the 'Structure and Criteria' page for general guidance.



Custom checks

Make sure you include the custom checks shown below, in your review.



Raw data check

Review the raw data. Download from the [materials page](#).



Image check

Check that figures and images have not been inappropriately manipulated.

Privacy reminder: If uploading an annotated PDF, remove identifiable information to remain anonymous.

Files

Download and review all files from the [materials page](#).

4 Figure file(s)

6 Raw data file(s)

2 Other file(s)

! Custom checks

DNA data checks

- ! Have you checked the authors [data deposition statement](#)?
- ! Can you access the deposited data?
- ! Has the data been deposited correctly?
- ! Is the deposition information noted in the manuscript?

Vertebrate animal usage checks

- ! Have you checked the authors [ethical approval statement](#)?
- ! Were the experiments necessary and ethical?
- ! Have you checked our [animal research policies](#)?

Field study

- ! Have you checked the authors [field study permits](#)?
- ! Are the field study permits appropriate?



Structure and Criteria

Structure your review

The review form is divided into 5 sections.
Please consider these when composing your review:

1. BASIC REPORTING
2. EXPERIMENTAL DESIGN
3. VALIDITY OF THE FINDINGS
4. General comments
5. Confidential notes to the editor

 You can also annotate this PDF and upload it as part of your review

When ready [submit online](#).

Editorial Criteria

Use these criteria points to structure your review. The full detailed editorial criteria is on your [guidance page](#).

BASIC REPORTING

-  Clear, unambiguous, professional English language used throughout.
-  Intro & background to show context. Literature well referenced & relevant.
-  Structure conforms to [Peerj standards](#), discipline norm, or improved for clarity.
-  Figures are relevant, high quality, well labelled & described.
-  Raw data supplied (see [Peerj policy](#)).

EXPERIMENTAL DESIGN

-  Original primary research within [Scope of the journal](#).
-  Research question well defined, relevant & meaningful. It is stated how the research fills an identified knowledge gap.
-  Rigorous investigation performed to a high technical & ethical standard.
-  Methods described with sufficient detail & information to replicate.

VALIDITY OF THE FINDINGS

-  Impact and novelty not assessed. Negative/inconclusive results accepted. *Meaningful* replication encouraged where rationale & benefit to literature is clearly stated.
-  Data is robust, statistically sound, & controlled.
-  Conclusions are well stated, linked to original research question & limited to supporting results.
-  Speculation is welcome, but should be identified as such.

Standout reviewing tips

3



The best reviewers use these techniques

Tip

Support criticisms with evidence from the text or from other sources

Example

Smith et al (J of Methodology, 2005, V3, pp 123) have shown that the analysis you use in Lines 241-250 is not the most appropriate for this situation. Please explain why you used this method.

Give specific suggestions on how to improve the manuscript

Your introduction needs more detail. I suggest that you improve the description at lines 57- 86 to provide more justification for your study (specifically, you should expand upon the knowledge gap being filled).

Comment on language and grammar issues

The English language should be improved to ensure that an international audience can clearly understand your text. Some examples where the language could be improved include lines 23, 77, 121, 128 – the current phrasing makes comprehension difficult.

Organize by importance of the issues, and number your points

1. Your most important issue
2. The next most important item
3. ...
4. The least important points

Please provide constructive criticism, and avoid personal opinions

I thank you for providing the raw data, however your supplemental files need more descriptive metadata identifiers to be useful to future readers. Although your results are compelling, the data analysis should be improved in the following ways: AA, BB, CC

Comment on strengths (as well as weaknesses) of the manuscript

I commend the authors for their extensive data set, compiled over many years of detailed fieldwork. In addition, the manuscript is clearly written in professional, unambiguous language. If there is a weakness, it is in the statistical analysis (as I have noted above) which should be improved upon before Acceptance.

Comparison of cryptobenthic reef fish communities among microhabitats in the Red Sea

Emily M Troyer^{Corresp., 1}, Darren J Coker¹, Michael L Berumen¹

¹ Red Sea Research Center, Division of Biological and Environmental Sciences and Engineering, King Abdullah University of Science and Technology, Thuwal, Saudi Arabia

Corresponding Author: Emily M Troyer
Email address: emily.troyer@kaust.edu.sa

Knowledge of community structure within an ecosystem is essential when trying to understand the function and importance of the system as well as when making management decisions related to this system. Within the larger ecosystem, microhabitats play an important role by providing inhabitants with a subset of available resources. On coral reefs, cryptobenthic fishes encompass many groups and make up an important proportion of the biodiversity. However, these fishes are relatively small and extremely cryptic, either behaviorally or visually and therefore are often overlooked. The largest family of cryptobenthic fishes is Gobiidae, a family currently containing more than 1600 species, although new species are continuously discovered. Many goby species are associated with a very specific microhabitat type, however most of this knowledge is limited to coral-dependent species. We examined the differences in fish community structure within three common reef microhabitats (live hard coral, dead coral rubble, and sand) using rotenone stations in the central Red Sea. Using a combination of morphological and genetic (COI barcoding) techniques, we identified 326 individuals representing 73 species spread across 17 families from the collections. The largest group collected was gobies, representing 232 individuals and 31 species. Goby assemblages in the three microhabitats were significantly different from each other – rubble microhabitats hosted the majority of collected gobies (69% of individuals), followed by live hard coral (20.6%), then sand (9.9%). These results provide essential baseline information about the ecology of understudied cryptobenthic fishes that can be used in future large-scale studies in the Red Sea region. Future reef assessments should also incorporate cryptobenthic fishes, as they are often ignored despite their potential functional importance.

Comparison of cryptobenthic reef fish communities among microhabitats in the Red Sea

Short title: Red Sea cryptobenthic fishes

Emily M. Troyer^{1*}, Darren J. Coker¹, Michael L. Berumen¹

¹Red Sea Research Center, Division of Biological and Environmental Sciences and Engineering,
King Abdullah University of Science and Technology, Thuwal 23955-6900, Saudi Arabia

*corresponding author: emily.troyer@kaust.edu.sa

ABSTRACT

Knowledge of community structure within an ecosystem is essential when trying to understand the function and importance of the system as well as when making management decisions related to this system. Within the larger ecosystem, microhabitats play an important role by providing inhabitants with a subset of available resources. On coral reefs, cryptobenthic fishes encompass many groups and make up an important proportion of the biodiversity. However, these fishes are relatively small and extremely cryptic, either behaviorally or visually and therefore are often overlooked. The largest family of cryptobenthic fishes is Gobiidae, a family currently containing more than 1600 species, although new species are continuously discovered. Many goby species are associated with a very specific microhabitat type, however most of this knowledge is limited to coral-dependent species. We examined the differences in fish community structure within three common reef microhabitats (live hard coral, dead coral rubble, and sand) using rotenone stations in the central Red Sea. Using a combination of morphological and genetic (COI barcoding) techniques, we identified 326 individuals representing 73 species spread across 17 families from the collections. The largest group collected was gobies, representing 232 individuals and 31 species. Goby assemblages in the three microhabitats were significantly different from each other – rubble microhabitats hosted the majority of collected gobies (69% of individuals), followed by live hard coral (20.6%), then sand (9.9%). These results provide essential baseline information about the ecology of understudied cryptobenthic fishes that can be used in future large-scale studies in the Red Sea region. Future reef assessments should also incorporate cryptobenthic fishes, as they are often ignored despite their potential functional importance.

INTRODUCTION

Habitat has frequently been shown to influence species abundances and distribution patterns in a range of ecosystems (Venier & Fahrig, 1996; Warren et al., 2001). Within global tropical regions, coral reefs provide complex habitats that support around 25% of marine fish species, despite occupying only 0.1% of ocean area (Spalding et al., 2001). On coral reefs, benthic composition and coral cover may influence the associated fish assemblages (Messmer et al., 2011; Chong-Seng et al., 2012; Komyakova, Munday, & Jones, 2013), with live coral habitats having a direct effect on many species through the provision of shelter and food (Bell & Galzin, 1984; Buchheim & Hixon, 1992; Cole, Pratchett & Jones, 2008; Coker, Wilson & Pratchett, 2014). In addition to biotic habitats, the physical structure and internal complexity of the reef has been shown to influence a number of reef fishes (Graham & Nash, 2013). Within coral reefs, different types of microhabitats exist, such as areas dominated by living hard corals, soft corals, rubble patches, macroalgae, or sandy areas. These habitats can offer a range of resources such as food and shelter for small fishes (Beukers & Jones, 1997; Depczynski & Bellwood, 2004,  Brooker, Munday & Ainsworth, 2010). For example, corals provide refuge spaces within the branches (Robertson & Sheldon, 1979), while the colony itself can provide shelter underneath for larger fishes (Kerry & Bellwood, 2012). In addition, degraded and structurally deteriorated reefs provide additional habitats that are inhabited by species that are better adapted to these habitat features (Ahmadia, Pezold & Smith, 2012; Coker, Graham & Pratchett, 2012). It is therefore expected that small and benthic-associated fishes would be influenced greatly by available habitats, and therefore understanding microhabitat requirements is essential for reef fishes.

On coral reefs, there are many groups of fishes (blennies, gobies, triplefins, etc.) that are

68 relatively small (< 50 mm), have a close association with the substrate, and have a cryptic nature
 69 (Depczynski & Bellwood, 2003). These fishes, termed “cryptobenthic reef fishes”, can be
 70 behaviorally cryptic by seeking out cracks and crevices in the reef in which to hide, or they can
 71 be visually cryptic, having coloration that matches the substrate where they live (Depczynski &
 72 Bellwood, 2003). Fast-growing and with naturally short lifespans, these fishes are an important
 73 functional group on coral reefs, contributing greatly to the transfer of energy through the food
 74 chain. For example, in the Great Barrier Reef up to 8% of the cryptobenthic fish population are
 75 consumed by predators every day (Depczynski & Bellwood, 2006). Despite their functional
 76 importance, they are difficult to sample due to their cryptic nature and therefore vastly
 77 understudied worldwide. Previous visual surveys have not been able to account for the majority
 78 of cryptobenthic fishes on a reef (Ackerman & Bellwood, 2000; Robertson & Smith-Vaniz,
 79 2008) and most surveys ignore them due to logistical and taxonomic difficulties. Thus, more
 80 targeted surveying techniques have recently been employed in order to gain a more accurate
 81 count of cryptobenthic reef fishes. Chemical ichthyocides, such as rotenone and clove oil, have
 82 proven to be an effective tool to collect small fishes and have been shown to reveal more cryptic
 83 species when compared to traditional visual surveys (Brock, 1982). Many groups of fish are
 84 considered to be cryptobenthic but within this group the largest (most speciose) is the family
 85 Gobiidae, which currently contains over 1600 species spread across 200 genera (Thacker, 2003;
 86 Tornabene et al., 2013). Gobies occupy a variety of habitats including corals, seagrass, sand, and
 87 rubble (Munday, Jones & Caley, 1997; Munday et al., 2002; Patzner et al., 2011). The majority
 88 of gobies are epibenthic or cryptobenthic, living on or just above the substrate. Gobies are small
 89 fish, with most attaining an adult size of less than 3 cm (Patzner et al., 2011). They will also have
 90 small home ranges, often no more than 0.25 to 2 m² (Depczynski & Bellwood, 2004). Their

91 small size, coupled with a small home range, has enabled gobies to make use of a multitude of
 92 different microhabitats. Many species of goby are habitat specialists and coral-obligate gobies
 93 have been highly studied (e.g., Munday, Jones & Caley, 1997, 2001; Munday, 2002). Other
 94 groups of cryptobenthic fishes should be studied to determine more broadly understand the
 95 importance of various habitats. 

96 Globally, knowledge of cryptobenthic fishes is still limited. A majority of studies have
 97 arisen from well-studied areas, such as the Great Barrier Reef and the Caribbean (see Ackerman
 98 & Bellwood, 2000, 2002; Harborne et al., 2012), but an important region that is greatly
 99 understudied when it comes to cryptobenthic fishes as well as conspicuous ones (Berumen et al.,
 100 2013), is the Red Sea. The Red Sea is a unique environment, with higher average temperatures
 101 and salinity compared to other regions.  It is recognized as a biodiversity hotspot with an
 102 estimated 14% of fish species in the Red Sea endemic to the region (DiBattista et al., 2016).
 103 Recent studies in this region show that cryptobenthic fish communities differ latitudinally and
 104 with distance from shore, and that habitat may be a driving factor (Coker et al., 2017).

105 This study aims to examine differences among the community of cryptobenthic reef
 106 fishes associated with three common reef microhabitats. Live hard corals, rubble patches, and
 107 sandy areas all represent habitats that cryptobenthic fishes are known to utilize (Depczynski &
 108 Bellwood, 2004; Ahmadi et al., 2015).  and these habitats may also be representative of different
 109 stages of reef degradation. It is hypothesized that cryptobenthic fish assemblages will
 110 significantly differ between these three microhabitats. This information will help us better
 111 understand abundance and distribution patterns within the region and provide baseline data about
 112 understudied cryptobenthic fishes and their functional role in the Red Sea.

113

MATERIALS AND METHODS

Study site

Fishes were sampled in May 2017 during daylight hours from the southern portion of a reef situated near the edge of the continental shelf (Al Fahal, 22° 13.6558 N, 38° 58.1853 E) (Fig. 1 A). The specific study site was located on the eastern side of the reef and thus is protected from the typical north-western wave patterns. Al Fahal was chosen as a sample site because it is a relatively large reef that hosts the variety of habitat types used in this study. This single reef (and wave exposure type) was selected to minimize any potential environmental variables, such as temperature, wave energy, current, and turbidity that could be introduced if sampling in varying parts of the reef (e.g., sides exposed to or sheltered from dominant wave action).

Microhabitat definitions

Three microhabitat types were sampled: live hard coral, dead coral rubble, and sand (Fig. 1 B-D). A hard coral microhabitat was defined as having at least 70% of the quadrat covered with hard coral; tabular *Acropora* corals were targeted. A rubble microhabitat was defined as a flat or gently sloping area (<10°) containing at least 50% coverage of dead hard coral rubble, rocks, or empty shells. A sand microhabitat was defined as a flat or gently sloping area (<10°) containing at least 90% sand coverage containing little or no rubble. For each of the three microhabitat types, five replicate quadrats were sampled, resulting in a total of 15 quadrats.

Percent cover of each microhabitat category was calculated using a grid point system. A 10 x10 grid (i.e., 100 total intercepting points) was overlaid onto a digital image of each quadrat. Each point in the image was categorized into the above microhabitat type. From these tallies, a total percent cover for each microhabitat type was estimated for each quadrat.

Rugosity of the sample quadrat was measured as a linear distance using a 1m chain (sensu Risk, 1972), wherein the chain is draped in a straight line through the quadrat following the vertical contour of the substrate. Subsequently, the 'overhead' view (i.e., two-dimensional horizontal length) of the chain is measured, and the rugosity value is calculated by dividing the length of the chain by the length of the 'overhead' view of the draped chain. A value of 1 would thus represent a flat surface, while higher values indicate a more three-dimensionally complex habitat. This method was repeated three times within each quadrat to provide a mean rugosity for each quadrat.

Fish community sampling

An ichthyocide (rotenone) was used to collect fish due to its high success rate in targeting cryptic species (Robertson & Smith-Vaniz, 2008). A rotenone mixture was prepared by mixing 500g of 4% rotenone powder (Consolidated Chemical Company) with 100ml 96% ethanol, 250ml liquid dishwashing detergent, and 100ml water (adapted from Ackerman & Bellwood, 2000).

Quadrats (1 m²) constructed from PVC pipe (25mm diameter, see Fig. 1) were placed at 10-15m depth onto a flat or gently sloping surface (<10°) of the reef in one of the three targeted microhabitats. A 4mm mesh net weighted down with chain was placed around the quadrat, enclosing the whole area. The rotenone mixture was gently squirted into the netted area until the whole area was covered. After waiting about five minutes for the rotenone to take effect, the net was removed and fishes were collected using handnets and tweezers (following Ackerman & Bellwood, 2000). Because some fish were observed escaping the enclosed quadrat (through the reef matrix) while the rotenone took effect, all deceased fishes directly around the sample area

were collected for consistency. After collecting the visible fishes, three divers intensively searched within each quadrat by lifting up any rubble or debris where hidden fishes could have settled. The quadrats were searched until no new fishes were found for a period of five minutes. Larger predatory fishes that came too close to the quadrat were chased away before they could consume any of the asphyxiated fishes.. 

Immediately after the dive, collected fishes were placed into an ice slurry to preserve coloration for photography (within ~ 2 hrs). Once photographed, total body length and standard length measurements (to the nearest mm) were taken. Fishes were then placed into individual labeled vials containing a solution of 96% ethanol for preservation. A tissue sample (pectoral or caudal fin) was collected from each individual for genetic analysis.

Fish sampling was done in accordance with the guidelines and procedures approved under the auspices of the King Abdullah University of Science and Technology (KAUST) Institutional Animal Use and Care Committee (IACUC) under approval number 17-04-004.

Genetic fish identification

Tissue samples were cleaned with 96% ethanol and gently patted dry before placed into 96-well plates containing 100µl 50mM NaOH. DNA was extracted from the samples using the HotSHOT protocol (95°C for 20 minutes, 4°C for 10 minutes) (Meeker et al., 2007). After extraction, 10µl of 1M Tris-Hcl (10%) was added to each well and mixed with pipettes.

80µl from each sample was transferred to a new plate for DNA amplification. A polymerase chain reaction (PCR) Qiagen Multiplex Mastermix containing Taq polymerase, dNTPs, MgCl₂ and reaction buffers was added to each well. Primers COI Universal Fish R2 5' ACT TCA GGG TGA CCG AAG AAT CAG AA 3' and F2 5' TCG ACT AAT CAT AAA GAT

ATC GGC AC 3' (Ward et al., 2005) were used to amplify the COI region of DNA. Each PCR well contained 6.25µl MasterMix, 4.25µl nuclease free water, 0.5µl forward primer, 0.5µl reverse primer, and 1µl DNA, for a total volume of 12.5µl. Thermocycling occurred with an initial denaturation period of 15 minutes at 95 °C, followed by 35 cycles of 30 seconds at 94 °C, 60 seconds at the annealing temperature of 45 °C, and 60 seconds at 72 °C, followed by a final extension period of 10 minutes at 72 °C. PCR products were visualized using a QIAxcel system. Following PCR amplification, 2.14µl ExoStar was added to each well to clean the PCR product. The mixture was incubated in the thermocycler for 60 minutes at 37 °C, then 15 minutes at 85 °C. Cleaned PCR products were sequenced via Sanger sequencing via ABI 3730xl sequencers in the KAUST Bioscience Core Lab.

In the event that a sample sequence did not yield a long enough strand of base pairs to be entered into a database (~500 bp), DNA was extracted again using a more precise Qiagen DNeasy Blood and Tissue kit. The PCR process was then repeated on those samples.

After sequencing was completed, sample sequences were checked against several sequence databases for potential matches. The National Center for Biotechnology Information (NCBI) GenBank (<https://www.ncbi.nlm.nih.gov/genbank/>) and Barcode of Life Data System (BOLD) (<http://www.boldsystems.org>) were used as public databases. A custom (in-house) Red Sea fish sequence database was also checked (see Coker et al., 2017; DiBattista et al., 2017; Isari et al., 2017).  A sequence was considered to be a good match for a species in the database if the

matched sequence was 98% similar or higher. Sequence matches were then checked against visual guides and morphological keys to double-check identity. In the event of a non-matching sequence, a sample was identified to species level, or as close as possible, with the use of keys that examined morphological features (see next section). When a species was unable to be

identified using the aforementioned methods, it was assigned an operational taxonomic unit (OTU) so as to still be included in the data analysis. OTUs were named for previous uploaded sequences in the databases if there was a match. In the event a sequence did not match a previously assigned OTU within one of the databases, a new OTU was assigned.

Morphological fish identification

With the use of keys, fishes that could not be identified genetically were assigned to a species, genus, or family level. The typical morphological characters that were useful in discriminating taxa included: counts of fin spines and soft rays, pelvic fin structure, cephalic pore counts, lateral line scale counts, as well as general meristics. Morphological characters were assessed using light dissection microscopy and an online image analysis tool for measuring morphometrics (Froese & Pauly, 2017). For a list of keys used, see Appendix S1.

Community analysis

For each quadrat, several metrics were calculated. Species richness was defined as the number of species or OTUs identified from each quadrat. Abundance was determined by the total number of individuals collected in each quadrat. Diversity (Shannon's diversity index, H') was calculated using the following formula: $H' = -\sum p_i \ln(p_i)$ (Shannon & Weaver, 1949), and then subsequently averaged within each microhabitat type. The fish communities present at each microhabitat type were plotted using non-metric multidimensional scaling (nMDS) using a Bray-Curtis resemblance matrix. Analyses were conducted using R version 3.4.0 (R Core Team, 2017) and the vegan package (Oksanen et al., 2017). Statistics used included analysis of variance (ANOVA) and permutational analysis of variance (PERMANOVA).

RESULTS

Fish communities

A total of 326 individuals representing 73 species and 17 families were collected from three microhabitat types (Appendix S2). The total number of all fishes collected at each quadrat ranged from 1 (Quadrat sand_4) to 65 (Quadrat rubble_3). Rubble quadrats had the highest average numbers of fish abundances (one-way ANOVA $F_{2,12}=11.59$, $p=0.002$), species richness ($F_{2,12}=6.78$, $p=0.011$), and diversity ($F_{2,12}=2.80$, $p=0.100$), followed by coral quadrats, and then sand quadrats (Fig. 2) and fish communities in all three microhabitats differed significantly (PERMANOVA $F_{2,12}=2.61$, $p<0.001$) (Fig. 3). Out of the 326 fish collected, the family Gobiidae comprised the majority, with a total of 232 individuals (71.1%) collected representing 31 species. The next most common families were Pseudochromidae, with 33 individuals (10.1%) collected representing 12 species, and Pomacentridae with 16 individuals (4.9%) representing 7 species.

Goby communities

The number of individual gobies collected at each microhabitat type followed patterns similar to that of all collected fishes. Abundance ranged from 0 gobies (Quadrat sand_4) to 46 gobies (Quadrat rubble_3). The most abundant goby was *Trimma avidori*, with a total of 44 individuals. This goby was collected in greatest abundance in both coral and rubble microhabitats. The second most abundant goby collected was *Callogobius bifasciatus* with 29 individuals collected in mainly rubble quadrats (Fig. 4). Of 31 species of goby collected, 11 species (35.4%) were represented by only a single individual, and the majority of collected species (64.5%) had less than 5 individuals collected.



252 Out of the coral quadrats, the most abundant goby collected was *T. avidori*. Both *C.*
 253 *bifasciatus* and *A. semipunctata* were the most abundant species found in rubble, and *Istigobius*
 254 *decoratus* was the most abundant species found in the sand microhabitat quadrats.

255 On average, rubble communities had the highest levels of goby abundance (one way
 256 ANOVA $F_{2,12}=24.61$, $p<0.001$), species richness ($F_{2,12}=14.95$, $p<0.001$), and diversity
 257 ($F_{2,12}=5.10$, $p=0.025$), followed by coral communities, and then sand communities (Fig. 2).
 258 Overall, goby community composition significantly differed among microhabitats
 259 (PERMANOVA $F_{2,12}=3.67$, $p<0.001$) (Fig. 3).

260 Of all the fish collected, 25.7% could not be confidently assigned to a species using either
 261 morphological or genetic techniques, so they were given an OTU. Several candidates for new
 262 species were identified from these OTUs; some of these are possibly Red Sea endemics. COI 
 263 sequences for new OTUs have been deposited to GenBank under accession numbers MG583518-
 264 MG583524, as well as sequences from *Bryaninops natans*, *Gobiodon reticulatus*, and *Trimma*
 265 *flavicaudatum*, all which currently have no COI sequence information on GenBank.

266

267 Habitat composition

268 Rugosity was highest on average in coral microhabitats (Fig. 1 E). Rubble and sand
 269 microhabitats were less rugose, with rubble quadrats having higher rugosity, on average, than
 270 sand (one-way ANOVA, $F_{2,10}=19.86$, $p<0.001$). Percent cover of each microhabitat type was
 271 highest for sand quadrats, with sand quadrats averaging 97.6% (± 0.66 SE) cover of sand. Coral
 272 microhabitat quadrats averaged 85.3% (± 4.83 SE) cover of all hard corals, with tabular
 273 *Acropora* averaging 56.2% cover (± 9.56 SE) within the coral microhabitat quadrats. Rubble
 274 microhabitat quadrats averaged 65.6% cover (± 4.43 SE) of rubble (Fig. 1 F).

275 DISCUSSION

276 Cryptobenthic fish communities, including gobies, are vastly understudied compared to other,
 277 more conspicuous groups, especially in the Red Sea. Fish communities from live hard coral,
 278 rubble, and sand microhabitats differed from each other by having differing levels of abundance,
 279 species richness, and diversity.  ~~Research is still just scratching the surface of cryptobenthic fish~~
 280 ~~ecology and this study found that habitat is a strong indicator of what kind of cryptobenthic fish~~
 281 ~~assemblages will be present on a reef in the Red Sea. Importantly, these results provide a~~
 282 ~~framework for future studies in the region to examine additional microhabitats in additional~~
 283 ~~locations.~~

284 The rubble microhabitats sampled yielded the highest levels of fish abundance, richness,
 285 and diversity. A possible explanation for this preference is the sediment load that is deposited
 286 onto the rubble. Runoff and detritus from more productive areas on a reef will eventually settle
 287 onto the substrate below, delivering more nutrients to the benthos (Crossman et al., 2001; Wilson
 288 et al., 2003; Depczynski & Bellwood, 2004; Goatley, González-Cabello & Bellwood, 2016).
 289 Cryptobenthic fishes will primarily feed upon microscopic prey such as epibenthic invertebrates,
 290 algae, or detritus (Depczynski & Bellwood, 2003), and this food is readily available on the
 291 substrate. Indeed, in the rubble microhabitats sampled, there was a presence of epilithic algae on
 292 many of the large pieces of rubble. *Asterropteryx semipunctata*, one of the most abundant species
 293 collected from the rubble microhabitats, is a known detritivore (Depczynski & Bellwood, 2003).
 294 Because cryptobenthic fishes are  dentary and will not forage far from their home areas
 295 (Depczynski & Bellwood, 2004), they must live in places that have a steady supply of food.

296 Coral microhabitats also yielded high levels of fish abundance, richness, and diversity.
 297 Coral quadrats sampled were characterized by a large percentage of live hard coral cover, with a

majority of the coral cover represented by tabular *Acropora*. These coral habitats also exhibited the highest measures of rugosity. There is a positive correlation between habitat complexity and fish abundance in both conspicuous fishes (Caley & St John, 1996; Friedlander & Parrish, 1998) as well as cryptobenthic fishes (Depczynski & Bellod, 2004). Complexity offers shelter from predators as well as lowered mortality rates for juveniles (Beukers & Jones, 1997). One of the most abundant fish collected from coral microhabitats was *Trimma avidori*. This goby is known to inhabit small overhangs or caves underneath coral structures (Herler & Hilgers, 2005), and it is likely that the large tabular *Acropora* present in the coral quadrats provided adequate shelter for these fishes.

Sand microhabitats exhibited the lowest levels of fish abundance, richness, and diversity. These areas were characterized by a lack of biotic structures and were overall quite barren. The sand quadrats were extremely flat with low rugosity and no complexity, and this may have been a contributing factor to the low levels of abundance, species richness, and diversity found within. Structural complexity has long been shown to influence fish abundance and diversity (Risk, 1972; Luckhurst & Luckhurst, 1978). Cryptic fishes rely on hiding places to avoid being preyed upon and unlike the coral habitats, the sand microhabitats had little, if any, areas to hide. However, sand habitats are not completely devoid of life. Many cryptobenthic specialists are able to thrive in sandy habitats due to less competition from other fishes and the ability to form internal tunnels within the substrate. Flatfish, such as flounders and soles, are able to blend in perfectly with the coloration and the extreme flatness of sand, and many wrasses are known to burrow underneath it to escape predation. These microhabitats may be further used by cryptobenthic fishes for foraging at night. Our daytime sampling regime precludes us for assessing this possibility.

The microhabitat types used in this study represent microhabitats nearly universal to coral reefs that are important to fish assemblages. However, it is important to note that the hard coral, rubble, and sand microhabitats in the Red Sea may not be functionally equivalent to those worldwide. For instance, coral species will vary from region to region, rubble size and detrital material can vary, and sand sediment grain size and composition may vary as well. Because the three microhabitats used in this study were found to be different from each other, it provides a good foundation for future research in this region. Future studies may address the coarseness of sand grains across multiple sand microhabitats or the size of rubble across multiple rubble microhabitats to determine if there are finer-scale differences that may determine microhabitat use among cryptobenthic fishes. Additional factors not considered here, such as exposure to wave action or distance from shore (Depczynski & Bellwood, 2005; Goatley, González-Cabello & Bellwood, 2016) may also warrant further investigation in the Red Sea.

Cryptobenthic fishes are an understudied group; hence there is a lack of public sequence information available for many species, especially gobies. For 232 collected gobies, using the 98% similarity threshold, only 128 (55.2%) gobies were able to be identified genetically using COI barcoding. The remaining 104 (44.8%) could not be identified using genetic methods, and were either identified using morphological characteristics or given an OTU. This gap in publicly available cryptobenthic fish sequences is a hindrance for identification that relies on genetic data. These difficulties with identifying cryptobenthic fishes highlight the need for a comprehensive sequence library that includes reference images.

The Red Sea is a known biological hotspot with high rates of endemism for many groups of species. An estimated 14% of Red Sea fishes are endemic (DiBattista et al., 2016), however these estimates may under-represent cryptic species. Out of 73 fish species, a total of 17 (23.2%)

endemic species from the Red Sea and Arabian Peninsula were found in this study. Four of those endemic species were gobies (see DiBattista et al., 2016 for checklist). It is also extremely likely that some of the unnamed goby OTUs are undescribed species, and these could possibly be endemic to the region. However, endemic status could only be more confidently asserted if supported by more sampling in adjacent regions. With the potential number of undiscovered cryptobenthic fishes in the region, the number of known endemics can be expected to increase in time.

CONCLUSIONS



This study provides baseline information on cryptobenthic fish communities in the central Red Sea and their microhabitat preferences. Three distinct microhabitats were found to host different communities of fishes, with different abundance, richness, and diversity levels. Rubble microhabitats were found to have the highest levels of abundance, richness, and diversity overall, followed by hard coral microhabitats, and finally sand microhabitats. We are only just beginning to understand the ecology of cryptobenthic fishes and such fine-scale habitat partitioning is likely to contribute to the high diversity found on a reef. On a single reef, there can be distinct microhabitat communities that are within meters of each other, yet support different fish assemblages. More data is needed to accurately quantify the cryptobenthic biodiversity of the region. Future larger-scale projects may find more evidence of microhabitat associations in the Red Sea. Because cryptobenthic fishes are estimated to compose up to half of all fishes on a reef and previous studies have found only a fraction of that proportion, more microhabitat types, reef parts, and depths should be sampled. By expanding the parameters of a cryptobenthic fish study, it is likely that knowledge of these cryptic fishes in the region will increase.



367 **ACKNOWLEDGEMENTS**

368 The authors would like to thank Michael Campbell for his mapmaking expertise, as well as
 369 Calder Atta, Royale Hardenstine, Alison Monroe, Tullia Terraneo, Matthew Tietbohl, and Sara
 370 Wilson for their assistance collecting fishes. We are also grateful to Calder Atta, Simon Brandl,
 371 and Luke Tornabene for their help in identifying several fish species. Fieldwork was supported
 372 by the KAUST Coastal and Marine Resources Core Laboratory.

REFERENCES

- Ackerman JL, Bellwood DR (2000) Reef fish assemblages: a re-evaluation using enclosed rotenone stations. *Mar Ecol Prog Ser* 206:227-237
- Ackerman JL, Bellwood DR (2002) Comparative efficiency of clove oil vs rotenone for sampling tropical reef fish assemblages. *J Fish Biol* 60:893-901
- Ahmadia, GN, Pezold FL, Smith DJ (2012) Cryptobenthic fish biodiversity and microhabitat use in healthy and degraded coral reefs in SE Sulawesi, Indonesia. *Mar Biodivers* 42:433-442
- Ahmadia GN, Sheard LJ, Pezold FL, Smith DJ (2012) Cryptobenthic fish assemblages across the coral reef-seagrass continuum in SE Sulawesi, Indonesia. *Aquat Biol* 16:125-135
- Bell JD, Galzin R (1984) Influence of live coral cover on coral-reef fish communities. *Mar Ecol Prog Ser* 15: 265-274
- Berumen ML, Hoey A, Bass W, Bouwmeester J, Catania D, Cochran JE, Khalil MT, Miyake S, Mughal MR, Spaet J (2013) The status of coral reef ecology research in the Red Sea. *Coral Reefs* 32:737-748
- Beukers JS, Jones GP (1997) Habitat complexity modifies the impact of piscivores on a coral reef fish population. *Oecologia* 114:50-59.
- Brock RE (1982) A critique of the visual census method for assessing coral reef fish populations. *B Mar Sci* 32:269-276
- Brooker RM, Munday PL, Ainsworth TD (2010) Diets of coral-dwelling fishes of the genus *Gobiodon* with evidence of corallivory. *J Fish Biol* 76:2578-2583
- Buchheim JR, Hixon MA (1992) Competition for shelter holes in the coral-reef fish *Acanthemblemaria spinosa* Metzelaar. *J Exp Mar Biol Ecol* 164:45-54

396 Caley JM, St John J (1996) Refuge availability structures assemblages of tropical reef fishes. *J*
 397 *Anim Ecol* 65:414-428

398 Chong-Seng KM, Mannering TD, Pratchett MS, Bellwood DR, Graham NAJ (2012) The
 399 influence of coral reef benthic condition on associated fish assemblages. *PLoS ONE* 7(8):
 400 e42167

401 Coker DJ, DiBattista JD, Sinclair-Taylor TH, Berumen ML (2017) Spatial patterns of
 402 cryptobenthic coral-reef fishes in the Red Sea. *Coral Reefs* DOI:10.1007/s00338-017-
 403 1647-9

404 Coker DJ, Graham NAJ, Pratchett MS (2012) Interactive effects of live coral and structural
 405 complexity on the recruitment of reef fishes. *Coral Reefs* 31:919-927

406 Coker DJ, Wilson SK, Pratchett MS (2014) Importance of live coral habitats for reef fishes. *Rev*
 407 *Fish Biol Fisher* 24:89-126

408 Cole AJ, Pratchett MS, Jones GP (2008) Diversity and functional importance of coral-feeding
 409 fishes on tropical coral reefs. *Fish and Fisheries* 9:286-307

410 Crossman DJ, Choat JH, Clements KD, Hardy T, McConochie J (2001) Detritus as food for
 411 grazing fishes on coral reefs. *Limnol Oceanogr* 46:1596-1605

412 Depczynski M, Bellwood DR (2003) The role of cryptobenthic reef fishes in coral reef
 413 trophodynamics. *Mar Ecol Prog Ser* 256:183–191

414 Depczynski M, Bellwood DR (2004) Microhabitat utilisation patterns in cryptobenthic coral reef
 415 fish communities. *Mar Biol* 145:455-463

416 Depczynski M, Bellwood DR (2005) Wave energy and spatial variability in community structure
 417 of small cryptic coral reef fishes. *Mar Ecol Prog Ser* 303:283-293

418 Depczynski M, Bellwood DR (2006) Extremes, plasticity, and invariance in vertebrate life

419 history traits: insights from coral reef fishes. *Ecology* 87: 3119-3127

420 DiBattista JD, Roberts MB, Bouwmeester J, Bowen BW, Coker DJ, Lozano-Cortes DF, Choat
 421 JH, Gaither MR, Hobbs JPA, Khalil MT, Kochzius M, Myers RF, Paulay G, Robizch
 422 VSN, Saenz-Agudelo P, Salas E, Sinclair-Taylor TH, Toonen RJ, Westneat MW,
 423 Williams ST, Berumen ML (2016) A review of contemporary patterns of endemism for
 424 shallow water reef fauna in the Red Sea. *J Biogeogr* 43:423-439

425 DiBattista JD, Coker DJ, Sinclair-Taylor TH, Stat M, Berumen ML, Bunce M (2017) Assessing
 426 the utility of eDNA as a tool to survey reef-fish communities in the Red Sea. *Coral Reefs*
 427 DOI:10.1007/s00338-017-1618-1

428 Friedlander AM, Parrish JD (1998) Habitat characteristics affecting fish assemblages on a
 429 Hawaiian reef. *J Exp Mar Biol Ecol* 224:1-30

430 Froese R, Pauly D (2017) FishBase. World Wide Web electronic publication. www.fishbase.org,
 431 version (06/2017)

432 Goatley CHR, González-Cabello A, Bellwood DR (2016) Reef-scale partitioning of
 433 cryptobenthic fish assemblages across the Great Barrier Reef, Australia. *Mar Ecol Prog*
 434 Ser 544:271-280

435 Graham NAJ, Nash KL (2013) The importance of structural complexity in coral reef ecosystems.
 436 *Coral Reefs* 32:315-326

437 Harborne AR, Jelks HL, Smith-Vaniz WF, Rocha LA (2012) Abiotic and biotic controls of
 438 cryptobenthic fish assemblages across a Caribbean seascape. *Coral Reefs* 31:977-990

439 Herler J, Hilgers H (2005) A synopsis of coral and coral-rock associated gobies (Pisces:
 440 Gobiidae) in the Gulf of Aqaba, northern Red Sea. *Aqua* 10:103-132

441 Isari S, Pearman JK, Casas L, Michell CT, Curdia J, Berumen ML, Irigoien X (2017) Exploring

the larval fish community of the central Red Sea with an integrated morphological and molecular approach. PloS ONE 12:e0182503

Kerry JT, Bellwood DR (2012) The effect of coral morphology on shelter selection by coral reef fishes. Coral Reefs 31:415-424

Komyakova V, Munday PL, Jones GP (2013) Relative importance of coral cover, habitat complexity and diversity in determining the structure of reef fish communities. PloS ONE 8:e83178

Luckhurst BE, Luckhurst K (1978) Analysis of the influence of substrate variables on coral reef fish communities. Mar Biol 49:317-323

Meeker ND, Hutchinson SA, Ho L, Trede NS (2007) Method for isolation of PCR-ready genomic DNA from zebrafish tissues. Biotechniques 43:610

Messmer V, Jones GP, Munday PL, Holbrook SJ, Schmitt RJ, Brooks AJ (2011) Habitat biodiversity as a determinant of fish community structure on coral reefs. Ecology 92:2285-2298

Munday PL (2002) Does habitat availability determine geographical-scale abundance of coral-dwelling fishes? Coral Reefs 21:105-116

Munday PL, Jones GP, Caley MJ (1997) Habitat specialisation and the distribution and abundance of coral-dwelling gobies. Mar Ecol Prog Ser 152:227-239

Munday PL, Jones GP, Caley MJ (2001) Interspecific competition and coexistence in a guild of coral-dwelling fishes. Ecology 82:2177-2189

Munday PL, Pierce SJ, Jones GP, Larson HK (2002) Habitat use, social organization and reproductive biology of the seawhip goby, *Bryaninops youngei*. Mar Freshwater Res

464 53:769-775

465 Oksanen J, Blanchet FG, Friendly M, Kindt R, Legendre P, McGlinn D, Minchin PR, O'Hara
 466 RB, Simpson GL, Solymos P, Stevens MHH, Szoecs E, Wagner H (2017) vegan:
 467 Community Ecology Package. R package version 2.4-3.
 468 <https://CRAN.R-project.org/package=vegan>

469 Patzner RA, Van Tassell JL, Kovačić M, Kapoor BG. Eds. (2011) *The Biology of Gobies*.
 470 Enfield, NH: Science Publishers

471 R Core Team (2017) R: A language and environment for statistical computing. R Foundation for
 472 Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>

473 Risk MJ (1972) Fish diversity on a coral reef in the Virgin Islands. *Atoll Res Bull* 153:1-7

474 Robertson DR, Sheldon JM (1979) Competitive interactions and the availability of sleeping sites
 475 for a diurnal coral reef fish. *J Exp Mar Biol Ecol* 40:285-298

476 Robertson DR, Smith-Vaniz WF (2008) Rotenone: an essential but demonized tool for assessing
 477 marine fish diversity. *Bioscience* 58:165-170

478 Shannon CW, Weaver W (1963) *Mathematical theory of communication*. University Illinois
 479 Press

480 Spalding MD, Ravilious C, Green EP (2001) *World atlas of coral reefs*. Univ. of California
 481 Press, Berkeley, USA

482 Thacker CE (2003) Molecular phylogeny of the gobioid fishes (Teleostei: Perciformes:
 483 Gobioidae). *Mol Phylogenet Evol* 26:354-368

484 Tornabene L, Ahmadi GN, Berumen ML, Smith DJ, Jompa J, Pezold F (2013) Evolution of
 485 microhabitat association and morphology in a diverse group of cryptobenthic coral reef
 486 fishes (Teleostei: Gobiidae: *Eviota*). *Mol Phylogenet Evol* 66:391-400

Venier LA, Fahrig L (1996) Habitat availability causes the species abundance-distribution relationship. *Oikos* 76: 564-570

Ward RD, Zemlak TS, Innes BH, Last PR (2005) DNA barcoding Australia's fish species. *Philos Trans R Soc Lond B Biol Sci* 360:1847-1857

Warren MS, Hill JK, Thomas JA, Asher J, Fox R, Huntley B, Roy DB, Telfer MG, Jeffcoate S, Harding P, Jeffcoate G, Willis SG, Greatorex-Davies JN, Moss D, Thomas CD (2001) Rapid responses of British butterflies to opposing forces of climate and habitat change. *Nature* 414:65-69

Wilson SK, Bellwood DR, Choat JH, Furnas MJ (2003) Detritus in the epilithic algal matrix and its use by coral reef fishes. *Oceanogr Mar Biol Annu Rev* 41:279-309

Figure 1

Microhabitat characteristics of sampled quadrats

Examples of 1m² quadrats sampled in the central Saudi Arabian Red Sea to assess cryptobenthic fish assemblages in each of three microhabitat types: coral (indicated by purple), rubble (orange), and sand (teal). Quadrats were sampled using rotenone at 10-15m depth. (A) Map of study site, with reef habitat indicated in grey. (B) Coral quadrat. (C) Rubble quadrat. (D) Sand quadrat. (E) Average rugosity for each microhabitat type. A value of 1 indicates a completely flat surface while higher values represent greater vertical complexity. (F) Percent cover of the target substrate for each quadrat for each microhabitat type. Lighter colors in the coral quadrats represent the proportion of the coral cover comprised by tabular *Acropora* (visible in panel B). (Photographs taken by EMT. Map created in ArcMap, version 10.3, by Michael Campbell using various mapping sources freely available through ESRI.)




Figure 2

Bar chart of community indices

Abundance (mean number of individuals per $\text{m}^2 \pm \text{SE}$), species richness (mean number of species per $\text{m}^2 \pm \text{SE}$), and diversity (H') (mean value per $\text{m}^2 \pm \text{SE}$) for all collected fish species and for family Gobiidae for each of the three microhabitat types sampled using rotenone stations (1m^2 quadrats, $n = 5$ quadrats per microhabitat type) in the central Saudi Arabian Red Sea.




Figure 3

nMDS plots of community similarities

nMDS plots using Bray-Curtis similarity of (A) all fish communities (i.e., relative abundances of all collected species) and (B) goby communities (i.e., relative abundances of individuals from the family Gobiidae) sampled using rotenone stations in each of the three microhabitat types in the central Saudi Arabian Red Sea. Solid points represent individual quadrats from each microhabitat type, which define the shaded minimum convex polygons for each habitat type. (Note that one quadrat from coral microhabitat and one quadrat from sand microhabitat were excluded from the goby nMDS due to an absence of gobies in these quadrats.) Ellipse size represents 95% confidence limits for each microhabitat type. Overall, total fish community composition significantly differs for each microhabitat (PERMANOVA $F_{2,12}=2.61$, $p<0.001$) and goby community composition significantly differs for each microhabitat (PERMANOVA $F_{2,12}=3.67$, $p<0.001$).





Figure 4

Goby species abundances

Rank-abundance plot of goby species sampled in the central Saudi Arabian Red Sea using 1m² rotenone stations in three microhabitat types (color coded). Vertical bars represent the total combined number of individuals sampled in all quadrats (n = 5 quadrats per microhabitat type) for all 31 goby species found in this study. Coloration of the vertical bars indicates the portion of individuals of each species found in the three microhabitat types.

