

Beetles in bamboo forests: community structure in a heterogeneous landscape of southwestern Amazonia (#9056)

1

First submission

Please read the **Important notes** below, and the **Review guidance** on the next page.
When ready [submit online](#). The manuscript starts on page 3.

Important notes

Editor and deadline

Dezene Huber / 14 Mar 2016

Files

1 Table file(s)

Please visit the overview page to [download and review](#) the files not included in this review pdf.

Declarations

Involves a field study on animals or plants.



Please in full read before you begin

How to review

When ready [submit your review online](#). The review form is divided into 5 sections. Please consider these when composing your review:

1. BASIC REPORTING

2. EXPERIMENTAL DESIGN

3. VALIDITY OF THE FINDINGS






4. General comments

5. Confidential notes to the editor





 You can also annotate this **pdf** and upload it as part of your review

To finish, enter your editorial recommendation (accept, revise or reject) and submit.





BASIC REPORTING

-  Clear, unambiguous, professional English language used throughout.
-  Intro & background to show context. Literature well referenced & relevant.
-  Structure conforms to [PeerJ standard](#), discipline norm, or improved for clarity.
-  Figures are relevant, high quality, well labelled & described.
-  Raw data supplied (See [PeerJ policy](#)).

EXPERIMENTAL DESIGN

-  Original primary research within [Scope of the journal](#).
-  Research question well defined, relevant & meaningful. It is stated how research fills an identified knowledge gap.
-  Rigorous investigation performed to a high technical & ethical standard.
-  Methods described with sufficient detail & information to replicate.

VALIDITY OF THE FINDINGS

-  Impact and novelty not assessed. Negative/inconclusive results accepted. *Meaningful* replication encouraged where rationale & benefit to literature is clearly stated.
-  Data is robust, statistically sound, & controlled.
-  Conclusion well stated, linked to original research question & limited to supporting results.
-  Speculation is welcome, but should be identified as such.

The above is the editorial criteria summary. To view in full visit <https://peerj.com/about/editorial-criteria/>

Beetles in bamboo forests: community structure in a heterogeneous landscape of southwestern Amazonia

Jennifer Jacobs, Rudolf von May, David H. Kavanaugh, Edward M. Connor

Amazonian bamboo forests dominated by large woody bamboo plants in the genus *Guadua* cover approximately 180,000 km² and represent a key resource for many organisms. In southwestern Amazonia, native bamboo forests differ in structure, biodiversity, and growth dynamics from other forest types in the region. However, with the exception of a few species in which habitat specialization or a strong habitat association has been demonstrated, little is known about how bamboo forests influence animal community structure. In an effort to understand more about the animal assemblages associated with Amazonian bamboo forests, we characterized the structure of ground-dwelling beetle assemblages living in bamboo forests and adjacent terra firme forests in a lowland rainforest site in Peru. We conducted intensive pitfall trap surveys in 13 bamboo habitat patches and 13 adjacent terra firme habitat patches to determine if there were differences in the abundance and richness of beetle species in these two habitat types. Additionally, given that southwestern Amazonia experiences distinct dry and wet seasons, we conducted our study during the dry and wet season of one year to account for differences in seasonality. We found a distinct beetle assemblage associated with each forest type, and identified a set of dominant species that significantly contributed to the distinctness in beetle community structure between bamboo and terra firme forest. The terra firme forest had a greater number of rare species than the bamboo forest. Several beetle species exhibited a strong association with the bamboo forest, including a large species of Scarabaeidae that appears to be specializing on bamboo. We also found marked differences in beetle assemblages between dry and wet seasons. Our results support the prediction that beetle community structure in bamboo forest differs from that of terra firme in terms of species richness, abundance, and composition. Bamboo-associated animal communities require more exploration and study, and must be included in regional conservation plans seeking to protect entire animal communities in southwestern Amazonia.

Beetles in bamboo forests: community structure in a heterogeneous landscape of southwestern Amazonia

Jennifer Jacobs^{1,4}, Rudolf von May², David H. Kavanaugh³, Edward F. Connor¹

¹ Department of Biology, San Francisco State University, 1600 Holloway Avenue, San Francisco, CA 94132

² Department of Ecology and Evolutionary Biology, University of Michigan, 830 North University, 2019 Kraus Natural Science Building, Ann Arbor, MI 48109-1048

³ Department of Entomology, California Academy of Sciences, 55 Music Concourse Drive, San Francisco, CA 94118

⁴ Corresponding author; e-mail: jmjacobs.berkeley@gmail.com

Abstract

Amazonian bamboo forests dominated by large woody bamboo plants in the genus *Guadua* cover approximately 180,000 km² and represent a key resource for many organisms. In southwestern Amazonia, native bamboo forests differ in structure, biodiversity, and growth dynamics from other forest types in the region. However, with the exception of a few species in which habitat specialization or a strong habitat association has been demonstrated, little is known about how bamboo forests influence animal community structure. In an effort to understand more about the animal assemblages associated with Amazonian bamboo forests, we characterized the structure of ground-dwelling beetle assemblages living in bamboo forests and adjacent terra firme forests in a lowland rainforest site in Peru. We conducted intensive pitfall trap surveys in 13 bamboo habitat patches and 13 adjacent terra firme habitat patches to determine if there were differences in the abundance and richness of beetle species in these two habitat types. Additionally, given that southwestern Amazonia experiences distinct dry and wet seasons, we conducted our study during the dry and wet season of one year to account for differences in seasonality. We found a distinct beetle assemblage associated with each forest type, and identified a set of dominant species that significantly contributed to the distinctness in beetle community structure between bamboo and terra firme forest. The terra firme forest had a greater number of rare species than the bamboo forest. Several beetle species exhibited a strong association with the bamboo forest, including a large species of Scarabaeidae that appears to be specializing on bamboo. We also found marked differences in beetle assemblages between dry and wet seasons. Our results support the prediction that beetle community structure in bamboo forest differs from that of terra firme in terms of species richness, abundance, and composition. Bamboo-associated animal communities require more exploration and study, and must be included in regional conservation plans seeking to protect entire animal communities in southwestern Amazonia.

Key words

Biodiversity, Coleoptera, *Guadua* bamboo, pitfall traps, terra firme

Introduction

Bamboo forests in southwestern Amazonia cover approximately 180,000 km² and represent the largest bamboo-dominated forest in the Neotropics (Nelson 1994; Griscom and Ashton 2003). These forests are primarily dominated by woody species in the genus *Guadua*, a native bamboo that has been present in the region since the pre-Holocene as suggested by fossil evidence (Olivier et al. 2008), or perhaps earlier periods as suggested by recent phylogenetic studies (Ruíz-Sánchez 2011). Many animal species use bamboo habitat for shelter, foraging, reproduction, or a combination of purposes (Emmons and Feer 1990; Louton et al. 1996; Kratter 1997; Dunnum and Salazar-Bravo 2004; Lebbin 2007; von May et al. 2008; von May et al. 2009a; 2009b; 2010; Jacobs and von May 2012; Jacobs et al. 2012), although most of these species also live in other forest types and only a few are considered to be bamboo specialists (Kratter 1997; Davidson et al. 2006a; 2006b; Lebbin et al. 2007).

The dynamics of bamboo forests is relatively well understood, but little is known about the structuring of animal communities associated with bamboo habitat. Most bamboo in southwestern Amazonia grows in patches ranging from less than a hectare up to tens or even thousands of hectares (Griscom and Ashton 2006). Those patches are typically surrounded by upland terra firme or, less frequently, by floodplain forest. Bamboo patches might flower synchronously and then die over large areas, and bamboo plants may or may not recolonize the same areas (Nelson 1994; Griscom and Ashton 2003; Silman 2003; Nelson and Bianchini 2005; Griscom and Ashton 2006; Olivier 2007; 2008). Thus large patches of bamboo may blink in and out of existence over time and space, presenting a dynamic, relatively short-lived resource for other organisms. In this context, it is relevant to investigate how animal communities are structured in bamboo forest compared to neighboring forest types. This question can be empirically addressed by conducting one or more surveys of species richness, composition and abundance in bamboo forest and neighboring forest habitat.


In an effort to understand more about animal communities associated with bamboo forests in southwestern Amazonia, our primary goal in this study was to quantitatively characterize the structure of ground-dwelling beetle assemblages living in bamboo forest and compare them with those of the adjacent terra firme forest. We selected beetles as focal taxa because they are taxonomically and trophically diverse, play vital roles in ecosystem maintenance, serve as an important food source for many vertebrates, and are relatively easy to collect (Didham 1997; Ward and Lariviere 2004). Previous inventories of particular beetle taxa in a variety of habitats in the southeastern portion of the Peruvian Amazon have included minimal sampling in bamboo forests (Pearson and Derr 1986; Erwin 1997; Larsen et al. 2006).




Until recently, with the exception of a small number of herbivorous beetles that are known to specialize on the *Guadua* plants themselves (T. Erwin, pers. comm.; Anderson 2008) and a large dynastine beetle strongly associated with bamboo (Jacobs et al. 2012), it was not known if other beetle species are closely associated with bamboo forest as a primary habitat, and whether distinct assemblages of beetles are associated with bamboo forests compared to the adjacent terra


firme. Because of the structural differences between these two habitats (Griscom and Ashton 2003; 2006), and the evidence of bamboo specialists in various animal taxa (see summary in Jacobs and von May 2012), we hypothesized that beetle assemblages in bamboo forest would differ from those of terra firme. We predicted that some beetle species would show preferences either for bamboo or terra firme. Additionally, because of seasonal rainfall variation in southeastern Peru, we expected to find differences in beetle community structure between dry and wet seasons. Based on previous findings regarding temporal patterns of insect communities in Neotropical rainforests (Janzen 1973; Pearson and Derr 1986; Richards and Windsor 2007), we hypothesized that beetle species richness and abundance would be greater in the wet season compared to the dry season.

Materials and methods

STUDY SITE – We conducted this study at the Los Amigos Research Center (12°34'07" S, 70°05'57" W; 270 m elev.), in Manu province, Madre de Dios region, southeastern Peru (Fig. 1). The Los Amigos Research center (hereafter referred to as LA) covers approximately 1,000 ha and borders the Los Amigos Conservation Concession, which covers 145,918 ha of undisturbed lowland rainforest including several forest types (von May et al. 2010). Annual rainfall is variable and ranges between 2700 and 3000 mm, and >80% of the rainfall in this region occurs during the October-May wet season. The dry season, which usually occurs between June and September and has precipitation below or around 100 mm/month, is relatively cooler than the wet season (<http://atrium-biodiversity.org>, accessed on 10 April 2014). The mean annual temperature ranges between 21°C and 26°C (Pitman 2008).

DESIGN OF STUDY – We selected four beetle families for analysis: Scarabaeidae, Carabidae, Histeridae, and Curculionidae. These families represent the majority of the individuals captured in our preliminary sampling. These families also span several functional groups, and include relatively large species which simplify uration and identification (Grimbacher and Stork 2009). ~~In addition, we targeted a subset of ground-dwelling and understory taxa from these families that would potentially be caught with pitfall traps.~~ In focusing on these families, we were attempting to collect information on decomposers, fruit, seed, and humus eaters, generalist ground foragers, and generalist predators.

We used a paired sampling design with 13 sites, each consisting of one patch of bamboo and an equal sized adjacent area of terra firme. We followed a proportional sampling approach (Schoereder et al. 2004) in which sampling effort was standardized with respect to patch area for bamboo forest and an equal area for terra firme pitfall trap/hectare). Thus, depending on the area of the patch, bamboo patches had different numbers of traps. The terra firme forest received the same number of traps as their “paired” bamboo patches. We sampled each site during two periods in the dry season (July and August 2006) and two periods in the wet season (January and March 2007) to evaluate the effect of seasonality.

SITE SELECTION We selected bamboo patches that contained dense stands of *Guadua weberbaueri*, few other plants, and leaf litter almost entirely composed of *G. weberbaueri* leaves to define a sample patch (Fig. 2). However, some patches also contained a second bamboo species, *Ichnanthus breviscrobis*. Though a few other intermittently distributed bamboo species

occur at LA, the upland terraces are dominated by *G. weberbaueri* (Olivier 2007). We selected sampling sites using Landsat images and vegetation maps used by previous researchers (Lebbin 2007; M. Tobler, pers. comm.), and ground-based investigations along the entire trail system at LA (>100km). We selected 13 dense patches of bamboo forest, which based on an initial power calculation, was sufficient for statistical analyses. Each bamboo patch and adjacent terra firme habitat (Fig. 3) was considered a site. We estimated the size of the bamboo patch using Arc GIS 9.1 and a pre-existing vegetation map of LA (Daniel Lebbin, pers. comm.). We also verified the GIS-derived measurements of bamboo patch size with ground observations. Every hectare within a selected bamboo patch, and adjacent terra firme, was allocated one pitfall trap, with patches ranging in size between approximately 1 and 25 ha.

SAMPLING METHODS – We used un-baited pitfall traps to sample beetle communities. Pitfall traps are commonly used to catch surface-active invertebrates and are simple, inexpensive, and yield high numbers of specimens (Ward et al. 2001; Work et al. 2002). The advantage of using un-baited pitfall traps is that they represent a passive capture method to evaluate habitat preference as opposed to trapping with baits which may draw animals from greater distances, potentially outside the bamboo patch. Each pitfall trap consisted of two plastic 16-ounce containers (12 cm diameter) stacked together and inserted into the soil so that the top of the containers was flush with the ground. Traps were filled with approximately 0.125 L of a mixture comprised of 50% water and 50% ethanol (95% solution). A roof composed of palm leaves propped on a stick platform was placed 30 cm above each trap in order to prevent the traps from being flooded by rainwater or filled by falling leaves and other objects. The leaves covered the trap from above but allowed plenty of room for insects to enter laterally.

At each site, traps were placed 10–30 m into the bamboo patch, depending on the patch size. We then placed pitfall traps 5 m apart in either a linear or semi-circular array, depending on the shape, size and accessibility of the bamboo patch. Because *Guadua* forms extremely dense thickets and each plant is covered with large spines, it was impossible to access all locations within each patch. We always maintained a 5 m distance between traps regardless of the type of trap array. Our spatial array of traps was relatively similar to trap arrays used in studies by Barbosa and Marquet (2002), Driscoll (2005), and Baker and Barmuta (2006) (Fig. 4). We followed exactly the same procedure for placing traps in the adjacent terra firme. We recorded the location of trap arrays with a Garmin 76 Map GPS in the center portion of the trap array in each habitat at every site. In the largest patch of bamboo (~ 25 ha), three separate transects of pitfall traps were established because the high density of bamboo culms made it impossible to place all 25 pitfall traps in one transect in the same location. Three separate transects were also established in the terra firme for that particular sampling site.

Pitfall traps were open for seven continuous days in both July and August of 2006 during the dry season, and seven continuous days in both January and March of 2007 during the wet season. Each seven-day period was considered a “sampling period”. During the interval between trap openings and closings, traps were monitored by visual inspection in the field for disturbance and functionality. If traps appeared dry, a small amount of ethanol was added. Traps were always examined following a large storm in case of flooding. For each sampling period, we used a total of 150 traps (75 for bamboo and 75 for terra firme), and with four sampling periods we employed a total of 600 traps. For the purpose of analyses and because we were interested in

seasonal patterns, data from July and August were pooled as “dry season” and data from January and March were pooled as “wet season”.

SAMPLE PROCESSING – Following trap collection specimens were cleaned, sorted, organized and preserved in jars containing 95% ethanol. Specimens were pinned, labeled, sorted to morpho-species, and entered into a database. Families, genera, species, and morpho-species were identified with help from a variety of taxonomic resources, including specialists at the California Academy of Sciences, Santa Barbara Natural History Museum, the Smithsonian Institution, and the University of Nebraska (see Acknowledgments). All other beetle specimens and arachnid specimens were kept and set aside for other researchers at the California Academy of Sciences and the Museo de Historia Natural San Marcos in Lima, Peru. Permits to conduct this work and collect specimens were issued by the Instituto Nacional de Recursos Naturales (INRENA), Peru (Research authorizations 053-2005-INRENA-IFFS-DCB, 23-2006-INRENA-IFFS-DCB, 67-2007-INRENA-IFFS-DCB, and 11-2008-INRENA-IFFS).

ANALYSES –To initially compare patterns of beetle species richness as a function of sampling effort between terra firme and bamboo forest, we generated species accumulation curves (EstimateS 7.5; Colwell 2005). We observed whether the confidence intervals of the curves overlapped to help determine if the communities of beetles in both habitats were different in terms of species richness. Although species accumulation curves are often used to determine the completeness of sampling when conducting an inventory (Longino and Colwell 1997; Longino et al. 2002), they are also helpful in gross comparison of communities in terms of diversity. Because we maintained exactly the same sampling effort in both habitats, we compared the curves directly and did not need to use richness estimators (Magurran 2004). If the curve of one beetle community in either forest type was lower in species richness than the other forest type, we interpreted that result as a community pattern, not an artifact of under-sampling.

COMPARISON OF BEETLE ASSEMBLAGES BETWEEN FOREST TYPES – To statistically test for overall differences in mean species richness and abundance between the two habitats, we used a paired *t*-test to compare the species richness and abundance of beetles from all families at each pair of sites, while pooling data from dry and wet seasons. We used log transformations to normalize the data for these analyses, and *t*-tests were conducted in SPSS v. 11.0 (Chicago, IL). We also used rank/abundance plots, also known as Whittaker plots, to illustrate contrasts or similarities in patterns of species richness and qualitatively assess the evenness of assemblages (Magurran 2004; Gardner et al. 2007), to compare species abundance distributions in the two habitats. Prior to creating the Whittaker plots, we normalized our data with log-transformations and pooled data from dry and wet seasons.

To characterize the structure of beetle assemblages found in the bamboo and terra firme, we used species composition and abundance data collected in each habitat. However, we first used a Mantel test to test for a correlation between species composition and geographic distance among sites. We used a Bray–Curtis similarity matrix and coupled this with a matrix of all pairwise distances (in meters) between sites. We used an Excel spreadsheet integrated with PopTools (<http://www.cse.csiro.au/poptools>) to perform Mantel tests using 1000 randomizations of the distance matrix.

To compare beetle community structure in bamboo versus terra firme, while pooling across seasons, we conducted a one-way analysis of similarity (ANOSIM; Clarke and Warwick 1994) using a Bray–Curtis similarity matrix. We used non-metric multidimensional scaling (nMDS; 100 restarts) plots to create graphical representations of ANOSIM results. The distance between points in nMDS plots is proportional to the compositional similarity of those points. Stress values of nMDS plots indicate the “goodness of fit” or quality of the test. Values beginning with zero (perfect fit) through 0.2 are considered effective for interpreting community data. All ANOSIM and nMDS analyses were conducted using a Bray–Curtis similarity matrix based on log-transformed data, with Primer 6.0 (Clarke and Gorley 2006). We also used the application SIMPER (in Primer 6.0), considering each collecting site as a sample and each forest type as a group, to determine which species best characterized each habitat. This procedure initially calculates the species that account for 90% of the abundance of all species analyzed, and from that list, determines the percentages that each species contributes to the dissimilarity between any two groups.

We conducted a second set of multivariate analyses on each beetle family individually to determine if community structure for every family differed depending on forest type. We compared beetle species composition and abundance in bamboo to that of terra firme for each family (ANOSIM). Data were pooled across seasons.

To compare patterns of abundance or presence/absence of individual species between forest types, we created matched rank-occurrence plots with all species that were represented by at least five individuals. We applied this criterion because it is common that samples of tropical forest invertebrates contain many uncommon species and a substantial percentage of singletons (Coddington et al. 2009). This graphical method allowed us to visualize the degree of overlap between forest types for every species, and illustrate which species were absent, rare, or common in our collection (Longino and Colwell 1997; Delsinne et al. 2008); the exclusion of rare species allowed us to better address differences driven by the most abundant species in the community.

COMPARISON BETWEEN WET AND DRY SEASON – We tested for differences in overall mean species richness and abundance between the dry and wet seasons, using paired *t*-tests. To compare species abundance distributions between dry and wet season, we created Whittaker plots. We identified and ordered the ten most abundant species in each season to compare the identity and rank of the most abundant species during the dry and wet seasons. For both analyses, we pooled data from bamboo and terra firme and used log-transformed data.

~~To understand how species composition and abundance differed between dry and wet seasons, each within bamboo and terra firme, we used a variety of analyses.~~ To test for seasonal differences in abundance within each forest type, we calculated the difference in number of individuals collected between dry and wet season at all sites for bamboo forest and for terra firme. We then applied a paired *t*-test to determine which forest type exhibited a greater change in beetle abundance between dry and wet season (SPSS v. 11.0, Chicago, IL). To test for differences in species turnover between seasons in bamboo and terra firme, we performed a similar procedure. We calculated how many species were unique between the dry and wet seasons for each site in the bamboo forest and in terra firme. We then applied a paired *t*-test to the number of unique species in each forest type.

To further explore the effects of habitat and season on beetle community composition, we performed both one-way and two-way ANOSIM analyses. To test for differences in beetle community composition between forest types in the wet and dry seasons individually, we performed one-way ANOSIM tests and associated nMDS plots. To simultaneously compare the effects of forest type and season on beetle community structure, we ran a two-way ANOSIM test using forest type and season as factors (we did not create an nMDS plot for the two-way ANOSIM because it is difficult to interpret given the more complex structure of the two-factor similarity matrix on which the nMDS analysis is based). The patterns of community structure are best visualized by looking at the one-way nMDS plots for the dry and wet seasons individually. In addition, we created matched rank-occurrence plots to compare patterns of abundance and presence/absence for individual species in the dry and wet seasons. For ANOSIM tests, nMDS plots, and matched rank-occurrence plots, we used species represented by at least five individuals.

Results

COMPARISON OF BEETLE ASSEMBLAGES BETWEEN FOREST TYPES – The beetle community in bamboo forest differed from that of terra firme in terms of species richness, abundance and composition. We collected a total of 189 species of beetles and 3752 individuals in our target families. In bamboo forest we trapped 120 species of beetles and 1539 individuals. In the terra firme we trapped 141 species of beetles and 2213 individuals. Forty-eight species were found only in bamboo forest, 69 species were found only in terra firme, and 72 species were found in both forest types. Overall, beetles from the family Scarabaeidae comprised the majority of all specimens captured, accounting for 83% of the total number of individuals and approximately 48% of all species captured. All families except Histeridae were more species rich in the terra firme by 38%, 33%, and 11%, respectively for Carabidae, Curculionidae, and Scarabaeidae. All families except Carabidae were more abundant in the terra firme by 50%, 49% and 45%, respectively for Curculionidae, Histeridae and Scarabaeidae (Table 1 and 2).

Overall, we found significant differences in the mean number of beetle species and individuals in bamboo versus terra firme using data pooled across seasons ($t_{12} = 2.712$, $p = 0.019$; $t_{12} = 2.088$, $p = 0.059$, respectively). The species accumulation curves illustrate that the number of species captured in terra firme was higher than that of bamboo forest, but overlapping confidence intervals indicate that the curves did not differ substantially (Fig. 5).

The species abundance distributions in the Whittaker plots for bamboo and terra firme (Fig. 6a) were similar, but terra firme exhibited higher abundances of the most abundant species and a surplus of rare species in comparison to bamboo forest. Though species abundance patterns were similar between bamboo and terra firme habitat, the order of the most dominant species differed between forest types. The ten most abundant species in order of rank abundance in bamboo forest (and displayed as letters on Fig. 6a) were not all the same species, nor were they in the same rank order as those in terra firme.

We did not detect any autocorrelation between differences in beetle assemblages and the distance between sites (Mantel test - Pearson $r = -0.07888$, $p = 0.225$). Therefore, we proceeded with multivariate analyses of beetle community structure.

We found significant differences in beetle assemblages between bamboo and terra firme pooling across seasons (ANOSIM Global $R = 0.160$, $p = 0.001$). An nMDS plot (Fig. 7) also suggests the presence of two distinct beetle communities in the bamboo and terra firme, though one terra firme sample from one of the smallest plots (1 ha) is clearly an outlier. A set of 24 beetle species are largely responsible for the dissimilarity of beetle community composition between bamboo and terra firme (Table 3). Four of these species were very abundant only in bamboo forests, and eight other species were very abundant only in terra firme. The remaining 12 species were collected in both habitats but in varied abundances.

For species represented by more than five individuals, matched rank-occurrence plots exhibit varying patterns of abundance and presence/absence in the two habitats (Fig. 8a). Many species exhibited greater abundances in only one forest type. Overall, we found a greater number of the most abundant species in the terra firme compared to the bamboo forest. Beetle species in the family Carabidae and Histeridae appear equally abundant in bamboo and terra firme whereas, Curculionidae and Scarabaeidae exhibited a greater number of species with higher abundances in the terra firme (Fig. 8a).

For two beetle families, community structure also differed between bamboo and terra firme. We detected significant differences in beetle assemblages between bamboo and terra firme for Scarabaeidae and Curculionidae (ANOSIM Global $R = 0.190$, $p = 0.002$; Global $R = 0.154$, $p = 0.003$, respectively), but not for Carabidae and Histeridae (Global $R = -0.030$, $p = 0.785$; Global $R = 0.044$, $p = 0.187$, respectively).

SEASONALITY AND FOREST TYPE—With data from bamboo and terra firme pooled, overall mean species richness and number of individuals did not significantly differ between dry and wet seasons (paired $t_{12} = 0.907$, $p = 0.382$; $t_{12} = 1.619$, $p = 0.132$, respectively). However, we collected 10% more individual beetles in the dry season and 35% more species in the wet season. Species abundance distributions were similar for the dry and wet season, but the rank-order of the most abundant species changed with the seasons and a greater number of rare species were detected in the wet season (Fig. 6b).

Seasonal differences in abundance and species turnover were greater in terra firme than in bamboo forest. We found a significant difference between the mean number of individuals collected in the dry versus wet seasons in terra firme, compared to that of bamboo forest, across all sites (paired $t_{12} = 2.951$, $p = 0.012$) (Fig. 9a). Similarly, we found a greater amount of species turnover between the dry versus wet season in terra firme compared to that of bamboo forest ($t_{12} = 4.366$, $p = 0.001$) (Fig. 9b).

Differences in beetle community structure between bamboo and terra firme was more pronounced in the wet season than in the dry season. We found significant differences in beetle community structure in the wet season (ANOSIM Global $R = 0.259$, $p = 0.001$, Figure 10a), with less distinction between forest types in the dry season (ANOSIM Global $R = 0.068$, $p = 0.070$,

Figure 10b). However, the nMDS plot from the dry season illustrates some separation in beetle communities between bamboo and terra firme. One terra firme sampling point that appears to be a distinct outlier (the same site that is an outlier when examining data pooled among seasons, Figure 5), may be influencing the results. The results of the two-way ANOSIM suggest that both forest type and season significantly affect beetle community structure, although the effect of seasonality was slightly stronger than that of forest type (season: Global $R = 0.298$, $p = 0.001$, forest type: Global $R = 0.163$, $p = 0.001$).

Matched rank-occurrence plots illustrate the variation in abundance for individual species, with more than five individuals captured, in the dry and wet seasons (Fig. 8b). Species in the family Carabidae were found primarily in the wet season, Curculionidae and Histeridae did not show strong seasonal variation, and we trapped more species of Scarabaeidae with higher abundances in the dry season. For a complete list of all species, where they were collected (forest type) and when they were collected (season) see [Online Resource 1](#).

Discussion

BEETLE ASSEMBLAGES IN BAMBOO VS. TERRA FIRME – Our results support the prediction that beetle community structure in bamboo forest differs from that of terra firme in terms of species richness, abundance, and composition. We found a greater number of beetle species (17.5%) and individuals (43.8%) in terra firme compared to bamboo forest, but nonetheless found high beetle richness (120 species) and abundance in bamboo forest. Though relative abundance distributions were similar between bamboo and terra firme, each community was characterized by a distinct set of dominant species and terra firme had a greater number of rare species. In analogous study systems comprised of two naturally-occurring, contrasting, and adjacent habitat types, most often in the form of forest and savanna, researchers have reported distinct beetle communities for each habitat (Kotze and Samways 2001; Spector and Ayzama 2003; Yu et al. 2007).

Our most striking results were the large number of beetle species found in bamboo, and that many species were only collected in bamboo forest and not in terra firme. Of the 120 beetle species we captured in the bamboo forest approximately 40% (48 species) were only collected in bamboo. Although further collecting might have yielded some of these species in terra firme, our results support the idea that bamboo is a rich habitat that contains a beetle community distinct from that of the adjacent terra firme. It is likely that some beetle species have evolved to specialize on bamboo forests as has been shown for other organisms (Kratter 1997; Davidson 2006; Kondo and Gullen 2004; Lebbin et al. 2007).

Each beetle family exhibited different patterns of richness, abundance and composition within bamboo and terra firme. Because beetles from the Scarabaeidae constituted the majority of species in our collection (48%), and the majority of individuals (83%), diversity patterns of the scarabs were the most influential on the overall results. Out of the 24 most abundant species that best characterized either bamboo or terra firme, the greatest number of species (19) were scarab beetles (Table 3). All families except Histeridae were more species-rich in terra firme. All families except Carabidae (equally abundant in both forest types) were more abundant in terra firme.

We found significant differences in beetle community structure comparing bamboo and terra firme only for Scarabaeidae and Curculionidae. For Carabidae, it is possible that a low number of individuals (103) compared to the number of species (25), in addition to the high number of singleton and doubleton species in the carabid dataset (20), yielded nonsignificant results in terms of community structure between habitats. Raw data for the Carabidae family suggests that there are differences in beetle assemblages from bamboo and terra firme as there was a large turnover in species between bamboo and terra firme.

We collected many beetle species from every family that were found only in bamboo forest. Fifty-four percent of the carabid beetle species that we collected in bamboo were only from bamboo, followed by 59% for the Curculionidae, 38% for the Histeridae, and 30% for the Scarabaeidae. In addition, we found the greatest number of singletons and doubletons in the Carabidae and Curculionidae which may be influencing the high degree of observed habitat affinity compared to the more abundant families of Histeridae and Scarabaeidae. Carabid and curculionid beetles may also be more sensitive to forest type. Additional, and more targeted sampling is necessary to elucidate habitat affinities for some beetle groups.

A number of studies have shown that invertebrate species richness is correlated with plant species richness (Siemann et al. 1998; Beals 2006; Schaffers et al. 2008). The high plant diversity of terra firme leads one to initially assume that terra firme habitat has more food resources than that of the neighboring bamboo forest, which may increase the number of herbivorous insects and other animals. The higher levels of plant-herbivore diversity could in turn increase animal diversity in higher trophic levels (Hunter and Price 1993; Brose 2003). Perhaps the fact that terra firme forest has more than double the number of tree species compared to adjacent bamboo-dominated forests (Griscom et al. 2007) accounts at least in part, for why we found more beetle species and individuals in terra firme (especially for Carabidae and Curculionidae). However, our results show that overall beetle species richness from terra firme is only 17.5% greater than that of bamboo forest – a proportional difference that, according to the hypothesis above, is smaller than would be expected based on tree diversity patterns. Though we did not inventory tree species in our sample sites, our results suggest that factors other than plant diversity are contributing to the relatively high diversity of beetles in bamboo forests.

For beetles in the families, Carabidae and Curculionidae, which are not often associated with dung or carrion, mechanistic hypotheses are more difficult. However, of our four target beetle families, these two families may be most strongly responding to the higher diversity of trees in the terra firme compared to bamboo forest. The majority of Neotropical carabid diversity occurs in the canopy of trees and the higher tree diversity of the terra firme may help explain the greater diversity of carabid beetles collected in this forest type. In addition, many curculionid beetles are plant, seed, or humus eaters, and thus may also be responding to the higher tree diversity of the terra firme. Although we collected a greater number of species of these two families in the terra firme, there was nonetheless high turnover in species composition between bamboo and terra firme.

SEASONALITY AND FOREST TYPE – Seasonality alone is playing a role in structuring beetle communities, though the effects are mixed for different beetle families. Overall, we found more species in the wet season and more individuals in the dry season. Differences in beetle

community structure between bamboo and terra firme were more pronounced in the wet season. Additionally, terra firme exhibited greater seasonal variation in the number of individuals captured and greater species turnover between seasons. Seasons in southwestern Peru are primarily defined by fluctuations in rainfall, and it is often assumed that increased rainfall will drive an increase in observed richness resulting from increased activity of most insect species (Wolda 1978; Novotny and Basset 1998).

We captured 35% more species in the wet season compared to the dry season, though many of the additional wet season species were singletons (i.e., the long tail in the species abundance distribution in Figure 6b). Based on previous studies (Pearson and Derr 1986; Devries et al. 1999; Richards and Windsor 2007), we also expected to observe higher abundances in the wet season versus the dry season. Interestingly, we captured 10% more individuals in the dry season compared to the wet season. However, it is important to acknowledge that the observed pattern of higher beetle abundance in the dry season is being driven primarily by scarab and histerid beetles – the most abundant families in our study. For the family Carabidae, 84% of all individuals were captured in the wet season. The abundance of curculionid beetles was approximately equal across dry and wet seasons.

While examining the effects of both seasonality and forest type on beetle community structure, our results, along with those from recent studies (Lucky et al. 2002; Grimbacher and Stork 2009), illustrate that temporal patterns of insects may be more complex than originally assumed. Overall, we found that season had a slightly stronger influence than forest type on beetle community structure when simultaneously analyzing the effects of habitat and season. However, when comparing beetle community structure in bamboo and terra firme separately for the dry and wet seasons, we observed that community structure was more highly defined in the wet season compared to the dry season.

In addition, there was more variation in the number of individuals captured, and greater species turnover, between the dry and wet seasons in terra firme compared to bamboo forest. Interestingly, while we found a larger difference in the number of individuals collected between dry and wet seasons in the terra firme, there was not a clear directional pattern. Thus, we collected a greater number of individuals at some sites in the wet season and a greater number of individuals at other sites in the dry season. In contrast, species richness was clearly higher in the wet versus the dry season for terra firme compared to bamboo forest (Figs. 9a and 9b). It is possible that bamboo forest is a more “predictable” habitat than terra firme in terms of food resources. Because terra firme has a much higher diversity of plants than bamboo forest, greater fluctuations in food resources from varying flowering and fruiting phenologies may occur throughout the year. Populations of beetle species that rely directly or indirectly on plant fruiting phenologies may be affected by this variation in terra firme. The potentially higher variation in beetle activity may explain why we observed more fluctuation in the number of individuals captured, and greater species turnover, between dry and wet seasons in terra firme. In contrast, bamboo plants from the genus *Guadua* do not experience the same annual phenological fluctuations compared to other tree species because they flower and set seed approximately every 30 years (Griscom et al. 2007).

Our results show that many beetle species are sensitive to seasonality and forest type. While a handful of beetle species were equally abundant in bamboo and terra firme, across wet and dry seasons, many were not. Other species, such as *Enema pan*, exhibited a strong preference for one forest type in only one season (Online Resource 1). After conducting additional investigations of *E. pan*, we discovered that this species lives in close association with the bamboo plants (Jacobs et al. 2012). These results are important in that this species is possibly impacting bamboo plants through feeding and nesting damage, and because of their large size (~100 mm), is potentially contributing considerably to the biomass of bamboo forest fauna (Fig. 11). In addition to *E. pan*, we found eight species of beetles that were captured in much greater numbers in the bamboo forest versus terra firme, suggesting that these species may also be associated with bamboo forest habitat (Online Resource 1). However, their biology is unknown. Bamboo-associated organisms are being discovered with increasing frequency (Jacobs & von May 2012) and more in-depth field work, throughout the dry and wet seasons, is needed to determine if and why particular insect species are associated with specific forest types. Understanding insect diversity patterns in *Guadua* bamboo forests is particularly important because insects are the food source for so many other animals, and this is especially true for the 19 known species of insectivorous, *Guadua*-specialist birds (Kratler 1997; Socolar et al. 2013).

Guadua bamboo forests are strikingly unique habitats because they occur as mono-dominant forest islands surrounded by hyper-diverse rainforest—providing an uncommon, spatially and biologically distinct habitat. Results from our study clearly show that bamboo forests in Peru maintain different communities of beetles compared to those of adjacent terra firme. Out of the total number of species that we collected in bamboo forest, 40% were collected only in bamboo forest. In addition, our results suggest that the species richness of beetles in terra firme is not considerably higher (17%) than that of bamboo forest—a surprising result given the substantially higher tree diversity in terra firme. Historically, bamboo forests in southwestern Amazonia have been regarded as species-poor, weedy habitats because of their relatively low plant diversity. It is encouraging to see that more research efforts are focusing on this important component of western Amazonian ecosystems. Bamboo-associated animal communities require more exploration and study, and must be included in regional conservation plans seeking to protect entire animal communities in southwestern Amazonia. Furthermore, with increased interest in harvesting natural bamboo or creating bamboo plantations for alternative timber sources, naturally-occurring bamboo forests may be at risk. In southwestern Amazonia, *Guadua* bamboo forests are also threatened by deforestation due to agriculture, illegal mining, and illegal logging. Thus, more studies focusing on the ecology of native bamboo forests and their associated fauna are of immediate and great importance.

Acknowledgments

We thank Jesus Ramos, Nigel Pitman, and the staff at Los Amigos Field Station for facilitating field work logistics. We thank Karina Ramirez for advice with permit applications. We thank Gerardo Lamas for support in the Entomology Department at the Museo de Historia Natural Universidad de San Marcos, Lima, Peru and the Entomology Department at the California Academy of Sciences. We also thank Trond Larsen, Terry Erwin, Michael Caterino, Kipling Will, Brett Ratcliffe, and Charles O'Brien for help with identifying beetle species. We

additionally thank Daniel Lebbin and Jan Olivier for help with describing, selecting and mapping bamboo forest patches at Los Amigos Field Station, and to Bruce Nelson for providing information on Guadua bamboo life history dynamics. And, we thank Don Driscoll for help with preliminary sampling and initial study design.

References

- Anderson RS (2008) A review of the Neotropical genus *Eucalandra* Faust, 1899 (Coleoptera; Curculionidae: Dryophthorinae). *Zootaxa* 1791:57-67
- Baker SC, Barmuta LA (2006) Evaluating spatial autocorrelation and depletion in pitfall-trap studies of environmental gradients. *J of Insect Cons* 10:269-276
- Barbosa O, Marquet, PA (2002) Effects of forest fragmentation on the beetle assemblage at the relict forest of Fray Jorge, Chile. *Oecologia* 132:296-306
- Beals ML (2006) Understanding community structure: a data-driven multivariate approach. *Oecologia* 150:484-495
- Blaum N, Seymour C, Rossmianith E, Schwager M, Jeltsch F (2009) Changes in arthropod diversity along a land use driven gradient of shrub cover in savanna rangelands: identification of suitable indicators. *Biodiversity Conserv* 18:1187-1199
- Brose U (2003) Bottom-up control of carabid beetle communities in early successional wetlands: mediated by vegetation structure of plant diversity? *Oecologia* 135:407-413
- Clarke KR (1993) Non-parametric multivariate analyses of changes in community structure. *Aus. J. Ecol.* 18:117-143
- Clarke KR, Gorley RN (2006) *PRIMER v6: User Manual/Tutorial*. PRIMER-E, Plymouth, UK
- Clarke KR, Warwick RM (1994) *Change in marine communities: an approach to statistical analyses and interpretation*. Primer-E. Natural Environment Research Council, Plymouth, UK
- Crump ML (1971) Quantitative analysis of the ecological distribution of a tropical herpetofauna. *Occasional Papers of the Museum of Natural History The University of Kansas* 3:1–62
- Coddington JA, Agnarsson I, Miller JA, Kuntner M, Hormiga G (2009) Undersampling bias: the null hypothesis for singleton species in tropical arthropod surveys. *J of Anim. Ecol.* 78:573-584
- Colwell RK (2005) *EstimateS: Statistical Estimation of Species Richness and Shared Species from Samples*, Version 7.5 <http://viceroy.eeb.uconn.edu/estimates>, accessed 3/15/09
- Davidson DW, Arias JA, Mann (2006a) An experimental study of bamboo ants in western Amazonia. *Insect Soc* 53:108-114

- Davidson DW, Castro-Delgado SR, Arias JA, Mann J (2006b) Unveiling a ghost of Amazonian rain forests: *Camponotus mirabilis*, engineer of *Guadua* bamboo. *Biotropica* 38(5): 653-666.
- Delsinne T, Leponce M, Laurence T, Braet Y, Roisin Y (2008) Rainfall influences ant sampling in dry forests. *Biotropica* 40(5): 590-596
- Dennis P, Young MR, Gordon IJ (1998) Distribution and abundance of small insects and arachnids in relation to structural heterogeneity of grazed, indigenous grasslands. *Ecol Entomol* 23:253-264
- Dennis P, Young MR, Howard CL, Gordon IJ (1997) The response of epigeal beetles (Col.: Carabidae, Staphylinidae) to varied grazing regimes on upland *Nardus stricta* grasslands. *J Appl Ecol* 24:433-443
- Devries PJ, Walla TR, Greeney HF (1999) Species diversity in spatial and temporal dimensions of fruit-feeding butterflies from two Ecuadorian rainforests. *Biol J Linn Soc* 68:333-353
- Didham RK (1997) The influence of edge effects and forest fragmentation on leaf litter invertebrates in central Amazonia. In: Laurance WF and Bierregaard Jr. RO (ed) *Tropical Forest Remnants: Ecology, Management, and Conservation of Fragmented Communities*. University of Chicago Press, Chicago pp 55-70
- Driscoll DA (2005) Is the matrix a sea? Habitat specificity in a naturally fragmented landscape. *Ecol Entomol* 30:8-16
- Dunnum JL, Salazar-Bravo J (2004) *Dactylomys boliviensis*. *Mamm Spec* 745:1-4
- Emmons LH, Feer F (1990) Neotropical rainforest mammals: A field guide. University of Chicago Press, Chicago
- Erwin TL (1997) Natural history of the carabid beetles at the BIOLAT Biological Station, Rio Manu, Pakitza, Peru, Supplement I. Additional records. In: Wilson DE, Sandoval A (ed) *Manu: The Biodiversity of Southeastern Peru*. The Smithsonian Institution Press, Washington D.C. pp 359-368
- Fine PVA, Mesones I, Coley P (2004) Herbivores promote habitat specialization by trees in Amazonian forests. *Science* 305:663-665
- Foster RB, Carr JL, Forsyth AB (1994) The Tambopata-Candamo reserved zone of southeastern Peru: a biological assessment. Rapid Assessment Program. Working Papers 6. Conservation International, Washington D.C.
- Gardner TA, Fitzherbert EB, Drewes RC, Howell KM, Caro T (2007) Spatial and temporal patterns of abundance and diversity of an east African leaf litter amphibian fauna. *Biotropica* 39(1):105-113

- Grimbacher PS, Stork NE (2009) Seasonality of a diverse beetle assemblage inhabiting lowland tropical rain forest in Australia. *Biotropica* 41(3):328-337
- Griscom BW, Ashton PMS (2003) Bamboo control of forest succession: *Guadua sarcocarpa* in Southeastern Peru. *Forest Ecol Manag* 175: 445-454
- Griscom BW, Ashton PMS (2006) A self-perpetuating bamboo disturbance cycle in a neotropical forest. *J Trop Ecol* 22:587-597
- Griscom BW, Daly DC, Ashton PMS (2007) Floristics of bamboo-dominated stands in lowland terra-firma forests of southwestern Amazonia. *J Torrey Bot Soc* 134(1):108-125
- Hunter MD, Price PW (1993) Playing chutes and ladders: heterogeneity and the relative roles of bottom-up and top-down forces in natural communities. *Ecology* 73:724-732
- Jacobs JM, von May R (2012) Forest of Grass: discovering biodiversity in the Amazon's bamboo jungles. *Natural History Magazine* Jan: 22–29
- Jacobs JM, von May R, Ratcliffe BC (2012) Observations on the life history of *Enema pan* (Fabricius) (Coleoptera: Scarabaeidae: Dynastinae) and its association with bamboo plants (*Guadua* Kunth: Poaceae: Bambusoideae) in southwestern Amazonia. *The Coleops Bull* 66(3): 253–260
- Janzen DH (1973) Sweep samples of tropical foliage insects: Effects of seasons, vegetation types, elevation, time of day, and insularity. *Ecology* 54:687-708
- Judziwickz EJ, Clark LG, Londoño X, Stern MJ (1999) *American Bamboos*. Smithsonian Institution Press, Washington D.C.
- Kratter AW (1997) Bamboo specialization by Amazonian birds. *Biotropica* 29(1):100-110
- Kondo T, Gullan PJ (2004) A new species of ant-tended soft scale of the genus *Cryptostigma* Ferris (Hemiptera: Coccidae) associated with bamboo in Peru. *Systematics, Morphology, and Physiology* Nov – Dec: 717-723
- Kotze DJ, Samways MJ (1999) Invertebrate conservation at the interface between the grassland matrix and natural Afromontane forest fragments. *Biodivers Conserv* 8:1339-1363
- Kotze DJ, Samways MJ (2001) No general edge effects for invertebrates at Afromontane forest/grassland ecotones. *Biodivers Conserv* 10:443-466
- Larsen TH, Lopera A, Forsyth A (2006) Extreme trophic and habitat specialization by Peruvian dung beetles. *The Coleops Bull* 60(4): 315-324

- Lawton JH (1983) Plant architecture and the diversity of phytophagous insects. *Annu Rev Entomol* 28:23-39
- Lebbin DJ (2006) Notes on birds consuming *Guadua* bamboo seeds. *Ornitol Neotropica* 17:602-612
- Lebbin DJ (2007) Habitat specialization among Amazonian birds: why are there so many bamboo specialists? Dissertation, Cornell University
- Lebbin DJ, Hosner PA, Anderson MJ, Valdez U, Tori WP (2007) First description of nest and eggs of the White-lined antbird (*Percnostola lophotes*), and breeding observations of poorly known birds inhabiting *Guadua* bamboo in southeastern Peru. *Bol SAO Vol. XVII*(2): 119-132
- Ledesma KJ (2003) Small mammal communities of bamboo forests in the Peruvian Amazon. Master's Thesis, Florida Atlantic University
- Longino JT, Coddington J, Colwell RK (2002) The ant fauna of a tropical rain forest: Estimating species richness three different ways. *Ecology* 83:689–702
- Longino JT, Colwell RK (1997) Biodiversity assessment using structured inventory: capturing the ant fauna of a tropical rain forest. *Ecological Applications* 7(4):1263-1277
- Louton J, Gelhaus J, Bouchard R (1996) The aquatic macrofauna of water-filled bamboo (Poaceae: Bambusoideae: *Guadua*) internodes in a Peruvian lowland tropical forest. *Biotropica* 28:228-242
- Lucky A, Erwin TL, Witman JD (2002) Temporal and spatial diversity and distribution of arboreal Carabidae (Coleoptera) in a western Amazonian rain forest. *Biotropica* 34(3):376-386
- Magurran AE (2004) *Measuring Biological Diversity*. Blackwell Publishing, UK
- Nelson BW (1994) Natural forest disturbance and change in the Brazilian Amazon. *Rem Sens Revs* 10:105-125
- Nelson BW, Bianchini MC (2005) Complete life cycle of southwest Amazon bamboos (*Guadua* spp) detected with orbital optical sensors. *Anais XII. Simpósio Brasileiro de Sensoriamento Remoto, INPE, Goiânia, Brasil* pp 1629-1636
- Novotny V, Basset Y (1998) Seasonality of sap-sucking insects (Auchenorrhyncha, Hemiptera) feeding on *Ficus* (Moraceae) in a lowland rain forest in New Guinea. *Oecologia* 115: 514-522
- Olivier J (2007) Etude spatio-temporelle de la distribution de bambous dans le Sud-ouest amazonien (Sud-Pérou). Histoire, dynamique et futur d'une végétation « monodominante » en forêt tropicale humide. Dissertation, Université Paul Sabatier, Toulouse, France

- Olivier J (2008) Gramíneas (Poaceae) bambusiformes del Río de Los Amigos, Madre de Dios, Perú. *Rev Peru Biol* 15(1):121-126
- Olivier J, Otto T, Roddaz M, Antoine PO, Londoño X, Clark LG (2008) First macrofossil evidence of a pre-Holocene thorny bamboo cf. *Guadua* (Poaceae: Bambusoideae: Bambuseae: guaduinae) in south-western Amazonia (Madre de Dios-Peru). *Rev Paleob and Palyno* 15(3):1-7
- Pearson DL, Derr JA (1986) Seasonal patterns of lowland forest floor arthropod abundance in southeastern Peru. *Biotropica* 18(3): 244-256
- Pitman NCA (2008) An overview of the Los Amigos Watershed, Madre de Dios, southeastern Peru. Amazon Conservation Association.
http://cicra.acca.org.pe/espanol/paisaje_biodiversidad/los-amigos-overview9.pdf. Accessed 12 March 2009
- Richards LA, Windsor DM (2007) Seasonal variation of arthropod abundance in gaps and the understory of a lowland moist forest in Panama. *J Trop Ecol* 23:169-176
- Ruíz-Sánchez E (2011) Biogeography and divergence time estimates of woody bamboos: insights in the evolution of Neotropical bamboos. *B Soc Bot Mex* 88:67-75
- Schaffers AP, Raemakers IP, Sýkora KV, Ter Braak CJF (2008) Arthropod assemblages are best predicted by plant species composition. *Ecology* 89(3):782-794
- Schoereder JH, Galbiati C, Ribas CR, Sobrinho TG, Sperber CF, DeSouza O, Lopes-Andrade C (2004) Should we use proportional sampling for species–area studies? *J Biogeogr* 31: 1219–1226
- Siemann E, Tilaman D, Haarstad J, Ritchie M (1998) Experimental tests of the dependence of arthropod diversity on plant diversity. *The American Naturalist* 152:738-750
- Socular SJ, Robinson SK, Terborgh J (2013) Bird diversity and occurrence of bamboo specialists in two bamboo die-offs in southeastern Peru. *The Condor* 115(2): 253-262
- Spector S, Aysama S (2003) Rapid turnover and edge effects in dung beetle assemblages (Scarabaeidae) at a Bolivian neotropical forest-savanna ecotone. *Biotropica* 35(3):394-404
- von May R, Medina-Müller M, Donnelly MA, Summers K (2008) The tadpole of the bamboo-breeding poison frog *Ranitomeya biolat* (Anura: Dendrobatidae). *Zootaxa* 1857:66–68
- von May R, Medina-Müller M, Donnelly MA, Summers K (2009a) Breeding-site selection by poison frogs (*Ranitomeya biolat*) in Amazonian bamboo forests: an experimental approach. *Can J Zoolog* 87(5):453–464
- von May R, Siu-Ting K, Jacobs JM, Medina-Müller M, Gagliardi G, Rodríguez LO, Donnelly MA (2009b) Species diversity and conservation status of amphibians in Madre de Dios, Perú. *Herpetol Cons and Biol* 4:14–29

von May R, Jacobs JM, Santa Cruz R, Valdivia J, Huamán J, Donnelly MA (2010) Amphibian community structure as a function of forest type in Amazonian Peru. *J Trop Ecol* 26(5): 509-519

Ward DF, Larivière M-C (2004) Terrestrial invertebrate surveys and rapid biodiversity assessment in New Zealand: Lessons from Australia. *NZ J Ecol* 28(1):151-159


Ward DF, New TR, Yen AL (2001) Effects of pitfall trap spacing on the abundance, richness and composition of invertebrate catches. *J Insect Conserv* 5(1):47-53

Wolda H (1978) Seasonal fluctuations in rainfall, food and abundance of tropical insects. *J Anim Ecol* 47(2):369-381

Work TT, Buddle CM, Korinus LM, Spence JR (2002) Pitfall trap size and capture of three taxa of litter-dwelling arthropods: implications for biodiversity studies. *Environ Entomol* 31(3):438-448

Yu DX, Luo TH, Zhou HZ, Yang J (2007) Distribution of Carabid beetles (Coleoptera: Carabidae) across a forest-grassland ecotone in southwestern China. *Community Ecosys Ecol* 36: 348-355

844



845
846
847
848
849
850
851
852
853
854

Fig 1. Map of study site in southeastern Peru. All collecting took place inside the Los Amigos Conservation Concession (LACC). Map courtesy of Karen Siu-Ting.





Fig. 2 Dense patch of *Guadua weberbaueri* bamboo habitat with trail cut through patch



Fig. 3 Terra firme habitat




Figure 1 Sampling scheme of paired sites in bamboo and terra firme forest. Each pair of forest types received the same number of pitfall traps in the same array. Pitfall trap arrays were always greater than 10 meters from the edge of a forest type and individual traps were always five meters apart




Fig. 5 Species accumulation curves. The number of species collected in bamboo and terra firme forests as a function of sampling effort (bars indicate 95% confidence intervals)




Fig. 6 Whitaker plots comparing rank abundances between (a) forest types and (b) seasons. Species are ranked according to their abundances. Each letter represents one species, and the ten most abundant species are ordered. For each plot, one letter represents the same species, and a new letter is assigned for every new species. However, the letters are not constant for both plots in that species D in plot (a) may not be species D in plot (b). The lettered order illustrates that the ten most abundant species are not the same for bamboo and terra firme, and dry and wet seasons




Fig. 7 nMDS plot illustrating beetle community structure between bamboo and terra firme forest with data from all families and seasons pooled. The plot exhibits distinct communities of beetles in the two forest types with some degree of overlap between assemblages. The arrow is pointing to one out-lying data point from the terra firme forest. This point represents one of the smallest collecting sites, Luisa 2, and may be influencing the results




Fig. 8 Matched rank/occurrence plots for **(a)** bamboo and terra firme forest and **(b)** dry and wet seasons. Each pair of positive (dark) and negative (hatched) bars represents the relative abundance of one species in each forest type or each season. The letter “A” column represents beetles from the family Carabidae, “B” represents Curculionidae, “C” represents Histeridae and “D” represents Scarabaeidae




Fig. 9 Plots expressing variation in numbers of individuals and numbers of unique species captured between dry and wet seasons, for bamboo forest compared to terra firme forest, at all sites. **(a)** Each bar represents the difference in number of individuals captured between the dry and wet season in one forest type. **(b)** Each bar represents the number of unique (not shared) species captured in the dry and wet season, in one forest type. The grey arrow in (b) is pointing to one of the smallest collecting sites, Louisa 2, and the only site where bamboo and terra firme forest both had the same number of unique species between dry and wet seasons. Overall, there was more variation in the numbers of individuals captured and greater species turnover between dry and wet seasons in the terra firme forest compared to bamboo forest





Fig. 10 nMDS plots comparing beetle community structure between bamboo and terra firme forest in (a) the wet season only and (b) the dry season only. There is a greater distinction in beetle community structure between bamboo and terra firme forest in the wet season compared to the dry season



Fig. 11 Individual of male *Enema pan* perching outside of his mound entrance at the base of a *Guadua weberbaueri* stem. *Enema pan* beetles construct these mounds connecting underground to approximately 1 meter long tunnels. When the mound is closed during the day, they stay hidden in their tunnels and feed on bamboo sap through the bark that they have shredded at the base of the bamboo stem. They open their mounds at night and perch at the entrance of their mound, presumably looking for mates or guarding the entrances to the mound.

Table 1. The number of species of beetle in each family, and in all families combined, collected in only bamboo forest, only terra firme forest, in both habitats, only in the dry season, only in the wet season, and both seasons. The 7th column lists the total number of species collected per family.



Richness	Bamboo only	Terra Firme only	Both habitats	Dry only	Wet only	Both seasons	Total no. species
Carabidae	7	12	6	7	15	4	25
Curculionidae	16	25	11	13	31	8	52
Histeridae	6	6	10	7	5	10	22
Scarabaeidae	19	26	45	14	29	47	90
All families	48	69	72	41	80	69	189

Table 2. The number of individuals from each beetle family, and all families combined, collected in bamboo forests, terra firme forests, dry season, wet season. The last column lists the total number of individuals collected within each family and all for all families combined.

Abundance	Bamboo	Terra Firme	Dry	Wet	Total no. individuals
Carabidae	51	52	16	87	103
Curculionidae	62	93	71	84	155
Histeridae	153	228	204	177	381
Scarabaeidae	1273	1840	1677	1436	3113
All families	1539	2213	1968	1784	3752

Table 3. Beetle species contributing 90% of the total number of individuals collected in bamboo and terra firme forest communities (SIMPER – PRIMER 6.0). The values represent the percent contribution of each species to the distinctness in beetle community structure between bamboo and terra firme forest. The largest numbers are associated with species that most greatly distinguish assemblages in one forest type or another.

Species	Bamboo	Terra Firme
Carabidae		
<i>Odontocheila cayennensis</i>	0	1.89
Curculionidae		
<i>Curculionidae 13</i>	0	0.96
Histeridae		
<i>Omolodes A</i>	6.71	8.35
<i>Phelister B</i>	2.1	5.62
<i>Operclipygus C</i>	1.66	0
Scarabaeidae		
<i>Enema pan</i>	14.8	0
<i>Scybalocanthos C</i>	12.63	8.51
<i>Canthidium A</i>	9.59	3.04
<i>Onthophagus xanthomerus</i>	9.4	10.96
<i>Canthidium F</i>	6.1	2.33
<i>Canthidium gerstaeckeri</i>	5.28	3.31
<i>Dichotomius ohausi</i>	5.03	6.51
<i>Dichotomius batesi</i>	4.87	0
<i>Dichotomius nr. lucasi</i>	3.57	3.34
<i>Canthonella D</i>	3.05	0
<i>Scybalocanthos D</i>	2.06	5.2
<i>Canthidium nr. deyrollei</i>	1.66	2.75
<i>Dichotomius lucasi</i>	1.58	3.09
<i>Ateuchus C</i>	0	7.21
<i>Ateuchus D</i>	0	3.11
<i>Canthidium batesi</i>	0	7.84
<i>Ceratocanthinae A</i>	0	1.4
<i>Deltochilum nr. komerecki</i>	0	1.19
<i>Eurysternus nov. stigilatus</i>	0	3.72