

# Standardizing microsatellite panels for individual identification of seabirds' Snow Petrel *Pagodroma nivea* and Wilson's Storm Petrel *Oceanites oceanicus* in Antarctica (#22989)

1

First submission

## Editor guidance

Please submit by **30 Jan 2018** for the benefit of the authors (and your \$200 publishing discount).



### Structure and Criteria

Please read the 'Structure and Criteria' page for general guidance.



### Custom checks

Make sure you include the custom checks shown below, in your review.



### Author notes

Have you read the author notes on the [guidance page](#)?



### Raw data check

Review the raw data. Download from the location [described by the author](#).



### Image check

Check that figures and images have not been inappropriately manipulated.

Privacy reminder: If uploading an annotated PDF, remove identifiable information to remain anonymous.

## Files

Download and review all files from the [materials page](#).

- 1 Figure file(s)
- 1 Table file(s)
- 2 Raw data file(s)

## ! Custom checks

### Vertebrate animal usage checks

- ! Have you checked the authors [ethical approval statement](#)?
- ! Were the experiments necessary and ethical?
- ! Have you checked our [animal research policies](#)?

### Field study

- ! Have you checked the authors [field study permits](#)?
- ! Are the field study permits appropriate?




# Structure and Criteria

## Structure your review

The review form is divided into 5 sections.  
Please consider these when composing your review:

1. BASIC REPORTING
2. EXPERIMENTAL DESIGN
3. VALIDITY OF THE FINDINGS
4. General comments
5. Confidential notes to the editor






 You can also annotate this PDF and upload it as part of your review

When ready [submit online](#).





## Editorial Criteria

Use these criteria points to structure your review. The full detailed editorial criteria is on your [guidance page](#).





### BASIC REPORTING

-  Clear, unambiguous, professional English language used throughout.
-  Intro & background to show context. Literature well referenced & relevant.
-  Structure conforms to [Peerj standards](#), discipline norm, or improved for clarity.
-  Figures are relevant, high quality, well labelled & described.
-  Raw data supplied (see [Peerj policy](#)).

### EXPERIMENTAL DESIGN

-  Original primary research within [Scope of the journal](#).
-  Research question well defined, relevant & meaningful. It is stated how the research fills an identified knowledge gap.
-  Rigorous investigation performed to a high technical & ethical standard.
-  Methods described with sufficient detail & information to replicate.

### VALIDITY OF THE FINDINGS

-  Impact and novelty not assessed. Negative/inconclusive results accepted. *Meaningful* replication encouraged where rationale & benefit to literature is clearly stated.
-  Data is robust, statistically sound, & controlled.
-  Conclusions are well stated, linked to original research question & limited to supporting results.
-  Speculation is welcome, but should be identified as such.

# Standout reviewing tips

3



The best reviewers use these techniques

## Tip

**Support criticisms with evidence from the text or from other sources**

## Example

*Smith et al (J of Methodology, 2005, V3, pp 123) have shown that the analysis you use in Lines 241-250 is not the most appropriate for this situation. Please explain why you used this method.*

**Give specific suggestions on how to improve the manuscript**

*Your introduction needs more detail. I suggest that you improve the description at lines 57- 86 to provide more justification for your study (specifically, you should expand upon the knowledge gap being filled).*

**Comment on language and grammar issues**

*The English language should be improved to ensure that an international audience can clearly understand your text. Some examples where the language could be improved include lines 23, 77, 121, 128 – the current phrasing makes comprehension difficult.*

**Organize by importance of the issues, and number your points**

1. Your most important issue
2. The next most important item
3. ...
4. The least important points

**Please provide constructive criticism, and avoid personal opinions**

*I thank you for providing the raw data, however your supplemental files need more descriptive metadata identifiers to be useful to future readers. Although your results are compelling, the data analysis should be improved in the following ways: AA, BB, CC*

**Comment on strengths (as well as weaknesses) of the manuscript**

*I commend the authors for their extensive data set, compiled over many years of detailed fieldwork. In addition, the manuscript is clearly written in professional, unambiguous language. If there is a weakness, it is in the statistical analysis (as I have noted above) which should be improved upon before Acceptance.*

# Standardizing microsatellite panels for individual identification of seabirds' Snow Petrel *Pagodroma nivea* and Wilson's Storm Petrel *Oceanites oceanicus* in Antarctica

Anant Pande<sup>1</sup>, Nidhi Rawat<sup>2</sup>, Kuppusamy Sivakumar<sup>1</sup>, Sambandan Sathyakumar<sup>1</sup>, Vinod B Mathur<sup>3</sup>, Samrat Mondol<sup>Corresp.</sup><sup>2</sup>

<sup>1</sup> Endangered Species Management, Wildlife Institute of India, Dehradun, Uttarakhand, India

<sup>2</sup> Animal Ecology and Conservation Biology, Wildlife Institute of India, Dehradun, Uttarakhand, India

<sup>3</sup> Director, Wildlife Institute of India, Dehradun, Uttarakhand, India

Corresponding Author: Samrat Mondol

Email address: samrat@wii.gov.in

Seabirds are known to be important indicators of marine ecosystems health. Procellariiformes are one of the most abundant seabird species distributed from warm tropical to cold temperate regions including Antarctica. With few long-term studies on breeding seabirds at the Antarctic continent, crucial biological parameters such as genetic variation, population genetic structure and past population demography is lacking for most of the commonly occurring species. Under the 'Biology and Environmental Sciences' component of the Indian Antarctic programme, long-term monitoring of Antarctic biodiversity is being conducted. In this paper, we describe a panel of 12 and 10 cross-species microsatellite markers for two relatively less studied seabird species in Antarctica, snow petrel *Pagodroma nivea* and Wilson's storm petrel *Oceanites oceanicus*, respectively. These loci showed high amplification success and moderate level of polymorphism in snow petrel (mean no. of alleles  $7.08 \pm 3.01$  and mean observed heterozygosity  $0.35 \pm 0.23$ ), but low polymorphism in Wilson's storm petrel (mean no. of alleles  $3.9 \pm 1.3$  and mean observed heterozygosity  $0.28 \pm 0.18$ ). The results demonstrate that these panels can unambiguously identify individuals of both species from various types of biological materials. This work forms a baseline for undertaking long-term genetic research of Antarctic seabird species and provides critical insights into their population genetics.

**Standardizing microsatellite panels for individual identification of seabirds' Snow  
Petrel *Pagodroma nivea* and Wilson's Storm Petrel *Oceanites oceanicus* in Antarctica**

**Running head: Individual identification of two Antarctic seabird species**


Anant Pande<sup>\*</sup>, Nidhi Rawat<sup>\*</sup>, Kuppusamy Sivakumar, Sambandan Sathyakumar, Vinod B.  
Mathur, Samrat Mondol<sup>#</sup>

Wildlife Institute of India, Chandrabani, Dehradun, Uttarakhand, India 248001

*\*equal contribution*

<sup>#</sup> Corresponding author: Samrat Mondol, Assistant Professor, Animal Ecology and  
Conservation Biology, Wildlife Institute of India, Chandrabani, Dehradun, Uttarakhand, India  
248001. Email: [samrat@wii.gov.in](mailto:samrat@wii.gov.in)

# 29 ABSTRACT

30 Seabirds are ~~known to be~~ important indicators of marine ecosystems health. Procellariiformes  
 31 are ~~one of the~~ most abundant seabird species distributed from warm tropical to cold temperate  
 32 regions including Antarctica. ~~With few long-term studies on breeding seabirds at the~~  
 33 ~~Antarctic continent, crucial biological parameters such as genetic variation, population~~  
 34 ~~genetic structure and past population demography is lacking for most of the commonly~~  
 35 ~~occurring species.~~ Under the ‘Biology and Environmental Sciences’ component of the Indian  
 36 Antarctic programme, long-term monitoring of Antarctic biodiversity is being conducted. In  
 37 this paper, we describe a panel of 12 and 10 ~~cross-species~~ microsatellite markers for two  
 38 relatively ~~less studied~~ seabird species in Antarctica, snow petrel *Pagodroma nivea* and  
 39 Wilson’s storm petrel *Oceanites oceanicus*, respectively. These loci showed high  
 40 amplification success and moderate level of polymorphism in snow petrel (mean no. of  
 41 alleles  $7.08 \pm 3.01$  and mean observed heterozygosity  $0.35 \pm 0.23$ ), but low polymorphism in  
 42 Wilson’s storm petrel (mean no. of alleles  $3.9 \pm 1.3$  and mean observed heterozygosity  
 43  $0.28 \pm 0.18$ ). The results demonstrate that these panels can unambiguously identify individuals  
 44 of both species from various types of biological materials. This work forms a baseline for  
 45 undertaking long-term genetic research of Antarctic seabird species and provides critical  
 46 ghts into their population genetics.

47

48

49

50 **Keywords:** Antarctic seabirds; Indian Antarctic programme; genetic monitoring;  
 51 Procellariiformes, Antarctic biodiversity.

52

# 53 INTRODUCTION

54 ~~Seabirds, being top predators,~~ maintain structure of marine food webs, regulate island and  
 55 marine ecosystem processes and act as indicators of marine ecosystem health (Lascelles *et al.*  
 56 2012; Paleczny *et al.* 2015). Their ~~natural~~ ability to fly over large distances, extreme life  
 57 history strategies (monogamy, slow reproduction, late sexual maturity), ~~natal~~ philopatry, high  
 58 visibility and dependence on land for breeding makes them ideal candidates for long-term  
 59 population level studies (Piatt *et al.* 2007). ~~A number of recent~~ studies focusing on seabird  
 60 population monitoring have highlighted the threatened status of seabirds ~~across the globe~~  
 61 (Croxall *et al.* 2012), especially in the southern ocean where seabird populations have  
 62 declined substantially over last few decades (Paleczny *et al.* 2015). ~~This has led to efforts~~  
 63 ~~focusing on understanding seabird population dynamics using interdisciplinary approaches to~~  
 64 ~~aid~~ conservation and management across their distribution range (Croxall *et al.* 2012; Taylor  
 65 and Friesen 2012).

66 ~~Among seabirds, order Procellariiformes includes Petrels, Shearwaters, Albatrosses, Storm~~  
 67 ~~Petrels and Diving Petrels representing the most widely distributed and abundant species~~  
 68 (Warham 1996). ~~In spite of their wide range~~ and large population sizes, long-term ecological  
 69 and genetic data exists for few of these species across the globe. In addition to several  
 70 ecological studies ~~on~~ Procellariiformes (Croxall *et al.* 2012), some recent studies have used  
 71 genetic data to address important biological parameters such as relatedness, population  
 72 structure, past population demography (e.g. see Gómez-Díaz *et al.* 2009 for Cory's  
 73 shearwater; Welch *et al.* 2012 for Hawaiian petrel) for species distributed in tropical and  
 74 **arctic** marine ecosystems. Research on ~~Procellariiformes' biology~~ is relatively limited in the  
 75 Southern Ocean ecosystem, especially in Antarctica because of its remoteness and associated  
 76 logistical difficulties. Despite few site-specific monitoring of some ~~Procellariiformes~~ on sub-  
 77 Antarctic islands (e.g. Brown *et al.* 2015 for giant petrels; Quillfeldt *et al.* 2017 for Antarctic

prion, thin-billed prion and blue petrel) and Antarctic coast (e.g. Barbraud & Weimerskirch, 2001 on snow petrel; Barbraud and Weimerskirch 2006 on multiple species; Techow *et al.* 2010 on giant petrels), long-term ecological as well as genetic research is sparse. Nunn and Stanley (1998) reported the phylogenetic relationships among procellariiform species' using a neighbour-joining approach, but within each family groups detailed population genetic information is lacking. Other preliminary studies have used Restriction Fragment Length Polymorphisms (RFLP) and allozymes to investigate genetic variation and extra-pair paternity in snow petrel as well as some other Procellariiformes (Jouventin and Viot 1985, Viot *et al.* 1993, Lorensten *et al.* 2000, Quillfeldt *et al.* 2001) in Antarctica.

The 'Biology and Environmental Sciences Programme' of Indian Scientific Expeditions to Antarctica has a specific focus on understanding distribution, abundance, population dynamics and genetics of Antarctic seabirds, including Procellariiformes. As part of this program, comprehensive ecological surveys were conducted between 2009-2016 to understand seabird and marine mammal ecology around Indian Antarctic research stations (Pande *et al.* 2017). Currently this programme is focused on generating baseline genetic data of breeding seabird species found around Indian area of operations in Antarctica, especially on Snow Petrel *Pagodroma nivea* and Wilson's Storm Petrel *Oceanites oceanicus*. Snow petrel is endemic to Antarctica and Southern Ocean with breeding distribution along Antarctic coast including some inland mountains and few sub-Antarctic islands (Croxall *et al.* 1995). On the other hand, Wilson's storm petrel has a much wider breeding distribution from Cape Horn to the Kerguelen Islands and coastal Antarctica. It also migrates to the mid-latitudes of the north Atlantic, north Indian and Pacific Ocean during non-breeding period (BirdLife International 2017). Effective monitoring of these species in the Indian Antarctic sector will require systematic information on their distribution, current population status and genetics. With the broad objective of assessing population genetic structure, we describe a



panel of cross-species microsatellite markers for individual identification of snow petrel and Wilson's storm petrel in Antarctica and sub-Antarctic islands. These tested panels will be of great help in understanding genetic variation, genetic relatedness and demographic history of both these species across their ranges.

## METHODS

### Permits and ethical clearances

All samples were collected under the 'Biology and Environmental Sciences' component (Letter no: NCAOR/ANT/ASPA/2014-15/01) of the Indian Scientific Expeditions to Antarctica with appropriate approvals from the Environment Officer, Committee for Environmental Protection (Antarctic Treaty Secretariat), National Centre for Antarctic and Ocean Research, Earth System Science Organisation, Ministry of Earth Sciences, Government of India, Goa, India.

### Study Area

Sampling was carried out at Larsemann hills, Prydz bay and Schirmacher oasis, Central Dronning Maudland (Figure 1); close to permanent Indian research stations in Antarctica Bharati (Larsemann hills) and Maitri (Schirmacher oasis). Distance between these two study areas is about 2,500 km. Larsemann hills (69° 20'S to 69° 30'S; 75° 55'E to 76° 30'E coordinates), are a group of islands in Prydz Bay located on the Ingrid Christensen Coast, Princess Elizabeth Land of east Antarctica. It comprises of variously sized islands and peninsulas, located halfway between the eastern extremity of the Amery Ice Shelf and the southern boundary of the Vestfold Hills. Schirmacher Oasis, Central Dronning Maudland (70° 44' to 70° 46' S and 11° 22' to 11° 54' E coordinates) is situated on the Princess Astrid coast about 120 km from the Fimbul ice shelf. Four species of seabirds (Adelie penguin *Pygoscelis adeliae*, south polar skua *Stercorarius maccormickii*, snow petrel and Wilson's

storm petrel) breed in the ice-free areas of Larsemann hills whereas only south polar skua breeds at Schirmacher oasis (Pande et al. 2017). Other key wildlife species found around these areas include emperor penguin (*Aptenodytes forsteri*), crabeater seal *Lobodon carcinophaga*, leopard seal *Hydrurga leptonyx*, Ross seal *Ommatophoca rossii*, Weddell seal *Leptonychotes weddellii* and orca *Orcinus orca*.

## Field Sampling

Sampling for this study was conducted as part of the ‘Antarctic Wildlife Monitoring Programme’ under the Indian Scientific Expedition to Antarctica (Expedition nos. 33, 34 and 35)’ during the austral summers (November-March) of 2013-14, 2014-15 and 2015-16. We adopted a systematic genetic sampling approach under the seabird nest monitoring protocol (see Pande et al., 2017) for snow petrel sample collection. First, identified nest sites with breeding snow petrel individuals were selected for genetic sampling. Subsequently, both non-destructive (buccal swabs and blood smears) and non-invasive (hatched eggshells and abandoned eggs) sampling approaches were used to collect biological materials from monitored nesting sites. During non-destructive sampling of snow petrel individuals, birds were carefully hand-captured at their nest cavities and buccal swabs or blood samples were collected. Blood samples were collected from bird’s brachial vein using 0.1 ml sterilized syringe needles and stored in an EDTA containing vial. All individuals were released within 60 seconds of capture. We could also collect few hatched eggshells and abandoned eggs from the nests. In addition, opportunistic sampling of snow petrel carcasses was also conducted. These dead animals were mostly predated by south polar skua or naturally dead. Systematic snow petrel sampling was conducted only at Larsemann hills. No nesting sites of snow petrel were found at Schirmacher oasis during field surveys but opportunistic sampling of carcasses was conducted.

Similarly, Wilson's storm petrel ~~tissue~~ samples were collected from monitored nesting sites at Larsemann ~~hills~~. All genetic samples of Wilson's storm petrel were collected opportunistically through carcass **tissue** collection, as capturing them was not possible due to their narrow nest cavities. No Wilson's storm petrel samples were collected from Schirmacher ~~oasis~~. Samples collected at **field** were stored at -20° C at respective Indian Antarctic research stations before being brought to Wildlife Institute of India, Dehradun for further laboratory analysis.

## Primer selection

~~As there is no published work available with nuclear DNA markers for snow petrel and no species-specific microsatellite markers are yet developed, we tested a panel of cross-species markers for individual identification of snow petrels. We selected a total of 15 microsatellite markers earlier developed for Hawaiian petrel *Pterodroma sandwichensis* (Nine markers from Welch and Fleischer 2011) and white-chinned petrel *Procellaria aequinoctialis* (Six markers from Techow and O'Ryan 2004). These markers were selected based on their polymorphic information content (number of alleles as well as expected heterozygosity) in the aforementioned species.~~

~~On the other hand, a set of cross-specific microsatellite markers developed for prion **species** has been tested in Wilson's storm petrel (Moodley *et al.* 2015). However, the study has reported very low amplification success rate. In this study, we also tested these 15 microsatellite loci for individual identification of Wilson's storm petrel.~~

## DNA extraction and primer standardization

We used **tissue** samples of snow petrel and Wilson's storm petrel for initial standardization and validation of microsatellite ~~panel~~. Genomic DNA was extracted in duplicate from all **tissue** samples using commercially available DNeasy Tissue kit (QIAGEN Inc.) ~~using a~~

~~modified approach~~. In brief, all samples were macerated with sterile blades independently, followed by overnight ~~complete~~ tissue digestion with 25  $\mu$ l proteinase-K. Post-digestion, extraction was performed using Qiagen animal tissue spin column protocol. DNA was eluted twice with 100  $\mu$ l of 1X TE and stored in  $-20^{\circ}\text{C}$  until further processing. Each set of 11 extractions was accompanied with one extraction control to monitor possible contamination.

All initial PCR standardizations were conducted using tissue DNA samples. Amplifications were carried out for each primer in 10  $\mu$ l reaction volumes containing 4  $\mu$ l Qiagen Multiplex PCR buffer mix (Qiagen Inc.), 0.2  $\mu$ M labeled forward primer, 0.2  $\mu$ M reverse primer, 4  $\mu$ M BSA and 2  $\mu$ l of 1:10 diluted DNA extract. The temperature regime included an initial denaturation ( $94^{\circ}\text{C}$  for 15 min); 35 cycles of denaturation ( $94^{\circ}\text{C}$  for 30 s), annealing ( $53$  or  $57^{\circ}\text{C}$  for 45 s) and extension ( $72^{\circ}\text{C}$  for 45 s); followed by a final extension ( $72^{\circ}\text{C}$  for 30 min). ~~Post-temperature standardization, primers with identical annealing temperatures was optimized for multiplex reactions with the same samples of both species (see Table 1).~~ Subsequently, all test samples were amplified with standardized parameters. During all amplifications, both extraction controls and PCR negative controls (one PCR negative every set of amplifications) were included to monitor ~~any possible~~ contamination. PCR products were visualized in 2% agarose gels, and genotyped using LIZ500 size standard in an automated ABI3500XL genetic analyzer. Microsatellite alleles were scored using program GENEMARKER (SOFTGENETICS Inc.) and allele bins for each locus were created from the data generated. We randomly re-genotyped 15% of each locus from different samples to check for reliable genotypes and estimated genotyping error rates.

## Data analysis

Average amplification success was calculated as the percent positive PCR for each locus, as described by Broquet *et al.* (2007) . Allelic dropout and false allele rates were quantified

manually as the number of dropouts or false alleles over the total number of amplifications, respectively (Broquet *et al.* 2007). We also calculated the Probability of Identity for siblings (PID)<sub>sibs</sub>, the probability of two individuals drawn from a population sharing the same genotype at multiple loci (Waits *et. al* 2001) using program GIMLET (Valière 2002). We tested the frequency of null alleles in our data set using FREENA (Chapuis & Estoup 2007), whereas summary statistics and tests for deviations from Hardy-Weinberg equilibrium were calculated for each locus using program ARLEQUIN v.3.1 (Excoffier and Schneider 2005).

## Results and discussion

We genotyped a total of 55 snow petrel and 24 Wilson's storm petrel samples to test and standardize the selected microsatellite markers. Snow petrel samples were selected from blood (n=1), buccal swab (n=2), carcass tissue (n=24) and eggshells (n=28) to test amplification success from different types of biological samples. Wilson's storm petrel samples were all from muscle tissue of individual carcasses collected in the field.

Of the 15 loci tested during the initial standardization, 12 loci showed amplification for snow petrel (loci Ptero2, Ptero6 and Ptero10 did not amplify), whereas only 10 loci successfully amplified for Wilson's storm petrel (loci Paequ2, Ptero2, Ptero6, Ptero8 and Ptero10 did not amplify) (see Table 1 for details). Subsequently, these panels of 12 and 10 loci were tested with all snow petrel and Wilson's storm petrel samples, respectively. For snow petrel, the loci varied from highly polymorphic (Paequ03- 12 alleles,  $H_o$ - 0.68) to less polymorphic (Paequ13- 4 alleles,  $H_o$ - 0.07), whereas for Wilson's storm petrel the loci were moderately polymorphic (Ptero07- 6 alleles,  $H_o$ -0.76) to less polymorphic (Paequ13- 2 alleles,  $H_o$ - 0.08) (see Table 1 for detailed summary statistics). We could not find any locus that deviated from the Hardy-Weinberg Equilibrium, and there was no evidence for linkage disequilibrium between any pair of loci. Overall, the amplification success ranged between 96.4-100% for

223 snow petrel and 91.7–100% for Wilson’s storm petrel; ~~and~~ allelic dropout rates were 0 –  
 224 3.6% and 0 – 8.3% for snow petrel and Wilson’s storm petrel respectively. The estimated  
 225 cumulative probability of identity assuming all individuals were siblings ( $PID_{(sibs)}$ ) was found  
 226 to be  $1.1 \times 10^{-3}$  for snow petrel and  $5.0 \times 10^{-3}$  for Wilson’s storm petrel. Average values for  
 227 observed and expected heterozygosity, number of alleles, allelic range sizes are presented in  
 228 Table 1. The frequency of null alleles across the loci was ~~observed to be~~ low in both ~~the~~ study  
 229 species (snow petrel -  $0.11 \pm 0.09$  and Wilson’s storm petrel -  $0.15 \pm 0.07$ ).

230 This paper is the first attempt to use nuclear microsatellite markers to individually identify  
 231 both snow petrel and Wilson’s storm petrel in Antarctica. Based on the results of this study  
 232 ( $PID_{(sibs)}$  value of  $1 \times 10^{-3}$ ), it can be ascertained that our standardized 12 microsatellite loci  
 233 panels are ~~sufficient enough~~ to differentiate among related individuals of snow petrel.  
 234 However, in case of Wilson’s storm petrel ( $PID_{(sibs)}$  value of  $5 \times 10^{-3}$ ) the statistical power is  
 235 ~~not enough~~, and additional loci need to be standardized to avoid any possible errors in case of  
 236 **population genetic** study of Wilson’s storm petrel. ~~Previously tested microsatellite loci by~~  
 237 Moodley *et al.* (2015), though not used for individual identification, could be used along with  
 238 the current panel of markers to increase the statistical power during individual identification  
 239 in Wilson’s storm petrel. Overall, our results show that both panels of loci provide  
 240 unambiguous individuals for respective seabird species in Antarctica.

## 241 Conclusion

242 Molecular genetic analysis ~~has become crucial in understanding~~ levels of genetic  
 243 differentiation, hybridisation and extinction risk in seabird populations (Taylor and Friesen,  
 244 2012). In **critical** ecosystems such as Antarctica, individual-level genetic data can be a  
 245 valuable tool to study evolution, adaptation, past events of diversifications and extinctions for  
 246 wide-ranging seabirds. However, genetic studies on species of flying seabirds such as snow

petrel and Wilson's storm petrel are generally lacking in comparison with charismatic species such as penguins. In this study, we could establish the efficacy of cross-species markers in individual identification of these two common Antarctic seabird species. We aim to continue this long-term genetic research under the current 'Antarctica Wildlife Monitoring Programme' by increasing spatio-temporal sampling efforts to understand the population structure, relatedness and other aspects and provide insights to seabird behaviour (monogamy, extra-pair paternity etc.) and evolution. This detailed genetic research will also aid long-term ecological monitoring of breeding seabird populations and help informed conservation management of these species in Antarctica.

## Acknowledgements

We thank the National Centre for Antarctic and Ocean Research, Ministry of Earth Sciences for providing all logistic support during the Indian Scientific Expeditions to Antarctica. We are grateful to respective expedition leaders and team member volunteers of 33<sup>rd</sup>, 34<sup>th</sup> & 35<sup>th</sup> Indian Scientific Expeditions to Antarctica for their assistance during field work. We thank A. Madhanraj and MEERCAT lab members for their help in laboratory analysis of samples. We sincerely thank Wildlife Forensics and Conservation Genetics Cell, CAMPA Cell, Research Coordinator and Dean, Wildlife Institute of India for their support. We thank the Wildlife Institute of India and DST-INSPIRE faculty Award to SM (Grant no: IFA12-LSBM-47) for financial support to this study.

## Figure legend

Figure 1: Seabird sampling locations in Antarctica. A) Schirmacher oasis, site of *Maitri* station B) Larsemann hills, site of *Bharati* station.

## References

Barbraud, C., & Weimerskirch, H. (2001). Contrasting effects of the extent of sea-ice on the

- 271 breeding performance of an Antarctic top predator , the Snow Petrel *Pagodroma nivea*.  
272 *Journal of Avian Biology*, 32, 297–302.
- 273 Barbraud, C., & Weimerskirch, H. (2006). Antarctic birds breed later in response to climate  
274 change. *Proceedings of the National Academy of Sciences of the United States of*  
275 *America*, 103, 6248–51.
- 276 BirdLife International (2017) Species factsheet: *Oceanites oceanicus*. Downloaded from  
277 <http://www.birdlife.org> on 17/11/2017.
- 278 Broquet, T., Ménard, N., & Petit, E. (2007). Noninvasive population genetics: A review of  
279 sample source, diet, fragment length and microsatellite motif effects on amplification  
280 success and genotyping error rates. *Conservation Genetics*, 8, 249–260.
- 281 Brown, R. M., Techow, N. M. S. M., Wood, A. G., & Phillips, R. A. (2015). Hybridization  
282 and Back-Crossing in Giant Petrels (*Macronectes giganteus* and *M. halli*) at Bird Island,  
283 South Georgia, and a Summary of Hybridization in Seabirds. *Plos One*, 10, e0121688.
- 284 Chapuis M. P., & Estoup, A. (2007) Microsatellite null alleles and estimation of population  
285 differentiation. *Molecular Biology and Evolution*, 24, 621–631.
- 286 Croxall, J.P., Steele, W.K., McInnes, S.J. & Prince, P. A. (1995). Breeding distribution of the  
287 Snow Petrel *Pagodroma nivea*. *Marine Ornithology*, 23, 69–99.
- 288 Croxall, J. P., Butchart, S. H. M., Lascelles, B. E. N., Stattersfield, A. J., Sullivan, B. E. N.,  
289 *et. al* (2012). Seabird conservation status—, threats and priority actions: a global  
290 assessment. *Bird Conservation International*, 22, 1–34.
- 291 Excoffier, L. & Lischer, H.E. L. (2010) Arlequin suite ver 3.5: A new series of programs to  
292 perform population genetics analyses under Linux and Windows. *Molecular Ecology*  
293 *Resources*, 10, 564-567.



- 294 Gómez-Díaz, E., González-Solís, J., & Peinado, M. A. (2009). Population structure in a  
295 highly pelagic seabird, the Cory's shearwater *Calonectris diomedea*: An examination of  
296 genetics, morphology and ecology. *Marine Ecology Progress Series*, 382, 197–209.
- 297 Jouventin, P., & Viot, C. R. (1985). Morphological and genetic variability of Snow Petrels  
298 *Pagodroma nivea*. *Ibis*, 127, 430–441.
- 299 Lascelles, B. G., Langham, G. M., Ronconi, R. A., & Reid, J. B. (2012). From hotspots to site  
300 protection: Identifying Marine Protected Areas for seabirds around the globe. *Biological*  
301 *Conservation*, 156, 5–14.
- 302 Lorensten, S., Amundsen, T., Anthonisen, K., & Lifjeld, J. (2000). Molecular evidence for  
303 extrapair paternity and female-female pairs in Antarctic petrels. *The Auk*, 117, 1042–  
304 1047.
- 305 Moodley, Y., Masello, J. F., Cole, T. L., Calderon, L., Munimanda, G. K., Thali, M. R. *et. al*  
306 (2015). Evolutionary factors affecting the cross-species utility of newly developed  
307 microsatellite markers in seabirds, *Molecular Ecology Resources*, 15, 1046–1058
- 308 Nunn, G. B., & Stanley, S. E. (1998). Body size effects and rates of cytochrome b evolution  
309 in tube-nosed seabirds. *Molecular Biology and Evolution*, 15, 1360–1371.
- 310 Paleczny, M., Hammill, E., Karpouzi, V., & Pauly, D. (2015a). Population Trend of the  
311 World's Monitored Seabirds, 1950-2010. *Plos One*, 10, e0129342.
- 312 Pande, A., Sivakumar, K., Sathyakumar, S., Kumar, R.S., Johnson, J.A., Mondol, S., &  
313 Mathur, V. B. (2017). Monitoring Wildlife and their Habitats in the Southern Ocean and  
314 Around Indian Research Stations in Antarctica. *Proceedings of the Indian National*  
315 *Science Academy*, 83, 483–496.
- 316 Piatt, J. F., Sydeman, W. J., & Wiese, F. (2007). Introduction: A modern role for seabirds as

- 317 indicators. *Marine Ecology Progress Series*, 352, 199–204.
- 318 Quillfeldt, P., Moodley, Y., Weimerskirch, H., Cherel, Y., Delord, K., *et. al* (2017). Does  
319 genetic structure reflect differences in non-breeding movements ? A case study in small ,  
320 highly mobile seabirds. *BMC Evolutionary Biology*, 17, 1-11.
- 321 Quillfeldt, P., Schmoll, T., Peter, H. U., Epplen, J. T., & Lubjuhn, T. (2001). Genetic  
322 Monogamy in Wilson’s Storm-Petrel. *The Auk*, 118, 242.
- 323 Taylor, S. A., & Friesen, V. L. (2012). Use of molecular genetics for understanding seabird  
324 evolution, ecology and conservation. *Marine Ecology Progress Series*, 451, 285–304.
- 325 Techow, N. M. S. M., & O’Ryan, C. (2004). Characterization of microsatellite loci in White-  
326 chinned Petrel (*Procellaria aequinoctialis*) and cross-amplification in six other  
327 procellariiform species. *Molecular Ecology Notes*, 4, 33–35.
- 328 Techow, N. M. S. M., O’Ryan, C., Phillips, R. A., Gales, R., Marin, M., *et. al* (2010).  
329 Speciation and phylogeography of giant petrels *Macronectes*. *Molecular Phylogenetics*  
330 *and Evolution*, 54, 472–487.
- 331 Valière, N. (2002) GIMLET: a computer program for analysing genetic individual  
332 identification data. *Molecular Ecology Notes* 2, 377–379.
- 333 Viot, C. R., Jouventin, P., & Bried, J. (1993). Population genetics of southern seabirds.  
334 *Marine Ornithology*, 21, 1-5.
- 335 Waits, L., Taberlet, P., & Luikart, G. (2001). Estimating the probability of identity among  
336 genotypes in natural populations: cautions and guidelines. *Molecular Ecology*, 10, 249–  
337 256.
- 338 Warham, J. (1996). *The behaviour, population biology and physiology of the Petrels*.  
339 Academic Press, 623 pp.

340 Welch, A. J., & Fleischer, R. C. (2011). Polymorphic microsatellite markers for the  
 341 endangered Hawaiian petrel (*Pterodroma sandwichensis*). *Conservation Genetics*  
 342 *Resources*, 3, 581–584.

343 Welch, A. J., Fleischer, R. C., James, H. F., Wiley, A. E., Ostrom, P. H., *et. al* (2012).  
 344 Population divergence and gene flow in an endangered and highly mobile seabird.  
 345 *Heredity*, 109, 19–28.

346

**Table 1** (on next page)

Table 1

Characterization of microsatellite loci genotyped in snow petrel and Wilsons's storm petrel individuals from Antarctica

**Table 1** Characterization of microsatellite loci genotyped in snow petrel and Wilson's storm petrel individuals from Antarctica

Species	Locus	Nucleotide Repeat nature	Dye	Product size range (bp)	Ta (°C)	Number of alleles	Ho	He	Allelic Range	PID (unrelated) Cumulative	PID (sibs) cumulative	Amplification success (%)	Allelic dropout (%)	Multiplex sets for PCR
Snow Petrel (n=55)	Ptero01	Di	PET	82-104	53	5	0.33	0.32	24	4.752 x 10 <sup>-01</sup>	7.072 x 10 <sup>-01</sup>	98.2	0	Set 1
	Ptero07	Tetra	FAM	177-217	53	8	0.53	0.66	48	7.599 x 10 <sup>-02</sup>	3.266 x 10 <sup>-01</sup>	98.2	3.6	
	Paequ03	Di	VIC	219-243	53	12	0.68	0.72	24	8.205 x 10 <sup>-03</sup>	1.362 x 10 <sup>-01</sup>	98.2	0	
	Ptero03	Di	FAM	165-177	53	4	0.10	0.23	22	4.917 x 10 <sup>-03</sup>	1.067 x 10 <sup>-01</sup>	100	0	
	Paequ08	Di	PET	215-223	53	4	0.16	0.18	8	3.291 x 10 <sup>-03</sup>	8.799 x 10 <sup>-02</sup>	100	0	
	Paequ02	Di	PET	180-200	53	7	0.03	0.30	30	1.642 x 10 <sup>-03</sup>	6.379 x 10 <sup>-02</sup>	98.2	1.8	Set 2
	Ptero04	Di	FAM	117-147	53	11	0.67	0.63	32	2.906 x 10 <sup>-04</sup>	3.059 x 10 <sup>-02</sup>	100	0	
	Ptero08	Tetra	VIC	181-221	53	11	0.49	0.73	52	2.742 x 10 <sup>-05</sup>	1.253 x 10 <sup>-02</sup>	96.4	0	
	Paequ10	Di	NED	159-183	53	7	0.20	0.56	12	6.217 x 10 <sup>-06</sup>	6.593 x 10 <sup>-03</sup>	98.2	0	
	Paequ13	Di	PET	144-150	57	4	0.07	0.44	6	2.189 x 10 <sup>-06</sup>	4.063 x 10 <sup>-03</sup>	100	0	Set 3
	Paequ07	Di	FAM	314-320	57	3	0.30	0.40	6	8.665 x 10 <sup>-07</sup>	2.625 x 10 <sup>-03</sup>	100	0	
	Ptero09	Tetra	FAM	161-189	57	9	0.67	0.72	28	1.041 x 10 <sup>-07</sup>	1.106 x 10 <sup>-03</sup>	100	0	
	Mean (SD)					7.08 (3.01)	0.35(0.23)	0.49(0.19)	24.5(14.5)					
Wilson's Storm Petrel (n=24)	Ptero01	Di	PET	165-177	53	4	0.17	0.44	12	3.65 x 10 <sup>-01</sup>	6.19 x 10 <sup>-01</sup>	100	0	Set 1
	Paequ10	Di	NED	181-191	53	4	0.38	0.64	10	7.44 x 10 <sup>-02</sup>	2.99 x 10 <sup>-01</sup>	100	0	
	Ptero07	Tetra	FAM	177-217	53	6	0.42	0.76	40	7.29 x 10 <sup>-03</sup>	1.18 x 10 <sup>-01</sup>	100	0	
	Paequ03	Di	VIC	219-235	53	5	0.21	0.39	16	2.86 x 10 <sup>-03</sup>	7.73 x 10 <sup>-02</sup>	100	0	
	Ptero03	Di	FAM	88-104	57	2	0.17	0.35	16	1.38 x 10 <sup>-03</sup>	5.37 x 10 <sup>-02</sup>	91.7	0	Set 2
	Ptero04	Di	FAM	127-139	57	4	0.38	0.52	12	4.30 x 10 <sup>-04</sup>	3.05 x 10 <sup>-02</sup>	100	0	
	Paequ13	Di	PET	146-148	57	2	0.08	0.5	2	1.62 x 10 <sup>-04</sup>	1.82 x 10 <sup>-02</sup>	100	8.3	
	Paequ08	Di	PET	219-227	51	3	0.21	0.47	8	6.04 x 10 <sup>-05</sup>	1.10 x 10 <sup>-02</sup>	100	0	Set 3
	Ptero09	Tetra	FAM	173-185	61	6	0.67	0.55	16	1.43 x 10 <sup>-05</sup>	5.88 x 10 <sup>-03</sup>	91.7	0	
	Paequ07	Di	FAM	312-318	51	3	0.08	0.16	6	1.03 x 10 <sup>-05</sup>	5.01 x 10 <sup>-03</sup>	100	4.2	
	Mean (SD)					3.9 (1.3)	0.28 (0.18)	0.48 (0.15)	13.8(9.7)					




# Figure 1(on next page)

Figure 1

Seabird sampling locations in Antarctica. A) Schirmacher oasis, site of Maitri station B) Larsemann hills, site of Bharati station







Indian Research Station

Sampling locations

North

Scale



Wilson's Storm Petrel



Snow Petrel