

Prevalence and extent of chronic periodontitis and its risk factors in a Portuguese subpopulation: a retrospective cross-sectional study and analysis of Clinical Attachment Loss

Vanessa Machado^{1,2}, João Botelho^{Corresp., 1,2}, António Amaral², Luís Proença², Ricardo Alves^{1,2}, João Rua², Maria Alzira Cavacas³, Ana Sintra Delgado², José João Mendes²

¹ Department of Periodontology, Clinical Research Unit, Centro de Investigação Interdisciplinar Egas Moniz (CiiEM), Instituto Universitário Egas Moniz, Almada, Portugal

² Clinical Research Unit, Centro de Investigação Interdisciplinar Egas Moniz (CiiEM), Instituto Universitário Egas Moniz, Almada, Portugal

³ Environmental Health, Centro de Investigação Interdisciplinar Egas Moniz (CiiEM), Instituto Universitário Egas Moniz, Almada, Portugal

Corresponding Author: João Botelho

Email address: jbotelho@egasmoniz.edu.pt

Objectives: To assess the prevalence and extent of chronic periodontitis, and its risk factors in a Portuguese subpopulation referred to periodontal examination.

Methods: This retrospective cross-sectional study used a subset of data from patients who sought dental treatment in a university dental clinic in the Lisbon metropolitan area. The sample consisted of 405 individuals (225 females/180 males), aged 20-90 years. All patients underwent a full-mouth periodontal examination and chronic periodontitis was defined as Clinical Attachment Loss (CAL) \geq 3mm affecting two or more teeth. Aggressive periodontitis cases were excluded from the analysis.

Results: Prevalence of chronic periodontitis was 83.5% (95% CI 80.4-86.6%). For these subjects, CAL \geq 3 mm affected 86.0% (95% CI 84.7-87.2) of sites and 83.7% (95% CI 81.7-85.6) of teeth, respectively. Mean CAL ranged from 3.6 to 4.3 mm, according to age. In the multivariate logistic regression model, smoking (OR=3.55, 95% CI 1.80-7.02) and older age (OR=8.70, 95% CI 3.66-20.69 and OR=4.85, 95% CI 2.57-9.16), for 65+ and 45-64 years old, respectively, were identified as risk indicators for CAL \geq 3mm.

Conclusions: This particular Portuguese adult subpopulation had a high prevalence of chronic periodontitis, with severe and generalized clinical attachment loss, and its presence was significantly associated with age and smoking. This data should serve to prepare future detailed epidemiological studies and appropriate public health programs.

Prevalence and extent of chronic periodontitis and its risk factors in a Portuguese subpopulation: a **retrospective** cross-sectional study and analysis of Clinical Attachment Loss

Vanessa Machado^{1,2}, João Botelho^{1,2}, António Amaral², Luís Proença², Ricardo Alves^{1,2}, João Rua², Maria Alzira Cavacas³, Ana Sintra Delgado², José João Mendes²

¹Department of Periodontology, Clinical Research Unit, Centro de Investigação Interdisciplinar Egas Moniz (CiiEM), Instituto Universitário Egas Moniz, Almada, Portugal

²Clinical Research Unit, Centro de Investigação Interdisciplinar Egas Moniz (CiiEM), Instituto Universitário Egas Moniz, Almada, Portugal

³Environmental Health, Centro de Investigação Interdisciplinar Egas Moniz (CiiEM), Instituto Universitário Egas Moniz, Almada, Portugal

Corresponding author:

João Botelho

Clinical Research Unit

Centro de Investigação Interdisciplinar Egas Moniz (CiiEM)

Egas Moniz Cooperativa de Ensino Superior, C.R.L.

Campus Universitário, Monte da Caparica, 2829 - 511 Caparica, Almada, Portugal

Phone: +351 212 946 737

e-mail: jbotelho@egasmoniz.edu.pt.

Abstract

Objectives: To assess the prevalence and extent of chronic periodontitis, and its risk factors in a Portuguese subpopulation referred to periodontal examination.

Methods: This **retrospective** cross-sectional study used a subset of data from patients who sought dental treatment in a university dental clinic in the Lisbon metropolitan area. The sample consisted of 405 individuals (225 females/180 males), aged 20-90 years. All patients underwent

a full-mouth periodontal examination and chronic periodontitis was defined as Clinical Attachment Loss (CAL) \geq 3 mm affecting two or more teeth. Aggressive periodontitis cases were excluded from the analysis.

Results: Prevalence of chronic periodontitis was 83.5% (95% CI 80.4-86.6%). For these subjects, CAL \geq 3 mm affected 86.0% (95% CI 84.7-87.2) of sites and 83.7% (95% CI 81.7-85.6) of teeth, respectively. Mean CAL ranged from 3.6 to 4.3 mm, according to age. In the multivariate logistic regression model, smoking (OR=3.55, 95% CI 1.80-7.02) and older age (OR=8.70, 95% CI 3.66-20.69 and OR=4.85, 95% CI 2.57-9.16), for 65+ and 45-64 years old, respectively, were identified as risk indicators for CAL \geq 3 mm.

Conclusions: This particular Portuguese adult subpopulation had high prevalence of chronic periodontitis, with severe and generalized clinical attachment loss, and its presence was significantly associated with age and smoking. This data should serve to prepare future detailed epidemiological studies and appropriate public health programs.

INTRODUCTION

Chronic periodontitis is an inflammatory disease characterized by a polymicrobial breakdown of host homeostasis and a progressive destruction of tooth-supporting structures,^{1,2} and its epidemiology and risk factors have been broadly studied.³⁻⁶

Periodontal diseases have a significant impact on oral health-related quality of life, especially with the worsening and extension of the disease in which it presents higher destructive consequences.⁷ There are important risk factors/indicators for periodontal disease such as alcohol,⁸ overweight and obesity,⁹ smoking⁶ and diabetes.¹⁰ Also, periodontitis can be a risk factor for several systemic diseases.¹⁰⁻¹⁸

Some European epidemiological studies have demonstrated the high prevalence of periodontitis among the populations.¹⁹⁻²³ However, data on the prevalence and risk factors for periodontal disease in the Portuguese population are still missing. According to the latest Portuguese Oral National Health Survey, the prevalence of periodontitis was 10.8% in adults and 15.3% in the elderly.²⁴ This nationwide survey used the Community Periodontal Index (CPI), with its

recognized limitations. To the best of our knowledge, there are no epidemiological studies that used full-mouth periodontal examination (FMPE) methodology to estimate the prevalence of periodontitis regarding Portuguese samples.

The aim of this study was to assess the prevalence, severity, and extent of chronic periodontitis through the full-mouth examination of CAL, and its association with sociodemographic, behavioral and environmental risk factors, in a Portuguese adult subpopulation, of a suburban area of the Lisbon Region, forwarded to periodontal examination.

MATERIAL AND METHODS

The study was conducted in accordance with the Declaration of Helsinki of 1975, as revised in 2013, and approved by the Ethics Committee of Egas Moniz (Ethical Application Ref: 595). A written informed consent was obtained from all participants during the first appointment. After the examination, the participants were informed of their periodontal status, and those with diagnosed periodontal diseases were advised to follow the proper treatment. This protocol followed the STrengthening the Reporting of OBservational studies in Epidemiology (STROBE) guidelines.²⁵

Study subjects

All participants were patients of Egas Moniz Dental Clinic (Almada, Portugal). This university clinic, located in the municipality of Almada, in Setúbal Peninsula (a NUTS III subregion, part of NUTS II Lisbon Region), provides dental health services to the general public.

At the first appointment, patients were submitted to a dental triage protocol, with the application of a self-reported health questionnaire and oral and dental examinations, to guide their treatment needs. Regarding periodontal triage, patients were assessed using the Periodontal Screening and Recording (PSR) procedure²⁶, and, if diagnosed with code 2 or higher, they were forwarded to a periodontology appointment.

Patient selection

This retrospective cross-sectional study analyzed patients who attended the dental clinic between September 2015 and March 2017. From a total of 3648 subjects who sought the first consultation in the university dental clinic during that period, 1501 (41%) patients were referred to the periodontology department, based on their triage status. From these, 459 attended a periodontal consultation and were considered for this study. Fifty-two participants were excluded due to incomplete questionnaires and periodontal data, and two subjects diagnosed with aggressive periodontitis. Hence, a final sample size of 405 subjects was obtained (11% of the total, 27% of the patients forwarded for periodontal treatment).

Health questionnaire

Before clinical examinations, all patients answered a general and oral health questionnaire that included information such as age, gender, educational level, employment status, general medical history and medication, smoking status and oral hygiene habits.

Clinical data

Periodontal status

Five well-trained and calibrated periodontists (R.A., J.C., C.I., F.J., L.A.) performed all dental and periodontal examinations. Periodontal examinations were performed using CDC/AAP full-mouth periodontal examination (FMEP) methodology.²⁷ We defined chronic periodontitis as defined as $CAL \geq 3$ mm affecting two or more teeth.²⁸ All permanent fully erupted teeth were examined, excluding third molars, retained roots, and implants. The evaluated parameters were: missing teeth, presence or absence of supragingival biofilm (SB), probing depth (PD), bleeding on probing (BOP), gingival recession (REC) and clinical attachment loss (CAL). SB and BOP were scored on four surfaces of each tooth (mesial, distal, buccal and lingual). At six sites per tooth (mesiobuccal, mid-buccal, distobuccal, mesiolingual, mid-lingual and distolingual), PD was measured as the distance from the cemento-enamel junction (CEJ) to the bottom of the pocket and REC as the distance from the CEJ to the free gingival margin, and this assessment was assigned a negative sign if the gingival margin was located coronally to the CEJ. CAL was

calculated as the algebraic sum of PD and REC. It was used a CP-12 SE (Hu-Friedy, Chicago, IL, USA).

Measurement reproducibility

Prior to the initiation of the study, all examiners were submitted to theoretical and practical training in a total of ten volunteer non-study patients suffering from moderate to severe periodontitis. The inter-examiner correlation coefficients, at subject level, ranged from 0.76 to 0.97 and between 0.91 and 0.99, for mean PD and mean CAL, respectively.

Covariates

Sociodemographic variables and several periodontal disease risk factors were selected as confounding variables. The selected variables were: age, gender, educational level, employment status, smoking status, Body Mass Index (BMI), time elapsed since the last dental appointment, consultation motive and oral hygiene habits.

Educational level was assessed as three categories: elementary (1-4 years), middle (5-12 years) and higher (> 12 years) education. Employment status of each participant was classified as: employed, unemployed or retired. Smoking status was defined as non-smoker or smoker. Active smokers were further divided into three categories: light smokers (< 10 cigarettes per day), medium smokers (10-20), heavy smokers (> 20). The height of the participants was measured in centimeters, using a hard ruler installed vertically and secured with a stable base. Weight was assessed in kilograms using mechanical scales. BMI was calculated as the ratio of the individual's body weight to the square of their height. Four BMI categories were defined using WHO criteria ²⁹: underweight ($\text{BMI} < 18.5 \text{ kg/m}^2$), normal weight ($\text{BMI} 18.5 - 24.9 \text{ kg/m}^2$), overweight ($\text{BMI} 25 - 29.9 \text{ kg/m}^2$) and obese ($\text{BMI} \geq 30 \text{ kg/m}^2$). The time elapsed since last dental consult was classified into five categories (never visited, less than one year, 1-2 years, 3-4 years, 5 years or over). Consultation motives were classified as routine, aesthetics, pain, functional or other. Oral hygiene habits were assessed by information on toothbrush frequency (2-3 times / daily, one time daily, 2-6 times/weekly) and dental floss use.

Data analysis

Data analysis was performed using IBM SPSS Statistics version 24.0 for Windows (Armonk, NY: IBM Corp.). Descriptive and inferential statistics methodologies were applied. In the latter, Mann-Whitney and Kruskal-Wallis tests were used to compare the clinical data as a function of the sociodemographic variables. Further, logistic regression analysis was used to model the relationship between chronic periodontitis and several risk indicators. Preliminary analyses were performed using univariate models. Next, a multivariate model was constructed for the outcome variable $CAL \geq 3$ mm. Only variables showing a significance $p \leq 0.25$ in the univariate model were included in the multivariate stepwise procedure. Predictor variables considered in this procedure were: age (years), smoking status, education (years), employment status, last dental visit and dental floss use. The contribution of each variable to the model was evaluated by Wald statistics. Interactions were also analyzed for all tested variables. The final reduced model was obtained with the following predictor variable categories: age (45-64 and ≥ 65 years) and smoking status (smoker). Odds ratio (OR) and 95% confidence intervals (95% CI) were calculated for both univariate and multivariate analyses. The level of statistical significance was set at 5%.

RESULTS

Table 1 shows the distribution of sociodemographic, behavioral, biometric and oral hygiene data in the studied sample. Ages ranged from 20 to 90 years. The sample had 55.6% of female patients. It is worth to mention that 65.2% of subjects did not smoke and active smokers were mainly medium smokers (66%), followed by light smokers (29%) and heavy smokers (5%). Regarding education and employment status 77.1% of subjects had ~~an~~ elementary or middle ~~level~~ education, and 51.9% of the subjects were employed. Approximately 59% were overweight and obese, and only 40% had normal values. Interestingly, 53.1% had a period of over one year without any dental visit and 1.2% never had a dental appointment, whereas functional complaint was the major consultation motive.

Table 2 shows the periodontal data of this sample according to age, gender, and smoking status. Subjects over 65 years of age had a significantly higher mean number of missing teeth and, in total, this **subpopulation** presented a mean loss of 8 teeth. Younger individuals (< 45 years of age) presented a significantly lower mean number of missing teeth, PD, REC, furcation lesions and teeth with mobility compared to older subjects. Male patients presented a significantly higher mean PD, deep periodontal pockets (≥ 5 mm) and teeth with furcation lesions than female. Compared to smokers, non-smokers had lower mean SB, PD and CAL, and less deep periodontal pockets.

Chronic periodontitis was diagnosed in 83.5% of the patients (Table 3), and subjects with chronic periodontitis had CAL ≥ 3 mm, ≥ 4 mm, ≥ 5 mm, ≥ 6 mm and ≥ 7 mm affecting, on average, 83.7%, 54.4%, 32.1%, 17.8% and 9.2% of their teeth, respectively (Table 4). Besides, the first lower molar was the most frequently missing tooth, while the lower canine was the least lost but the most severely affected tooth (Fig. 1).

In the logistic regression analysis, similar results were observed in the univariable (Table 5) and multivariable models (Table 6). In the multivariable analysis, smoking (OR=3.55, 95% CI 1.80-7.02) and older age (OR=8.70, 95% CI 3.66-20.69 and OR=4.85, 95% CI 2.57-9.16), for 65+ and 45-64 years old, respectively, were identified as risk indicators for CAL ≥ 3 mm (table 5). Chronic periodontitis was not significantly associated with the remaining variables.

DISCUSSION

This **retrospective** cross-sectional study assessed the periodontal status of forwarded adult subjects who sought dental treatment in a Portuguese university dental clinic, that is located in the metropolitan area of Lisbon. This area has over 2.8 million inhabitants and is the largest Portuguese metropolitan area ³⁰. This university dental clinic is an important reference dental center in the Lisbon Region and receives patients from all social strata. The absence of complete socioeconomic data constitutes a limitation of this study. Unfortunately, over 70% of patients (data not shown) refused to provide socioeconomic status information.

The results of this **retrospective** study can't be compared with previous investigations performed in Portugal because in these it was applied the CPITN methodology.^{24,31-33} This is the first FMPE protocol used in a Portuguese population and provides direct evidence for estimating periodontal

status and results in a better representation of the population.²⁷ Although FMPE methodology can result in an overestimation of periodontal treatment needs among young adults,²¹ the partial-mouth examination can miscalculate the prevalence of periodontitis in almost 50% of the population.²⁷ The overall results demonstrate that this referred subpopulation had a high prevalence of chronic periodontitis (79.3%, 95% CI 77.5-88.1%), and severe extensity of periodontal destruction among the affected subjects (83.7%, 95% CI 81.7-85.6%).

This investigation study design is not an epidemiological study per se, but rather an observational study of patients who were forwarded to a periodontology consultation. Thus, we were only able to estimate the prevalence and extent of our referred subpopulation. However, these results underline the fact that the majority of patients attended the periodontal consultation already in a state of advanced periodontal destruction and only a small percentage appeared in the early stages or healthy. Still, a disturbing percentage of patients did not attend periodontal consultations despite the triage referral with approximately 69% missing or unchecking the appointment.

Regarding tooth loss, the most frequently missed teeth were the lower first molars and the less missed were the lower canines, as with recent European data.^{21,22} Additionally, lower canines and incisors were the most affected teeth with CAL and the lower molars the less. The lower arch presented more periodontal destruction than the upper, and the teeth with more severe CAL levels in the upper arch were the canines.

Concerning periodontal parameters, unlike PD, CAL severity increased with age and can be related to the increase of gingival recession with aging.³⁴ As in the literature,¹⁹⁻²³ age was confirmed in the multivariate analysis as a risk indicator for chronic periodontitis for 45-64 years old (OR=4.85, 95% CI 2.57-9.16) and 65+ years old (OR=8.70, 95% CI 3.66-20.69). However, it's important to highlight that, in the majority of CAL thresholds of subjects with the disease, 45-64 years old group presented worse results for prevalence of chronic periodontitis, while 65+ years old group had worse levels of periodontal destruction extent.

Smoking was strongly associated with chronic periodontitis (OR = 3.55, 95% CI 1.80-7.02). Previous studies reported OR values ranging between 2 and 9 of having periodontitis.^{21-23,35-38} Despite not accounting for lifetime smoking exposure, we stratified current smokers according to

the number of cigarettes smoked although it was not significantly associated with the severity and progression of the periodontal disease.

Several studies found that obesity was associated with an increased risk of periodontitis.³⁹⁻⁴¹ Besides that, Suvan et al.⁴² concluded that overweight/obese individuals are more likely to suffer from periodontitis compared to normal weight individuals. Although our results show that overweight and obesity have no impact on the aggravation of periodontitis, we emphasize that more than half of this **subpopulation** was overweight or obese, in agreement with the latest national IAN-AF Food and Activity Survey.⁴³

In the past, several epidemiological surveys reported that people with lower educational level had higher prevalence and severity of periodontal disease.^{20,21,23,44,45} However, other studies have indicated that this impact cannot be seen in a singular way but in a multifactorial view.^{3,46} Our results show that despite middle education had significance in the univariable model (OR = 2.09 (95% CI 1.05-4.13), p = 0.035), when analyzed in a multivariable model it had no impact on the probability of having chronic periodontitis.

CONCLUSION

This specific subpopulation of individuals referred to periodontal examination in a university dental clinic of the Lisbon region presented high prevalence and severe extent of chronic periodontitis. Age and smoking were identified as risk indicators for chronic periodontitis **in this referred subpopulation**. Within the limitations of this study, these results highlight the importance of developing appropriate public health programs to educate the Portuguese population about the burden of periodontal diseases.

ACKNOWLEDGMENTS

We thank Júlio Guilherme for his substantial support **for** database development and management during this study.

262 REFERENCES

- 263 1. Pihlstrom BL, Michalowicz BS, Johnson NW. Periodontal diseases. *Lancet*.
264 2005;366(9499):1809–20.
- 265 2. Darveau RP. Periodontitis: a polymicrobial disruption of host homeostasis. *Nat Rev*
266 *Microbiol*. 2010;8(7):481–90.
- 267 3. Albandar JM. Global risk factors and risk indicators for periodontal diseases. *J*
268 *Periodontol* 2000. 2002;29:177–206.
- 269 4. Albandar JM, Rams TE. Global epidemiology of periodontal diseases: an overview.
270 *Periodontol* 2000. 2002;29(1):7–10.
- 271 5. Brett PM, Zygogianni P, Griffiths GS, Tomaz M, Parkar M, D’Aiuto F, Tonetti M.
272 Functional gene polymorphisms in aggressive and chronic periodontitis. *J Dent Res*.
273 2005;84(12):1149–53.
- 274 6. Burt B. Position paper - Epidemiology of periodontal diseases. *J Periodontol*.
275 2005;76(8):1406–19.
- 276 7. Buset SL, Walter C, Friedmann A, Weiger R, Borgnakke WS, Zitzmann NU. Are
277 periodontal diseases really silent? A systematic review of their effect on quality of life. *J*
278 *Clin Periodontol*. 2016;43(4):333–44.
- 279 8. Wang J, Lv J, Wang W, Jiang X. Alcohol consumption and risk of periodontitis: A meta-
280 analysis. *J Clin Periodontol*. 2016;43(7):572–83.
- 281 9. Keller A, Rohde JF, Raymond K, Heitmann BL. Association Between Periodontal Disease
282 and Overweight and Obesity: A Systematic Review. *J Periodontol*. 2015;86(6):766–76.
- 283 10. Preshaw PM, Alba AL, Herrera D, Jepsen S, Konstantinidis A, Makrilakis K, Taylor R.
284 Periodontitis and diabetes: a two-way relationship. *Diabetologia*. 2012;55(1):21–31.
- 285 11. Bahekar AA, Singh S, Saha S, Molnar J, Arora R. The prevalence and incidence of
286 coronary heart disease is significantly increased in periodontitis: A meta-analysis. *Am*
287 *Heart J*. 2007;154(5):830–7.
- 288 12. Humphrey LL, Fu R, Buckley DI, Freeman M, Helfand M. Periodontal disease and
289 coronary heart disease incidence: A systematic review and meta-analysis. *J Gen Intern*

- Med. 2008;23(12):2079–86.
13. Nibali L, Tatarakis N, Needleman I, Tu Y-K, D’Aiuto F, Rizzo M, Donos N. Association Between Metabolic Syndrome and Periodontitis: A Systematic Review and Meta-analysis. *J Clin Endocrinol Metab.* 2013;98(3):913–20.
14. Lafon A, Pereira B, Dufour T, Rigouby V, Giroud M, Béjot Y, Tubert-Jeannin S. Periodontal disease and stroke: A meta-analysis of cohort studies. *Eur J Neurol.* 2014;21(9):1155–61.
15. Leira Y, Seoane J, Blanco M, Rodríguez-Yáñez M, Takkouche B, Blanco J, Castillo J. Association between periodontitis and ischemic stroke: a systematic review and meta-analysis. *Eur J Epidemiol.* 2017;32(1):43–53.
16. Fuggle NR, Smith TO, Kaul A, Sofat N. Hand to mouth: A systematic review and meta-analysis of the association between rheumatoid arthritis and periodontitis. *Front Immunol.* 2016;7(MAR):1–10.
17. Maisonneuve AP, Amar S, Lowenfels AB. Periodontal Disease , Edentulism and Pancreatic Cancer : A Meta Analysis. *Ann Oncol.* 2017;28(5):985–95.
18. Papageorgiou SN, Hagner M, Nogueira AVB, Franke A, Jäger A, Deschner J. Inflammatory bowel disease and oral health: systematic review and a meta-analysis. *J Clin Periodontol.* 2017;44(4):382–93.
19. Bouchard P, Boutouyrie P, Mattout C, Bourgeois D. Risk assessment for severe clinical attachment loss in an adult population. *J Periodontol.* 2006;77(3):479–89.
20. Bourgeois D, Bouchard P, Epidemiology MC. Epidemiology of periodontal status in dentate adults in. *J Periodontal Res.* 2007;42(2):219–27.
21. Aimetti M, Perotto S, Castiglione A, Mariani GM, Ferrarotti F, Romano F. Prevalence of periodontitis in an adult population from an urban area in North Italy : findings from a cross-sectional epidemiological survey. *J Clin Periodontol.* 2015;(42):622–31.
22. Schutzhold S, Kocher T, Biffar R, Hoffmann T, Schmidt C, Micheelis W, Jordan R, Holtfreter B. Changes in prevalence of periodontitis in two German population-based studies. *J Clin Periodontol.* 2015;(42):121–30.

23. Holde GE, Oscarson N, Trovik TA, Tillberg A, Jönsson B. Periodontitis Prevalence and Severity in Adults: A Cross-Sectional Study in Norwegian Circumpolar Communities. *J Periodontol.* 2017;1–17.
24. DGS. III Estudo de Prevalência das Doenças Orais. 2015.
25. von Elm E, Altman DG, Egger M, Pocock SJ, Gøtzsche PC, Vandenbroucke JP. The strengthening the reporting of observational studies in epidemiology (STROBE) statement: Guidelines for reporting observational studies. *Int J Surg.* 2014;12(12):1495–9.
26. Landry RG, Jean M. Periodontal Screening and Recording (PSR) Index: Precursors, utility and limitations in a clinical setting. *Int Dent J.* 2002;52(1):35–40.
27. Eke PI, Page RC, Wei L, Thornton-Evans G, Genco RJ. Update of the Case Definitions for Population-Based Surveillance of Periodontitis. *J Periodontol.* 2012;83(12):1449–54.
28. Susin C, Haas AN, Valle PM, Oppermann R V., Albandar JM. Prevalence and risk indicators for chronic periodontitis in adolescents and young adults in south Brazil. *J Clin Periodontol.* 2011;38(4):326–33.
29. Kopelman PG. Obesity as a medical problem. *Nature.* 2000;404(6778):635–43.
30. Área Metropolitana de Lisboa. Municípios AML [Internet]. 2018. p. (accessed 29 January 2018). Available from: <https://www.aml.pt/index.php>
31. Freitas E. A baseline study of periodontal conditions among youth and adults in portugal. Vol. 11, Community dental oral epidemiology. 1983. p. 250–4.
32. Marques MD, Teixeira-Pinto A, da Costa-Pereira A, Eriksen HM. Prevalence and determinants of periodontal disease in Portuguese adults: results from a multifactorial approach. *Acta Odontol Scand.* 2000;58(5):201–6.
33. Petersen PE, Ogawa H. The global burden of periodontal disease: Towards integration with chronic disease prevention and control. *Periodontol 2000.* 2012;60(1):15–39.
34. Eke PI, Dye BA, Wei L, Thornton-Evans GO, Genco RJ. Prevalence of Periodontitis in Adults in the United States: 2009 and 2010. *J Dent Res.* 2012;91(10):914–20.
35. Bergström J, Preber H. Tobacco Use as a Risk Factor. *J Periodontol.* 1994;65(5):545–50.

36. Bergström J. Periodontitis and Smoking: An Evidence-Based Appraisal. *J Evid Based Dent Pract.* 2006;6(1):33–41.
37. Kinane DF, Stathopoulou PG, Papapanou PN. Periodontal diseases. *Nat Rev Dis Prim.* 2017;3:17038.
38. Silva-Boghossian CM, Luiz RR, Colombo AP V. Periodontal Status, Sociodemographic, and Behavioral Indicators in Subjects Attending a Public Dental School in Brazil: Analysis of Clinical Attachment Loss. *J Periodontol.* 2009;80(12):1945–54.
39. Nishida N, Tanaka M, Hayashi N, Nagata H, Takeshita T, Nakayama K, Morimoto K, Shizukuishi S. Determination of smoking and obesity as periodontitis risks using the classification and regression tree method. *J Periodontol.* 2005;76(6):923–8.
40. Saito T, Shimazaki Y, Kiyohara Y, Kato I, Kubo M, Iida M, Yamashita Y. Relationship between obesity, glucose tolerance, and periodontal disease in Japanese women: The Hisayama study. *J Periodontal Res.* 2005;40(4):346–53.
41. Dalla Vecchia CF, Susin C, Ro CK, Oppermann RV, Albandar JM. Overweight and Obesity as Risk Indicators for Periodontitis in Adults. *J Periodontol.* 2005;76(10):1721–8.
42. Suvan JE, Petrie A, Nibali L, Darbar U, Rakmanee T, Donos N, D’Aiuto F. Association between overweight/obesity and increased risk of periodontitis. *J Clin Periodontol.* 2015;42(8):733–9.
43. IAN-AF. Inquérito Alimentar Nacional e de Atividade Física 2015-2016. 2016.
44. Albandar JM, Brunelle JA, Kingman A. Destructive periodontal disease in adults 30 years of age and older in the United States, 1988-1994. *J Periodontol.* 1999;70(1):13–29.
45. Krstrup U, Erik Petersen P. Periodontal conditions in 35-44 and 65-74-year-old adults in Denmark. *Acta Odontol Scand.* 2006;64(2):65–73.
46. Geyer S, Schneller T, Micheelis W. Social gradients and cumulative effects of income and education on dental health in the fourth German oral health study. *Community Dent Oral Epidemiol.* 2010;38(2):120–8.

Figure 1

Percentage of subjects with the respective tooth present and by thresholds of CAL (mm), at each specific position, for all teeth in all quadrants

The black lines indicate the separation by each quadrant. (Dark blue) percentage of missing teeth, (Blue) percentage of teeth with less than 3mm of CAL, (pink) percentage of teeth with 3-4 mm of CAL, (yellow) percentage of teeth over 4 mm of CAL




Table 1(on next page)

Sociodemographic, behavioral, biometric and oral hygiene data (N = 405).

1 Table 1 - Sociodemographic, behavioural, biometric and oral hygiene data (*N* = 405)

2

3

Variable		<i>n</i> (%)
Gender	Female	225 (55.6)
	Male	180 (44.4)
Age (years)	20 - 44	90 (22.2)
	45 - 64	217 (53.6)
	≥ 65	98 (24.2)
Smoking status	Smoker	141 (34.8)
	Non-smoker	264 (65.2)
Active smokers (cigarettes per day) (<i>n</i> =141)	Light (< 10)	41 (29.1)
	Medium (10-20)	93 (66.0)
	Heavy (>20)	7 (5.0)
Education	Elementary	157 (38.8)
	Middle	155 (38.3)
	Higher	93 (23.0)
Employment status	Employed	210 (51.9)
	Unemployed	63 (15.6)
	Retired	132 (32.6)
BMI (kg/m ²)	< 18.5	5 (1.2)
	18.5 - 24.9	162 (40.0)

	25.0 - 29.9	159 (39.3)
	≥ 30	79 (19.5)
Last dental visit	< 1 year	185 (45.7)
	1 - 2 years	57 (14.1)
	3 - 4 years	75 (18.5)
	≥ 5 years	83 (20.5)
	Never	5 (1.2)
Consultation motive	Routine	125 (30.9)
	Aesthetics	35 (8.6)
	Pain	73 (18.0)
	Functional	157 (38.8)
	Other	15 (3.7)
Dental floss usage	Yes	141 (34.8)
	No	264 (65.2)
Toothbrush frequency	2 - 3 times / daily	313 (77.3)
	1 time / daily	75 (18.5)
	2 - 6 times / weekly	17 (4.2)

BMI (kg/m²) – Body Mass Index (kilogram/meter²)

4
5
6
7

Table 2(on next page)

Periodontal clinical data (presented as mean \pm standard deviation) as a function of gender, age and smoking status (N = 405).

Table 2 - Periodontal clinical data (presented as mean \pm standard deviation) as a function of gender, age and smoking status ($N = 405$).

		SB (%)	BOP (%)	PD (mm)	REC (mm)	Missing teeth (n)	Teeth w/ mobility (n)	Teeth w/ furcation lesions (n)	Deep periodontal pockets (≥ 5 mm) (n)	CAL			
										Total	≥ 3 mm (%)	≥ 5 mm (%)	≥ 7 mm (%)
Gender	Female	34.6 \pm 22.8	9.1 \pm 12.6	3.1 \pm 0.7 *	1.0 \pm 0.9	8.5 \pm 5.9	5.2 \pm 5.0	0.4 \pm 0.8 *	15.0 \pm 18.8 *	4.0 \pm 1.2	77.6 \pm 19.8 *	33.2 \pm 24.8	11.1 \pm 15.8 *
	Male	37.4 \pm 23.6	11.3 \pm 15.7	3.3 \pm 0.8 *	1.0 \pm 0.9	8.1 \pm 5.6	4.4 \pm 4.2	0.5 \pm 0.9 *	20.1 \pm 19.8 *	4.3 \pm 1.5	81.5 \pm 17.7 *	38.4 \pm 27.1	14.8 \pm 18.9 *
Age (years)	20 - 44	33.3 \pm 20.9	9.9 \pm 12.3	3.1 \pm 0.7	0.6 \pm 0.7 **	5.3 \pm 5.0 **	4.0 \pm 4.7 **	0.2 \pm 0.5 **	18.4 \pm 19.9	3.6 \pm 1.2 **	72.5 \pm 21.2 **	25.6 \pm 24.3 **	7.6 \pm 15.1 **
	45 - 64	35.4 \pm 23.5	10.1 \pm 14.6	3.3 \pm 0.8	1.0 \pm 0.9 **	8.5 \pm 5.5 **	5.4 \pm 5.0 **	0.5 \pm 0.9 **	19.1 \pm 21.2	4.3 \pm 1.3 **	81.5 \pm 18.2 **	38.5 \pm 25.9 **	13.9 \pm 17.4 **
	≥ 65	39.0 \pm 24.1	10.1 \pm 14.5	3.1 \pm 0.7	1.2 \pm 1.0 **	10.7 \pm 6.0 **	4.5 \pm 3.6 **	0.6 \pm 0.9 **	12.3 \pm 12.8	4.3 \pm 1.4 **	80.8 \pm 17.2 **	38.0 \pm 25.5 **	14.7 \pm 18.3 **
Smoking status	Smoker	38.7 \pm 23.9 *	8.6 \pm 14.2	3.4 \pm 0.8 *	1.1 \pm 1.0	8.2 \pm 5.8	5.3 \pm 5.2	0.4 \pm 0.8	22.5 \pm 22.0 *	4.5 \pm 1.4 *	85.5 \pm 16.1 *	42.9 \pm 28.6 *	15.8 \pm 19.6 *
	Non-smoker	34.3 \pm 22.6 *	10.8 \pm 13.9	3.1 \pm 0.7 *	0.9 \pm 0.8	8.4 \pm 5.8	4.6 \pm 4.3	0.5 \pm 0.9	14.5 \pm 17.3 *	3.9 \pm 1.3 *	76.0 \pm 19.6 *	31.6 \pm 23.5 *	11.1 \pm 15.8 *
Total		35.8 \pm 23.1	10.1 \pm 14.1	3.2 \pm 0.8	1.0 \pm 0.9	8.3 \pm 5.8	4.8 \pm 4.6	0.4 \pm 0.9	17.3 \pm 19.4	4.1 \pm 1.3	79.3 \pm 19.0	79.3 \pm 19.0	79.3 \pm 19.0

SB – Supragingival Biofilm, BOP – Bleeding on Probing, PD – Pocket Depth, REC – Recession, CAL – Clinical Attachment Loss

* Mann-Whitney test ($p < 0.05$), ** Kruskal-Wallis test ($p < 0.05$)

Table 3(on next page)

Percentage of patients, with 95% confidence interval (95% CI), by threshold of CAL (mm), severity and age group (years).

1 Table 3 - Percentage of patients, with 95% confidence interval (95% CI), by threshold of CAL (mm), severity and age group (years)

2

CAL (mm)	Subjects with chronic periodontitis								All subjects							
	20-44 (n = 59)		45-64 (n = 190)		≥ 65 (n = 89)		Total (n = 338)		20-44 (n = 90)		45-64 (n = 217)		≥ 65 (n = 98)		Total (n = 405)	
	%	95% CI	%	95% CI	%	95% CI	%	95% CI	%	95% CI	%	95% CI	%	95% CI	%	95% CI
Prevalence (patients)																
≥ 3	100	100.0-100.0	100	100.0-100.0	100	100.0-100.0	100	100.0-100.0	65.6	56.1-75.1	87.6	83.5-91.7	90.8	85.3-96.3	83.5	80.4-86.6
≥ 4	42.4	30.0-54.8	62.1	55.7-68.6	59.6	49.7-69.5	58.0	53.4-62.6	27.8	18.8-36.8	54.4	48.3-60.5	54.1	44.6-63.6	48.4	44.2-52.6
≥ 5	20.3	10.2-30.4	30.0	23.9-36.1	25.8	17.0-34.6	15.4	12.0-18.8	13.3	6.5-20.1	26.3	20.9-31.7	23.5	15.4-31.6	12.8	10.0-15.6
≥ 6	8.5	1.5-15.5	13.2	8.7-17.7	12.4	5.8-19.0	12.1	9.0-15.2	5.6	1.0-10.2	11.5	7.6-15.4	11.2	5.2-17.2	10.1	7.6-12.6
≥ 7	5.1	0.0-10.6	0.5	0.0-1.4	5.6	1.0-10.2	5.0	3.0-7.1	3.3	0.0-6.9	4.1	1.7-6.5	5.1	0.9-9.3	4.2	2.5-5.9

3

CI- Confidence Interval, CAL – Clinical Attachment Loss

4

Table 4(on next page)

Percentage, with 95% confidence interval (95% CI), of sites (prevalence) and affected teeth (extent), by threshold of CAL (mm), severity and age group (years).

Table 4 - Percentage, with 95% confidence interval (95% CI), of sites (prevalence) and affected teeth (extent), by threshold of CAL (mm), severity and age group (years)

CAL (mm)	Subjects with chronic periodontitis								All subjects							
	20-44 (n = 59)		45-64 (n = 190)		≥ 65 (n = 89)		Total (n = 338)		20-44 (n = 90)		45-64 (n = 217)		≥ 65 (n = 98)		Total (n = 405)	
	%	95% CI	%	95% CI	%	95% CI	%	95% CI	%	95% CI	%	95% CI	%	95% CI	%	95% CI
Prevalence (sites)																
≥ 3	85.3	82.7-87.4	86.9	85.3-88.5	84.4	81.7-87.1	86.0	84.7-87.2	72.5	68.0-76.9	81.5	79.0-83.9	80.8	77.4-84.2	79.3	77.5-81.2
≥ 4	57.0	51.6-62.4	61.3	58.1-64.5	59.2	54.5-63.9	60.0	57.6-62.4	41.5	35.8-47.1	55.2	51.7-58.7	55.3	50.2-60.4	52.2	49.6-54.8
≥ 5	36.9	30.7-43.0	43.3	39.8-46.8	41.2	36.1-46.3	41.6	39.0-44.2	25.6	20.5-30.7	38.5	35.1-42.0	38.0	33.0-43.1	35.5	33.0-38.1
≥ 6	21.9	16.4-27.4	27.7	24.5-30.9	26.8	22.2-31.4	26.5	24.1-28.9	14.7	10.6-18.8	24.5	21.5-27.5	24.6	20.1-29.1	22.4	20.2-24.5
≥ 7	11.7	7.0-16.4	15.8	13.2-18.3	16.1	12.2-20.1	15.2	13.2-17.1	7.6	4.4-10.8	13.9	11.6-16.2	14.7	11.0-18.4	12.7	11.0-14.4
Extent (affected teeth)																
≥ 3	82.1	77.6-86.6	85.0	82.5-87.7	81.7	77.6-85.7	83.7	81.7-85.6	62.1	55.5-68.7	77.3	73.7-80.9	77.0	72.3-81.7	73.9	71.1-76.6
≥ 4	49.3	41.9-56.7	56.1	51.9-60.3	54.0	47.8-60.2	54.4	51.3-57.5	33.5	27.1-39.9	49.8	45.5-54.0	49.5	43.2-55.8	46.1	43.0-49.2
≥ 5	25.0	17.8-32.1	33.4	29.2-37.6	33.8	27.8-39.8	32.1	29.0-35.2	16.0	10.9-21.2	29.5	25.6-33.4	30.8	25.0-36.5	26.8	24.0-29.6
≥ 6	12.3	6.8-17.8	18.6	15.2-22.0	19.5	14.3-24.7	17.8	15.2-20.3	7.9	4.2-11.6	16.3	13.2-19.3	17.7	12.8-22.6	14.7	12.6-16.9
≥ 7	5.8	2.1-9.4	10.0	7.5-12.4	9.9	5.8-14.1	9.2	7.4-11.1	3.7	1.3-6.0	8.7	6.5-10.8	9.0	5.2-12.8	7.6	6.1-9.2

CI- Confidence Interval, CAL – Clinical Attachment Loss

Table 5(on next page)

Univariate logistic regression analysis of sociodemographic, behavioural, anthropometric and oral hygiene variables for the outcome variable $CAL \geq 3 \text{ mm}$ ($N = 405$)

1 Table 5 - Univariate logistic regression analysis of sociodemographic, behavioural, anthropometric and oral hygiene variables for the
2 outcome variable CAL ≥ 3 mm ($N = 405$)
3

Predictor variables		OR (95% CI)	<i>p</i>
Gender	Female	1	-
	Male	1.32 (0.77-2.26)	0.310
Age (years)			< 0.001
	20 - 44	1	-
	45 - 64	3.70 (2.04-6.69)	< 0.001
	≥ 65	5.20 (2.31-11.70)	< 0.001
Smoking status	Smoker	2.06 (1.11-3.81)	0.021
	Non-smoker	1	-
Education (years)			0.107
	1 - 4	1.40 (0.74-2.66)	0.298
	5 - 12	2.09 (1.05-4.13)	0.035
	> 12	1	-
Employment status			0.246
	Employed	1	-
	Unemployed	1.67 (0.74-3.77)	0.219
	Retired	1.54 (0.84-2.81)	0.163
BMI (kg/m ²)			0.699
	< 18.5	1	-

	18.5 - 24.9	1.06 (0.11-9.79)	0.961
	25.0 - 29.9	1.48 (0.16-13.82)	0.732
	≥ 30	1.40 (0.14-13.59)	0.774
Last dental visit			0.026
	< 1 year	1	-
	1 - 2 years	1.39 (0.54-3.57)	0.493
	3 - 4 years	0.42 (0.22-0.81)	0.009
	>= 5 years	0.97 (0.46-2.03)	0.930
	Never	0.24 (0.04-1.54)	0.134
Consultation motive			0.806
	Routine	1	-
	Aesthetics	0.72 (0.29-1.80)	0.483
	Pain	1.09 (0.50-2.35)	0.834
	Functional	1.24 (0.66-2.36)	0.502
	Other	1.39 (0.29-6.60)	0.680
Dental floss use	Yes	1	-
	No	1.66 (0.97-2.82)	0.063
Toothbrush frequency			0.803
	2 - 3 times / daily	1	-
	1 time / daily	1.27 (0.63-2.56)	0.508
	2 - 6 times / weekly	-	0.998

4 BMI (kg/m²) – Body Mass Index (kilogram/meter²), CI- Confidence Interval, OR – Odds Ratio

Table 6(on next page)

Multivariate logistic regression analysis (final reduced model) (*) for the outcome variable $CAL \geq 3 \text{ mm}$ (N = 405).

Table 6 - Multivariate logistic regression analysis (final reduced model) (*) for the outcome variable CAL \geq 3 mm ($N = 405$)

Predictor variables		CAL \geq 3 mm	
		OR (95% CI)	<i>p</i>
Age (years)	20-44	1	-
	45 - 64	4.85 (2.57-9.16)	< 0.001
	\geq 65	8.70 (3.66-20.69)	< 0.001
Smoking status	Non-smoker	1	-
	Smoker	3.55 (1.80-7.02)	< 0.001

CI- Confidence Interval, OR – Odds Ratio, CAL – Clinical Attachment Loss

(*) The model was statistically significant, $\chi^2(3) = 39.507$, $p < 0.001$, explained 15.7% (Nagelkerke R^2) of the variance and correctly classified 83.5% of cases.