



# Trends in bacterial and fungal communities in ant nests observed with Terminal-Restriction Fragment Length Polymorphism (T-RFLP) and Next Generation Sequencing (NGS) techniques—validity and compatibility in ecological studies

Stafva Lindström<sup>1,2,3</sup>, Owen Rowe<sup>4,5,6</sup>, Sari Timonen<sup>6</sup>, Liselotte Sundström<sup>1,2,3</sup> and Helena Johansson<sup>1,2</sup>

<sup>1</sup> Centre of Excellence in Biological Interactions, Department of Biosciences, University of Helsinki, Helsinki, Finland

<sup>2</sup> Tvärminne Zoological Station, University of Helsinki, Hanko, Finland

<sup>3</sup> Organismal and Evolutionary Biology, University of Helsinki, Helsinki, Finland

<sup>4</sup> Umeå Marine Sciences Centre, Umeå University, Hörnefors, Sweden

<sup>5</sup> Department of Ecology and Environmental Science, Umeå University, Umeå, Sweden

<sup>6</sup> Department of Microbiology, University of Helsinki, Helsinki, Finland

## ABSTRACT

Microbes are ubiquitous and often occur in functionally and taxonomically complex communities. Unveiling these community dynamics is one of the main challenges of microbial research. Combining a robust, cost effective and widely used method such as Terminal Restriction Fragment Length Polymorphism (T-RFLP) with a Next Generation Sequencing (NGS) method (Illumina MiSeq), offers a solid alternative for comprehensive assessment of microbial communities. Here, these two methods were combined in a study of complex bacterial and fungal communities in the nest mounds of the ant *Formica exsecta*, with the aim to assess the degree to which these methods can be used to complement each other. The results show that these methodologies capture similar spatiotemporal variations, as well as corresponding functional and taxonomical detail, of the microbial communities in a challenging medium consisting of soil, decomposing plant litter and an insect inhabitant. Both methods are suitable for the analysis of complex environmental microbial communities, but when combined, they complement each other well and can provide even more robust results. T-RFLP can be trusted to show similar general community patterns as Illumina MiSeq and remains a good option if resources for NGS methods are lacking.

Submitted 14 February 2018

Accepted 1 July 2018

Published 20 July 2018

Corresponding author

Stafva Lindström,  
stafva.lindstrom@helsinki.fi

Academic editor

Hannah Buckley

Additional Information and  
Declarations can be found on  
page 15

DOI 10.7717/peerj.5289

© Copyright

2018 Lindström et al.

Distributed under

Creative Commons CC-BY 4.0

OPEN ACCESS

**Subjects** Ecology, Microbiology, Molecular Biology

**Keywords** NGS, T-RFLP, Fungi, Bacteria, Ants, *Formica exsecta*, Illumina MiSeq, Spatiotemporal trends, Microbial communities

## INTRODUCTION

Microbes are the most numerous organisms on earth (*Bertrand et al., 2011*). Their small size and the difficulty of cultivating many (if not most) species has spurred the development of molecular techniques to study the taxonomy, community structure and functions of natural microbial communities. Several techniques such as Automated Ribosomal Intergenic Spacer Analysis (ARISA) (*Ramette, 2009*), Temperature Gradient Gel Electrophoresis (TGGE), Denaturing Gradient Gel Electrophoresis (DGGE) (*Nocker, Burr & Camper, 2007*) and Terminal Restriction Fragment Length Polymorphism (T-RFLP) (*Liu et al., 1997; Kent et al., 2003; Cao et al., 2013*) have been developed for the assessment of complex microbial communities. The semi-quantitative properties (*Blackwood et al., 2003; Aiken, 2011*), high reproducibility, low cost (*Thies, 2007*) and the technical and analytical straightforwardness of T-RFLP has cemented it as one of the leading fingerprinting methods for several decades (*Schütte et al., 2008*). T-RFLP has successfully been used to evaluate the effects of spatial and temporal variation in the structure of microbial communities (*Schütte et al., 2008; Robinson et al., 2009; Barnes et al., 2016*). However, to obtain taxonomic identity of microbial community members, T-RFLP is usually combined with time-consuming and expensive preparation and sequencing of clone libraries (*Van Elsas & Boersma, 2011*).

Recent advances in high-throughput DNA sequencing techniques have made NGS (Next Generation Sequencing) methods such as Illumina MiSeq/HiSeq, IonTorrent and 454 pyrosequencing attractive alternatives for analysis of microbial communities (*Teeling & Glöckner, 2012*). In contrast to T-RFLP, sequence data generated by these methods reveal taxonomic identity to the extent that designated sequence data is available (*Weissbrodt et al., 2012; De La Fuente et al., 2014; Prakash et al., 2014*). However, NGS datasets can be extensive, require more complex bioinformatic interpretation and the protocols and methods used for microbial NGS analysis are far from standardized (*Amend, Seifert & Bruns, 2010; Tedersoo et al., 2015; Balint et al., 2016*). Complementing T-RFLP with NGS techniques can, however, offer a convenient way of generating and analyzing metagenomic microbial data, including the addition of taxonomic information achieved by NGS sequencing, to the T-RFLP fingerprints. This is especially useful for longitudinal studies, where substantial T-RFLP-fingerprinting data, including data generated prior to the NGS era, are available. Furthermore, different environments facilitate and promote microbiomes of different composition and diversity and the analysis should be carefully adapted to the specific study at hand (*Buttigieg & Ramette, 2014; Faust et al., 2015; Weiss et al., 2016*). The few studies to date on bacteria, indicate that the patterns and trends in data generated by the two techniques are largely congruent (rumen bacteria (*De La Fuente et al., 2014*), bacteria in Antarctic soil (*Van Dorst et al., 2014*), bacteria in tropical soil (*Supramaniam et al., 2016*)). Studies that utilize both T-RFLP and NGS data to explore fungal communities appear absent from the literature, as do studies comparing both bacterial and fungal communities using both techniques.

Ants are frequent in most terrestrial environments (*Vander Meer, 2012*), often being referred to as ecosystem engineers due to their skills in manipulating the abiotic and biotic properties of soil (*Dauber, Schroeter & Wolters, 2001; Jurgensen et al., 2008*). While

building their nests, the bioturbation and the selective picking of appropriate construction materials create a soil chemistry that differs significantly from the one of the background soil (Kilpeläinen *et al.*, 2007; Dostál *et al.*, 2005). The distinct nest environment is clearly reflected by the nest microbiome: the nests of several mound building ant species contain a substantially higher microbial biomass (Dauber, Schroeter & Wolters, 2001), more numerous and diverse fungal colonies (Duff *et al.*, 2016) and significantly different microbial assemblages (Boots *et al.*, 2012), than those of the reference soils.

The aim of this study was to evaluate the congruence of Illumina MiSeq and T-RFLP data, based on the bacterial 16s and fungal ITS areas of the rRNA genes from complex environmental communities. To do this, the microbial communities inside the nests of the mound building ant *Formica exsecta* (Nylander, 1846) were sampled over a two-year period. The bacterial and fungal communities in the *F. exsecta* mounds have not been studied previously, but here molecular microbial data of active nest mounds is reported, generated with both methodologies. The relative abundance patterns, diversities and sampling efficiencies generated by the two molecular techniques were compared, and the similarity of the functional, spatial, and temporal patterns were evaluated. The extent to which T-RFLP can be complemented with the taxonomic information gained from the NGS data was also assessed.

## MATERIALS & METHODS

### Ant species, study site and sampling

Ants of the genus *Formica* are common in Northern Eurasia (Collingwood, 1979; Czechowski, Radchenko & Czechowska, 2002; Goropashnaya *et al.*, 2007). Most species construct perennial above-ground mounds from litter and soil particles gathered from the surrounding forest floor (Jurgensen *et al.*, 2008). The ant *Formica exsecta* used in this study is common (Douwes *et al.*, 2012), and inhabits meadows, sunny woodland openings, and forest clearings (Sundstrom, Chapuisat & Keller, 1996). The study sites are located on islands in the south-western archipelago of Finland, (Fig. S1) near Tvärminne zoological station (59°84'196"N, 23°20'182"E), where populations of *F. exsecta* have been monitored and studied since 1994 (e.g., Sundstrom, Chapuisat & Keller, 1996; Sundström, Keller & Chapuisat, 2003; Vitikainen, Haag-Liautard & Sundström, 2011). The islands represent typical biotopes of the SW coast of Finland, where granite outcrops, pine spinneys, and dry meadows alternate with intermixed lush patches of vegetation. The soil is classified as leptosol (<http://www.fao.org/soils-portal/soil-survey/soil-classification/world-reference-base/en/soil>) of varying thickness due to the topographic factors. Scots pine and Norway spruce dominate, together with ericaceous shrubs and species of *Deschampsia* and *Festuca*. Sampling was approved by the land owner (Tvärminne Zoological station), and no endangered or protected species were included in this study. Sampling was carried out once a month in May, June, July, August and September in both 2013 and 2014. Four nest mounds were sampled, two per island (F12 and F120 on Furuskär, J30 and J40 on Joskär, (Fig. S1)) on each consecutive sampling occasion, totalling 40 samples. The samples (~0.2 l) were collected by hand from the inside of the mound at the depth of 10–15 cm

using sterile gloves and placed in clean plastic bags. Samples were stored at  $-80^{\circ}\text{C}$  until extraction of DNA.

### DNA extraction and PCR amplification for T-RFLP

DNA was extracted from a  $\sim 0.25$  g subsample of nest mound material using the PowerSoil® DNA Isolation Kit (MoBio Laboratories Inc., Carlsbad, CA, USA) according to the manufacturer's instructions, with one exception being the use of TissueLyser II (Qiagen, Valencia, CA, USA) for 3 min at 20,000 rpm during the cell lysis phase. DNA was eluted in 100  $\mu\text{l}$  of elution buffer, and the same DNA extractions were used for the T-RFLP analysis and Illumina MiSeq sequencing as described below.

Sequences from the fungal ribosomal ITS2 region were amplified with the TAMRA-tagged forward primer fITS7 (GTGA(A/G)TCATCGAATCTTTG, [Ihrmark et al., 2012](#)), and the reverse primer ITS4 (TCCTCCGCTTATTGATATGC, [White et al., 1990](#)). PCR reactions were performed in a final volume of 15  $\mu\text{l}$ , containing 0.75  $\mu\text{l}$  of each 10  $\mu\text{M}$  primer, 3.0  $\mu\text{l}$  GC 5  $\times$  buffer, 0.3  $\mu\text{l}$  of 10 mM dNTP's, 0.15  $\mu\text{l}$  of Phusion™ DNA polymerase (Thermo Fisher, Waltham, MA, USA), 0.75  $\mu\text{l}$  of DNA template and 8.7  $\mu\text{l}$  Milli-Q water. The PCR conditions were 30 s at  $98^{\circ}\text{C}$ ; followed by 34 cycles of 10 s at  $98^{\circ}\text{C}$ , 30 s at  $54^{\circ}\text{C}$ , and 30 s at  $72^{\circ}\text{C}$ , followed by a final 5 min elongation step at  $72^{\circ}\text{C}$ . The PCRs were performed in duplicates and pooled into one product. Sequences from the bacterial 16S rRNA region were amplified with the FAM-tagged forward primer 27F (AGAGTTTGATC(A/C)TGGCTCAG, [Weisburg et al., 1991](#); [Chung et al., 2004](#)) and the reverse primer 1387R (GGGCGG(A/T)GTGTACAAGGC, [Wade et al., 1998](#)). PCR reactions were performed as described above, with the exception of the step in cycling before final elongation that was 40 s for bacteria.

Pooled duplicate PCR products from bacteria or fungi were purified with Agencourt AMPure XP beads (1 DNA:1.2 AMPure, Beckman Coulter) and digested with the restriction endonucleases HaeIII and MspI (Thermo Fisher, Waltham, MA, USA). The reaction volumes were 10  $\mu\text{l}$  purified PCR product, 1  $\mu\text{l}$  (10U) of each enzyme, 2  $\mu\text{l}$  10  $\times$  FastDigest buffer and 17  $\mu\text{l}$  nuclease free water. Samples were digested for 5 min at  $37^{\circ}\text{C}$ . The labelled terminal restriction fragments (T-RFs) were separated on an ABI-sequencer (model 3730; Applied Biosystems, Foster City, CA, USA) with the internal size standard GeneScan 500 ROX (Thermo Fisher, Waltham, MA, USA) and analysed using GeneMapper v.5 (Applied Biosystems, Foster City, CA, USA).

### Analysis of T-RFLP data

Peak data of T-RFs within the range of 70–400 bp and above a threshold of units of fluorescence of 100 for bacteria, and 70 for fungi were extracted from GeneMapper and imported into T-rex (<http://trex.biohpc.org>) for filtering of noise (removal of peaks not considered to comprise unique T-RFs), alignment and binning ([Culman et al., 2009](#)). Exact matches were used, i.e., only single base errors ( $\pm 1$  bp), and height of peaks ([Dickie & Fitzjohn, 2007](#); [Aiken, 2011](#); [Fredriksson, Hermansson & Wilén, 2014](#)) were used to determine the relative abundance. The peak data produced by the two enzymes (HaeIII and MspI) were filtered, aligned and binned in T-rex separately and pooled together ([De La](#)

*Fuente et al., 2014*). The only instances where the data from the two enzymes were analysed separately was for matching the experimental T-RFs with the virtual T-RFs (described below) and for assessing the level of correlation between the ranked relative abundances of T-RFs and OTUs.

### **Illumina MiSeq library preparation and sequencing**

The fungal ITS region was amplified with the primers fITS7 (GTGA(A/G)TCATCGAATCTTTG, *Ihrmark et al., 2012*) and ITS4 (TCCTCCGCT-TATTGATATGC, *White et al., 1990*) modified with partial Illumina TruSeq adaptor sequence. An initial PCR was performed with Phusion Hot Start II polymerase (Thermo Fischer) with a denaturation step of 98 °C for 30 s, followed by 15 cycles at 98 °C for 10 s, 65 °C for 30 s, 72 °C for 10 s followed by a final extension for 5 min at 72 °C. PCR samples were purified with Exonuclease I (Thermo Scientific, Waltham, MA, USA) and Thermosensitive Alkaline Phosphatase (FastAP; Thermo Scientific, Waltham, MA, USA). A second PCR was performed on the template products with full-length TruSeq P5 and index-containing P7 adapters. Cycling conditions were the same as the first library preparation PCR, but with an increased number of cycles (18). DNA libraries were quantified with Qubit (Invitrogen, Carlsbad, CA, USA) and library quality assessed with a Bioanalyzer 2100 (Agilent, Santa Clara, CA, USA), before pooling to equimolar amounts and submitted to sequencing. Due to technical requirements, the same reverse primer as for the T-RFLP could not be used. The bacterial 16s rRNA region was amplified using the primers 27F (AGAGTTTGATC(A/C)TGGCTCAG, *Weisburg et al., 1991*; *Chung et al., 2004*) and pD' (GTATTACCGCGGCTGCTG, *Edwards et al., 1989*). Libraries were otherwise constructed using the protocol described above. Sequencing was carried out using an Illumina MiSeq v2 600 cycle kit in paired-end mode at the DNA Sequencing and Genomics Laboratory, Institute of Biotechnology, University of Helsinki.

### **Illumina MiSeq—bioinformatics data analysis**

Read filtering and OTU clustering (at 97% identity) was performed using UPARSE v. 8.1 (*Edgar, 2013*; *Edgar & Flyvbjerg, 2015*), [Table S1](#). Database SILVAv123 (*Quast et al., 2013*) was used as reference database for alignment of the bacterial sequences, and UNITE v7 (*Kõljalg et al., 2013*) for the fungal sequences. For taxonomic classification of the bacterial OTUs, the reference database RDP16s training set v.14 (*Wang et al., 2007*) was used, and for the fungal OTUs the RDP ITS Warcup training set v.4 (*Deshpande et al., 2016*) was used. The queries against the databases were done by using the RDP Naïve Bayesian Classifier with bootstrap cut-off at 80% (*Wang et al., 2007*). Singletons, doubletons, and sequences not identified to the level of kingdom were removed prior to further analysis. The unprocessed sequences are available at NCBI Sequence Read Archive, Bioproject number [PRJNA399258](#).

### **Species diversity, effective sampling estimates, and functional organisation**

The two molecular methods were compared based on species accumulation curves, performed in the statistical framework R using the function `specaccum` in package `vegan` v.

2.4-5 (Oksanen *et al.*, 2011), and Good's estimate (Good, 1953) of sampling coverage. The curves visualize the level of sampling effort as a function of the accumulated number of samples with a more saturated curve indicating a better sampling effort. Good's estimate indicates the proportion of the population that has been captured by the sampling or sequencing (Good, 1953). Furthermore, the number of T-RFs and OTUs were counted, and the mean value of the total fungal and bacterial diversity was estimated by calculating respective Shannon-Wiener's diversity index ( $H'$ ) on the T-RFs and OTUs using the diversity function in R, package vegan (Oksanen *et al.*, 2011).

The functional organization ( $F_o$ ) of bacteria and fungi obtained with the two molecular methods, was compared by estimating Pareto-Lorenz curves (performed in JMP v.11, SAS Institute Inc.). To assess the degree of functional organization, the cumulative normalized number of T-RFs or OTUs were plotted against their respective cumulative normalized height (T-RFs) or sequence abundance (OTUs) (Marzorati *et al.*, 2008). This always creates a convex curve as the values are positive by default. The deviation of the curve from the 45° diagonal line (representing perfect evenness with no functional organization) indicates the degree of functional organization (Mertens, Boon & Verstraete, 2005; Wittebolle *et al.*, 2009). The  $F_o$  value is determined from the  $y$ -axis where the curve intercepts with the 20%  $x$ -axis line. A  $F_o$  value of 25% represents a community of high evenness with no distinct structure in terms of species dominance. A community at the  $F_o$  value of 45% is more functionally organized due to lower evenness, and a value of 80% for the  $F_o$  stands for a highly specialized community, dominated by a low number of species on which the functional stability depends. The level of  $F_o$  could also be defined as the community's ability to rebuild itself to the level of functionality it had before a disturbance (Marzorati *et al.*, 2008).

To further assess similarity between the datasets produced by T-RFLP and Illumina sequencing, the T-RFs were rarefied without replacement to the lowest peak height, and OTUs to the lowest number of reads (Table S1), before Bray-Curtis inter-sample dissimilarity matrices were generated. These matrices were then subjected to principal coordinates analysis (PCoA), and a PERMANOVA (a one-way nonparametric permutational multivariate analysis of variance, (Anderson, 2001), with 999 permutations was used to test the effects of year, month, nest and island on the matrices. To ensure that the results for the bacterial OTUs were robust (Morton *et al.*, 2017), PCoAs were also performed on Bray-Curtis matrices constructed from log transformed data, and on Morisita and Jaccard distance matrices. The uniformity of the T-RFLP and the OTU matrices was tested with Mantel's tests (method 'pearson', significance based on 999 permutations). Rarefying and the analyses described above were carried out in R with package vegan (Oksanen *et al.*, 2011). Finally, the relative abundance patterns of the (non-rarefied) T-RFs and the OTUs were compared, with the expectation that both techniques would show similar proportions of abundances. The T-RFs (enzymes analysed separately), and the equivalent number of OTUs were ranked from the highest to the lowest (T-RFs according to their height, and the OTUs according to their read abundance), and the correlations (Pearson's  $\rho$ ) tested by  $t$ -test, were performed in JMP v.11 (SAS Institute Inc., Cary, NC, USA).


## Virtual T-RFs

To match the Illumina OTUs and their identities with the experimental T-RFs, a virtual restriction was performed on the 200 most abundant OTUs of each bacteria and fungi. These subsets were chosen based on the outcome of the correlation test, and the number of sequences they covered. The OTU sequences were first aligned (ClustalX2, [Larkin et al., 2007](#)) to confirm the coverage of the entire targeted 16s region (bacteria) or the ITS2 region (fungi) and the starting points of the primers were identified. Virtual T-RFs were generated with the Webcutter on-line tool (<http://rna.lundberg.gu.se/cutter2/>), for the two enzymes separately. The experimental T-RFs that were generated were matched with virtual T-RFs in Excel, and taxonomic identities were derived from the classification of the Illumina MiSeq sequences. A taxa was considered as being represented by the experimental T-RFs only if restriction sites for both enzymes were detected on the Illumina OTU sequences.

## RESULTS

The T-RFLP-analysis generated 129 (HaeIII) and 120 (MspI) unique bacterial T-RFs ( $n = 37$ ), and 102 (HaeIII) and 124 (MspI) fungal T-RFs ( $n = 36$ ), respectively ([Table 1](#)). The Illumina MiSeq sequencing yielded in total (i.e., for all samples) 1,896,920 bacterial reads (16S sequences), and 3,274,825 fungal reads (ITS2 sequences), of high quality ( $n = 38$  for both bacteria and fungi) ([Table S1](#) and [SI\\_Bioinformatics](#)). These clustered into 4,699 unique bacterial and 2,315 unique fungal OTUs ([Table 1](#)). The Good's estimate of coverage for the T-RF data suggested that 97.9% of the bacterial, and 93% of the fungal diversity had been captured by the method, whereas the coverage for the OTUs was 99.9% and 99.4%, respectively ([Table 1](#)). The species accumulation curves ([Fig. 1](#)) showed a similar pattern, with the curves for the T-RFs being generally less flattened towards the end of their curves, particularly so in the fungal data. Only the bacterial OTU communities reached asymptote, suggesting high sampling efficiency ([Fig. 1](#)). The mean Shannon-Wiener diversity index for the T-RFs was lower than for the OTUs for both the bacteria (3.35 vs. 5.69) and fungi (2.71 vs. 3.48) ([Table 1](#)). The Pareto-Lorenz curves suggested a high  $F_0$ , indicating a high functional organization, for both the bacterial and fungal communities, regardless of the method used ([Fig. 2](#)). In the bacterial data, 20% of the T-RFs and OTUs accounted for 85% and 87% of the species abundance, respectively ([Fig. 2A](#)). The corresponding values for the fungal T-RFs and OTUs projected 73% and 95%, respectively ([Fig. 2B](#)).

The PCoAs indicated clustering of bacterial communities within islands and nests, in both the T-RF and OTU data, but the clustering was more pronounced in the OTU data ([Figs. 3A](#) and [3B](#)). The horseshoe pattern in the bacterial NGS data remained in the PCoA analysis when it was repeated with log-transformed data, and also when the analysis was performed on alternative distance matrices ([Figs. S2A–S2E](#)), suggesting a true pattern rather than an artefact. A clustering by month ([Fig. S3](#)) was indicated for the T-RFs, which seemed mainly driven by the values of September. No such indication was seen for the OTUs. The PERMANOVA confirmed the effect of month for the bacterial T-RFs. The PERMANOVA suggested compliance between the two methods, showing significant effects of island and nest for both the T-RF and OTU data ([Table 2](#)). The effect of year



**Figure 1** Species accumulation curves for bacterial T-RFs (A) and OTUs (B), and fungal T-RFs (C) and OTUs (D). The line represents the actual sampling, the grey area depicts the standard deviation and the box plot shows the species richness based on linear interpolation of random permutations.

Full-size [DOI: 10.7717/peerj.5289/fig-1](https://doi.org/10.7717/peerj.5289/fig-1)

**Table 1** The number of T-RFs and OTUs obtained, Shannon-Wiener diversity index ( $H'$ ) and Good's estimate of sampling coverage.


	T-RFs	SD	T-RFs HaeIII	T-RFs MspI	OTUs	SD
<b>Bacteria</b>						
No. of	249		129	120	4,699	
Mean no. of (per sample)	64	27			1,743	599
Mean value of Shannon-Wiener $H'$	3.35	0.43			5.69	0.67
Good's estimate	98.0%		90.6%	91.1%	99.9%	
<b>Fungi</b>						
No. of	226		102	124	2,315	
Mean no. of (per sample)	32	20			482	14
Mean value of Shannon-Wiener $H'$	2.71	0.76			3.48	0.78
Good's estimate	95.0%		74.0%	93.0%	99.4%	

**Notes.**

T-RFs,  $n = 37$  (bacteria) and  $n = 36$  (fungi); OTUs,  $n = 38$  for both bacteria and fungi.

was not significant, neither for the T-RFs nor the OTUs (Table 2). The PCoAs from the fungal T-RF and OTU data indicated clustering across both island and nest (Figs. 3C and 3D), with the fungal OTU data showing a much more pronounced clustering than the T-RF data (Figs. 3C and 3D). The PERMANOVA showed significant effects of island and nest for both T-RFs and OTUs in the fungal data, suggesting full compliance between the two methods for all the tested effects (Table 2). The Mantel tests showed moderate to






**Figure 2** Functional organization ( $F_o$ ) according to the Pareto-Lorenz curves of the bacterial (A) and fungal (B) T-RFs and OTUs. The continuous lines represent the T-RFs and the dashed lines the OTUs. The value of  $F_o$  equals the value projected at the  $y$ -axis where the critical 20% at the  $x$ -axis (thin continuous line) intercepts with the Pareto-Lorenz curves. The dotted line shows the  $45^\circ$  slope, representing perfect evenness. A  $F_o$  value of 25% represents a community of high evenness with no distinct structure in terms of species dominance. A community at the  $F_o$  value of 45% is more functionally organized due to its evenness, and a value of 80% for the  $F_o$  stands for a highly specialized community, dominated by a low number of species on which the functional stability depends.

Full-size DOI: [10.7717/peerj.5289/fig-2](https://doi.org/10.7717/peerj.5289/fig-2)

good correspondence between the T-RF and the OTU data, being significantly correlated (Pearson's  $\rho = 0.743$ ,  $P \leq 0.001$ , and  $0.574$ ,  $P \leq 0.001$ ) for the bacterial and the fungal data sets, respectively.

The relative abundances of the T-RFs, and the equivalent number of OTUs, ranked highest to lowest (encompassing 52% (HaeIII) and 50% (MspI) of the bacterial, and 78% and 81% of the fungal OTU sequences), were highly and significantly correlated (Pearson's  $\rho = 0.949$  (HaeIII), and  $0.849$  (MspI), respectively) for the bacterial data, and  $0.953$  and  $0.904$ , respectively for the fungal data;  $P \leq 0.001$  in all cases; Figs. 4A–4D). Of all the bacterial OTUs, 89% were assigned identities at the phylum, and 53% at the family level (Table 3), whereas the corresponding values for the fungal OTUs were 85%, and 39%, respectively (Table 3, Figs. S4A–S4B).

The 200 most abundant bacterial OTUs, on which the virtual restriction was performed, comprised approximately 60% of all bacterial OTU sequences. Taxonomic information (based on the taxonomic identification of the Illumina OTUs) was available for 66% of the 200 bacterial OTUs down to the level of family. The enzyme HaeIII successfully cut nearly all (99%) of the most abundant 200 OTUs, whereas MspI cut 86%. Matching virtual T-RF with the T-RFs generated from soil samples returned 93 taxonomic items, with restriction patterns from both enzymes (Table S2). Seventeen different restriction patterns were found, three of which were shared between several phyla, whereas 14 patterns were unique at the order level, and six of these at the family level (Table S2). No taxonomic information was available for three of the restriction patterns below the class level, and one pattern returned phylum only. Four of the six identified families (Burkholderiaceae, Caulobacteraceae,



**Figure 3** PCoA of bacterial T-RFs (A) and OTUs (B), and fungal T-RFs (C) and OTUs (D). Year is indicated by colour and nest by shape of symbols as indicated in the graph.


Full-size [DOI: 10.7717/peerj.5289/fig-3](https://doi.org/10.7717/peerj.5289/fig-3)

**Table 2** PERMANOVA test of the temporal (year and month) and spatial (island and nest) effects on the Bray-Curtis distances of the bacterial and fungal T-RF and OTU data.

Test effect	T-RFs				OTUs			
	<i>F</i>	<i>df</i>	<i>R</i> <sup>2</sup>	<i>p</i>	<i>F</i>	<i>df</i>	<i>R</i> <sup>2</sup>	<i>p</i>
<b>Bacteria</b>								
Year	1.08	36	0.030	0.336	0.49	37	0.014	0.873
Island	2.50	36	0.067	0.021	6.84	37	0.160	0.001
Nest	3.58	36	0.093	0.001	4.20	37	0.104	0.005
Month	2.55	36	0.068	0.020	1.32	37	0.035	0.210
<b>Fungi</b>								
Year	1.12	35	0.032	0.284	0.85	37	0.023	0.555
Island	2.99	35	0.081	0.001	7.46	37	0.172	0.001
Nest	4.65	35	0.120	0.001	7.31	37	0.169	0.001
Month	0.94	35	0.027	0.558	1.01	37	0.027	0.378

Nocardioideae and Streptomycetaceae) belonged to the bacterial families ranked as the most abundant according to the Illumina MiSeq data (Table 3).

The corresponding fungal data, based on the 200 most abundant fungal OTUs, on which the virtual restriction was performed, comprised approximately 87% of all fungal OTU sequences. Taxonomic information down to the family level was available for 50% of the fungal OTUs. A virtual HaeIII restriction site was found in 97%, and an MspI restriction site in 93% of the fungal OTUs. Matching the virtual T-RFs with the T-RFs generated from



**Figure 4** Correlation (Pearson  $\rho$ ) of T-RFs with the equivalent number of the most abundant OTUs, both ranked highest to lowest. (A) HaeIII, (B) MspI generated bacterial T-RFs, (C) and (D) HaeIII and MspI generated fungal T-RFs respectively.

Full-size DOI: [10.7717/peerj.5289/fig-4](https://doi.org/10.7717/peerj.5289/fig-4)

nest mound soil samples returned 41 taxonomic items, with restriction patterns from both enzymes. These taxonomic items represent 14 distinct restriction patterns, all of which were unique at the level of phylum (Table S3). Nine unique patterns were identified at the order level, and 8 patterns were further identified to the family level. Four patterns were identified to phylum only. Of the identified families, the Tremellales\_incertae\_sedis and Venturiaceae were among the most abundant ones according to the Illumina MiSeq data (Table 3).

## DISCUSSION

In this study, T-RFLP fingerprinting and Illumina sequencing were used to analyze the bacterial and fungal communities in the nest mounds of the ant *F. exsecta*. The results show that the combination of the two approaches is well suited to study microbial communities of high complexity, and that they provide similar, yet complementary information. With both methods, the microbial communities showed high and similar functional organization, and similar spatiotemporal patterns across both nest mounds and sampling occasions.

**Table 3** The number of the Illumina MiSeq OTUs assigned to the different taxonomic levels. The five most abundant bacterial phyla and families, together with the four fungal phyla present in the data and the five most abundant fungal families.

Taxonomic level	Nbr of OTUs assigned	
	Bacteria	Fungi
Kingdom	4,699	2,315
Phylum	4,163	1,963
	The five most abundant bacterial phyla:	The four fungal phyla present in data:
	Proteobacteria	Ascomycota
	Actinobacteria	Basidiomycota
	Bacteroidetes	Mucoromycota
	Acidobacteria	Chytridiomycota
	TM7 candidate phylum	
Class	3,836	1,326
Order	3,159	1,105
Family	2,486	908
	The five most abundant bacterial families:	The five most abundant fungal families:
	Streptomycetaceae	Herpotrichiellaceae
	Acetobacteraceae	Sporidiobolales_Incertae_sedis
	Nocardioidaceae	Venturiaceae
	Caulobacteraceae	Tremellales_Incertae_sedis
	Burkholderiaceae	Mortierellaceae
Genus	2,031	725

Comparisons between fungi and bacteria also revealed that bacterial communities were more comprehensively sampled than fungal communities.

The number of T-RFs obtained in this study were similar in magnitude to other comparable studies of soil microbial communities in polar, pasture, meadow, agricultural and forest soils, both for bacteria (*Blackwood et al., 2003; Frey et al., 2009; Anderson et al., 2011; Van Dorst et al., 2014*), and fungi (*Schwarzenbach, Enkerli & Widmer, 2007; Boots et al., 2012; Van Dorst et al., 2014*). Similarly, the number of microbial sequences, the sum of refined OTUs, and the resultant unique OTUs identified in this study, were in good correspondence with recent NGS studies of soil bacteria (*Lazzaro, Hilfiker & Zeyer, 2015; Supramaniam et al., 2016*) and fungi (*Meiser, Bálint & Schmitt, 2014*). In general, the number of defined, unique OTUs was much higher than the number of respective T-RFs (approximately 36 times the bacterial, and 19 times the fungal T-RFs). Nevertheless, when ranked per relative abundance, the high Pearson correlation suggest similar proportions of the most abundant T-RFs and OTUs, for both bacteria and fungi. This does not automatically mean that the most abundant T-RFs correspond with the most abundant OTUs or vice versa, but it indicates that the most abundant taxa were obtained by both techniques, in similar relative proportion.

Comparisons between the sampling coverage, i.e., Good's estimates, and the species accumulation curves of fungi and bacteria, revealed that bacterial communities were more comprehensively sampled than fungal communities. Notably, the asymptote was not reached for neither fungi nor bacteria using T-RFLP, suggesting undersampling by this technique (Zhou *et al.*, 2008). Only a small fraction of microbial diversity is sampled with this (and similar) methods (Jankowski, Schindler & Horner-Devine, 2014), resulting in low detection sensitivity and flatter species accumulation curves. Comparatively, the asymptote was reached for bacteria, and the curve better saturated for fungi using NGS, highlighting the power of this method to detect rarer taxa. Consequently, the diversity estimates were also lower for fungi than for bacteria. Technical factors likely to contribute to the observed differences in sampling success include the length of sequenced fragments, the choice of genes sequenced, and primer specificity. The technical requirement of using different primer pairs for both methods, resulted in fragments of differing length. The bacterial Illumina sequences were ~500 bp (Timonen *et al.*, 2017), and the T-RFLP sequences ~1,200 bp (Muehling *et al.*, 2016) long, respectively, whereas the fungal target sequence was 280–310 bp long (Ihrmark *et al.*, 2012) for both methods. Thus the longer fragment can be expected to provide more phylogenetic information, leading to a finer tuned OTU clustering. For the T-RFLP differentiation, the longer fragment could provide an increased number of enzyme cutting sites, which could partly explain the higher yield of the bacterial T-RFs compared to the fungal T-RFs. In addition, the bacterial 16S gene is highly conserved, and successfully targets most bacterial phyla (Větrovský *et al.*, 2013). Developing fungal primers with similar attributes has proven more challenging (Martin & Rygiel, 2005; Asemaninejad *et al.*, 2016), as fungal markers fulfilling the criteria of being of suitable length and specificity, yet not discriminating against any fungal taxa are not available (Lindahl *et al.*, 2013). This is further compounded by the fact that a substantial proportion of the fungal ITS sequences in the International Nucleotide Sequence Database are incomplete, with fragmentary regions in the ITS1 and ITS2 subsections (Nilsson *et al.*, 2009). From a biological standpoint, it is nevertheless possible that the fungal communities are less diverse, owing to abiotic factors (De Vries & Shade, 2013; Wagg *et al.*, 2014).

Illumina sequencing and T-RFLP fingerprinting captured the same patterns of functional organization in the nest mounds of *F. exsecta*. The Pareto-Lorenz curves suggested a high functional organization of the microbial communities inside the nest mounds. This suggests that the number of dominant species within the community is relatively low (Wittebolle *et al.*, 2009), given that a high functional organization signals numerical dominance by a few species (Mertens, Boon & Verstraete, 2005; Marzorati *et al.*, 2008; Wittebolle *et al.*, 2008; Gonzalez-Gil & Holliger, 2011). This agrees with our finding that 50% of the fungal sequences, and 60% of the bacterial sequences were found among the 200 most abundant OTUs. Such communities are often specialized, but potentially susceptible to disturbance, requiring a long recovery time for regaining their functionality. Conversely, a community of high species evenness is expected to cope better with environmental disturbances (Marzorati *et al.*, 2008). The high level of functional organization suggests that the microbial communities in the nest mounds may be specialized and strongly affected by the habitat with which they are associated (Edwards *et al.*, 2011; Boots & Clipson, 2013;

*Cai et al., 2014*). Further study to confirm an existent influence of the *F. exsecta* nest environment on the bacterial and fungal communities, compared with the effect of a reference substrate, would be required.

Both techniques also captured similar community patterns in bacteria and fungi. The results suggest that microbial communities were primarily structured at the level of nest mounds, with a more distinct clustering in the OTU data. Structure at the level of island was also found, but this effect may be compromised by the low sample size, and a more extensive study with a larger number of sampling sites would be required to confirm this effect. In this study, the higher sequencing depth (i.e., higher number of OTUs generated) likely contributed to the slightly lower correspondence between NGS and T-RFLP matrices, compared to that found in e.g., *Pilloni et al. (2012)* and *De La Fuente et al. (2014)*. A higher sequencing depth is likely to reveal rarer taxa, which would not have been defined in earlier studies, and are inaccessible to methods such as T-RFLP.

Although even greater taxonomic details for the T-RFs, and a better agreement between the two methods could be achieved by the use of additional enzymes (*Dunbar, Ticknor & Kuske, 2001; Edwards & Turco, 2005*), the overall taxonomic information obtained was comparable to those reported for extensive T-RFLP studies involving clone libraries (*Axelrood et al., 2002; Youssef & Elshahed, 2009; Lin et al., 2011*). The correlation between the relative abundance of T-RFs, and the equivalent number of the most abundant OTUs was significant both for bacteria and fungi, which corroborates the congruence between the methods. The outcome of the correlation served as a workable guideline for the number of OTUs (200) that were forwarded to the *in silico* T-RF cuttings. For both bacteria and fungi, the virtual T-RFLP generated restriction patterns from the majority of this subset of OTU sequences obtained by the Illumina sequencing. The bacterial OTUs were assigned to 13 different phyla, with roughly half (52%) being assigned to the phyla of Proteobacteria and Actinobacteria, both of which were also captured by the matching of T-RFs. At lower taxonomic levels, T-RFs and the virtual restriction cutting patterns corresponded to the taxonomic families of Burkholderiaceae, Caulobacteraceae, Nocardioideae and Streptomycetaceae. In the fungal data, the Illumina sequencing captured four phyla, whereas, based on the virtual restriction, the T-RFLP only captured two phyla, the Ascomycota and Basidiomycota. At lower taxonomic levels, the virtual restriction captured the families Tremellales incertae sedis, and Venturiaceae, both of which were the most abundant fungal families captured by the Illumina sequencing.

The bacterial and fungal taxa captured by our analysis are abundant in soil environments (*Acosta-Martínez et al., 2008; Buée et al., 2009; Anderson et al., 2011; Voglmayr et al., 2011; Lanzén et al., 2015; DeAngelis et al., 2015*). Furthermore, the bacterial genera belonging to Burkholderiaceae have also been shown to be associated with ant genera such as, *Camponotus* (*He et al., 2011*), *Cephalotes* (*Russell et al., 2009; Kautz et al., 2013*), and *Tetraponera* (*Van Borm et al., 2002*). Several genera representing both Proteobacteria and Actinobacteria, and sequences matching *Burkholderia* were also present in the transcriptomic data of *F. exsecta* (*Johansson et al., 2013*). The two fungal families that were captured by both T-RFLP and Illumina (Tremellales incertae sedis and Venturiaceae) are not currently defined in the literature as having any association with ants. However, the

literature on fungi associated with ants is focused solely on the fungus-farming ants and the restricted number of fungal species they cultivate, or the few insect pathogenic fungi studied in connection with ants (*Hughes et al., 2004*).

## CONCLUSIONS

Thorough assessment of complex microbial communities is challenging, and the use of multiple methods is often required in order to achieve a comprehensive outcome. This study describes certain steps at which technical choices during the process have substantial impact on the outcome, pinpointing some plausible reasons for non-conformity between the two methods. Overall, this study shows that both T-RFLP and Illumina sequencing are suitable for analysis of the fungal and bacterial communities in nest mounds of ants, but when the two techniques are combined, it can provide an even more robust dataset. Furthermore, this study validates the use of both techniques when addressing the topic of microbial communities in complex environments such as ant nest mounds, both independently or to support findings within comparable studies. In particular, when comprehensive longitudinal studies based on T-RFLP exist, supplementing the fingerprinting data with NGS is a practical solution. Although refining the results gained by the two methods into a reliable synthesis requires testing of overall compliance, T-RFLP can be trusted to reveal the same general community patterns as NGS (Illumina MiSeq), and is a good option if resources for NGS are limited.

## ACKNOWLEDGEMENTS

We thank our technician Heini Ali-Kovero for her valuable input during the laboratory assays, Minna-Maarit Kytöviita for her kind assistance in identifying the soil type and typical plants of the sampling site, Claire Morandin for her valuable help with the bioinformatics and Hanna Sinkko for all her statistical support.

## ADDITIONAL INFORMATION AND DECLARATIONS

### Funding

This work was funded by the Academy of Finland (grant numbers 252411 and 284666). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

### Grant Disclosures

The following grant information was disclosed by the authors:  
Academy of Finland: 252411, 284666.

### Competing Interests

The authors declare there are no competing interests.

### Author Contributions

- Stafva Lindström conceived and designed the experiments, performed the experiments, analyzed the data, prepared figures and/or tables, authored or reviewed drafts of the paper, approved the final draft.
- Owen Rowe and Sari Timonen conceived and designed the experiments, authored or reviewed drafts of the paper, approved the final draft.
- Liselotte Sundström conceived and designed the experiments, contributed reagents/materials/analysis tools, authored or reviewed drafts of the paper, approved the final draft.
- Helena Johansson conceived and designed the experiments, performed the experiments, authored or reviewed drafts of the paper, approved the final draft.

### Field Study Permissions

The following information was supplied relating to field study approvals (i.e., approving body and any reference numbers):

Field experiments were approved by Helsinki University, Tvärminne Zoological station.

### DNA Deposition

The following information was supplied regarding the deposition of DNA sequences:

The sequences are deposited at NCBI Sequence Read Archive, project number [PRJNA399258](https://www.ncbi.nlm.nih.gov/sra/PRJNA399258).

### Data Availability

The following information was supplied regarding data availability:

The seq data and the t-rflp data are uploaded as [Supplemental Files](#).

### Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.5289#supplemental-information>.

## REFERENCES

- Acosta-Martínez V, Dowd S, Sun Y, Allen V. 2008.** Tag-encoded pyrosequencing analysis of bacterial diversity in a single soil type as affected by management and land use. *Soil Biology and Biochemistry* **40**:2762–2770 DOI [10.1016/j.soilbio.2008.07.022](https://doi.org/10.1016/j.soilbio.2008.07.022).
- Aiken JT. 2011.** Terminal restriction fragment length polymorphism for soil microbial community fingerprinting. *Soil Science Society of America Journal* **75**:102–111 DOI [10.2136/sssaj2008.0088](https://doi.org/10.2136/sssaj2008.0088).
- Amend AS, Seifert KA, Bruns TD. 2010.** Quantifying microbial communities with 454 pyrosequencing: does read abundance count? *Molecular Ecology* **19**:5555–5565 DOI [10.1111/j.1365-294X.2010.04898.x](https://doi.org/10.1111/j.1365-294X.2010.04898.x).
- Anderson MJ. 2001.** Permutation tests for univariate or multivariate analysis of variance and regression. *Canadian Journal of Fisheries and Aquatic Sciences* **58**:626–639 DOI [10.1139/f01-004](https://doi.org/10.1139/f01-004).



- Anderson CR, Condrón LM, Clough TJ, Fiers M, Stewart A, Hill RA, Sherlock RR. 2011.** Biochar induced soil microbial community change: implications for biogeochemical cycling of carbon, nitrogen and phosphorus. *Pedobiologia* 54:309–320 DOI 10.1016/j.pedobi.2011.07.005.
- Asemaninejad A, Weerasuriya N, Gloor GB, Lindo Z, Thorn RG, Kauserud H. 2016.** New primers for discovering fungal diversity using nuclear large ribosomal DNA. *PLOS ONE* 11:e0159043 DOI 10.1371/journal.pone.0159043.
- Axelrood PE, Chow ML, Radomski CC, McDermott JM, Davies J. 2002.** Molecular characterization of bacterial diversity from British Columbia forest soils subjected to disturbance. *Canadian Journal of Microbiology* 48:655–674 DOI 10.1139/w02-059.
- Balint M, Bahram M, Murat Eren A, Faust K, Fuhrman JA, Orn Lindahl B, O'hara RB, Opik M, Sogin ML, Unterseher M, Tedersoo L. 2016.** Millions of reads, thousands of taxa: microbial community structure and associations analyzed via marker genes FEMS microbiology reviews advance access. *FEMS Microbiology Reviews* 40:686–700 DOI 10.1093/femsre/fuw017.
- Barnes CJ, Van der Gast CJ, Burns CA, McNamara NP, Bending GD. 2016.** Temporally variable geographical distance effects contribute to the assembly of root-associated fungal communities. *Frontiers in Microbiology* 7:195 DOI 10.3389/fmicb.2016.00195.
- Bertrand J-C, Caumette P, Lebaron P, Matheron R, Normand P, Sime-Ngando T. 2011.** *Environmental microbiology: fundamentals and applications*. Dordrecht: Springer Science Business Media B.V. DOI 10.1007/978-94-017-9118-2.
- Blackwood CB, Marsh T, Kim S, Paul EA. 2003.** Terminal restriction fragment length polymorphism data analysis for quantitative comparison of microbial communities terminal restriction fragment length polymorphism data analysis for quantitative comparison of microbial communities. *Applied and Environmental Microbiology* 69:926–932 DOI 10.1128/AEM.69.2.926.
- Boots B, Clipson N. 2013.** Linking ecosystem modification by the yellow meadow ant (*Lasius flavus*) to microbial assemblages in different soil environments. *European Journal of Soil Biology* 55:100–106 DOI 10.1016/j.ejsobi.2013.01.002.
- Boots B, Keith AM, Niechoj R, Breen J, Schmidt O, Clipson N. 2012.** Unique soil microbial assemblages associated with grassland ant species with different nesting and foraging strategies. *Pedobiologia* 55:33–40 DOI 10.1016/j.pedobi.2011.10.004.
- Buée M, Reich M, Murat C, Morin E, Nilsson RH, Uroz S, Martin F. 2009.** 454 Pyrosequencing analyses of forest soils reveal an unexpectedly high fungal diversity. *New Phytologist* 184:449–456 DOI 10.1111/j.1469-8137.2009.03003.x.
- Buttigieg PL, Ramette A. 2014.** A guide to statistical analysis in microbial ecology: a community-focused, living review of multivariate data analyses. *FEMS Microbiology Ecology* 90:543–550 DOI 10.1111/1574-6941.12437.
- Cai H, Jiang H, Krumholz LR, Yang Z. 2014.** Bacterial community composition of size-fractionated aggregates within the phycosphere of cyanobacterial blooms in a eutrophic freshwater lake. *PLOS ONE* 9(8):e102879 DOI 10.1371/journal.pone.0102879.

- Cao Y, Van De Werfhorst LC, Dubinsky EA, Badgley BD, Sadowsky MJ, Andersen GL, Griffith JF, Holden PA. 2013. Evaluation of molecular community analysis methods for discerning fecal sources and human waste. *Water Research* 47:6862–6872 DOI 10.1016/j.watres.2013.02.061.
- Chung J, Ha E-SS, Park H-RR, Kim S. 2004. Isolation and characterization of Lactobacillus species inhibiting the formation of Streptococcus mutans biofilm. *Oral Microbiology and Immunology* 19:214–216 DOI 10.1111/j.0902-0055.2004.00137.x.
- Collingwood C. 1979. The Formicidae (Hymenoptera) of Fennoscandia and Denmark. *Fauna Entomologica Scandinavica* 8:1–174.
- Culman SW, Bukowski R, Gauch HG, Cadillo-Quiroz H, Buckley DH. 2009. T-REX: software for the processing and analysis of T-RFLP data. *BMC Bioinformatics* 10:171 DOI 10.1186/1471-2105-10-171.
- Czechowski W, Radchenko A, Czechowska W. 2002. *The ants (Hymenoptera, Formicidae) of Poland*. Wilcza, Warsaw: Museum of Institute of Zoology.
- Dauber J, Schroeter D, Wolters V. 2001. Species specific effects of ants on microbial activity and N-availability in the soil of an old-field. *European Journal of Soil Biology* 37:259–261 DOI 10.1016/S1164-5563(01)01094-9.
- DeAngelis KM, Pold G, Topcuoglu BD, Van Diepen LTA, Varney RM, Blanchard JL, Melillo J, Frey SD. 2015. Long-term forest soil warming alters microbial communities in temperate forest soils. *Frontiers in Microbiology* 6:104 DOI 10.3389/fmicb.2015.00104.
- De La Fuente G, Belanche A, Girwood SE, Pinloche E, Wilkinson T, Newbold CJ. 2014. Pros and cons of ion-torrent next generation sequencing versus terminal restriction fragment length polymorphism T-RFLP for studying the rumen bacterial community. *PLOS ONE* 9(7):e101435 DOI 10.1371/journal.pone.0101435.
- Deshpande V, Wang Q, Greenfield P, Charleston M, Porrás-Alfaro A, Kuske CR, Cole JR, Midgley DJ, Tran-Dinh N. 2016. Fungal identification using a Bayesian classifier and the Warcup training set of internal transcribed spacer sequences. *Mycologia* 108:1–5 DOI 10.3852/14-293.
- De Vries FT, Shade A. 2013. Controls on soil microbial community stability under climate change. *Frontiers in Microbiology* 4:1–16 DOI 10.3389/fmicb.2013.00265.
- Dickie IA, Fitzjohn RG. 2007. Using terminal restriction fragment length polymorphism (T-RFLP) to identify mycorrhizal fungi: a methods review. *Mycorrhiza* 17:259–270 DOI 10.1007/s00572-007-0129-2.
- Dostál P, Březnová M, Kozlíčková V, Herben T, Kovář P. 2005. Ant-induced soil modification and its effect on plant below-ground biomass. *Pedobiologia* 49:127–137 DOI 10.1016/j.pedobi.2004.09.004.
- Douwes P, Abenius J, Cederberg B, Wahlstedt U. 2012. *Nationalnyckeln till Sveriges flora och fauna. Steklar: Myror-getingar. Hymenoptera: Formicidae-Vespidae*. Uppsala: ArtDatabanken, SLU.
- Duff LB, Urichuk TM, Hodgins LN, Young JR, Untereiner WA. 2016. Diversity of fungi from the mound nests of *Formica ulkei* and adjacent non-nest soils. *Canadian Journal of Microbiology* 62:562–571 DOI 10.1139/cjm-2015-0628.

- Dunbar J, Ticknor LO, Kuske CR. 2001.** Phylogenetic specificity and reproducibility and new method for analysis of terminal restriction fragment profiles of 16S rRNA genes from bacterial communities phylogenetic specificity and reproducibility and new method for analysis of terminal restriction. *Applied and Environmental Microbiology* 67:190–197 DOI 10.1128/AEM.67.1.190.
- Edgar RC. 2013.** UPARSE: highly accurate OTU sequences from microbial amplicon reads. *Nature Methods* 10:996–998 DOI 10.1038/nmeth.2604.
- Edgar RC, Flyvbjerg H. 2015.** Error filtering, pair assembly and error correction for next-generation sequencing reads. *Bioinformatics* 31:3476–3482 DOI 10.1093/bioinformatics/btv401.
- Edwards A, Anesio AM, Rassner SM, Sattler B, Hubbard B, Perkins WT, Young M, Griffith GW. 2011.** Possible interactions between bacterial diversity, microbial activity and supraglacial hydrology of cryoconite holes in Svalbard. *The ISME Journal* 5:150–160 DOI 10.1038/ismej.2010.100.
- Edwards U, Rogall T, Blöcker H, Emde M, Böttger EC, Bottger EC, Blocker H, Emde M. 1989.** Isolation and direct complete nucleotide determination of entire genes. Characterization of a gene coding for 16S ribosomal RNA. *Nucleic Acids Research* 17:7843–7853 DOI 10.1093/nar/17.19.7843.
- Edwards IP, Turco RF. 2005.** Inter- and intraspecific resolution of nrDNA TRFLP assessed by computer-simulated restriction analysis of a diverse collection of ectomycorrhizal fungi. *Mycological Research* 109:212–226 DOI 10.1017/S0953756204002151.
- Faust K, Lahti L, Gonze D, De Vos WM, Raes J. 2015.** Metagenomics meets time series analysis: unraveling microbial community dynamics. *Current Opinion in Microbiology* 25:56–66 DOI 10.1016/j.mib.2015.04.004.
- Fredriksson N, Hermansson M, Wilén B-M. 2014.** Impact of T-RFLP data analysis choices on assessments of microbial community structure and dynamics. *BMC Bioinformatics* 15:360 DOI 10.1186/s12859-014-0360-8.
- Frey B, Kremer J, Rüdts A, Sciacca S, Matthies D, Lüscher P. 2009.** Compaction of forest soils with heavy logging machinery affects soil bacterial community structure. *European Journal of Soil Biology* 45:312–320 DOI 10.1016/j.ejsobi.2009.05.006.
- Gonzalez-Gil G, Holliger C. 2011.** Dynamics of microbial community structure of and enhanced biological phosphorus removal by aerobic granules cultivated on propionate or acetate. *Applied and Environmental Microbiology* 77:8041–8051 DOI 10.1128/AEM.05738-11.
- Good IJ. 1953.** The population frequencies of species and the estimation of population parameters. *Biometrika* 40:237–264 DOI 10.1093/biomet/40.3-4.237.
- Goropashnaya AV, Fedorov VB, Seifert B, Pamilo P. 2007.** Phylogeography and population structure in the ant *Formica exsecta* (Hymenoptera, Formicidae) across Eurasia as reflected by mitochondrial DNA variation and microsatellites. *Annales Zoologici Fennici* 2450:462–474.
- He H, Chen Y, Zhang Y, Wei C. 2011.** Bacteria associated with gut lumen of camponotus japonicus Mayr. *Environmental Entomology* 40:1405–1409 DOI 10.1603/EN11157.

- Hughes WOH, Thomsen L, Eilenberg J, Boomsma JJ. 2004. Diversity of entomopathogenic fungi near leaf-cutting ant nests in a neotropical forest, with particular reference to *Metarhizium anisopliae* var. *anisopliae*. *Journal of Invertebrate Pathology* 85:46–53 DOI 10.1016/j.jip.2003.12.005.
- Ihrmark K, Bödeker ITM, Cruz-Martinez K, Friberg H, Kubartova A, Schenck J, Strid Y, Stenlid J, Brandström-Durling M, Clemmensen KE, Lindahl BD. 2012. New primers to amplify the fungal ITS2 region—evaluation by 454-sequencing of artificial and natural communities. *FEMS Microbiology Ecology* 82:666–677 DOI 10.1111/j.1574-6941.2012.01437.x.
- Jankowski K, Schindler DE, Horner-Devine MC. 2014. Resource availability and spatial heterogeneity control bacterial community response to nutrient enrichment in lakes. *PLOS ONE* 9:e86991 DOI 10.1371/journal.pone.0086991.
- Johansson H, Dhaygude K, Lindström S, Helanterä H, Sundström L, Trontti K. 2013. A metatranscriptomic approach to the identification of microbiota associated with the ant *Formica exsecta*. *PLOS ONE* 8(11):e79777 DOI 10.1371/journal.pone.0079777.
- Jurgensen MF, Finér L, Domisch T, Kilpeläinen J, Punntila P, Ohashi M, Niemelä P, Sundström L, Neuvonen S, Risch AC. 2008. Organic mound-building ants: their impact on soil properties in temperate and boreal forests. *Journal of Applied Entomology* 132:266–275 DOI 10.1111/j.1439-0418.2008.01280.x.
- Kautz S, Rubin BER, Russell JA, Moreau CS. 2013. Surveying the microbiome of ants: comparing 454 pyrosequencing with traditional methods to uncover bacterial diversity. *Applied and Environmental Microbiology* 79:525–534 DOI 10.1128/AEM.03107-12.
- Kent AD, Smith DJ, Benson BJ, Triplett W, Triplett EW. 2003. Web-based phylogenetic assignment tool for analysis of terminal restriction fragment length polymorphism profiles of microbial communities. *Applied and Environmental Microbiology* 69:6768–6776 DOI 10.1128/AEM.69.11.6768.
- Kilpeläinen J, Finér L, Niemelä P, Domisch T, Neuvonen S, Ohashi M, Risch AC, Sundström L. 2007. Carbon, nitrogen and phosphorus dynamics of ant mounds (*Formica rufa* group) in managed boreal forests of different successional stages. *Applied Soil Ecology* 36:156–163 DOI 10.1016/j.apsoil.2007.01.005.
- Köljalg U, Nilsson RH, Abarenkov K, Tedersoo L, Taylor AFS, Bahram M, Bates ST, Bruns TD, Bengtsson-Palme J, Callaghan TM, Douglas B, Drenkhan T, Eberhardt U, Dueñas M, Grebenc T, Griffith GW, Hartmann M, Kirk PM, Kohout P, Larsson E, Lindahl BD, Lücking R, Martín MP, Matheny PB, Nguyen NH, Niskanen T, Oja J, Peay KG, Peintner U, Peterson M, Pöldmaa K, Saag L, Saar I, Schüßler A, Scott JA, Senés C, Smith ME, Suija A, Taylor DL, Telleria MT, Weiss M, Larsson KH. 2013. Towards a unified paradigm for sequence-based identification of fungi. *Molecular Ecology* 22:5271–5277 DOI 10.1111/mec.12481.
- Lanzén A, Epelde L, Garbisu C, Anza M, Martín-Sánchez I, Blanco F, Mijangos I. 2015. The community structures of prokaryotes and fungi in mountain pasture soils are highly correlated and primarily influenced by pH. *Frontiers in Microbiology* 6:1321 DOI 10.3389/fmicb.2015.01321.

- Larkin MA, Blackshields G, Brown NP, Chenna R, McGettigan PA, McWilliam H, Valentin F, Wallace IM, Wilm A, Lopez R, Thompson JD, Gibson TJ, Higgins DG. 2007. Clustal W and Clustal X version 2.0. *Bioinformatics* 23:2947–2948 DOI 10.1093/bioinformatics/btm404.
- Lazzaro A, Hilfiker D, Zeyer J. 2015. Structures of microbial communities in alpine soils: seasonal and elevational effects. *Frontiers in Microbiology* 6:1330 DOI 10.3389/fmicb.2015.01330.
- Lin Y Te, Jangid K, Whitman WB, Coleman DC, Chiu CY. 2011. Soil bacterial communities in native and regenerated perhumid montane forests. *Applied Soil Ecology* 47:111–118 DOI 10.1016/j.apsoil.2010.11.008.
- Lindahl B, Nilsson RH, Tedersoo L, Abarenkov K, Carlsen T, Pennanen T, Stenlid J. 2013. Fungal community analysis by high-throughput sequencing of amplified markers—a user’s guide. *New Phytologist* 199:288–299.
- Liu WT, Marsh TL, Cheng H, Forney LJ. 1997. Characterization of microbial diversity by determining terminal restriction fragment length polymorphisms of genes encoding 16S rRNA. *Applied and Environmental Microbiology* 63:4516–4522.
- Martin KJ, Rygielwicz PT. 2005. Fungal-specific PCR primers developed for analysis of the ITS region of environmental DNA extracts. *BMC Microbiology* 5:28 DOI 10.1186/1471-2180-5-28.
- Marzorati M, Wittebolle L, Boon N, Daffonchio D, Verstraete W. 2008. How to get more out of molecular fingerprints: practical tools for microbial ecology. *Environmental Microbiology* 10:1571–1581 DOI 10.1111/j.1462-2920.2008.01572.x.
- Meiser A, Bálint M, Schmitt I. 2014. Meta-analysis of deep-sequenced fungal communities indicates limited taxon sharing between studies and the presence of biogeographic patterns. *New Phytologist* 201:623–635 DOI 10.1111/nph.12532.
- Mertens B, Boon N, Verstraete W. 2005. Stereospecific effect of hexachlorocyclohexane on activity and structure of soil methanotrophic communities. *Environmental Microbiology* 7:660–669 DOI 10.1111/j.1462-2920.2004.00735.x.
- Morton JT, Toran L, Edlund A, Metcalf JL, Lauber C, Knight R. 2017. Uncovering the horseshoe effect in microbial analyses. *Systems* 2:e00166–16 DOI 10.1128/mSystems.00166-16.
- Muehling M, Poehlein A, Stuhr A, Voitel M, Daniel R, Schlömann M. 2016. Reconstruction of the metabolic potential of acidophilic sideroxydants strains from the metagenome of an microaerophilic enrichment culture of acidophilic iron-oxidizing bacteria from a pilot plant for the treatment of acid mine drainage reveals metabolic. *Frontiers in Microbiology* 7:1–16 DOI 10.3389/fmicb.2016.02082.
- Nilsson RH, Ryberg M, Abarenkov K, Sjökvist E, Kristiansson E. 2009. The ITS region as a target for characterization of fungal communities using emerging sequencing technologies. *FEMS Microbiology Letters* 296:97–101 DOI 10.1111/j.1574-6968.2009.01618.x.
- Nocker A, Burr M, Camper AK. 2007. Microbial ecology genotypic microbial community profiling: a critical technical review. *Microbial Ecology* 54:276–289 DOI 10.1007/s00248-006-9199-5.

- Oksanen J, Blanchet F, Kindt R, Legendre P, Minchin P, Al E. 2011. Vegan: community ecology package. R package version 2.0-1.
- Pilloni G, Granitsiotis MS, Engel M, Lueders T. 2012. Testing the limits of 454 pyrotag sequencing: Reproducibility, quantitative assessment and comparison to T-RFLP fingerprinting of aquifer microbes. *PLOS ONE* 7:e40467 DOI 10.1371/journal.pone.0040467.
- Prakash O, Pandey PK, Kulkarni GJ, Mahale KN, Shouche YS. 2014. Technicalities and glitches of Terminal Restriction Fragment Length Polymorphism (T-RFLP). *Indian Journal of Microbiology* 54:255–261 DOI 10.1007/s12088-014-0461-0.
- Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, Peplies J, Glöckner FO. 2013. The SILVA ribosomal RNA gene database project: improved data processing and web-based tools. *Nucleic Acids Research* 41:590–596 DOI 10.1093/nar/gks1219.
- Ramette A. 2009. Quantitative community fingerprinting methods for estimating the abundance of operational taxonomic units in natural microbial communities. *Applied and Environmental Microbiology* 75:2495–2505 DOI 10.1128/AEM.02409-08.
- Robinson CH, Szaro TM, Izzo AD, Anderson IC, Parkin PI, Bruns TD. 2009. Spatial distribution of fungal communities in a coastal grassland soil. *Soil Biology and Biochemistry* 41:414–416 DOI 10.1016/j.soilbio.2008.10.021.
- Russell JA, Moreau CS, Goldman-Huertas B, Fujiwara M, Lohman DJ, Pierce NE. 2009. Bacterial gut symbionts are tightly linked with the evolution of herbivory in ants. *Proceedings of the National Academy of Sciences of the United States of America* 106:21236–21241 DOI 10.1073/pnas.0907926106.
- Schütte UME, Abdo Z, Bent SJ, Shyu C, Williams CJ, Pierson JD, Forney LJ. 2008. Advances in the use of terminal restriction fragment length polymorphism (T-RFLP) analysis of 16S rRNA genes to characterize microbial communities. *Applied Microbiology and Biotechnology* 80:365–380 DOI 10.1007/s00253-008-1565-4.
- Schwarzenbach K, Enkerli J, Widmer F. 2007. Objective criteria to assess representativity of soil fungal community profiles. *Journal of Microbiological Methods* 68:358–366 DOI 10.1016/j.mimet.2006.09.015.
- Sundstrom L, Chapuisat M, Keller L. 1996. Conditional manipulation of sex ratios by ant workers: a test of kin selection theory. *Science* 274:993–995 DOI 10.1126/science.274.5289.993.
- Sundström L, Keller L, Chapuisat M. 2003. Inbreeding and sex-biased gene flow in the ant *Formica exsecta*. *Evolution; International Journal of Organic Evolution* 57:1552–1561 DOI 10.1111/j.0014-3820.2003.tb00363.x.
- Supramaniam Y, Chong C-W, Silvaraj S, Tan IK-P. 2016. Effect of short term variation in temperature and water content on the bacterial community in a tropical soil. *Applied Soil Ecology* 107:279–289 DOI 10.1016/j.apsoil.2016.07.003.
- Tedersoo L, Ramirez KS, Nilsson RH, Kaljuvee A, Kõljalg U, Abarenkov K, Orgiazzi A, Dunbar M, Panagos P, Groot G, Lemanceau P, Pante E, Schoelinck C, Puillandre N, Quince C, Lanzen A, Curtis T, Davenport R, Hall N, Head I, Whitlock M, Critescu M, Wiczorek J, Bloom D, Guralnick R, Blum S, Döring M, Robertson

- V, Yilmaz P, Kottmann R, Field D, Knight R, Cole J, Amaral-Zettler L, Meiser A, Balint M, Schmitt I, Kõljalg U, Nilsson R, Abarenkov K, Tedersoo L, Taylor A, Bahram M, Ratnasingham S, Hebert P, Abarenkov K, Tedersoo L, Nilsson R, Velak K, Saar I, Veldre V. 2015. Standardizing metadata and taxonomic identification in metabarcoding studies. *GigaScience* 4:34 DOI 10.1186/s13742-015-0074-5.
- Teeling H, Glöckner FO. 2012. Current opportunities and challenges in microbial metagenome analysis—A bioinformatic perspective. *Briefings in Bioinformatics* 13:728–742 DOI 10.1093/bib/bbs039.
- Thies JE. 2007. Soil microbial community analysis using terminal restriction fragment length polymorphisms. *Soil Science Society of America Journal* 71:579–591 DOI 10.2136/sssaj2006.0318.
- Timonen S, Sinkko H, Sun H, Sietiö OM, Rinta-Kanto JM, Kiheri H, Heinonsalo J. 2017. Ericoid roots and mycospheres govern plant-specific bacterial communities in boreal forest humus. *Microbial Ecology* 73:939–953 DOI 10.1007/s00248-016-0922-6.
- Van Borm S, Buschinger A, Boomsma JJ, Billen J. 2002. Tetraponera ants have gut symbionts related to nitrogen-fixing root-nodule bacteria. *Proceedings of the Royal Society B-Biological Sciences* 269:2023–2027 DOI 10.1098/rspb.2002.2101.
- Van Dorst J, Bissett A, Palmer AS, Brown M, Snape I, Stark JS, Raymond B, McKinlay J, Ji M, Winsley T, Ferrari BC. 2014. Community fingerprinting in a sequencing world. *FEMS Microbiology Ecology* 89:316–330 DOI 10.1111/1574-6941.12308.
- Van Elsas JD, Boersma FGH. 2011. A review of molecular methods to study the microbiota of soil and the mycosphere. *European Journal of Soil Biology* 47:77–87 DOI 10.1016/j.ejsobi.2010.11.010.
- Vander Meer R. 2012. Ant interactions with soil organisms and associated semiochemicals. *Journal of Chemical Ecology* 38:728–745 DOI 10.1007/s10886-012-0140-8.
- Větrovský T, Baldrian P, Pijl A, De Hollander M, Kowalchuk G. 2013. The variability of the 16S rRNA gene in bacterial genomes and its consequences for bacterial community analyses. *PLOS ONE* 8:e57923 DOI 10.1371/journal.pone.0057923.
- Vitikainen E, Haag-Liautard C, Sundström L. 2011. Inbreeding and reproductive investment in the ant *Formica exsecta*. *Evolution* 65:2026–2037 DOI 10.1111/j.1558-5646.2011.01273.x.
- Voglmayr H, Mayer V, Maschwitz U, Moog J, Djieto-Lordon C, Blatrix R. 2011. The diversity of ant-associated black yeasts: insights into a newly discovered world of symbiotic interactions. *Fungal Biology* 115:1077–1091 DOI 10.1016/j.funbio.2010.11.006.
- Wade JR, Marchesi JR, Sato T, Weightman AJ, Martin TA, Fry JC, Hiom SJ, Dymock D, Wade WG. 1998. Design and evaluation of useful bacterium-specific PCR primers that amplify genes coding for bacterial 16S rRNA. *Applied and Environmental Microbiology* 64:795–799.
- Wagg C, Bender SF, Widmer F, Van der Heijden MGA. 2014. Soil biodiversity and soil community composition determine ecosystem multifunctionality. *Proceedings of the National Academy of Sciences of the United States of America* 111:5266–5270 DOI 10.1073/pnas.1320054111.

- Wang Q, Garrity GM, Tiedje JM, Cole JR. 2007.** Naive Bayesian classifier for rapid assignment of rRNA sequences into the new bacterial taxonomy. *Applied and Environmental Microbiology* 73:5261–5267 DOI [10.1128/AEM.00062-07](https://doi.org/10.1128/AEM.00062-07).
- Weisburg WG, Barns SM, Pelletier DA, Lane DJ. 1991.** 16S ribosomal DNA amplification for phylogenetic study. *Journal of Bacteriology* 173:697–703 DOI [10.1128/JB.173.2.697-703.1991](https://doi.org/10.1128/JB.173.2.697-703.1991).
- Weiss S, Van Treuren W, Lozupone C, Faust K, Friedman J, Deng Y, Xia LC, Xu ZZ, Ursell L, Alm EJ, Birmingham A, Cram JA, Fuhrman JA, Raes J, Sun F, Zhou J, Knight R. 2016.** Correlation detection strategies in microbial data sets vary widely in sensitivity and precision. *The ISME Journal* 10:1669–1681 DOI [10.1038/ismej.2015.235](https://doi.org/10.1038/ismej.2015.235).
- Weissbrodt DG, Shani N, Sinclair L, Lefebvre G, Rossi P, Maillard J, Rougemont J, Holliger C. 2012.** PyroTRF-ID: a novel bioinformatics methodology for the affiliation of terminal-restriction fragments using 16S rRNA gene pyrosequencing data. *BMC Microbiology* 12:1471–2180.
- White TJ, Bruns T, Lee S, Taylor J. 1990.** *PCR protocols: a guide to methods and application*. San Diego: Academic Press.
- Wittebolle L, Marzorati M, Clement L, Balloi A, Daffonchio D, De Vos P, Heylen K, Verstraete W, Boon N. 2009.** Initial community evenness favours functionality under selective stress. *Nature* 458:623–626 DOI [10.1038/nature07840](https://doi.org/10.1038/nature07840).
- Wittebolle L, Vervaeren H, Verstraete W, Boon N. 2008.** Quantifying community dynamics of nitrifiers in functionally stable reactors. *Applied and Environmental Microbiology* 74:286–293 DOI [10.1128/AEM.01006-07](https://doi.org/10.1128/AEM.01006-07).
- Youssef NH, Elshahed MS. 2009.** Diversity rankings among bacterial lineages in soil. *The ISME Journal* 3:305–313 DOI [10.1038/ismej.2008.106](https://doi.org/10.1038/ismej.2008.106).
- Zhou J, Kang S, Schadt CW, Garten CT. 2008.** Spatial scaling of functional gene diversity across various microbial taxa. *Proceedings of the National Academy of Sciences of the United States of America* 105:7768–7773 DOI [10.1073/pnas.0709016105](https://doi.org/10.1073/pnas.0709016105).