

# Genetic diversity of Microsporidia in the circulatory system of endemic amphipods from different locations and depths of ancient Lake Baikal

Mariya Dimova<sup>1</sup>, Ekaterina Madyarova<sup>1,2</sup>, Anton Gurkov<sup>1,2</sup>, Polina Drozdova<sup>1</sup>, Yulia Lubyaga<sup>1,2</sup>, Elizaveta Kondrateva<sup>1</sup>, Renat Adelshin<sup>1,3</sup>, Maxim Timofeyev<sup>Corresp. 1</sup>

<sup>1</sup> Irkutsk State University, Irkutsk, Russia

<sup>2</sup> Baikal Research Centre, Irkutsk, Russia

<sup>3</sup> Irkutsk Anti-Plague Research Institute of Siberia and Far East, Irkutsk, Russia

Corresponding Author: Maxim Timofeyev  
Email address: m.a.timofeyev@gmail.com

Endemic amphipods (Amphipoda, Crustacea) of the most ancient and large freshwater Lake Baikal (Siberia, Russia) are a highly diverse group comprising >15 % of all known species of continental amphipods. The extensive endemic biodiversity of Baikal amphipods provides the unique opportunity to study interactions and possible coevolution of this group and their parasites, such as Microsporidia. In this study, we investigated microsporidian diversity in the circulatory system of 22 endemic species of amphipods inhabiting littoral, sublittoral and deep-water zones in all three basins of Lake Baikal. Using molecular genetic techniques, we found microsporidian DNA in two littoral (*Eulimnogammarus verrucosus*, *E. cyaneus*), two littoral/sublittoral (*Pallasea cancellus*, *E. maritiji*) and two sublittoral/deep-water (*Acanthogammarus lappaceus longispinus*, *A. victorii maculosus*) endemic species. Twenty sequences of the small subunit ribosomal (SSU) rDNA were obtained from the haemolymph of the six endemic amphipod species sampled from 0–60 m depths at the Southern Lake Baikal's basin (only the Western shore) and at the Central Baikal. They form clusters with similarity to *Enterocytozpora*, *Cucumispora*, *Dictyocoela*, and several unassigned Microsporidia sequences, respectively. Our sequence data show similarity to previously identified microsporidian DNA from inhabitants of both Lake Baikal and other water reservoirs. The results of our study suggest that the genetic diversity of Microsporidia in haemolymph of endemic amphipods from Lake Baikal does not correlate with host species, geographic location or depth factors but is homogeneously diverse.

**Genetic diversity of Microsporidia in the circulatory system of endemic amphipods from  
different locations and depths of ancient Lake Baikal**

Mariya Dimova<sup>1</sup>, Ekaterina Madyarova<sup>1,2</sup>, Anton Gurkov<sup>1,2</sup>, Polina Drozdova<sup>1</sup>, Yulia Lubyaga<sup>1,2</sup>,  
Elizaveta Kondrateva<sup>1</sup>, Renat Adelshin<sup>1,3</sup>, Maxim Timofeyev<sup>1</sup>

<sup>1</sup> Irkutsk State University, Irkutsk, Russia

<sup>2</sup> Baikal Research Centre, Irkutsk, Russia

<sup>3</sup> Irkutsk Anti-Plague Research Institute of Siberia and Far East, Irkutsk, Russia

Corresponding Author:

Maxim Timofeyev<sup>1</sup>

Email address: m.a.timofeyev@gmail.com

# Abstract

Endemic amphipods (Amphipoda, Crustacea) of the most ancient and large freshwater Lake Baikal (Siberia, Russia) are a highly diverse group comprising >15 % of all known species of continental amphipods. The extensive endemic biodiversity of Baikal amphipods provides the unique opportunity to study interactions and possible coevolution of this group and their parasites, such as Microsporidia. In this study, we investigated microsporidian diversity in the circulatory system of 22 endemic species of amphipods inhabiting littoral, sublittoral and deep-water zones in all three basins of Lake Baikal. Using molecular genetic techniques, we found microsporidian DNA in two littoral (*Eulimnogammarus verrucosus*, *E. cyaneus*), two littoral/sublittoral (*Pallasea cancellus*, *E. marituji*) and two sublittoral/deep-water (*Acanthogammarus lappaceus longispinus*, *A. victorii maculosus*) endemic species. Twenty sequences of the small subunit ribosomal (SSU) rDNA were obtained from the haemolymph of the six endemic amphipod species sampled from 0–60 m depths at the Southern Lake Baikal's basin (only the Western shore) and at the Central Baikal. They form clusters with similarity to *Enterocytozpora*, *Cucumispora*, *Dictyocoela*, and several unassigned Microsporidia sequences, respectively. Our sequence data show similarity to previously identified microsporidian DNA from inhabitants of both Lake Baikal and other water reservoirs. The results of our study suggest that the genetic diversity of Microsporidia in haemolymph of endemic amphipods from Lake Baikal does not correlate with host species, geographic location or depth factors but is homogeneously diverse.

# Introduction

Lake Baikal is the deepest (1643 m) and oldest lake in the world (25–30 million years); it is the largest (by volume; 23,000 km<sup>3</sup>) reservoir of clean fresh water, containing approximately 20 % of the available world's liquid fresh water (Martens, 1997; Yoshii, 1999). Lake Baikal has specific abiotic characteristics that distinguish it from all the other freshwater bodies in the world: high oxygen content throughout the entire water column, stable low water temperatures with long seasonal ice coverage on the lake's surface, and super-oligotrophic conditions. The environmental conditions of the lake's open-water and deep-water zones have remained close to their current state for the last 2–4 million years (Kozhova and Izmet'eva, 1998). As a unique ecosystem with exceptionally high degrees of biological diversity and endemism, Lake Baikal was designated a UNESCO World Heritage Site. To date, 2595 animal species from Lake Baikal have been described, approximately 80 % of which are endemic (Timoshkin, 2001).

Among macroinvertebrates, the highest number of species in the lake is presented by amphipods. The diversity of amphipods in Lake Baikal is very high: currently, 276 species and 78 subspecies (over 10 % of the total species diversity of fauna of Lake Baikal; over 15 % of all recognized amphipod species from fresh or inland waters world-wide), grouped in 7 families and 41 genera, have been found in the lake (Väinölä et al., 2008; Takhteev, 2015). Amphipods inhabit all depths of Baikal, including littoral (0-20 m), sublittoral (20-70 m), supra-abyssal (70-250 m) and abyssal (250-1640 m); the last two depths are typically grouped as a deep-water zone (Kozhov, 1962). The age and geographical isolation of this group create excellent opportunities for exploring the diversity and evolution of host-parasite relationships.

Microsporidia are a diverse phylum of eukaryotic parasites, which are a sister clade to the Fungi kingdom and infect a wide range of hosts from invertebrates to humans (Keeling and Fast, 2002; Smith, 2009; Issi et al., 2012; Gismondi et al., 2012; Stentiford et al., 2013; Bojko et al.,

2016). The number of described species of Microsporidia is more than 1300, and the various species belong to approximately 187 genera, 50 of which are found in aquatic arthropods (Stentiford et al. 2013; Weiss and Becnel, 2014). Some microsporidian species may influence the sex ratio of arthropod populations (Terry et al., 1998; Ironside et al., 2003; Ironside and Alexander, 2015), the behaviour of the host, and the host population dynamics (Dunn and Smith, 2001). For example, a selective sexual behaviour is described for amphipod species *Gammarus duebeni* Liljeborg, 1852. The males of this species infected by Microsporidia are only able to hold down infected females for breeding (Dunn et al., 2006). Microsporidia can be transmitted horizontally, vertically or both ways (Terry et al., 2004; Smith, 2009). They have been found to infect muscles, gonads, intestinal walls, hepatopancreas, haemocytes, and other organs (Kelly and Anthony, 1979; Weiss and Becnel, 2014).

Despite the fact that amphipods endemic to Lake Baikal offer great possibility to study the basic principles of coevolution of the host-parasite system and distribution of parasites among evolutionary close species, only a few papers have been published on the microsporidian diversity in the lake's fauna. The study of Microsporidia in Baikal amphipods began in 1967 when *Nosema kozhovi* was found in intestinal epithelium of *Brandtia latissima lata* (Dyb., 1874) (Lipa, 1967). The first molecular genetic studies of Microsporidia were performed in the 21st century. Six endemic and one cosmopolitan microsporidian species (*Dictyocoela duebenum*) were detected in the Baikal amphipod *Gmelinoides fasciatus* (Stebbing, 1899; Kuzmenkova et al., 2008), and 100 new gene isolates of Microsporidia in 31 species of amphipods of Lake Baikal were discovered by molecular methods (Smith et al., 2008). A recent study provided the first glimpse at horizontal distribution and exchange of Microsporidia between Baikal littoral and non-Baikal amphipods, which indicated frequent introductions of the parasites into Lake Baikal

ecosystem and high homogeneity of Microsporidia between species of endemic hosts in the coastal zone (Ironside and Wilkinson, 2017).

However, most previous studies utilized the whole individuals or soft amphipod tissues for amplification of microsporidian DNA, which does not rule out the possibility of contamination. To avoid this problem, our group recently conducted a study using molecular genetic techniques (SSU rDNA sequencing) to detect Microsporidia in circulatory system of several endemic amphipod species from Lake Baikal (Madyarova et al., 2015). Searching directly in the haemolymph minimizes the possibility of identifying SSU rDNA of Microsporidia located on exoskeleton, in gut lumen or inside parasites of the amphipods and can guarantee that the found Microsporidia are parasitic directly to the amphipod species studied. Moreover, using haemolymph provides a unified way of treating amphipods of very different sizes, including small species and young individuals.

Spatial variations, especially vertical distribution, in biodiversity of Microsporidia in Lake Baikal amphipods remain poorly investigated. Thus, the aim of the current study was to search for diversity of Microsporidia in the haemolymph of endemic amphipods inhabiting littoral, sublittoral and deep-water zones of the lake in all three basins of Baikal: Southern, Central and Northern.

## Materials and Methods

### *Sampling and location*

To study microsporidian biodiversity, the haemolymph of 22 amphipod species from 11

points of Lake Baikal was used (Table S1). The following seven species were collected at depths of 0-1 m using a hand net in the period from 2011 to 2016: *Eulimnogammarus verrucosus*, *E. cyaneus*, *E. maritiji*, *E. vitatus*, *E. maaki*, *E. viridis olivaceus*, *Pallasea cancelus*. The other amphipod species were sampled from depths of 20-200 m: *Acanthogammarus victorii maculosus*, *A. flavus*, *A. lappaceus longispinus*, *A. godlewskii*, *A. brevispinus*, *A. reicherii*, *Carinurus belkini*, *Garjajewia cabanisi*, *Propachygammarus maximus*, *Ommatogammarus carneolus melanophthalmus*, *Parapallasea lagovskii*, *P. puzyllii*, *P. brandtia*, *P. kesslerii*, *P. grubii*. The latter group of species was collected during the expeditions in 2014 and 2016 using a trawl, dredge and deep-water traps with rotten fish installed at a depth of 200 m. In this work, only benthic amphipod species were studied. The chosen amphipod species are not endangered or protected.

Species determination was performed according to the identification key (Bazikalova, 1945; Takhteev and Didorenko, 2015).

The collections cover all three of Lake Baikal's basins: the Southern (Listvyanka, Bolshie Koty, Port Baikal), the Central (Olkhon Island, Kharauz, Ushkany Islands) and the Northern (Kotelnikovskiy Cap), including both the Western (Listvyanka, Bolshie Koty, Port Baikal) and Eastern (Kharauz) shores (Fig. 1). Around Olkhon Island, five sampling points were screened: Burkhan Cape, Cape Umysh-Tame, Olkhon Gate, Kharantsy Island and Peschanka Bay. The Lake Baikal's basins and the Angara River act as natural geographic barriers for benthic amphipods (Mashiko, 2000; Gomanenko et al., 2005). No specific permissions were required for sampling of amphipods in these locations.

# *Haemolymph extraction*

Haemolymph extraction was performed from live amphipods in a laboratory using a glass capillary. Each sample subsequently used for DNA extraction (volumes ranging from 50 to 100 µl) contained haemolymph collected from one or pooled from several individuals of the same species. The number of individuals per sample depended on the size of the amphipods (1–50 individuals; see Table S1 for details). Isolated haemolymph samples were stored in liquid nitrogen.

# *DNA extraction and PCR*

Total DNA extraction from individual or pooled haemolymph was performed using the “Riboprep” reagent kit (AmpliSens, Russia). The concentration and purity of the isolated DNA was determined using the UV spectrophotometer UNICO 2802 (UNICO, USA).

The small subunit of ribosomal DNA (SSU rDNA) of the Microsporidia was selected as a molecular phylogenetic marker. Amplification was performed in the nested PCR with universal primers for Microsporidia: V1f: 5'-CACCAGGTTGATTCTGCCTGAC-3' (Weiss et al., 1994); 1342r: 5'-ACGGGCGGTGTGTACAAAGAACAG-3' (McClymont et al., 2005); 18sf: 5'-GTTGATTCTGCCTGACGT-3' (Baker et al., 1995); and 981r: 5'-TGGTAAGCTGTCCCGCGTTGAGTC-3' (MacNeil et al., 2003). Each PCR was performed in a Gradient Thermocycler (Biometra, Germany) in a volume of 25 µl and contained 1X PCR buffer, 6.25 pmol of each dNTP, 2.5 U SynTaq DNA polymerase, 6.25 pmol MgCl<sub>2</sub>, 10 pmol primers, 3 µl of DNA template, and deionized water. The conditions for the first round were 95°C for 5 min, followed by 40 cycles of 95°C for 30 sec, 55°C for 30 sec, and 72°C for 90 sec, and then 72°C



for 7 min. Conditions for the second round of PCR were 95°C for 5 min, followed by 35 cycles of 95°C for 30 sec, 52°C for 30 sec, and 72°C for 1 min, followed by 72°C for 7 min.

The PCR products of the second round were visualized in 1 % agarose gels, and products of the expected size were excised from the gel and purified by ethanol and sodium acetate precipitation (Sambrook et al., 1989). Sequencing of the amplified DNA fragments was performed using Genetic Analyzer 3500 xL (Applied Biosystems, Japan) with the BigDye Terminator Cycle Sequencing kit v.3.1 (Applied Biosystems, USA).

# *Nucleotide sequence analysis, phylogenetic reconstruction and statistical analysis*

The obtained sequences were identified with NCBI nucleotide BLAST with default parameters against the nt database (last accessed 2018/05/17). The best B hit belonging to a sample identified in other studies was chosen to describe the samples (Table S2).

The sequences were aligned with the MAFFT 7.397 E-INS-i strategy (Katoh et al., 2009) and then trimmed with Gblocks (Castresana et al., 2000) accessed through the Phylogeny.fr web interface (Dereeper et al., 2008) with all options for less stringent selection checked. The resulting alignment (Alignment S1) contained 514 bases. Nucleotide substitution models were evaluated with jModelTest 2.1.10 (Guindon and Gascuel, 2003; Darriba et al., 2012). The best model according to Bayesian information criterion (BIC) score was TrN+G with GTR+G falling very close behind. As the formed model was not available in RAxML, GTR+G was used for all calculations. The phylogenetic tree was reconstructed with BEAST 2.5 (Bouckaert et al., 2014) for 10,000,000 generations with sampling every 1,000<sup>th</sup> tree. After this run, ESS value was close to 150, indicating acceptable convergence. In addition, we reconstructed the tree with RAxML

8.2.11 (Stamatakis, 2014) under the GTRGAMMA model and 5,000 rapid bootstrap generations. RAxML bootstrap results were reanalysed with the tree obtained with BEAST to get comparable support rates for the nodes. Raw trees obtained with BEAST and RAxML are shown in Tree S1 and Tree S2, respectively. The results were visualized in FigTree 1.4.2 (<http://tree.bio.ed.ac.uk/software/figtree/>) and Dendroscope, and Fig. 2 was prepared with the ggtree package (Yu et al., 2017) for R (R Core team, 2017) and Inkscape (inkscape.org).

Simpson's diversity index was calculated using the diversity function (option *index* set to *invsimpson*) of the vegan package (Dixon, 2003) for R. Fisher's exact test was implemented with the stats package for R.

## Results

We screened the haemolymph of 22 species (Table S1) of endemic amphipods from Lake Baikal and found microsporidian DNA in two littoral (*E. verrucosus*, *E. cyaneus*), two littoral/sublittoral (*P. cancellus*, *E. marituji*) and two sublittoral/deep-water species (*A. lappaceus longispinus*, *A. victorii maculosus*). A total of over 1,000 individual amphipods were analysed (Table S1). Twenty nucleotide sequences of SSU rDNA belonging to the *Cucumispora*, *Dictyocoela*, *Enterocytopora*-like groups and unassigned Microsporidia were obtained from the haemolymph of the six endemic amphipod species sampled from depths of 0-60 m at Southern Lake Baikal's basin (only the Western shore) and at Central Baikal (Table S2). Only one microsporidian isolate has been identified in each positive pooled haemolymph sample. The phylogenetic tree (Fig. 2) reconstructed using these and some other published microsporidian sequences (Alignment S1) shows clustering of the DNA identified in the current study to four

groups.

Three microsporidian DNA sequences were found in the haemolymph of *E. cyaneus* (sampled at 0-1 m in the Southern Baikal), and one was found in *A. victorii maculosus* (sampled at 53-60 m in the Central Baikal). These four sequences are presented as a single cluster on the phylogenetic tree marked as *Cucumispora*-like (C) and are similar to taxonomically defined *Cucumispora* isolates (Ovcharenko et al., 2010; Bojko et al., 2015, 2017b) as well as microsporidian sequences found earlier in some other endemic amphipods of Lake Baikal, such as *P. cancellus* (KM977842, KM977843, KM977844, KM977845, KM977846; Madyarova et al., 2015), *Acanthogammarus victorii* (FJ756173), *Garjajewia cabanisii* (FJ755959) and *Pallaseopsis kessleri* (FJ756022).

The other microsporidian DNA from one haemolymph sample of *P. cancellus*, three samples of *E. verrucosus* (both sampled at 0-1 m in the Southern Baikal) and two of *A. lappaceus longispinus* (sampled at 25-35 m in the Central Baikal) were clustered with those of Microsporidia of the recently defined (Bacela-Spychalska et al., 2018) genus *Dictyocoela* (D), namely *D. duebenum* (Grabner et al., 2015) and *D. berillonum*. Representatives of *Dictyocoela* were found earlier in an endemic Baikal amphipod *A. lappaceus* (KM977839; Madyarova et al., 2015).

Two more sequences amplified from *Eulimnogammarus verrucosus* haemolymph clustered with other isolates of the *Enterocytopora*-like clade (E), such as *Enterocytopora artemiae* (JX915760; Rode et al., 2013a), *Globulispora mitoportans* (KT762153; Vávra et al., 2016) and recently described *Parahepatospora carcini* (KX757849; Bojko et al., 2017a). A similar sequence was also found in an endemic Baikal amphipod *Micruropus wahlui* (FJ756186).

Finally, some isolates detected in the littoral species *E. verrucosus* (sampled at 0–1 m) from four different habitats, Listvyanka, Bolshie Koty, Port Baikal (Southern Baikal) and Burkhan Cape (Central Baikal), clustered with other microsporidian sequences previously amplified from Baikal endemic amphipods (FJ756111 from *Dorogostaiskia parasitica* and FJ755992 from *Odontogammarus calcaratus*). This cluster is designated as unclassified Microsporidia (uM).

As we have data for seven different locations, we have calculated Simpson's diversity index for the groups of amphipods defined according to the phylogenetic tree and these locations (Fig. S1). Our samples are not big enough to draw any conclusions, as we quite predictably saw higher diversity in locations with higher numbers of analysed pools, but further studies may help clarify this issue.

## Discussion

The microsporidian DNA sequences found in the haemolymph of amphipods from Lake Baikal during this study belong to four clusters: *Cucumispora*-, *Dictyocoela*-, *Enterocytoispora*-like, as well as some unclassified Microsporidia. Some other sequences belonging to the *Cucumispora*-like and *Dictyocoela*-like clusters, as well as to unclassified Microsporidia were also identified in our previous work (Madyarova et al., 2015). Apart from these similarities, there were also some differences, as in the earlier work we also identified representatives of the genus *Nosema* (N) in Baikal amphipods *E. verrucosus* from Listvyanka (Madyarova et al., 2015), but in the current study, these microsporidian species were not found at all.

Moreover, in winter 2013-2014 (Table S1) *E. verrucosus* from Listvyanka showed

relatively high infection (~30 % of pools) by Microsporidia of groups *Nosema/Vairimorpha*- (NV), *Dictyocoela*- (D), *Enterocytoispora*-like (E) and unclassified Microsporidia (uM), while in summer 2016 only one infected pool (~3 %) was identified in this location and corresponded to an unclassified Microsporidia (uM). We have compared the largest samples with exact Fisher's test (5 out of 18 vs. 1 out of 33) and found that the difference was statistically significant ( $p = 0.017$ ). Sampling of Microsporidia during summer and autumn (2015 and 2016) at other points (Port Baikal, Bolshie Koty and Burkhan Cape) also demonstrated relatively low infection (no more than 13 % of pools) only by unclassified Microsporidia. It is interesting to note that two highest infection rates belonged to pools sampled in winter, the reproduction season of *E. verrucosus* (Bazikalova, 1941). Generally, by comparing the percentage of infected haemolymph pools of different amphipod species from chosen sampling points (Table S1), we can conclude that the proportion of infected individuals can vary significantly.

The species diversity of the obtained samples is also interesting in the context of worldwide diversity of microsporidian parasites of amphipods. Microsporidian SSU rDNA nucleotide sequences similar to the C group in our analysis (Fig. 2) were also found in *Gammarus chevreuxi* Sexton, 1913 from the Avon River in the UK (AJ438962) (Terry et al., 2004). The identified sequences of the D group are similar to the *Dictyocoela duebenum* isolate 775 (FN434091), which was previously found in *Gammarus duebeni* from Iceland (Krebes et al., 2010), and the *Dictyocoela muelleri* isolate (AJ438955) found in *Gammarus duebeni celticus* from Ireland (Terry et al., 2004). The *Enterocytoispora*-like sequences are similar to the sequences of *Enterocytoispora artemiae* found in *Artemia franciscana monica* from Mono Lake, USA (Rode et al., 2013b). These results may suggest that microsporidian distribution in Baikal amphipods is similar to that in amphipods and other crustaceans from other water bodies.

Concerning the spatial and host distribution of Microsporidia in Baikal amphipods, we should note that similar and identical sequences of microsporidian DNA were found in amphipods inhabiting different depths and basins of the lake. Two microsporidian sequences from *P. cancellus* (KM977844 and KM977846) were identical to sequences from completely different amphipod species *G. cabanisi* (FJ755959) and *D. parasitica* (FJ756113) (Table S2; Fig. 2). Almost identical DNA of Microsporidia that belong to the group C were found at sampling depths 0-1 m of Southern Baikal (KT956054, KR080135, KT956053 from *E. cyaneus* and KM977844 from *P. cancellus*) and depths of 54-60 m of Central Baikal (KT956063 from *A. victorii maculosus*). Similar sequences of Microsporidia of the group D are found at depths of 0-1 m (*E. marituji*, *E. verrucosus* and *P. cancellus* from the Southern Baikal) and at 25-35 m (*A. lappaceus longispinus* from the Central Baikal). Additionally, amphipods of species *E. verrucosus* from Southern and Central Baikal were infected by microsporidia with almost identical SSU rDNA belonging to some unclassified Microsporidia (uM). So, similar microsporidian isolates infect different hosts at different depths and locations. There are several possible explanations for this fact. Amphipods are omnivorous animals and can feed on other amphipods of different or the same species. Moreover, Baikal amphipods are characterized by high vertical and horizontal mobility: a significant part of the analysed species (*E. marituji*, *P. cancellus*, *A. lappaceus longispinus* and others) can be found in a range of depths of at least several dozen meters and can participate in nocturnal vertical migrations observed for many benthic animals in Baikal (Bazikalova, 1945; Karnaukhov et al., 2016); also, in Lake Baikal there are eurybathic scavenger amphipods (like *Ommatogammarus flavus* and *O. albinus*), found in depth ranges of several hundred meters (Bazikalova, 1945), which may serve as additional “carriers” of infections in the benthic community.

Thus, the results of our study do not suggest any link between the genetic diversity of Microsporidia found in circulatory system of endemic amphipods from Lake Baikal and host species, geographic location or depth. This result corroborates previous studies (Madyarova et al., 2015; Ironside & Wilkinson, 2017).

## Conclusions

The sequence isolates that we have identified during our study belong to four clusters and show similarity to previously identified microsporidian DNA from inhabitants of both Lake Baikal and other water reservoirs. No specificity in distribution of Microsporidia was identified depending on host amphipod species and their spatial relations, which may indicate high connectivity of individuals within species and between species of Lake Baikal amphipods. However, search for Microsporidia in amphipods at greater depths (200-1600 m) is still necessary to confirm the observed homogeneity in distribution of these parasites.

## Acknowledgements

We would like to thank Dr Denis Axenov-Gribanov, Dr Daria Bedulina, Prof. Vadim Takhteev and the crew of the research vessel “Prof. M. Kozhov” for assistance in the selection and fixation of samples, identifying species of amphipods and their valuable advice. We are extremely grateful to Dr Jamie Bojko and an anonymous peer reviewer for their helpful comments on the manuscript.

# References

- Bacela-Spychalska, K., Wróblewski, P., Mamos, T., Grabowski, M., Rigaud, T., Wattier, R., ...  
& Ovcharenko, M., 2018. Europe-wide reassessment of *Dictyocoela* (Microsporidia)  
infecting native and invasive amphipods (Crustacea): molecular versus ultrastructural  
traits. Scientific reports, 8, 8945.
- Baker, M. D., Vossbrinck, C. R., Didier E. S., Maddox J. V., Shadduck J. A., 1995. Small  
subunit ribosomal DNA phylogeny of various microsporidia with emphasis on AIDS-  
related forms. Journal of Eukaryotic Microbiology, 42, 564–570.
- Bazikalova, A. Y. 1945. Amphipods of Lake Baikal. Proceedings of Baikal Limnological Station,  
11, 1-440.
- Bojko, J., Dunn, A.M., Stebbing, P.D., Ross, S.H., Kerr, R.C. and Stentiford, G.D., 2015.  
*Cucumispora ornata* n. sp. (Fungi: Microsporidia) infecting invasive ‘demon shrimp’  
(*Dikerogammarus haemobaphes*) in the United Kingdom. Journal of invertebrate  
pathology, 128, 22-30.
- Bojko, J., Clark, F., Bass, D., Dunn, A.M., Stewart-Clark, S., Stebbing, P.D. and Stentiford, G.D.,  
2017a. Parahepatospora carcini n. gen., n. sp., a parasite of invasive Carcinus maenas  
with intermediate features of sporogony between the Enterocytozoon clade and other  
Microsporidia. Journal of invertebrate pathology, 143, 124-134.
- Bojko, J., Bącela-Spychalska, K., Stebbing, P. D., Dunn, A. M., Grabowski, M., Rachalewski, M.  
Stentiford, G. D., 2017b. Parasites, pathogens and commensals in the "low-impact" non-  
native amphipod host *Gammarus roeselii*. Parasites & Vectors, 10, 1-15.



328 Bouckaert, R., Heled, J., Kühnert, D., Vaughan, T., Wu, C-H., Xie, D., Suchard, M. A., Rambaut,  
329 A., & Drummond, A. J. (2014). BEAST 2: A software platform for Bayesian  
330 evolutionary analysis. *PLoS Computational Biology*, 10, e1003537.

331 Castresana, J., 2000. Selection of conserved blocks from multiple alignments for their use in  
332 phylogenetic analysis. *Molecular Biology and Evolution*, 17, 540-552.

333 Darriba, D., Taboada, G.L., Doallo, R., Posada, D., 2012. jModelTest 2: more models, new  
334 heuristics and parallel computing. *Nature Methods*, 9, 772.

335 Dereeper, A., Guignon, V., Blanc, G., Audic, S., Buffet, S., Chevenet, F., Dufayard, J.F.,  
336 Guindon, S., Lefort, V., Lescot, M. and Claverie, J.M., 2008. Phylogeny. fir: robust  
337 phylogenetic analysis for the non-specialist. *Nucleic acids research*, 36, W465-W469.

338 Dixon, P., 2003. VEGAN, a package of R functions for community ecology. *Journal of*  
339 *Vegetation Science*, 14(6), pp.927-930.

340 Dunn, A.M., Smith, J.E., 2001. Microsporidian life cycles and diversity: the relationship between  
341 virulence and transmission. *Microbes and Infection*, 3, 381-388.

342 Dunn, A.M., Hogg, J.C., Hatcher, M.J., 2006. Transmission and burden and the impact of  
343 temperature on two species of vertically transmitted microsporidia. *International Journal*  
344 *for Parasitology*, 36, 409-414.

345 Gismondi, E., Rigaud, T., Beisel, J. N., Cossu-Leguille, C., 2012. Microsporidia parasites disrupt  
346 the responses to cadmium exposure in a gammarid. *Environmental Pollution*, 160, 17-23.

347 Gomanenko, G. V., Kamaltynov, R. M., Kuzmenkova, Z. V., Berenos, K., Sherbakov, D. Y.,  
348 2005. Population structure of the Baikalian amphipod *Gmelinoides fasciatus* (Stebbing).

349 Russian Journal of Genetics, 41, 907-912.

350 Grabner, D. S., Weigand, A. M., Leese, F., Winking, C., Hering, D., Tollrian, R., Sures, B., 2015.

351 Invaders, natives and their enemies: distribution patterns of amphipods and their

352 microsporidian parasites in the Ruhr Metropolis, Germany. Parasites & vectors, 8, 419.

353 Guindon, S. and Gascuel, O., 2003. A simple, fast and accurate method to estimate large

354 phylogenies by maximum-likelihood. Systematic Biology, 52, 696-704.

355 Issi, I. V., Tokarev, Y. S., Seliverstova, E. V., Voronin V. N., 2012. Taxonomy of *Neoperezia*

356 *chironomi* and *Neoperezia semenovaiae* comb. nov. (Microsporidia, Aquasporidia):

357 Lessons from ultrastructure and ribosomal DNA sequence data. European Journal of

358 Protistology, 48, 17-29.

359 Ironside, J. E., Alexander, J., 2015. Microsporidian parasites feminise hosts without paramyxean

360 co-infection: support for convergent evolution of parasitic feminisation. International

361 journal for parasitology, 45, 427-433.

362 Ironside, J. E., Dunn, A. M., Rollinson, D., Smith, J. E., 2003. Association with host

363 mitochondrial haplotypes suggests that feminizing microsporidia lack horizontal

364 transmission. Journal of evolutionary biology, 16, 1077-1083.

365 Ironside, J.E., Wilkinson, T.J., 2017. Accumulation and exchange of parasites during adaptive

366 radiation in an ancient lake. International Journal for Parasitology, S0020-

367 7519(17)30349-1.

368 Karnaukhov, D. Y., Bedulina, D. S., Kaus, A., Prokosov, S. O., Sartoris, L., Timofeyev, M. A.,

369 Takhteev, V. V., 2016. Behaviour of Lake Baikal amphipods as a part of the night

- 370           migratory complex in the Kluevka settlement region (south-eastern Baikal).
- 371           *Crustaceana*, 89, 419-430.
- 372   Kato, K., Asimenos, G., Toh, H., 2009. Multiple alignment of DNA sequences with MAFFT.
- 373           Bioinformatics for DNA sequence analysis, 537, 39-64.
- 374   Keeling, P. J., Fast, N. M., 2002. Microsporidia: biology and evolution of highly reduced
- 375           intracellular parasites. Annual Reviews in Microbiology, 56, 93-116.
- 376   Kelly, J. F., Anthony, D. W., 1979. Susceptibility of spores of the microsporidian *Nosema*
- 377           *algerae* to sunlight and germicidal ultraviolet radiation. Journal of invertebrate
- 378           pathology, 34, 164-169.
- 379   Kozhov, M. M., 1962. Biology of Lake Baikal. Publishing house of Academy of Sciences of the
- 380           USSR, Moscow.
- 381   Kozhov, M. M., Kozhova O. M., Izmet'eva L. R. Izmet'eva L. R., 1998. Lake Baikal: evolution
- 382           and biodiversity. Backhuys Publishers.
- 383   Krebs, L., Blank, M., Frankowski, J., Bastrop, R., 2010. Molecular characterisation of the
- 384           Microsporidia of the amphipod *Gammarus duebeni* across its natural range revealed
- 385           hidden diversity, wide-ranging prevalence and potential for co-evolution. Infection,
- 386           Genetics and Evolution, 10, 1027-1038.
- 387   Kuzmenkova, Z.V., Sherbakov D.U., Smith J.E., 2008. Diversity of microsporidia, that parasitize
- 388           on baikal amphipods Gmelinoides fasciatus from different populations. The bulletin of
- 389           Irkutsk State University. Series "Biology, Ecology", 1, 56-61.
- 390   Lipa, J. J., 1967. *Nosema kozhovi* sp. n., a new microsporidian parasite of *Brandtia lata lata*

391 (Crustacea, Gammaridae) of Baical Lake. Acta protozool, 5, 93-97.

392 MacNeil, C., Dick, J. T., Hatcher, M. J., Terry, R. S., Smith, J. E., Dunn, A. M., 2003. Parasite-  
 393 mediated predation between native and invasive amphipods. Proceedings of the Royal  
 394 Society of London B: Biological Sciences, 270, 1309-1314.

395 Madyarova, E. V., Adelshin, R. V., Dimova, M. D., Axenov-Gribanov, D. V., Lubyaga, Y. A.,  
 396 Timofeyev, M. A., 2015. Microsporidian parasites found in the hemolymph of four  
 397 baikalian endemic amphipods. PLOS One, 10, e0130311.

398 Martens, K., 1997. Speciation in ancient lakes. Trends in Ecology & Evolution, 12, 177–182.

399 Mashiko, K., 2000. Insights into the mechanism of speciation in gammarid crustaceans of Lake  
 400 Baikal using a population-genetic approach. Advances in Ecological Research, 31, 221-  
 401 235.

402 McClymont, H. E., Dunn, A. M., Terry, R. S., Rollinson, D., Littlewood, D. T. J., Smith, J. E.,  
 403 2005. Molecular data suggest that microsporidian parasites in freshwater snails are  
 404 diverse. International journal for parasitology, 35, 1071-1078.

405 Ovcharenko, M.O., Bacela, K., Wilkinson, T., Ironside, J.E., Rigaud, T. and Wattier, R.A., 2010.  
 406 *Cucumispora dikerogammari* n. gen. (Fungi: Microsporidia) infecting the invasive  
 407 amphipod *Dikerogammarus villosus*: a potential emerging disease in European rivers.  
 408 Parasitology, 137(2), pp.191-204.

409 R Core Team, 2017. R: A language and environment for statistical computing. R Foundation for  
 410 Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>.

411 Rode, N. O., Landes, J., Lievens, E. J., Flaven, E., Segard, A., Jabbour-Zahab, R., ... &

412 Lenormand, T., 2013a. Cytological, molecular and life cycle characterization of  
413 *Anostracospora rigaudi* ng, n. sp. and *Enterocytozpora artemiae* ng, n. sp., two new  
414 microsporidian parasites infecting gut tissues of the brine shrimp *Artemia*. *Parasitology*,  
415 140(9), 1168-1185.

416 Rode, N. O., Lievens, E. J., Segard, A., Flaven, E., Jabbour-Zahab, R., Lenormand, T., 2013b.  
417 Cryptic microsporidian parasites differentially affect invasive and native *Artemia spp.*  
418 *International journal for parasitology*, 43, 795-803.

419 Sambrook, J., Fritsch, E. F., Maniatis, T., 1989. *Molecular cloning: a laboratory manual* (No. Ed.  
420 2). Cold Spring Harbor laboratory press.

421 Smith, J. E., 2009. The ecology and evolution of microsporidian parasites. *Parasitology*, 136,  
422 1901-1914.

423 Smith, J. E., Yang Q., Kamaltynov R. M., Sherbakov D. Y., 2008. Evolutionary interactions  
424 between microsporidia and their hosts: Lessons from an ancient lake. 41st Annual  
425 Meeting of the Society for Invertebrate Pathology and 9th International Conference on  
426 *Bacillus thuringiensis*. August 3-7. University of Warwick, Coventry, United Kingdom.

427 Stamatakis A., 2014. RAxML version 8: a tool for phylogenetic analysis and post-analysis of  
428 large phylogenies. *Bioinformatics*, 30, 1312-1313.

429 Stentiford, G. D., Feist, S. W., Stone, D. M., Bateman, K. S., Dunn, A. M., 2013. Microsporidia:  
430 diverse, dynamic, and emergent pathogens in aquatic systems. *Trends in parasitology*,  
431 29, 567-578.

432 Takhteev, V. V., Berezina, N. A., Sidorov, D. A., 2015. Checklist of the Amphipoda (Crustacea)

- 433 from continental waters of Russia, with data on alien species. *Arthropoda Selecta*, 24,  
434 335-370.
- 435 Terry, R. S., Smith, J. E., Dunn, A. M., 1998. Impact of a novel, feminising microsporidium on  
436 its crustacean host. *Journal of Eukaryotic Microbiology*, 45, 497-501.
- 437 Terry, R.S., Smith, J.E., Sharpe, R.G., Rigaud, T., Littlewood, D. T.J., Ironside, J.E., 2004.  
438 Widespread vertical transmission and associated host sex-ratio distortion within the  
439 eukaryotic phylum Microspora. *Proceedings of the Royal Society of London B:*  
440 *Biological Sciences*, 271, 1783-1789.
- 441 Timoshkin, O.A., 2001. Index of animal species inhabiting Lake Baikal and its catchment area,  
442 book first. Nauka, Novosibirsk.
- 443 Väinölä, R., Witt, J. D. S., Grabowski, M., Bradbury, J. H., Jazdzewski, K., Sket, B., 2008.  
444 Global diversity of amphipods (Amphipoda; Crustacea) in freshwater. *Hydrobiologia*,  
445 595, 241–255.
- 446 Vávra, J., Hyliš, M., Fiala, I., & Nebesářová, J. (2016). *Globulispora mitoportans* ng, n. sp.,  
447 (Opisthosporidia: Microsporidia) a microsporidian parasite of daphnids with unusual  
448 spore organization and prominent mitosome-like vesicles. *Journal of invertebrate*  
449 *pathology*, 135, 43-52.
- 450 Weiss, L. M., 1994. Utility of microsporidian rRNA in diagnosis and phylogeny: a review. *Folia*  
451 *parasitologica*, 41, 81-90.
- 452 Weiss, L. M., Becnel, J. J., 2014. *Microsporidia: Pathogens of Opportunity*. John Wiley & Sons.
- 453 Yoshii, K., 1999. Stable isotope analyses of benthic organisms in Lake Baikal. *Hydrobiologia*,

454           411, 145–159.

455   Yu G, Smith D, Zhu H, Guan Y, Lam TT (2017). ggtree: an R package for visualization and

456           annotation of phylogenetic trees with their covariates and other associated data.

457           Methods in Ecology and Evolution, 8, 28-36

458

# Figure legends

460

461 **Figure 1.** Sampling sites for amphipod collection in Lake Baikal. Grey dots correspond to sites  
462 where no infected amphipods were found, while black dots mark places where microsporidian  
463 DNA was identified in amphipod haemolymph. The bathymetric map of Lake Baikal was taken  
464 from <http://users.ugent.be/~mdbatist/intas/intas.htm> and modified with CorelDRAW.

465 **Figure 2.** Bayesian phylogenetic tree of Microsporidia identified in amphipods of Lake Baikal  
466 and some other crustaceans. The tree contains the following clusters:  
467 Enterocytozoonidae&Hepatosporidae-like (EH), *Enterocytozoon*- (E), *Cucumispora*- (C),  
468 *Dictyocoela*- (D), *Nosema-Vairimorpha*-like (NV), and some unclassified Microsporidia (uM),  
469 *Nosema* (N). Branch lengths are drawn to scale. Grey semi-circles mark nodes with bootstrap  
470 support >70%, and red semi-circles mark those with posterior probability >0.7. Baikal shapes  
471 mark microsporidian sequences amplified from Lake Baikal amphipods, and samples obtained in  
472 this work are in blue font. The alignment used to build this tree is presented in Alignment S1.

473

# Supplementary materials

475 **Supplemental Table S1.** All analysed samples of haemolymph of amphipods collected in Lake  
476 Baikal.

477 **Supplemental Table S2.** All microsporidian DNA identified in haemolymph of amphipods  
478 collected in Lake Baikal. It contains sequences obtained both in the current study and in  
479 (Madyarova et al., 2015). Columns A-H describe samples obtained in our work, while columns



480 H-M describe their closest their closest hits in the NCBI database. E-values equal to 0 represent  
481 values lower than  $10^{-200}$ .

482 **Supplemental Alignment S1.** Raw alignment of microsporidian DNA sequences used for  
483 construction of the phylogenetic tree on Fig. 2.

484 **Supplemental Tree S1.** Bayesian tree obtained from the Alignment S1.

485 **Supplemental Tree S2.** Maximum likelihood bootstrap support for the tree obtained from the  
486 Alignment S1.


487 **Supplemental Figure S1.** Inverse Simpson's diversity index for each location and number of  
488 samples for each location where microsporidian DNA was detected. The vertical axis features  
489 the number of microsporidian isolates.

# Figure 1

Sampling sites for amphipod collection in Lake Baikal.

Grey dots correspond to sites where no infected amphipods were found, while black dots mark places where microsporidian DNA was identified in amphipod haemolymph. The bathymetric map of Lake Baikal was taken from





<http://users.ugent.be/~mdbatist/intas/intas.htm> and modified with CorelDRAW.



## Figure 2 (on next page)

Bayesian phylogenetic tree of Microsporidia identified in amphipods of Lake Baikal and some other crustaceans.

The tree contains the following clusters: Enterocytozoonidae&Hepatosporidae-like (EH), *Enterocytozoon*- (E), *Cucumispora*- (C), *Dictyocoela*- (D), *Nosema-Vairimorpha*-like (NV), and some unclassified Microsporidia (uM), *Nosema* (N). Branch lengths are drawn to scale. Grey semi-circles mark nodes with bootstrap support >70%, and red semi-circles mark those with posterior probability >0.7. Baikal shapes mark microsporidian sequences amplified from Lake Baikal amphipods, and samples obtained in this work are in blue font. The alignment used to build this tree is presented in Alignment S1.

 Bootstrap > 70%  
 Posterior probability > 0.7  
 Baikal samples  
 Samples obtained in this work

0.04

