

Anatomical notes and discussion of the first described aetosaur *Stagonolepis robertsoni* (Archosauria: Suchia) from the Upper Triassic of Europe, and the use of plesiomorphies in aetosaur biochronology (#25967)

1

First submission

Editor guidance

Please submit by **20 Apr 2018** for the benefit of the authors (and your \$200 publishing discount).



Structure and Criteria

Please read the 'Structure and Criteria' page for general guidance.



Raw data check

Review the raw data. Download from the [materials page](#).



Image check

Check that figures and images have not been inappropriately manipulated.

Privacy reminder: If uploading an annotated PDF, remove identifiable information to remain anonymous.

Files

Download and review all files from the [materials page](#).

11 Figure file(s)



Structure your review

The review form is divided into 5 sections.

Please consider these when composing your review:

1. BASIC REPORTING

2. EXPERIMENTAL DESIGN

3. VALIDITY OF THE FINDINGS

4. General comments

5. Confidential notes to the editor

 You can also annotate this PDF and upload it as part of your review

When ready [submit online](#).

Editorial Criteria

Use these criteria points to structure your review. The full detailed editorial criteria is on your [guidance page](#).

BASIC REPORTING

-  Clear, unambiguous, professional English language used throughout.
-  Intro & background to show context. Literature well referenced & relevant.
-  Structure conforms to [PeerJ standards](#), discipline norm, or improved for clarity.
-  Figures are relevant, high quality, well labelled & described.
-  Raw data supplied (see [PeerJ policy](#)).

EXPERIMENTAL DESIGN

-  Original primary research within [Scope of the journal](#).
-  Research question well defined, relevant & meaningful. It is stated how the research fills an identified knowledge gap.
-  Rigorous investigation performed to a high technical & ethical standard.
-  Methods described with sufficient detail & information to replicate.

VALIDITY OF THE FINDINGS

-  Impact and novelty not assessed. Negative/inconclusive results accepted. *Meaningful* replication encouraged where rationale & benefit to literature is clearly stated.
-  Data is robust, statistically sound, & controlled.
-  Conclusions are well stated, linked to original research question & limited to supporting results.
-  Speculation is welcome, but should be identified as such.

Standout reviewing tips

3



The best reviewers use these techniques

Tip

Support criticisms with evidence from the text or from other sources

Example

Smith et al (J of Methodology, 2005, V3, pp 123) have shown that the analysis you use in Lines 241-250 is not the most appropriate for this situation. Please explain why you used this method.

Give specific suggestions on how to improve the manuscript

Your introduction needs more detail. I suggest that you improve the description at lines 57- 86 to provide more justification for your study (specifically, you should expand upon the knowledge gap being filled).

Comment on language and grammar issues

The English language should be improved to ensure that an international audience can clearly understand your text. Some examples where the language could be improved include lines 23, 77, 121, 128 – the current phrasing makes comprehension difficult.

Organize by importance of the issues, and number your points

1. Your most important issue
2. The next most important item
3. ...
4. The least important points

Please provide constructive criticism, and avoid personal opinions

I thank you for providing the raw data, however your supplemental files need more descriptive metadata identifiers to be useful to future readers. Although your results are compelling, the data analysis should be improved in the following ways: AA, BB, CC

Comment on strengths (as well as weaknesses) of the manuscript

I commend the authors for their extensive data set, compiled over many years of detailed fieldwork. In addition, the manuscript is clearly written in professional, unambiguous language. If there is a weakness, it is in the statistical analysis (as I have noted above) which should be improved upon before Acceptance.

Anatomical notes and discussion of the first described aetosaur *Stagonolepis robertsoni* (Archosauria: Suchia) from the Upper Triassic of Europe, and the use of plesiomorphies in aetosaur biochronology

William G Parker^{Corresp., 1}, Jeffrey W Martz²

¹ Division of Science and Resource Management, Petrified Forest National Park, Petrified Forest, Arizona, United States

² College of Sciences and Technology, University of Houston, Downtown, Houston, Texas, United States

Corresponding Author: William G Parker

Email address: William_Parker@nps.gov

Stagonolepis robertsoni was the first named aetosaurian from the Late Triassic of Scotland. Known mostly from a series of natural molds from two localities, the osteology of this taxon has been difficult to interpret. Detailed work on this material in the late 1950s resulted in a monograph that set the standard for the understanding of aetosaurians, making *Stagonolepis robertsoni* the best known aetosaurian; however, little has been done with this material since. Reanalysis of this material shows that despite the limitations of the material, the early 1960s reconstruction work depicts the preserved characteristics faithfully, especially in the skull. The first cervical rib is extremely anteroposteriorly elongate as in *Alligator*, a character not previously recognized in aetosaurians. Diapophyseal and zygapophyseal vertebral laminae are present in the cervical and trunk vertebrae. The ilium is autapomorphic with distinct pre- and post-processes of the iliac blade. The osteoderms differ from North and South American material that has been ascribed to the genus and those assignments are based on plesiomorphies within Aetosauria, such as a radial ornamentation and a posteriorly located and medially offset dorsal eminence. Biostratigraphic correlations using taxonomic conclusions based on plesiomorphic characters should not be used. The holotype specimen of *S. robertsoni* is currently diagnostic in part because ventral osteoderms are not known for many aetosaurian taxa and the surface ornamentation of randomly distributed, closely packed oblong pits found in *S. robertsoni* is unique for Aetosauria.

Anatomical notes and discussion of the first described aetosaur *Stagonolepis robertsoni*
(Archosauria: Suchia) from the Upper Triassic of Europe, and the use of plesiomorphies in
aetosaur biochronology.

William G. Parker^{1*}, Jeffrey W. Martz²

¹Division of Science and Resource Management, Petrified Forest National Park, Petrified Forest,
Arizona 86028

²Department of Natural Sciences, University of Houston-Downtown, Houston, Texas 77002

*Corresponding Author: William_Parker@nps.gov

ABSTRACT

Stagonolepis robertsoni was the first named aetosaurian from the Late Triassic of Scotland. Known mostly from a series of natural molds from two localities, the osteology of this taxon has been difficult to interpret. Detailed work on this material in the late 1950s resulted in a monograph that set the standard for the understanding of aetosaurians, making *Stagonolepis robertsoni* the best known aetosaurian; however, little has been done with this material since. Reanalysis of this material shows that despite the limitations of the material, the early 1960s reconstruction work depicts the preserved characteristics faithfully, especially in the skull. The first cervical rib is extremely anteroposteriorly elongate as in *Alligator*, a character not previously recognized in aetosaurians. Diapophyseal and zygapophyseal vertebral laminae are present in the cervical and trunk vertebrae. The ilium is autapomorphic with distinct pre- and post-processes of the iliac blade. The osteoderms differ from North and South American material that has been ascribed to the genus and those assignments are based on plesiomorphies within Aetosauria, such as a radial ornamentation and a posteriorly located and medially offset dorsal eminence. Biostratigraphic correlations using taxonomic conclusions based on plesiomorphic characters should not be used. The holotype specimen of *S. robertsoni* is currently diagnostic in part because ventral osteoderms are not known for many aetosaurian taxa and the surface ornamentation of randomly distributed, closely packed oblong pits found in *S. robertsoni* is unique for Aetosauria.

INTRODUCTION

Aetosaurs are exclusively Late Triassic quadrupedal, heavily armored pseudosuchians known globally with the exception of some parts of Gondwana: South Africa, Antarctica, and Australia. The most recent overview of this clade is by Desojo et al. (2013). The first described aetosaurian fossil was christened *Stagonolepis robertsoni* for a series of ventral osteoderms. Because the unit they were discovered in was thought to be the Devonian ‘Old Red Sandstone’, these osteoderms were considered to represent a large ganoid fish (Agassiz 1844). Subsequent discoveries (detailed in Walker 1961) demonstrated that these remains belonged instead to a stem pseudosuchian, and that the rocks they were found in were instead more likely Triassic in age (Huxley 1859, 1869; 1875; 1877). The original materials worked on by Louis Agassiz and T.H. Huxley, as well as additional discoveries made in the 1920s and 1930s, were the subject of an extensive monograph (Walker 1961). This work was influential because it finally solidified that aetosaurians were a globally distributed Late Triassic group distinct from phytosaurs.

Although since that time the only material described for *Stagonolepis robertsoni* is a braincase (MCZD 2-4; Gower & Walker 2002), *S. robertsoni* is still considered to be the ‘model’ aetosaurian (Desojo et al. 2013) and the taxon is featured prominently in phylogenetic analyses (e.g., Parrish 1994; Heckert, Hunt, & Lucas 1996; Heckert & Lucas 1999; Parker 2007; Parker 2016a). However, since 1961 the taxon has never been adequately re-evaluated in terms of the great strides that have been made in our understanding of aetosaurian morphology and systematics, particularly the recognition that armor surface ornament, width-length ratios, and lateral osteoderm morphology are diagnostic to clades, and in some cases, to species (Long & Ballew 1985; Long & Murry 1995; Heckert & Lucas 2000; Parker 2007; Martz & Small 2006). Other named taxa (e.g., *Calyptosuchus wellesi*, *Aetosauroides scagliai*) have been assigned to the genus *Stagonolepis* based on similarity of the dorsal paramedian osteoderm ornamentation (e.g.,

Long and Murry, 1995; Heckert & Lucas 2002); however, these assignments have been questioned by other workers (e.g., Desojo & Ezcurra 2011; Parker 2018).

The purpose of this paper is the reanalyze referred material of *Stagonolepis robertsoni* present at the Natural History Museum, London and evaluate it in recent context based on our current understanding of aetosaurian anatomy. The material is not completely redescribed here as the original descriptions by Walker (1961) and Gower & Walker (2002) are still quite adequate, instead we focus on characteristics whose taxonomic significance was not fully recognized by Walker given the understanding of aetosaurians at the time of his monograph. These newly recognized characteristics are useful for examining the phylogenetic relationships of *S. robertsoni* and other aetosaurians.

MATERIALS

The materials of *Stagonolepis robertsoni* at the Natural History Museum, London that were available to Huxley (1859, 1875, 1877) consist only of a couple of now chipped casts of a left femur (NHMUK R581) and a paramedian osteoderm and phalange (NHMUK R582). The rest of the material are in other institutions such as the Geological Survey Museum, London and the Elgin Museum in Scotland (Walker 1961; Walker undated and unpublished notes at the NHMUK). More prominent materials consist almost entirely of large blocks of a highly indurated sandstone that preserve natural molds of the various bones. Painted casts of these were created in 1885 and presented to the museum by the Reverend George Gordon. These specimens allow study of portions of the bones in two dimensions, and were available for study by Charles Camp in 1935 (Camp unpublished notes at the UCMP, 1935) and Alick Walker in the 1950s (Walker 1961). Walker (1961) also created a series of PVC casts (e.g., NHMUK 4787) from the

original sandstone molds that preserved more detail and allowed for 3D visualization of the bones. These PVC casts are still in the NHMUK collections although many have deteriorated.

For this study the following blocks/specimens were examined and photographed: 

MCZD2 (5 parts), partial skull and articulated cervical osteoderms; NHMUK R581, cast of left femur; NHMUK R582, cast of paramedian osteoderm and phalange; NHMUK R4784a, cast of slab with basioccipital and associated articulated series of cervical and trunk vertebrae, including the cervical ribs, scapulocoracoid, partial humerus; NHMUK R4785a, cast of slab with osteoderms and ribs; NHMUK R4786a, cast of slab with anterior caudal centra and ribs in articulation, distal end of left tibia; NHMUK R4787; Sandstone block containing natural mold of much of a skull plus associated PVC cast; NHMUK R4787a, cast of specimen showing the lower portion of the left side of a good skull, especially the mandible; NHMUK R4789a, cast of slab with ribs, osteoderms, right ilium, partial maxilla; NHMUK R4790a, cast of slab with osteoderms, right ilium, ischium; NHMUK R4797a, cast with lower limb bones and appendicular osteoderms; NHMUK R4799, sandstone block with impression of an anterior cervical vertebra; NHMUK R8586, cast of EM38R, left side of internal surface of snout (figured by Huxley, 1877); NHMUK R27404, cast of the holotype specimen that consists of a segment of the articulated ventral carapace; NHMUK R36392, sandstone slab with osteoderms, including six articulated left lateral osteoderms; NHMUK R36394; sandstone slab with impressions of articulated ventral osteoderms;

An attempt was made to use the CT scanner at the NHMUK to scan the negative space in block NHMUK R4787 to get a three-dimensional view of the skull, but unfortunately that scanner could not penetrate the hard sandstone matrix of the block 

121 **SYSTEMATIC PALEONTOLOGY**

122 **Archosauria Cope 1869 sensu Gauthier & Padian 1985**

123 **Pseudosuchia Zittel 1887-1890 sensu Gauthier & Padian 1985**

124 **Aetosauria Marsh 1884 sensu Parker 2007**

125 **Desmotosuchia sensu Parker 2016a**

126 **Stagonolepidinae sensu Heckert & Lucas 2000**

127 ***Stagonolepis* Agassiz 1844**

128 ***Stagonolepis robertsoni* Agassiz 1844**

129 **(Figs. 1-7, 9, 11)**

130 1844 *Stagonolepis robertsoni*: Agassiz, p. 139, pl. XXXI, figs. xiii, xiv

131 1859 *Stagonolepis robertsoni*: Huxley, p. 440, pl. XIV, figs. 1-3

132 1877 *Stagonolepis robertsoni*: Huxley, p. 1, pl. I-X.

133 1902 *Staganolepis* [sic] *robertsoni*: Huene, p. 54, figs. 62-67, 72, 73.

134 1908 *Stagonolepis robertsoni*: Huene, p. 392, figs. 347-348.

135 1936 *Stagonolepis robertsoni*: Huene, p. 207, fig. 3.

136 1942 *Stagonolepis robertsoni*: Huene, p. 223, figs. 45-49.

137 1961 *Stagonolepis robertsoni*: Walker, p. 103, figs. 2-23, 24b, 25b, pl. 9-12.

138 1976 *Stagonolepis robertsoni*: Krebs, p. 40, figs. 3, 4, 9, 10d, 12, 15, 16, 17c-e, 19d-e,

139 20d-e, 26b, 27.

140 1978 *Staganolepis* [sic]: Bonaparte, p. 300, figs. 137b, 138.

141 1986 *Stagonolepis*: Parrish, p. 8, fig. 6, 14c3.

142 1988 *Stagonolepis*: Carroll, p. 273, figs. 13.15, 13.16.

143 1988 *Stagonolepis*: Fraser, p. 132, fig. 5b.

- 144 1991 *Stagonolepis*: Sereno, p. 11, fig. 10, 27f.
 145 1996 *Stagonolepis*: Lucas & Heckert, p. 57, fig. 4.
 146 2000 *Stagonolepis robertsoni*: Heckert & Lucas, 2000, p. 1552, figs. 4c, e.
 147 2001 *Stagonolepis robertsoni*: Lucas & Heckert, p. 719, figs. 2, 3.
 148 2002 *Stagonolepis robertsoni*: Gower & Walker, p. 7, figs. 1-4, 6.
 149 2010 *Stagonolepis robertsoni*: Sulej, p. 878, figs. 8a, 9f.
 150 2011 *Stagonolepis robertsoni*: Desojo & Ezcurra, p. 599, figs. 3d-f, 7b.
 151 2013 *Stagonolepis robertsoni*: Desojo et al., p. 207, figs. 3e-f, 4g, 5a-h, 6a-f, 7c-f.
 152 2016a *Stagonolepis robertsoni*: Parker, p. 32, figs. 1, app. B, fig. 11j.

153

154 **Holotype** – ELGNM 27R, impression of a segment of the ventral carapace (Agassiz
 155 1844). NHMUK R27404 is a negative cast of this specimen.

156 **Referred Material** – see Materials for a list of NMHUK specimens.

157 **Occurrence** – Lossiemouth Sandstone Formation, Morayshire, Scotland, U.K (Walker
 158 1961).

159 **Age** – Late Triassic, Late Carnian to Early Norian (Benton & Walker 2010) 

160 **Diagnosis** – *Stagonolepis robertsoni* is diagnosed by the following autapomorphies:

161 ventral osteoderms rectangular with randomly arranged, oblong pits* first cervical vertebrae 

162 with elongate cervical ribs that extend back to the position of the 4th cervical vertebra*; trunk

163 paramedian osteoderms lack ornament along the posterior edge in the region of the dorsal

164 eminence  posterior process of iliac blade forms an acute angled tip and the anterior process of

165 the iliac blade is anteroposteriorly short, dorsoventrally thin, and ventrally hooked*; pubis with

166 two obturator foraminae* (convergent in *Scutarx deltatylus*) 

167 *Stagonolepis robertsoni* can also be distinguished by the following combination of
 168 character states: premaxilla with four, possibly five teeth in “*Stagonolepis*” *kae* and
 169 *Paratypothorax andressorum* differs from the three in *Stenomyti huangae* and the absence of
 170 premaxillary teeth in *Desmatosuchus smalli*); premaxillary tip expanded laterally as in
 171 “*Stagonolepis*” *olenkae*, *Desmatosuchus smalli*, and *Neoaetosauroides engaeus* (absent in
 172 *Aetosaurus ferratus*, *Stenomyti huangae*; *Paratypothorax andressorum*); distinct ridge on lateral
 173 side of maxilla beneath antorbital fossa as in *Paratypothorax andressorum* and *Stenomyti*
 174 *huangae* (this ridge is absent in *Desmatosuchus smalli*, *Longosuchus meadei*, *Neoaetosauroides*
 175 *engaeus*, and “*Stagonolepis*” *olenkae*); long axis of the jugal anterodorsally inclined as in
 176 *Desmatosuchus spurensis* and *Longosuchus meadei* (the ventral margin of the jugal is level in
 177 *Aetosaurus ferratus*, *Paratypothorax andressorum*, and *Stenomyti huangae*); teeth thecodont
 178 with a swollen base and non-recurved tip as in *Desmatosuchus smalli* (differs from
 179 *Aetosauroides scagliai* and *Aetosaurus ferratus* Parabasisphenoid elongate with
 180 anteroposteriorly separated basal tubera and basiptyergoid processes as in *Neoaetosauroides*
 181 *engaeus*, *Aetosaurus ferratus*, and *Desmatosuchus smalli* (differs from *Scutarx deltatylus*;
 182 *Paratypothorax andressorum*, *Tecovasuchus chatterjeei*, and *Desmatosuchus spurensis*);
 183 mandible “slipper-shaped” with ventrolateral portion of splenial visible in lateral view beneath
 184 the ventral margin of the dentary as all aetosaurs except for *Aetosauroides scagliai* and
 185 *Typothorax coccinarum*; posterior cervical vertebrae with ventral keel (as in *Aetosauroides*
 186 *scagliai*, *Calyptosuchus wellsi*, *Sierritasuchus macalpini*, and *Scutarx deltatylus* keels absent in
 187 *Longosuchus meadei* and *Desmatosuchus spurensis* trunk paramedian osteoderms with a
 188 length/width ratio of about 2.5:1 as in *Desmatosuchus spurensis*, *Longosuchus meadei*, and
 189 *Aetobarbakinoides brasiliensis* (differs from the much wider paramedians of *Typothorax*

coccinarum and *Paratypothorax andressorum* raised anterior bar on osteoderms as in all non-desmotosuchin aetosaurs; anteromedial and anterolateral projections of the anterior bar present on the trunk osteoderms as in all non-desmotosuchin aetosaurs; anterolateral projection of the anterior bar not elongate as in “*Stagonolepis*” *olenkae*, *Aetosaurus ferratus*, and *Typothorax coccinarum* (differs from *Adamanasuchus eisenhardtae* and *Scutarx deltatylus*); dorsal eminence of trunk paramedian osteoderms situated on the posterior osteoderm margin and offset medially (as in most aetosaurs except *Desmotosuchus* and *Paratypothorax andressorum*); dorsal surface ornament of the paramedian and lateral osteoderms anastomosing (interconnected series of radiating ridges surrounding subrounded and subrectangular pits) lateral osteoderms are nearly equant with distinct lateral and dorsal flanges as in all non-desmotosuchin aetosaurs; lateral osteoderms lack pronounced horns or spines as in most aetosaurs outside of Typothoracisinae and Desmotosuchini.

DESCRIPTION

Skull

Block NHMUK R4787a is a cast of the lower portion of a skull in semi-articulation including much of the lower jaw, quadrate, portions of the palate and maxilla, and the premaxilla (Figure 1A). These elements represent the left side of the skull, so the cast provides an internal (medial) view. The semi-articulated condition allows for the determination of the skull length, which from the retroarticular process to the tip of the premaxilla is about 240 mm. The dentigerous elements show the presence of at least eight dentary, four maxillary, and four premaxillary teeth. Alick Walker’s PVC cast of this specimen shows even more details (Figure 1B) including the upper portions of the skull and the braincase, especially the parabasisphenoid. Thus block NHMUK R4787 is the natural mold of a nearly complete skull of *Stagonolepis robertsoni* (Walker 1961).

213 Details of this specimen include: 1) a possible 5th premaxillary tooth crown, or alternatively the
 214 tip of the right premaxilla in articulation with the left; 2) a portion of the right squamosal and
 215 impressions of the skull roof, 3) the left quadrate in articulation with the left articular, and 4)
 216 impressions of the braincase, especially the basisphenoid including the left and right basitubera
 217 and basiptyergoid processes, and the cultriform process is also preserved.

218 A cast of the right maxilla from NHMUK R4787 (Figure 2) shows that overall the
 219 element is more slender than that of “*Stagonolepis*” *olenkae* (Sulej 2010) with at least six alveoli,
 220 but the anterior portion is covered. A very distinct transverse ridge is present on the lateral
 221 surface along the anterior and ventral borders of the antorbital fossa, making the fossa extremely
 222 pronounced; this ridge also occurs in *Stenomyti huangae* (Small and Martz, 2013) and
 223 *Paratypothorax andressorum* (Schoch and Desojo, 2016) but is not present in “*Stagonolepis*”
 224 *olenkae* (Sulej 2010), *Desmotosuchus* (Case, 1922; Small, 2002), *Longosuchus* (Parrish, 1994),
 225 or *Neoatesauroides* (Desojo and Báez, 2007). The medial side of the maxillary body is marked
 226 by an elongate medial shelf that articulates with the palatal bones (Walker 1961). The
 227 articulations with the lacrimal and jugal are each complex, with regions of overlap between the
 228 two bones; this is also the case at least in *Longosuchus meadei* (J. Martz, unpublished notes), but
 229 this region is rarely found sufficiently disarticulated in aetosaurs to establish the relationships
 230 between these elements. Finally, a pneumatic accessory cavity is present on the medial side of
 231 the ascending process as described for *Desmotosuchus smalli* (Small 2002).

232 NHMUK R8586 is a cast of ELGNM 38R (Figure 3), which was figured by Huxley
 233 (1877) and features the left side of the internal portion of the snout, including the premaxilla,
 234 maxilla, and nasals. The premaxilla measures 63 mm in length, bears four teeth with an
 235 edentulous anterior portion typical for aetosaurians; four or more teeth also occur in *Aetosaurus*

236 *ferratus* (Schoch, 2007) “*Stagonolepis*” *olenkae* (Sulej, 2010), and *Paratypothorax andressorum*
 237 (Schoch and Desojo, 2016) whereas *Stenomyti huangae* only possesses three (Small and Martz,
 238 2013), and *Desmatosuchus* lacks any (Small, 2002). The maxilla bears six teeth as preserved, but
 239 it is missing the posterior portion. The external naris is 22 mm at its deepest point and 72 mm in
 240 length, but missing the posteriormost section. The nasal has a pronounced ridge on the medial
 241 edge, which at a position just dorsal to the 3rd premaxillary tooth migrates ventrally to the ventral
 242 margin of the nasal where it contacts the premaxilla. The anterior tip of the premaxilla bears a
 243 prominent ridge that divides the element into a flat surface that slopes into the external naris, and
 244 a second triangular area that slopes anteroventrally (Figure 3). This ridge is the anterior
 245 expansion and also occurs in other aetosaurians such as *Desmatosuchus smalli* (Small 2002),
 246 *Neoaetosauroides engaeus* (PVL 4363), and differs from aetosaurians such as *Aetosaurus*
 247 *ferratus* (Schoch 2007); *Aetosauroides scagliai* (PVL 2073), and *Typothorax coccinarum* (YPM
 248 58121). The premaxilla bears a small dorsal protuberance above the first tooth position that
 249 extends dorsally into the external naris, which also occurs in “*Stagonolepis*” *olenkae* (Sulej
 250 2010) and *Desmatosuchus smalli* (Small 2002); however, in “*S.*” *olenkae* it is dorsal to the
 251 second tooth position (Antczak 2016) and *D. smalli* has an edentulous premaxilla (Small 2002).
 252 There is only a slight swelling in this position in *Stenomyti huangae* (Small and Martz, 2013).

253 The best-preserved skull material is MCZD 2, which consists of seven small blocks that
 254 fit together to present much of the skull and the anterior section of the neck (Walker 1961:figs.
 255 26-29; Gower & Walker 2002:fig. 1). The material consists of well-preserved bone and is not a
 256 natural mold as is most of the *S. robertsoni* material. This specimen has previously been
 257 described in great detail so this will not be duplicated here (Walker 1961; Gower & Walker
 258 2002). However, notable is that the basitubera and basiptyergoid processes are widely separated

anteroposteriorly from each other so that the parabasisphenoid is elongate supporting what can be observed in NHMUK R4787. *Neoaetosauroides engaeus* (PVL 5698; Desojo & Báez 2007), *Aetosaurus ferratus* (Schoch 2007) and the basal taxon *Aetosauroides scagliai* (PVSJ 326) also have an elongate parabasisphenoid so that is the plesiomorphic state within Aetosauria. This differs significantly from the condition in “*S. olenkae*”, where the basitubera and basiptyergoid processes nearly contact (Sulej, 2010: figs. 1d, f) and thus the parabasisphenoid is anteroposteriorly short. In *Scutarx deltatylus* (Parker, 2016b: fig. 7) and *Desmatosuchus spurensis* (UMMP 7476) the parabasisphenoid is also anteroposteriorly short; whereas the condition in *Calyptosuchus wellsi* is unknown (Parker 2018).

Cervical vertebrae and ribs

In block NHMUK R4784a, which is a cast created from a natural mold and represents the postcranial skeleton of the skull NHMUK R4787 (Walker 1961), the occipital condyle and left paroccipital process of the skull are present (Figure 4A). If the cervical count is nine (as in *Desmatosuchus spurensis*) the four anterior cervicals are missing (including the axis and atlas); however, cervical ribs are present for three of these positions. Very striking are two very elongate posteriorly projecting cervical ribs underlying the ventral surfaces of the two more posteriorly positioned ribs (Figure 4A). The elongate ribs (left and right sides) originate where the axis/atlas would be located and are very similar to the greatly elongate cervical ribs found in *Alligator* (Reese 1915). These exposed ribs measure 85mm in length (the ends are covered), more than three times the lengths of the other exposed cervical ribs. The elongate ribs were not noted by Walker (1961) and have not previously been described for any aetosaur. The axis/atlas and third cervical are present in MCZD 2, but unfortunately are very poorly preserved; however, this block also preserves a very elongate first cervical rib.

The first well-preserved cervical vertebra in NHMUK R4784a is the 8th (Walker 1961), which is visible in right lateral view and preserves details of the centrum and much of the neural arch and transverse process (Figure 4B). A disarticulated cervical rib is present across the centrum and probably does not belong to this vertebra. The centrum measures 25 mm in length and the ventral surface is strongly keeled. Keels occur on the ventral surface of the cervical vertebrae in *Aetosauroides* (Desojo and Ezcurra, 2011) and *Scutarx* (Parker 2016) while the ventral surface of the cervicals lacks keels in *Longosuchus* (Long and Murry, 1995) and *Desmatosuchus* (Case, 1922; Parker, 2018). The condition may be variable in *Typothorax*; ventral keels are absent in the cervicals of the small specimen described by Martz (2002) but present in at least some larger specimens (Heckert et al., 2010) suggesting there may be allometric or ontogenetic variability. The parapophysis in the 8th cervical of NHMUK R4784a is low on the anterior rim of the centrum, but not completely at the base. The transverse process projects laterally and slightly ventrally, is 25 mm long, and bears a flaring sub-rectangular head in lateral view. A distinct posterior centrodiapophyseal lamina (Wilson 1999) stretches from the base of the transverse process to the posterior portion of the neurocentral suture. This is the “T-beam” structure described by Case (1922) for the posterior cervical and dorsal vertebrae of *Desmatosuchus spurensis* and described as present in *Stagonolepis robertsoni* (Walker, 1961) and *Typothorax* (Martz, 2002) and also occurs in *Paratypothorax* (Martz et al., 2013, fig. 9D). The right prezygapophysis is present but not enough is present to tell if a hyposphene was present. The neural spine is present, but missing the apex.

The next three vertebrae are preserved in articulation (Figure 4B). The parapophysis on the 9th cervical is still situated on the centrum, but slightly higher than its position in the preceding vertebra. The centrum is more elongate as well measuring 31 mm. Pre- and

postzygadiapophyseal laminae (Wilson 1999) are visible on the 9th and 10th presacrals (Figure 4B). The 10th presacral is transitional between the cervical and dorsal series as previously described for *Desmotosuchus spurensis* (Case 1922; Parker 2008). The centrum is more elongate than the previous vertebra with a length of 36mm. However, there is still a slight ventral keel as in the other cervicals, and although the parapophysis has migrated upwards onto the base of the neural arch, it is not located on the arch itself as in the dorsal vertebrae. The neurocentral suture appears to be open.

The 11th presacral is the first true dorsal vertebra as the parapophysis would now be situated on the posteroventral surface of the transverse process, unfortunately this can't be seen clearly as it is broken away (Walker 1961:fig. 7i). The centrum has a length of 39 mm and is unkeeled. The 12th presacral is present, but covered by broken ribs. The cervical and trunk vertebrae of *S. robertsoni* lack oval depressions on the lateral sides just below the neurocentral sutures as in *Aetosauroides scagliai* (Desojo & Ezcurra 2011).

Caudal Vertebrae

NHMUK R4799b is a PVC cast of an isolated anterior caudal vertebra providing more details of the neural arch and spine (Walker 1961: figs. 10c-e). The broad transverse processes (left equals 79 mm) do not extend ventral to the base of the centrum; the postzygapophyses are oriented at 45 degrees above horizontal. The centrum is blocky, with equant width and height of about 40 mm, but the entire vertebral height is 112 mm, with the neural spine contributing 40 mm to this measurement. Spinopostzygapophyseal laminae (Wilson 1999) are present, as is the expanded neural spine table.

Osteoderms

Numerous osteoderms of *Stagonolepis robertsoni* are preserved as natural molds; however, often they only produce partial casts. Walker (1961) discussed the osteoderms in what we now consider superficial terms, therefore we redescribe one each of the best preserved dorsal paramedian and lateral osteoderms. The dorsal trunk paramedian osteoderm (NHMUK 4790a; Figure 5A) is from the left side and has a surface patterning of what we call anastomosing, an interlaced network of high ridges surrounding circular and elongate pits closer to the posterior plate margin, and elongate, but irregular grooves on the anterior portion of the osteoderm. This is similar to the pattern in *Aetosaurus ferratus*, *Aetosauroides scagliai*, *Neoaetosauroides engaeus*, “*Stagonolepis*” *olenkae*, and *Paratypothorax andressorum* that the ornamentation is non-radial posteromedially, but elongate and radial anteriorly along the full length of the anterior bar. However, in *S. robertsoni*, the ornamentation on the posterolateral region of the paramedians is distinctly non-radial and faint to absent along the entire posterior margin, whereas in the other taxa the paramedians possess very elongate grooves in the posterolateral portion of the paramedian osteoderm that are nearly parallel to the posterior margin.

This anastomosing ornamentation in *S. robertsoni* radiates from an elongate, but narrow, raised dorsal eminence that contacts the posterior osteoderm margin as in most non-desmatosuchine taxa that possess an eminence; in desmatosuchines, the eminence often contacts the posterior margin (Parker, 2018; Parker and Martz, 2010). This eminence is rounded rather than distinctly pyramidal as in *Typothorax* (Martz, 2002) and *Longosuchus* (Parker and Martz, 2010), and offset slightly medial to the center of the osteoderm as in nearly all aetosaur trunk paramedians. The anterior portion of the osteoderm bears a raised, transverse, smooth strip of bone called the anterior bar (Long & Ballew 1985) that bears anterolateral projection and anteromedial projections as in most non-desmatosuchine aetosaurs. The anterior bar maintains an

even width across the lateral portion of the osteoderm, but thins significantly medially before expanding again at the anteromedial projection. We term this distinct medial thinning, ‘scalloping’ (Parker, 2016b). This feature occurs in several other aetosaurians including *Aetosauroides scagliai* (PVL 2073); *Calyptosuchus wellsi* (UCMP 126844; Parker 2018), *Scutarx deltatylus* (Parker 2016b), and *Paratypothorax andressorum* (SMNS 5721; unpublished data). The medial edge is straight and the lateral edge slightly sinuous in dorsal view. In posterior view the osteoderm is moderately flexed. 

The lateral trunk osteoderm (NHMUK 4789a; Figure 5B) is from the right side based on the presence of a distinct beveling of the anteromedial corner of the anterior bar, which represents an articulation surface for the anterolateral process of the adjacent paramedian osteoderm. The bar is thin but stretches across the entire anterior margin of the osteoderm. The osteoderm is trapezoidal in dorsolateral view and a ridge-like dorsal eminence that contacts the posterior margin divides the osteoderm into distinct dorsal and lateral flanges. The dorsal flange is roughly trapezoidal in dorsal view, whereas the lateral flange is slightly larger and sub-rectangular in dorsolateral view. The surface ornamentation is anastomosing and very faint in the posterior portion of the osteoderm, as seen in the paramedian. The osteoderm is slightly flexed ventrally and the angle between the two flanges is obtuse. In dorsolateral view the lateral margin is gently rounded. The medial margin is angled posteromedially, corresponding with the shape of the adjacent paramedian osteoderm.

The ventral osteoderms (NHMUK R27404 [negative cast of holotype ELGNM 27R]; NHMUK R4789a; Figure 6A, B) are rectangular in dimension, bear an anterior bar, and possess an ornamentation of randomly arranged oblong pits, the ‘drops’ which Agassiz (1844) used to formulate the genus name. These differ from what is seen in *Calyptosuchus wellsi* (UCMP

27225; Parker 2018), *Aetosauroides scagliai* (MCP13-a-b-PV; Desojo & Ezcurra 2011) and *Typothorax coccinarum* (Martz, 2002; Heckert et al. 2010). These taxa also have rectangular ventral osteoderms, but the ornamentation arrangement is more radial and consists of very elongate furrows and ridges, even in *Typothorax*, where paramedian osteoderm ornamentation is distinctly pitted. *Coahomasuchus kahleorum* (Heckert & Lucas 1999) and *Aetosaurus ferratus* (Schoch 2007) preserve more equant, overlapping ventral osteoderms; but the surface ornament on these specimens is poorly preserved and difficult to comprehend. Where they are preserved; however, they appear to consist of finer pits, less densely packed, but nonetheless have a strong radial distribution as in *Coahomasuchus chathamensis* (Heckert, Fraser & Schneider 2017) and the previous taxa. The ventral osteoderms of *S. robertsoni* also differ from *Stenomyti huangae* where the osteoderms are subrounded rather than square and broadly separated rather than overlapping (Small & Martz 2013).

Ilium

Described in detail by Walker (1961) the ilium of *Stagonolepis robertsoni* is autapomorphic. The shapes and sizes of the processes of the iliac blade are particularly diagnostic for aetosaurian taxa. *S. robertsoni* possesses an elongate postacetabular blade that ends in an acute angled end in medial view (Figure 7). Most other aetosaurians have a squared-off end of the postacetabular blade such as *Aetosauroides scagliai* (PVSJ 326; PVL 2073), *Calypotosuchus wellesi* (UCMP 32422), *Longosuchus meadei* (TMM 31185-40) and *Neoaetosauroides engaeus* (PVL 3525). *Typothorax coccinarum* (UMCP 35255) has a postacetabular blade that has a squared off end and is very short, barely extending past the posterior edge of the ischiadic peduncle.

The preacetabular blade of *S. robertsoni* differs from almost all aetosaurians in that it is very short (does not extend to the anterior margin of the pubic peduncle, very narrow, and ventrally hooked (Figure 7). Most aetosaurians have preacetabular blades that extend anteriorly to the edge of the pubic peduncle, and a thick and triangular in lateral view, such as *Aetosauroides scagliai* (PVL 2073), *Desmatosuchus spurensis* (UMMP 7476), and *Calyptosuchus wellsi* (UCMP 32422). *Neoaetosauroides engaeus* (PVL 3525) and *Typothorax coccinarum* (UCMP 35255) both have thin preacetabular blades and the element in *T. coccinarum* is very strongly hooked ventrally; however, in both taxa the postacetabular blades are much more elongate than in *S. robertsoni*, even extending past the anterior margin of the public peduncle.

Two ilia similar to that of *S. robertsoni* are TMM-31100-1 and UMMP 7322. TMM-31100-1, assigned to *Lucasuchus hunti* by Long & Murry (1995) has the elongate and acute angled tip of the postacetabular blade, as well as a narrower and slightly ventrally hooked preacetabular blade; however, the preacetabular blade is more anteroposteriorly elongate than that of *S. robertsoni* extending to the margin of the pubic peduncle. UMMP 7322, referred to *Desmatosuchus spurensis* by Long & Murry (1995) also has an acutely angled postacetabular blade end and a relatively narrow and slightly ventrally hooked preacetabular blade, but again the preacetabular blade is more elongate, in this case extending slightly anterior to the anterior margin of the public peduncle.

Overall the ilium most resembles that of *Aetosaurus ferratus* (Schoch, 2007) with a short, narrow and ventrally hooked preacetabular blade, but the posterior acetabular blade end is not as acutely angled in medial view.

Pubis

An odd characteristic of the pubis of *Stagonolepis robertsoni* is the presence of two large pubic foramina, the uppermost which represents the obturator foramen (Walker 1961). This was unique until the discovery of the same feature in the pubis of *Scutarx deltatylus* (Parker 2016b). The pubis is poorly known in many aetosaurs; however, *Desmatosuchus spurensis* (MNA V9300) has only a single opening.

PHYLOGENY

The most inclusive recent phylogenetic analysis of the Aetosauria is that of Parker (2016a). That study is the only analysis to include both species of *Stagonolepis* (*S. robertsoni*; “*S.* *olenkae*”), and tests the relationships of other species that have historically been considered to belong to *Stagonolepis* (*Calyptosuchus wellsi*; *Aetosauroides scagliai*) (Heckert & Lucas 1999, 2000, 2002). That analysis analyzed relationships between 26 in-group taxa utilizing 83 characters (Parker 2016a). *Stagonolepis robertsoni* is recovered in a clade (Stagonolepidinae with *Polesinosuchus aurelioi*) within Desmatosuchia and as the sister taxon to Desmatosuchinae (Figure 8). “*Stagonolepis*” *olenkae* and *Calyptosuchus wellsi* are recovered within Desmatosuchinae, whereas *Aetosauroides scagliai* is recovered outside of Stagonolepididae as in other recent studies (Desojo et al. 2012; Heckert et al. 2015).

Stagonolepis sensu Sulej (2010) is found to be paraphyletic, with *S. robertsoni* and *Polesinosuchus* forming a sister clade to “*S.* *olenkae*” and all other desmatosuchines. The previous published differences between *S. robertsoni* and “*S.* *olenkae*” all appear to be in the cranium (Sulej 2010; Antczak 2016; Parker 2016a); however, many sections of the published description of the skull of “*S.* *olenkae*” (Sulej 2010) are nearly verbatim to those published by Walker (1961) for *S. robertsoni*, so it is difficult to determine what material is actually being described in the Sulej paper.

Antczak (2016) used newly referred material from the Krasiejów quarry to hypothesize that the two taxa may be conspecific, with some of the cranial differences between *S. robertsoni* and “*S. olenkae*” representing individual variation. Further support for this hypothesis was cited by Antczak (2016) as coming from the postcranial analysis of “*S. olenkae*” by Lucas, Spielmann & Hunt (2007) who also argued that “*S. olenkae*” was a synonym of *S. robertsoni*. However, this synonymy is based on plesiomorphies and not a detailed determination of apomorphies in the material. Moreover, the findings of Antczak (2016) suggest that the skull roof (ZPAL AbIII/466/17) proposed by Sulej (2010) may not serve adequately as a holotype for *S. olenkae*, making comparisons between the two taxa problematic.

Comparison of the paramedian osteoderms of the two taxa demonstrate that although both have similar width/length ratios, and anterior bars, and radial patterning, the patterning is quite distinct with that of “*S. olenkae*” consisting of more closely packed elongate ridges and grooves (Figure 9). Nonetheless, this comparison is based on a single published osteoderm of “*S. olenkae*” (Lucas, Spielmann & Hunt 2007:fig. 4a) and more osteoderm material is needed to further support any proposed differences. Thus, a full description of the postcrania and osteoderms of “*S. olenkae*” is required to further examine potential differences between these two species.

Another recent phylogenetic analysis (Figure 10; Hoffman, Heckert & Zanno 2018) of the Aetosauria that also builds on progressive analyses from Heckert & Lucas (1999), Parker (2007), Desojo, Ezcurra & Kischlat (2012), Heckert et al (2015), and Schoch & Desojo (2016), but does not include all currently known taxa (e.g., “*Stagonolepis*” *olenkae*) also recovers *S. robertsoni* in Stagonolepidinae as the sister taxon to Desmatosuchinae (=Desmatosuchini of Parker 2016a). Interestingly with the exclusion of “*S. olenkae*” from the analysis, the sister taxon

to *S. robertsoni* is *Calyptosuchus (Stagonolepis) wellsi*. Thus presently there is strong agreement between the phylogenetic position of *S. robertsoni* in these studies as more closely related to Desmotosuchinae than to Typothoracinae (Figures 7, 9; Parker 2016a; Hoffman, Heckert & Zanno 2018).

DISCUSSION

The first formal diagnosis of *S. robertsoni* (Heckert & Lucas 2000:1556) was based on plesiomorphies that do not even comprise a unique combination of characters within Aetosauria and was intended to only diagnose the genus. This conservative assessment of the taxon has allowed other specimens (e.g., *Aetosauroides scagliai*, *Calyptosuchus wellsi*) from other areas (North and South America) to be easily assigned to the genus, mainly for the purpose of building a global terrestrial vertebrate biostratigraphy for the Upper Triassic (e.g., Lucas & Heckert 1996; Heckert & Lucas 1999, 2000, 2002; Lucas, Spielmann, & Hunt 2007; Lucas 2017). However, detailed comparison demonstrates that these taxa all bear unique combinations of characters that allow them to be differentiated (e.g., Desojo & Ezcurra 2011; Parker 2018). Assignment of specimens to genera can be subjective and based upon the taxonomic philosophy of the researcher. We have argued elsewhere (Parker 2018) that utilizing monotypic genera in aetosaurian work can aid in removing some of the ambiguity regarding use of these taxa in broader scale studies where genera are often used as a proxy for species. Diagnoses should be apomorphy based, at the species level, or provide unique combinations of characters. Thus it is important that holotype specimens be reexamined utilizing discrete apomorphies in a phylogenetic context (Nesbitt, Irmis & Parker 2007; Irmis et al. 2007; Nesbitt & Stocker 2008).

This is especially true when taxa are being utilized for intercontinental biostratigraphic correlations (Irmis et al. 2010). For another example of how taxa may be lumped based on

plesiomorphies for biostratigraphic convenience, a recent discussion on global biostratigraphy for the Late Triassic proposes a new hypothesis that *Neoaetosauroides engaeus* is a junior subjective synonym of *Aetosaurus ferratus* (Lucas 2017). Support is given based on the width/length ratios and ornamentation patterns of the paramedian osteoderms (both symplesiomorphies for Aetosauria based on their presence in the non-stagonolepidid *Aetosauroides scagliai*) as well as an interpretation of similar sutural patterns of the skull based on published figures (Lucas 2017:369). However, this interpretation ignores several key characters such as the presence of a laterally expanded premaxillary tip in *Neoaetosauroides engaeus* (Desojo & Báez 2007), which is apomorphic for Aetosauria (Parker 2016a; Hoffman, Heckert & Zanno 2018) and absent in *Aetosaurus ferratus* (Schoch 2007) as well as *Aetosauroides scagliai*, which polarizes the character distribution. Among other character differences with *A. ferratus*, *N. engaeus*, also lacks a well-developed antorbital fossa. Thus, synonymy between the two taxa is based on overall general similarity and symplesiomorphy rather than discrete synapomorphies, and currently unsupported. The practice of using plesiomorphic characters to assign specimens and to synonymize taxa has long been out of favor in vertebrate paleontology and should no longer be acceptable (e.g., Sereno 1990; Padian, Lindberg & Polly 1994; see further discussion in Nesbitt & Stocker 2008). Data that utilize this approach should be strongly suspect and not used for broader scale studies until the synapomorphy distributions among included taxa are fully evaluated and supported.

Status of the holotype of *Stagonolepis robertsoni* Agassiz, 1844

The holotype of *Stagonolepis robertsoni* is the impression of a fragment of the ventral carapace of a single specimen (ELGNM 27R) that shows several partial rows and columns of imbricated predominantly rectangular osteoderms (Figure 11; Huxley 1877:pl.1, fig. 1). These

osteoderms have an anterior bar and a surface pattern of numerous drop shaped pits radiating from the osteoderm center; hence the name *Stagonolepis*, which means “drop scale”. Since the initial descriptions by Huxley (1877) and Walker (1961) subsequent authors have assigned other species to *Stagonolepis robertsoni* based on similarities of the dorsal paramedian osteoderms (e.g., Murry & Long 1989; Long & Murry 1995; Lucas & Heckert 2001; Heckert & Lucas 2002; Lucas et al. 2007b). However, none of these authors have addressed the diagnostic status of the type specimen.

Ventral osteoderms are known from several other aetosaurian taxa including *Coahomasuchus kahleorum*, *Calypotosuchus wellsi*, *Scutarx deltatylus*, *Aetosaurus ferratus* and even the non-aetosaurian *Revueltosaurus callenderi*. However, as described above presently the ornamentation in the ventral osteoderms of the holotype specimen (ELGNM 27R) of *S. robertsoni* differs from that of other aetosaurians in that it consists of randomly arranged oblong pits separated by a latticework of thin ridges. These pits are tightly packed and cover the majority of the osteoderm surface. This differs from what is seen in other aetosaurian taxa where the ornamentation consists of narrow grooves and oblong pits radiating from a central point on the osteoderm. Presently *Stagonolepis robertsoni* is currently valid based on this character as well as the unique combination of characters listed and discussed above.

CONCLUSIONS

Stagonolepis robertsoni is the oldest named aetosaurian and as such has long served as the standard for aetosaurian osteology. Despite this status the material is difficult to work with because of its preservation and because the only sites where *S. robertsoni* fossils have been recovered are no longer active. However, the holotype specimen, although fragmentary, is

presently diagnostic and Walker's (1961) description is extremely faithful to the existing material and still serves as the basis for our understanding of this taxon.

In this contribution we redescribe and note several features that have recently become important characters for the purpose of comparing aetosaurians taxonomically and phylogenetically. *S. robertsoni* has a paramedian osteoderm morphology that bears a unique combination of characters including a raised anterior bar with anteromedial, anterior, and anterolateral processes (projections), a 'scalloped' anterior edge of the anterior bar medial to the anterior process, a short anterolateral process, a posteriorly placed, pyramidal dorsal eminence, and an anastomosing pattern of pits and grooves radiating from the eminence that lacks elongate nearly parallel grooves. Furthermore, the posterior portion of the paramedian osteoderms are devoid of ornament. *S. robertsoni* is also the only aetosaurian to preserve and extremely elongate first cervical rib, which possibly is an autapomorphy of the taxon. Posterior cervical vertebrae are keeled ventrally and bear diapophyseal and zygapophyseal laminae. The parabasisphenoid is anteroposteriorly elongate with significant separation between the basitubera and the basiptyergoid processes. The ventral osteoderms have an autapomorphic ornamentation.

These characters serve to differentiate *S. robertsoni* from all other aetosaurs include those who are or have been historically assigned to the same genus including "*Stagonolepis*" *olenkae*, *Calyptosuchus wellsi*, and *Aetosauroides scagliai*. These assignments were made based on general similarity and symplesiomorphies rather than synapomorphies, a practice that is widely discouraged in vertebrate paleontology.

INSTITUTIONAL ABBREVIATIONS

ELGNM Elgin Museum; Elgin, Scotland

MCP Museo de Ciencias e Tecnología, Porto Alegre, Brazil

- 559 **MACN** Museo Museo Argentino de Ciencias Naturales ‘Bernardino Rivadavia’, Buenos
- 560 Aires, Argentina;
- 561 **MCZD** Marischal College, Zoology Department, Aberdeen, Scotland
- 562 **MNA** Museum of Northern Arizona, Flagstaff, Arizona, USA
- 563 **NHMUK** Natural History Museum, United Kingdom
- 564 **PEFO** Petrified Forest National Park, Petrified Forest, Arizona, USA
- 565 **PVL** Paleontología de Vertebrados, Instituto “Miguel Lillo,” San Miguel de Tucumán,
- 566 Argentina
- 567 **PVSJ** División de Paleontología de Vertebrados del Museo de Ciencias Naturales y
- 568 Universidad Nacional de San Juan, San Juan, Argentina
- 569 **TMM** Texas Vertebrate Paleontology Collections, University of Texas, Austin, Texas,
- 570 USA
- 571 **UCMP** University of California Museum of Paleontology, Berkeley, CA, USA
- 572 **UMMP** University of Michigan Museum of Paleontology, Ann Arbor, Michigan, USA
- 573 **YPM** Yale Peabody Museum of Natural History, New Haven, Connecticut, USA
- 574 **ZPAL** Institute of Paleobiology, Polish Academy of Sciences, Warsaw, Poland
- 575

576 **ACKNOWLEDGEMENTS**

577 Thank you to Sandra Chapman (NHMUK), Lorna Steele (NHMUK), David Gower
578 (NHMUK), Julia Desojo (MACN), Mark Goodwin (UCMP), Kevin Padian (UCMP), the late
579 Jaime Powell (PVL); Ricardo Martinez (PVSJ), David & Janet Gillette (MNA), and Matthew
580 Smith (PEFO) for access to specimens and data under their control as well as discussions. An
581 earlier version of the manuscript was completed under the partial requirements of a dissertation
582 at the University of Texas by the senior author. This is Petrified Forest National Park

Paleontological Contribution #56. The work presented here is that of the authors and does not represent the views or opinions of the United States Government.

REFERENCES

Agassiz L. 1844. Monographie des poissons fossiles du vieux grès rouge ou Système Dévonien (Old Red Sandstone) des Isles Britanniques et de Russie. Jent et Gassman, Neuchâtel, 171 pp.

Antczak M. 2016. Late Triassic aetosaur (Archosauria) from Krasiejów (SW Poland): new species or an example of individual variation? *Geological Journal* **51**:779-788.

Benton MJ, Walker AD. 2011. *Saltopus*, a dinosauriform from the Upper Triassic of Scotland. *Earth and Environmental Transactions of the Royal Society of Edinburgh* **101**:285-299.

Bonaparte JF. 1978. El Mesozoico de America del Sur y sus tetrapodos. *Opera Lilloana* **26**:1-596.

Camp CL. 1930. A study of the phytosaurs with description of new material from Western

North America. *Memoirs of the University of California* **10**:1-174. 

Carroll RL. 1988. *Vertebrate Paleontology and Evolution*. W.H. Freeman and Company, New York, 698 pp.

Case EC. 1922. New reptiles and stegocephalians from the Upper Triassic of western Texas. *Carnegie Institution of Washington Publication* **321**:1-84.

Cope ED. 1869. Synopsis of the extinct Batrachia, Reptilia, and Aves of North America. *Transactions of the American Philosophical Society* **14**:1-252.

Desojo JB, Báez AM. 2007. Cranial morphology of the Late Triassic South American archosaur *Neoaeosauroides engaeus*: evidence for aetosaurian diversity. *Palaeontology* **50**:267-276.

- Desojo JB, Ezcurra MD. 2011.** A reappraisal of the taxonomic status of *Aetosauroides* (Archosauria, Aetosauria) specimens from the Late Triassic of South America and their proposed synonymy with *Stagonolepis*. *Journal of Vertebrate Paleontology* **31**:596-609.
- Desojo JB, Ezcurra MD, Kischlat EE. 2012.** A new aetosaur genus (Archosauria: Pseudosuchia) from the early Late Triassic of southern Brazil. *Zootaxa* **3166**:1-33.
- Desojo JB, Heckert AB, Martz JW, Parker WG, Schoch RR, Small BJ, Sulej T. 2013.** Aetosauria: a clade of armored pseudosuchians from the Upper Triassic continental beds. In: Nesbitt SJ, Desojo J B, Irmis RB, eds. *Anatomy, Phylogeny, and Palaeobiology of Early Archosaurs and their Kin*. Geological Society, London, Special Publications 379. The Geological Society Publishing House, Bath, 203-239.
- Fraser NC. 1988.** Latest Triassic vertebrates and their biostratigraphy. *Modern Geology* **13**:125-140.
- Gauthier JA, Padian K. 1985.** Phylogenetic, functional, and aerodynamic analyses of the origin of birds and their flight. In: Hecht MK, Ostrom JH, ViohlG, Wellnhofer P. eds. *The Beginning of Birds: Proceedings of the International Archaeopteryx Conference, Eichstätt, 1984*. Freunde des Jura-Museums Eichstätt, Eichstätt, 185-197.
- Gower DJ, Walker AD. 2002.** New data on the braincase of the aetosaurian archosaur (Reptilia: Diapsida) *Stagonolepis robertsoni* Agassiz; pp. 7-23 in Norman D. B. and D. J. Gower (eds.), *Archosaurian Anatomy and Paleontology. Essays in Memory of Alick D. Walker*. *Zoological Journal of the Linnaean Society* **136**. The Linnaean Society of London, London.
- Heckert AB, Lucas SG. 1999.** A new aetosaur (Reptilia: Archosauria) from the Upper Triassic of Texas and the phylogeny of aetosaurs. *Journal of Vertebrate Paleontology* **19**:50-68.

- 630 **Heckert AB, Lucas SG. 2000.** Taxonomy, phylogeny, biostratigraphy, biochronology,
631 paleobiogeography, and evolution of the Late Triassic Aetosauria (Archosauria:
632 Crurotarsi). *Zentralblatt für Geologie und Paläontologie Teil I 1998 Heft 11–12*:1539-
633 1587.
- 634 **Heckert AB, Lucas SG. 2002.** South American occurrences of the Adamanian (Late Triassic:
635 latest Carnian) index taxon *Stagonolepis* (Archosauria: Aetosauria) and their
636 biochronological significance. *Journal of Paleontology* **76**:852-863.
- 637 **Heckert AB, Hunt AP, Lucas SG. 1996.** Redescription of *Redondasuchus reseri*, a Late
638 Triassic aetosaur (Reptilia: Archosauria) from New Mexico (U.S.A.), and the
639 biochronology and phylogeny of aetosaurs. *Geobios* **29**:619-632.
- 640 **Heckert AB, Schneider VP, Fraser NC, Webb RA. 2015.** A new aetosaur (Archosauria:
641 Suchia) from the Upper Triassic Pekin Formation, Deep River Basin, North Carolina, U.
642 S. A., and its implications for early aetosaur evolution. *Journal of Vertebrate*
643 *Paleontology* **35**:e881831.
- 644 **Heckert AB, Fraser NC, Schneider VP. 2017.** A new species of *Coahomasuchus* (Archosauria,
645 Aetosauria) from the Upper Triassic Pekin Formation, Deep River Basin, North Carolina.
646 *Journal of Paleontology* **91**:162-178.
- 647 **Huene F. 1902.** Übersicht über die reptilien der Trias. *Geologische und Palaeontologische*
648 *Abhandlungen n. F. 6*:1-84.
- 649 **Huene F. 1908.** Die dinosaurier der europäischen Triasformation mit Berücksichtigung der
650 aussereuropäischen Vorkommnisse. *Geologische und Palaeontologische Abhandlungen*
651 *Suppl-Bd. 1*:322-351.
- 652 **Huene F. 1936.** The constitution of the Thecodontia. *American Journal of Science* **32**:207-217.

- 653 **Huene F.1942.** *Die fossilen Reptilien des südamerikanischen Gonwanalandes. Ergebnisse der*
654 *Sauriergrabung in Südbrasilien 1928/1929.* C.H. Beck, München.
- 655 **Hillis DM, Bull JJ. 1993.** An empirical test of bootstrapping as a method for accessing
656 confidence in phylogenetic analysis. *Systematic Biology* **42**:182-192.
- 657 **Huxley TH. 1859.** On the *Stagonolepis robertsoni* (Agassiz) of the Elgin Sandstone; and on the
658 recently discovered footmarks in the Sandstones of Cummingstone. *Proceedings of the*
659 *Geological Society* **15**:440-460.
- 660 **Huxley TH. 1869.** On *Hyperodapedon*. *Quarterly Journal of the Geological Society, London*
661 **25**:138-152.
- 662 **Huxley TH. 1875.** On *Stagonolepis robertsoni*, and on the evolution of the crocodylia. *Quarterly*
663 *Journal of the Geological Society, London* **31**:423-438.
- 664 **Huxley TH. 1877.** The crocodilian remains found in the Elgin sandstones, with remarks on the
665 ichnites of Cummingstone. *Memoirs of the Geological Survey of the United Kingdom*
666 **3**:1-58.
- 667 **Irmis RB, Parker WG, Nesbitt SJ, Liu J. 2007.** Early ornithischian dinosaurs: the Triassic
668 record. *Historical Biology* **19**:3-22.
- 669 **Irmis RB, Martz JW, Parker WG, Nesbitt SJ. 2010.** Re-evaluating the correlation between
670 Late Triassic terrestrial vertebrate biostratigraphy and the GSSP-defined marine stages.
671 *Albertiana* **38**:40-52.
- 672 **Krebs B. 1976.** Pseudosuchia. In: Kuhn O, ed. *Handbuch der Paläoherpetologie, Volume 13.*
673 Gustav Fischer Verlag, Stuttgart, 40-98.
- 674 **Long RA, Ballew KL. 1985.** Aetosaur dermal armor from the late Triassic of southwestern
675 North America, with special reference to material from the Chinle Formation of Petrified

- Forest National Park. In: Colbert EH, Johnson RR, eds. *The Petrified Forest Through the*
Ages, 75th Anniversary Symposium November 7, 1981. Museum of Northern Arizona
Bulletin **54**. Museum of Northern Arizona Press, Flagstaff, 45-68.
- Long RA, Murry PA. 1995.** Late Triassic (Carnian and Norian) tetrapods from the southwestern
United States. *New Mexico Museum of Natural History and Science Bulletin* **4**:1–254.
- Lucas SG. 2017.** Late Triassic terrestrial tetrapods: biostratigraphy, biochronology, and biotic
events. In: Tanner LH, ed., *The Late Triassic World, Topics in Geobiology* **46**, Springer
International Publishing AG, Cham, Switzerland, 351-405.
- Lucas SG, Heckert AB. 1996.** Late Triassic aetosaur biochronology. *Albertiana* **17**:57-64.
- Lucas SG, Heckert AB. 2001.** The aetosaur *Stagonolepis* from the Upper Triassic of Brazil and
its biochronological significance. *Neues Jahrbuch für Geologie und Paläontologie,*
Monatshefte **2001**:719–732.
- Lucas SG, Spielmann JA, Hunt AP. 2007.** Biochronological significance of Late Triassic
tetrapods from Krasiejów, Poland. In: Lucas SG, Spielmann JA, eds., *The Global*
Triassic. New Mexico Museum of Natural History and Science Bulletin **41**. New Mexico
Museum of Natural History and Science, Albuquerque, 248-258.
- Lucas SG, Hunt AP, Heckert AB, Spielmann, JA. 2007b.** Global Triassic tetrapod
biostratigraphy and biochronology: 2007 status. In: Lucas SG, Spielmann JA, eds. *The*
Global Triassic. New Mexico Museum of Natural History and Science Bulletin **41**. New
Mexico Museum of Natural History and Science, Albuquerque, 229-240.
- Lydekker R. 1887.** The fossil Vertebrata of India. *Records of the Geological Survey of India*
20:51–80.
- Marsh OC. 1884.** The classification and affinities of dinosaurian reptiles. *Nature* **31**:68–69.

- Martz JW, Small BJ. 2006.** *Tecovasuchus chatterjeei*, a new aetosaur (Archosauria: Aetosauria) from the Tecovas Formation (Upper Triassic, Carnian) of Texas. *Journal of Vertebrate Paleontology* **26**:308–320.
- Murry PA, Long RA. 1989.** Geology and paleontology of the Chinle Formation, Petrified Forest National Park and vicinity, Arizona and a discussion of vertebrate fossils of the southwestern Upper Triassic. In: Lucas SG and Hunt AP, eds. *Dawn of the Age of Dinosaurs in the American Southwest*. New Mexico Museum of Natural History, Albuquerque, 29–64.
- Nesbitt SJ, Stocker MR. 2008.** The vertebrate assemblage of the Late Triassic Canjilon Quarry (northern New Mexico, U.S.A.), and the importance of apomorphy-based assemblage comparisons. *Journal of Vertebrate Paleontology* **28**:1063–72.
- Nesbitt SJ, Irmis RB, Parker WG. 2007.** A critical re-evaluation of the Late Triassic dinosaur taxa of North America. *Journal of Systematic Palaeontology* **5**:209–243.
- Padian K, Lindberg DR, Polly PD. 1994.** Cladistics and the fossil record: the uses of history. *Annual Review of Earth and Planetary Science*, **22**, 63-91.
- Parker WG. 2007.** Reassessment of the aetosaur “*Desmatosuchus*” *chamaensis* with a reanalysis of the phylogeny of the Aetosauria (Archosauria: Pseudosuchia). *Journal of Systematic Palaeontology* **5**:1–28.
- Parker WG. 2008.** Description of new material of the aetosaur *Desmatosuchus spurensis* (Archosauria: Suchia) from the Chinle Formation of Arizona and a revision of the genus *Desmatosuchus*. *PaleoBios* **28**:1–40.

- Parker WG. 2016a.** Revised phylogenetic analysis of the Aetosauria (Archosauria: Pseudosuchia); assessing the effects of incongruent morphological character sets. *PeerJ* **4**:e1583; DOI 10.7717/peerj.1583.
- Parker WG. 2016b.** Osteology of the Late Triassic aetosaur *Scutarx deltatylus* (Archosauria: Pseudosuchia). *PeerJ* **4**:e2411; DOI 10.7717/peerj.2411.
- Parker WG. 2018.** Redescription of *Calypotosuchus (Stagonolepis) wellsi* (Archosauria: Pseudosuchia: Aetosauria) from the Late Triassic of the southwestern United States with a discussion of genera in vertebrate paleontology. *PeerJ* **6**:e4291; DOI 10.7717/peerj.4291.
- Parrish JM. 1986.** Locomotor adaptations in the hind limb and pelvis of the Thecodontia. *Hunteria* **1**:3-35.
- Parrish JM. 1994.** Cranial osteology of *Longosuchus meadei* and the phylogeny and distribution of the Aetosauria. *Journal of Vertebrate Paleontology* **14**:196-209.
- Schoch RR. 2007.** Osteology of the small archosaur *Aetosaurus* from the Upper Triassic of Germany. *Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen* **246**:1–35.
- Schoch RR, Desojo JB. 2016.** Cranial anatomy of the aetosaur *Paratypothorax andressorum* Long & Ballew, 1985, from the Upper Triassic of Germany and its bearing on aetosaur phylogeny. *Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen* **279**:73–95.
- Sereno PC. 1990.** Grades and clades in dinosaur systematics. In: Carpenter K, Currie PJ, eds., *Dinosaur Systematics: approaches and perspectives*. Cambridge University Press, Cambridge, United Kingdom, 9-20.

- Sereno PC. 1991.** Basal archosaurs: phylogenetic relationships and functional implications. *Society of Vertebrate Paleontology Memoir* 2:1-53.
- Small BJ. 2002.** Cranial anatomy of *Desmatosuchus haplocerus* (Reptilia: Archosauria: Stagonolepididae). *Zoological Journal of the Linnean Society* **136**:97-111.
- Small BJ, Martz JW. 2013.** A new aetosaur from the Upper Triassic Chinle Formation of the Eagle Basin, Colorado, USA. In: Nesbitt SJ, Desojo JB, Irmis RB, eds., *Anatomy, Phylogeny and Palaeobiology of Early Archosaurs and their Kin. Geological Society, London, Special Publications* **379**: The Geological Society Publishing House, Bath, 393-412
- Sulej T. 2010.** The skull of an early Late Triassic aetosaur and the evolution of the stagonolepidid archosaurian reptiles. *Zoological Journal of the Linnean Society* **158**:860-881.
- Walker AD. 1961.** Triassic Reptiles from the Elgin Area: *Stagonolepis*, *Dasygnathus*, and their allies. *Philosophical Transactions of the Royal Society of London* **244**:103–204.
- Wilson JA. 1999.** Vertebral laminae in sauropods and other saurischian dinosaurs. *Journal of Vertebrate Paleontology* **19**:639-653.
- Zittel KAv. 1887–1890.** *Handbuch der Palaeontologie. 1. Abteilung: Paläozoologie, Band 3, Vertebrata (Pisces, Amphibia, Reptilia, Aves)*. Druck and Verlag von R. Oldenbourg, München and Leipzig, 900 pp.

FIGURES

Figure 1. Casts of bones of *Stagonolepis robertsoni*. A) NHMUK R4787a, 1885 cast which represents much of the lower portion of a skull. B) NHMUK R4787, PVC cast of the same

specimen that shows more details of the braincase and dorsal portion of the skull. Scale bars equal 1 cm. Abbreviations: **an**, angular; **art**, articular; **bpt**, basipterygoid process; **bs**, parabasisphenoid; **bt**, basal tuber; **d**, dentary; **fr**, frontals; **l**, left; **mx**, maxilla; **mpr**, area of the medial pharyngeal recess; **pa**, parietal; **pmx**, premaxilla; **pra**, prearticular; **pro**, prootic; **pt**, pterygoid; **qu**, quadrate; **r**, right; **sa**, surangular. Elements from the left side have a prefix of **l**.; elements of the right side have a prefix of **r**.

Figure 2. NHMUK R4787, cast of a right maxilla of *Stagonolepis robertsoni* in lateral A) and medial B) views. Scale bar equals 1 cm. Abbreviations: **a**., articulation with listed element; **alv**, alveoli; **aof**, antorbital fenestra; **ap.m**, ascending process of the maxilla; **ju**, jugal; **la**, lacrimal; **ms**, medial shelf; **na**, nasal; **pac**, pneumatic accessory cavity; **pmx**, premaxilla; **tr**, transverse ridge.

Figure 3. Cast of the anterior part of the right side of the skull of *Stagonolepis robertsoni* in lateral view. Scale bar equals 1 cm. Abbreviations: **en**, external naris; **mx**, maxilla; **na**, nasal; **nr**, nasal ridge; **pmr**, premaxillary ridge; **pmx**, premaxilla.

Figure 4. NHMUK 4784a, casts of articulated sections of the presacral vertebral column in lateral view including the rear of the skull through the fourth cervical position (A), and the seventh through the twelfth presacral vertebrae (B). Scale bars equal 1 cm. Abbreviations: **cvr**, cervical rib; **k**, ventral keel; **oc**, occipital condyle of the basicranium; **parp**, parapophysis; **pcdl**, posterior centrodiapophyseal lamina.

Figure 5. Casts of osteoderms of *Stagonolepis robertsoni*. A, left dorsal trunk paramedian (NHMUK R4790a) in dorsal view; B, right dorsal trunk lateral (NHMUK R4789a) in dorsolateral view. Scale bar equals 1 cm. Abbreviations: **a.**, articulation with listed element; **ab**, anterior bar; **alp**, anterolateral projection; **amp**, anteromedial projection; **de**, dorsal eminence; **df**, dorsal flange; **lf**, lateral flange; **sc**, scalloping of anterior bar margin.

Figure 6. Ventral osteoderms of *Stagonolepis robertsoni*. A) close-up of NHMUK R27404 (positive cast of EM 27R, the holotype specimen of *Stagonolepis robertsoni* Agassiz 1844) in ventrolateral view showing the detail of the overlapping osteoderms; B) NHMUK R4787a, referred ventral osteoderm in ventral view.

Figure 7. NHMUK R4789. Cast of medial side of right ilium in medial view. Scale bar = 2 cm. Abbreviations: **arsr2**, articulation for sacral rib 2; **ip**, ischiadic peduncle; **ost**, osteoderm; **poab**, postacetabular blade; **pp**, pubic peduncle; **prab**, preacetabular blade.

Figure 8. Phylogenetic analysis of the Aetosauria. The reduced strict consensus tree from Parker 2016a, with all named clades. Decay indices and bootstrap values are shown for all nodes, with bootstrap values under 70% (the confidence threshold of Hillis & Bull, 1993) shown in red.

Figure 9. Dorsal trunk paramedian osteoderms of *Stagonolepis*. A) NHMUK R4787a, left dorsal trunk paramedian in dorsal view; B) PAN ZPAL AbIII 57011, left dorsal trunk paramedian in dorsal view. Scale bars = 1 cm.

811 Figure 10. Alternate phylogenetic hypothesis of the Aetosauria of Hoffman, Heckert, and Zanno
812 (2018) with clade names and decay indices for each node.

813

814 Figure 11. Cast of EM 27R, the holotype specimen of *Stagonolepis robertsoni* Agassiz
815 1844. A series of imbricated osteoderms from the ventral trunk region.

816

Figure 1

Casts of bones of *Stagonolepis robertsoni*.

(A) NHMUK R4787a, 1885 cast which represents much of the lower portion of a skull. (B) NHMUK R4787, PVC cast of the same specimen that shows more details of the braincase and dorsal portion of the skull. Scale bars equal 1 cm. Abbreviations: **an**, angular; **art**, articular; **bpt**, basipterygoid process; **bs**, parabasisphenoid; **bt**, basal tuber; **d**, dentary; **fr**, frontals; **l**, left; **mx**, maxilla; **mpr**, area of the medial pharyngeal recess; **pa**, parietal; **pmx**, premaxilla; **pra**, prearticular; **pro**, prootic; **pt**, pterygoid; **qu**, quadrate; **r**, right; **sa**, surangular. Elements from the left side have a prefix of **l.**; elements of the right side have a prefix of **r.**



Figure 2

NHMUK R4787, cast of a right maxilla of *Stagonolepis robertsoni*

Lateral (A) and medial (B) views. Scale bar equals 1 cm. Abbreviations: **a.**, articulation with listed element; **alv**, alveoli; **aof**, antorbital fenestra; **ap.m**, ascending process of the maxilla; **ju**, jugal; **la**, lacrimal; **ms**, medial shelf; **na**, nasal; **pac**, pneumatic accessory cavity; **pmx**, premaxilla; **tr**, transverse ridge.




Figure 3

Skull of *Stagonolepis robertsoni*. 

Cast of the anterior part of the right side of the skull of *Stagonolepis robertsoni* in lateral view. Scale bar equals 1 cm. Abbreviations: **en**, external naris; **mx**, maxilla; **na**, nasal; **nr**, nasal ridge; **pmr**, premaxillary ridge; **pmx**, premaxilla.



Figure 4

Presacral vertebrae of *Stagonolepis robertsoni*.

NHMUK 4784a, casts of articulated sections of the presacral vertebral column in lateral view including the rear of the skull through the fourth cervical position (A), and the seventh through the twelfth presacral vertebrae (B). Scale bars equal 1 cm. Abbreviations: **cvr**, cervical rib; **k**, ventral keel; **oc**, occipital condyle of the basicranium; **parp**, parapophysis; **pcdl**, posterior centrodiapophyseal lamina.





Figure 5

Casts of osteoderms of *Stagonolepis robertsoni*.

(A) left dorsal trunk paramedian (NHMUK R4790a) in dorsal view; (B) right dorsal trunk lateral (NHMUK R4789a) in dorsolateral view. Scale bar equals 1 cm. Abbreviations: **a.**, articulation with listed element; **ab**, anterior bar; **alp**, anterolateral projection; **amp**, anteromedial projection; **de**, dorsal eminence; **df**, dorsal flange; **lf**, lateral flange; **sc**, scalloping of anterior bar margin.





Figure 6

Ventral osteoderms of *Stagonolepis robertsoni*.

(A) close-up of NHMUK R27404 (positive cast of EM 27R, the holotype specimen of *Stagonolepis robertsoni* Agassiz 1844) in ventrolateral view showing the detail of the overlapping osteoderms; (B) NHMUK R4787a, referred ventral osteoderm in ventral view. Scale bar = 1 cm.




Figure 7

Ilium of *Stagonolepis robertsoni*

NHMUK R4789. Cast of medial side of right ilium in medial view. Scale bar = 2 cm.

Abbreviations: **arsr2**, articulation for sacral rib 2; **ip**, ischiadic peduncle; **ost**, osteoderm; **poab**, postacetabular blade; **pp**, pubic peduncle; **prab**, preacetabular blade.




Figure 8

Phylogenetic Analysis of the Aetosauria.

The reduced strict consensus tree from Parker (2016a), with all named clades. Decay indices and bootstrap values are shown for all nodes, with bootstrap values under 70% (the confidence threshold of Hillis & Bull, 1993) shown in red.




Figure 9

Dorsal trunk paramedian osteoderms of *Stagonolepis*.

(A) NHMUK R4787a, left dorsal trunk paramedian in dorsal view; (B) PAN ZPAL AbIII 57011, left dorsal trunk paramedian in dorsal view. Scale bars = 1 cm.



Figure 10

Alternate phylogenetic analysis of the Aetosauria.

Alternate phylogenetic hypothesis of the Aetosauria of Hoffman, Heckert, and Zanno (2018) with clade names and decay indices for each node.




Figure 11

Holotype specimen of *Stagonolepis robertsoni*.

Cast of EM 27R, the holotype specimen of *Stagonolepis robertsoni* Agassiz 1844, a series of imbricated osteoderms from the ventral trunk region.

