

1 **High definition video loggers provide new insights into**
2 **behaviour, physiology, and the oceanic habitat of a marine**
3 **predator, the yellow-eyed penguin**

4
5 Thomas Mattern^{1,2}, Mike McPherson³, Ursula Ellenberg^{4,2}, Yolanda van
6 Heezik¹, Philip J. Seddon¹

7 ¹ Department of Zoology, PO Box 56, University of Otago, Dunedin, NZ

8 ² Global Penguin Society, Marcos Zar 2716, Puerto Madryn (9120), Chubut, Argentina.

9 ³ CTNova Ltd, Leighton Buzzard, Bedfordshire, England

10 ⁴ Department of Ecology, Environment and Evolution, La Trobe University, Melbourne, Australia

11 Corresponding author: Thomas Mattern¹

12 Email address: t.mattern@eudypates.net

Abstract

Camera loggers are increasingly used to examine behavioural aspects of free-ranging animals. However, often video loggers are deployed with a focus on specific behavioural traits utilizing small cameras with a limited field of view, poor light performance and video quality. Yet rapid developments in consumer electronics provide new devices with much improved visual data allowing a wider scope for studies employing this novel methodology. We developed a camera logger that records full high-definition (HD) video through a wide-angle lens, providing high resolution footage with a greater field of view than other camera loggers. The main goal was to assess the suitability of this type of camera for the analysis of various aspects of the foraging ecology of a marine predator, the yellow-eyed penguin (*Megadyptes antipodes*) in New Zealand. Frame-by-frame analysis allowed accurate timing of prey pursuits and time spent over certain seafloor types. The recorded video footage showed that prey species were associated with certain seafloor types, revealed different predator evasion strategies by benthic fishes, and highlighted varying energetic consequences for penguins pursuing certain types of prey. Other aspects that could be analysed were the timing of breathing intervals between dives and observed exhalation events during prey pursuits, a previously undescribed behaviour. Screen overlays facilitated analysis of flipper angles and beat frequencies throughout various stages of the dive cycle. Flipper movement analysis confirmed decreasing effort during descent phases as the bird gained depth, and that ascent was principally passive. Breathing episodes between dives were short (<1 s) while the majority of the time was devoted to subsurface scanning with a submerged head. Video data recorded on free-ranging animals not only provide a wealth of information recorded from a single deployment but also necessitate new approaches with regards to analysis of visual data. Here, we demonstrate the diversity of information that can be gleaned from video logger data, if devices with high video resolution and wide field of view are utilized.

Introduction

Examining the at-sea behaviour of marine animals has long been a challenging endeavour. Direct visual observations of behaviour are almost impossible, especially when most of it happens under the ocean's surface. In recent decades, advances in telemetry technologies and the emergence of bio-logging hardware have provided the means to track marine animals and reveal their foraging behaviour in great detail. Starting in the 1970s with rather crude location estimates and limited data quality recorded by unwieldy devices that could only be used on large animals, advancements in micro-electronics have resulted in ever smaller and more accurate loggers to pinpoint an animal's position to within a few metres and record their diving depths with oceanography-grade precision (Wilmers et al., 2015). New technologies such as accelerometers and gyroscopes further refined methods to study marine habitat use (e.g. Noda et al. 2014). Yet, placing dive metrics into a complex behavioural and environmental context can be difficult; ideally a reference framework based on direct observations is used to match up dive metrics and actual behaviours (e.g. Moreau et al. 2009; Volpov et al. 2016). So, the original dilemma of having to make direct observations of marine animal behaviours still persists. Animal-borne video recorders offer the means to overcome this problem.

In recent years animal-borne camera systems have made it possible to log in situ observations of behaviour from the animal's point of view (Moll et al., 2007). For example, deployment of light-weight video cameras on flying birds provided new perspectives on prey pursuit in falcons (Kane & Zamani, 2014) and revealed how albatrosses use the presence of killer whales to locate prey (Sakamoto et al., 2009). No other animal group has been more subject to deployment of video recording devices in recent years than marine animals. By overcoming the observational barrier at sea, video loggers are providing copious amounts of novel data that range from identification of feeding strategies (Ponganis et al., 2000; Takahashi et al., 2008) and previously unknown food sources (Thiebot et al., 2017), to social interactions such as group foraging (Sutton, Hoskins & Arnould, 2015) or kleptoparasitism (Handley & Pistorius, 2015). Video data

also offers the means to calibrate other bio-logging data (Watanabe & Takahashi, 2013; Gómez-Laich et al., 2015).

What most of these studies have in common is their focus on specific behavioural traits while providing limited information about the environment the behaviours occurred in. This is principally due to limitations of the video hardware used, which has to be small and light-weight so as to not overly impede the study animal's movement capabilities (Ludynia et al., 2012) and hence behaviour. As a result, video quality (i.e. image resolution and field of view/FOV) is sacrificed in favour of smaller cameras (e.g. Watanabe & Takahashi, 2013; Gómez-Laich et al., 2015; Thiebot et al., 2016, 2017). However, with the rise in popularity of action cams on the consumer market, new video devices have recently become available with high definition video capabilities and wide-angle optics, suitable for deployment even on smaller marine animals such as penguins. This leap in quality has significant implications for the study of marine animals as it not only allows more accurate monitoring of a wide-range of aspects of behaviours such as specific pursuit strategies and capture efficiency, as well as prey identification and interactions with other species, but also provides new opportunities for the visual analysis of the environment the animals use. This is particularly relevant in species that forage at the seafloor where video data can provide extensive information about the benthic habitat (Watanuki et al., 2008).

The ~~Y~~yellow-eyed penguin (*Megadyptes antipodes*) in New Zealand is known to be a benthic forager (Mattern et al., 2007) that feeds primarily on demersal fish species (van Heezik & Davis, 1990; Moore et al., 1995). It has been suggested that this strategy might come at the expense of reduced behavioural flexibility, with subsequent vulnerability to changes in the marine environment (Mattern et al., 2007). In particular, degradation of seafloor ecosystems in the wake of commercial bottom fisheries are suspected to influence ~~Y~~yellow-eyed penguin foraging success and population developments (Browne et al., 2011; Mattern et al., 2013). While the species' at-sea movement and diving behaviour has ~~yes~~ been subject to a number of studies in the

90 past decades (Moore et al., 1995; Mattern et al., 2007, 2013), information about their benthic
91 habitat is very limited.

92 To assess the extent to which penguin behaviour and foraging success correlates with the
93 composition of the benthic habitat, we developed a camera logger that records full high-
94 definition (HD) videos through wide-angle lenses. The main focus of our study was to assess the
95 suitability of the device for the visual analysis of penguin prey pursuit behaviour and
96 characteristics of the benthic ecosystem. However, the deployment revealed far more
97 information than was anticipated. The video data provided novel insights into physiological
98 aspects of the penguin's diving activities and allowed us to draw conclusions about prey capture
99 techniques. In this paper, we summarise our findings, demonstrate analytical approaches to
100 evaluate animal-borne video data, and highlight the multi-disciplinary potential of wide-angle,
101 full HD video loggers.

102 **Materials and methods**

103 **Study site and species**

104 The ~~Y~~yellow-eyed penguin, classified as "Endangered" by the IUCN Redlist (BirdLife
105 International, 2012), is one of five penguin species endemic to the New Zealand region and
106 occurs on the sub-Antarctic Auckland and Campbell Islands as well as the south-eastern
107 coastlines of New Zealand's South Island and Stewart Island (Seddon, Ellenberg & van Heezik,
108 2013). This study was carried out at the Boulder Beach complex, Otago Peninsula, South Island,
109 New Zealand (45.90°S, 170.56°E). Penguins from this site have been subject to foraging studies
110 that have suggested substantial impact of bottom trawling activities on the ~~Y~~yellow-eyed
111 penguins' at-sea movements (Ellenberg & Mattern, 2012; Mattern et al., 2013).

112 **Video logger & deployment**

113 We developed a high-definition video logger (dimensions LxWxH, 89x41x21mm; weight: 78 g)
114 which is combined with a time-depth recorder (TDR, 31x12x11mm, 6.5g ; AXY-depth,
115 Technosmart Ltd. Italy) and a GPS logger (modified, epoxy encased i-gotU, GT-120, Mobile

116 Action Technology Inc., Taiwan, 31x22x11mm, 12 g). The latter two devices were combined into
 117 a single unit by gluing the AXY-depth to the longer side of the GPS device. Camera and logger
 118 combination were then attached individually in line to the lower back of the penguin using
 119 adhesive tape (Wilson et al., 1997). Additional drag of the devices was principally limited to the
 120 cameras frontal area (Bannasch, Wilson & Culik, 1994).

121 The camera logger consisted of a modified Mobius action-cam with a 130° wide-angle lens
 122 (www.mobius-actioncam.com). To achieve the smallest and lightest device possible, the camera
 123 electronics, video sensor and lens were removed from the casing and the battery replaced with a
 124 1200 mAh Lithium Polymer battery to extend recording time. A small bespoke timer board was
 125 developed to allow the camera to be fired at a pre-determined time. Connections were provided
 126 to allow programming logger start time and also to access the camera's USB port for managing
 127 camera setting, extracting the video data and recharging the battery. The board was isolated
 128 electrically to prevent the contacts from shorting as sea-water is conductive. Activation of the
 129 interface was achieved using a Hall-effect device. An Arduino-based interface was developed to
 130 allow the current date/time and logger start time to be set. The camera was programmed to
 131 record video data at a resolution of 1920x1080 pixels (1080p) at a frame rate of 30 frames per
 132 second. Video data were recorded in H.264 MPEG4 format and stored on a 32GB MicroSD card.
 133 The camera was programmed to start recording at 11 am the following day when it was
 134 assumed that the penguin had completed its travel phase and arrived at its foraging destination.
 135 The camera operated from the programmed start time until the battery fell below the minimum
 136 operating voltage of the camera (ca. 2-4 hours). The device was recovered when the penguin
 137 returned from its foraging trip; data were then downloaded through the camera's USB interface.

138 Since the logger stores video data as a series of full frame images ('progressive scan'), it was
 139 possible to conduct a frame-by-frame analysis to accurately time components of the bird's
 140 behaviour – i.e. breathing intervals, flipper beat frequencies and amplitudes – as well as time
 141 spent over certain benthic habitats. Video analysis was conducted in professional editing
 142 software (Adobe Premiere Pro CS 6, Adobe Systems Inc., San Jose, CA, USA) which allows the

143 quick and precise backward and forward navigation of the video material using the keyboard
144 (“scrubbing”) and provides the option to display frame number in the preview timer.

145 The video logger was deployed on a breeding male ~~Y~~yellow-eyed penguin tending two chicks on
146 17 December 2015. Deployment occurred at the penguin’s nest on the evening of 17 December.
147 The bird was removed from the nest and placed in a cloth bag to reduce stress. The
148 instrumentation procedure lasted around 20 minutes after which the penguin was released back
149 on its nest. The bird left on a single foraging trip on 18 December before the device was
150 recovered on 19 December; the penguin continued to breed normally after the deployment.

151 **Failure to record GPS data**

152 Upon device recovery it became apparent that the GPS logger did not record any data after the
153 camera had started operating. It has since become evident, that the Mobius action-cam
154 generates significant electromagnetic interference which prevented the GPS logger from
155 functioning properly. This can be rectified by wrapping the camera with electrical shielding
156 tape; however, in our case the lack of shielding resulted in failure to record GPS data.

157 **Analysis of behaviours & habitat**

158 For detailed analysis of behaviours, we randomly selected 12 **dive cycles** from the 46 dive cycles
159 recorded (i.e. one fourth of all dives) independent from prevalent behaviours exhibited during
160 these dives. This was due to the labour-intensive frame-by-frame analysis necessary for several
161 of the behaviours. Future analyses are ideally conducted automatically using machine learning
162 algorithms to reduce analysis time and increase accuracy (e.g. Valletta et al., 2017).

163 **Prey pursuits & capture.** We defined the beginning of a prey pursuit as the moment when the
164 penguin markedly accelerated while swimming along the seafloor; the end was reached when
165 the penguin decelerated again to its previous cruise speed (if no prey was caught), or when the
166 prey item was swallowed completely. Acceleration and deceleration were associated with
167 temporary blurring of the video footage due to irregular body movement, allowing for exact

Commented [A1]: Define dive cycle. You didn't answer the reviewer's question about this.

168 timing of prey pursuits. Where possible, prey species were identified from frames providing a
169 clear view of the prey item.

170 **Benthic habitat**

171 For all dives, the benthic habitat was classified according to sediment type (fine sand, coarse
172 sand with shell fish fragments, gravel), sediment structure (flat, sediment ripples) and
173 composition of the epibenthic communities. For the latter, we used a presence/absence
174 approach with easy-to-identify epibenthic species brittlestars (*Ophiuroidea*), anthozoans
175 (anemones and soft corals), and horse mussels (*Atrina zelandica*), within a 30-frame time
176 window. Figure 1 provides a photographic overview of the different habitat characteristics used.
177 Future deployments with a functional GPS logger can be used for more elaborate analysis of the
178 benthic habitat, e.g. the creation of biodiversity indices.

179 Beyond prey and habitat interactions, the video data offered the opportunities to analyse
180 various physiological aspects of the penguin's behaviour.

181 **Flipper movements.** During dives, flipper beat frequencies (beats per minute, BPM) were
182 determined by counting the number of frames required to complete one flipper beat cycle,
183 beginning the count when the flipper angle reached its maximum upward inclination and ending
184 with the frame prior to the subsequent maximum upward inclination. In the video editing
185 software, we overlaid a template indicating 10, 30, 50, 70 and 90-degree angles radiating from
186 the base of the flippers on the video data (<https://vimeo.com/179414575>). This allowed us to
187 visually determine maximum amplitude of each flipper beat to the nearest 5°.

188 **Surface breathing & underwater exhalation.** We timed breathing events when the penguin
189 was at the surface following a dive. Noting frame numbers when the bird raised its head out of
190 the water before lowering below the surface again made it possible to determine the times the
191 penguin was able to respire (<https://vimeo.com/179414575#t=145>). Additional observations
192 of exhalations during the dive were noted.

193 A selection of edited video clips demonstrating the various behaviours and habitat types
194 described above can be accessed via <https://vimeo.com/album/4103142>.

195 **Dive data analysis**

196 Dive data recorded by the TDR at 1 s intervals and depth resolution of ~0.1m were analysed
197 following methods described in detail in Mattern et al. (2007). Dives were classified as pelagic or
198 benthic dives using dive profile characteristics, where near horizontal bottom phases with little
199 vertical variance as well as consistent maximum dive depths on consecutive dives were used as
200 cues for diving along the seafloor. This approach was validated by recorded video data. The TDR
201 also recorded tri-axial accelerometer data which has yet to be analysed.

202 Statistical analysis was carried out in R 3.4.2 (R Core Team, 2014). Correlations were examined
203 as linear models (Pearson's correlation). Comparisons were conducted as simple t-tests
204 accounting for unequal variances (Welch's t-Test, Ruxton, 2006).

205 **Permits**

206 This study was approved by the Animal Ethics Committee of the University of Otago (UOO AEC
207 69/15) and field experiments conducted under research permits issued by the New Zealand
208 Department of Conservation (45799-FAU).

209 **Results**

210 **Foraging trip length, diving events and video coverage**

211 The day following camera deployment, the penguin performed a 10.7 hour-long foraging trip.
212 The first dive event was recorded at 5:30 hrs and the last event concluded at 16:10 hrs. The bird
213 performed 286 dives of which 159 dive profiles matched the criteria for benthic dives (Figure 2).
214 Median dive depth reached during benthic dives was 54.4m (range: 4.8-62.1 m, n=159) whereas
215 the majority of pelagic dives occurred in the upper 10m of the water column (median: 7.8m,
216 range: 0.5-31.7 m, n=127); camera footage confirmed these to be principally travelling
217 behaviour (<https://vimeo.com/179414642>). For the first 3½ hours of the foraging trip (05:30-

218 09:00 hrs) the bird performed mainly pelagic dives, indicating primarily travelling behaviour
219 towards its main foraging grounds; ~~Y~~ellow-eyed penguins are known to exhibit high individual
220 site fidelity with regards to foraging locations (Moore, 1999; Mattern et al., 2007). Between
221 09:00 and 16:00 hrs the bird principally devoted its time to benthic diving while shallow dives
222 dominated the remaining 10 minutes of the foraging trip (Figure 2).

223 **Video coverage & quality**

224 The camera operated continuously from 11:00:22 hrs to 13:01:43 hrs. Due to frame loss
225 representing a mean 1.6 seconds of footage when video data were written to the file every 3
226 minutes, total length of the recorded footage amounted to 2 hours 8 seconds. Forty-six complete
227 dives were video recorded which corresponds to 16% of all dive events; of these 32 dives were
228 benthic dives. However, dives were longer during the middle of the day so that camera footage
229 covered 25% of the trip's cumulative dive time. The video quality proved to be significantly
230 better than that recorded with other animal-borne camera deployed on penguins to date
231 (<https://vimeo.com/268905870>). The light sensitivity of the camera was adequate to record
232 clear images at dive depths close to 70m and, combined with the large field of view, facilitated
233 detailed frame-by-frame analysis.

234 **Prey pursuits & capture**

235 A total of 20 prey pursuits were recorded at the seafloor (Figure 3). Fourteen of these resulted in
236 successful capture of either opalfish (*Hemerocoetes monopterygius*, 10 specimens) or blue cod
237 (*Parapercis colias*, 2 specimens); prey species could not be identified during two captures, but
238 the penguin's searching behaviour and ease of ingestion suggested these were opalfish and we
239 include them with opalfish captures below and in Figure 3. All of these prey pursuits occurred at
240 the sea floor with the penguin swimming very close to the bottom
241 (<https://vimeo.com/179414724>). During the camera operation time, the penguin spent 5.7
242 minutes on prey pursuit, which corresponds to 19% of the total time the bird foraged along the
243 seafloor (29.9 minutes) and 6% of its total dive time (89.9 minutes). The penguin spent most of
244 its active prey pursuit on opalfish (total 3.8 minutes, 12 events), 0.7 minutes were used to

capture blue cod (2 events), and 1.2 minutes of prey pursuit did not result in successful prey capture (Figure 3).

Two main prey pursuit strategies became apparent that were associated with prey species. When catching opalfish, the penguin would glide closely above the seafloor, sometimes briefly accelerating before starting to hover over a certain spot while repeatedly pecking at the substrate until the prey item was captured (<https://vimeo.com/179414724>). During encounters with blue cod prolonged pursuits ensued during which fish zigzagged at a fast pace along the seafloor (<https://vimeo.com/179414724#t=2m46s>). In one instance the fish was caught as it appeared to seek shelter at the base of a horse mussel protruding from the substrate (<https://vimeo.com/179414724#t=2m55s>). An unsuccessful prey pursuit of blue cod ended with the fish escaping under what appeared to be a half-buried back plate of a dishwasher (<https://vimeo.com/179414777>). A third blue cod encounter occurred just seconds after a successful capture of an opalfish; it seems likely that the resulting prolonged bottom time and oxygen-demanding prey pursuits drove the penguin to carry the fish to the surface at an almost vertical angle as indicated sun disc's central position in the frame; the fish was ultimately dropped at the surface (<https://vimeo.com/179414724#t=3m07s>).

Benthic habitat

During the video logger's operating time, the penguin spent 29.9 minutes foraging along the seafloor. The majority of the penguin's bottom time (90%) was spent over coarse sand, whereas time spent over fine sand (7%) and gravel (0.9%) was negligible (Figures 1 & 3). Two thirds of the bottom time (65.9%) was spent over sand ripples, the remaining time (34.1%) the bird foraged over flat ground. Brittle stars and anthozoans were present in most areas visited by the penguin with the former being present in 22.5 mins (75%) of the benthic video footage while the latter occur for a total of 17.9 mins (60%). Horse mussels were present for a total of 9.3 minutes (31%) of the bottom time.

Prey encounters were associated with certain benthic habitat types. All prey encounters occurred over coarse sand although the sediment structure differed depending on prey species.

Opalfish were principally encountered on sediment ripples (93.6% of the total prey pursuit time, <https://vimeo.com/179414724>), while flat bottom habitat played a more important role during blue cod pursuits (52.8% of pursuit time, <https://vimeo.com/179414724#t=2m32s>). With regards to epibenthic characteristics, brittle stars and anemones were present during the majority of the prey pursuit times for both fish species (Figure 3). However, horse mussels were present only during blue cod pursuits (81.4% of pursuit time).

Commented [A2]: 52.8% of blue cod pursuit time?

Commented [A3]: Of blue cod pursuit time?

Flipper movements

When descending to the sea floor the penguin propelled itself with fast, strong flipper strokes that got progressively slower and less pronounced with time and, thus, increasing depth (Pearson correlation – flipper amplitude: $R^2 = 0.69857$, $F_{1,363}=791.8$, $p<0.001$, BPM: $R^2=0.1324$, $F_{1,363}=55.2$, $p<0.001$, Figure 4a&b; <https://vimeo.com/179414575>). In contrast, ascending was principally passive with the penguin using its natural buoyancy to return to the surface, occasionally aided by a few strokes in the early stages of the ascent, decreasing beat frequency (flipper amplitude: $R^2=0.01065$, $F_{1,74}=0.4986$, $p=0.4988$; BPM: $R^2=0.2725$, $F_{1,76}=28.547$, $p<0.001$, Figure 4c&d) and no observable flipper movements towards the end of the dive (<https://vimeo.com/179414575#t=1m49s>). Despite differences in flipper movement between the two transit phases of a dive, the vertical velocities recorded by the TDR did not differ significantly (mean descent velocity: 1.45 ± 0.28 m/s, mean ascent velocity: 1.36 ± 0.57 m/s, $n=159$ dives, Welch's t-test: $t_{232}=1.73$, $p=0.09$).

During the bottom phase flipper amplitudes showed no correlation with relative bottom time (flipper amplitude: $R^2=0.001$, $F_{1,479}=0.6765$, $p=0.4245$ (Figure 4c), likely owing to the fact that bottom phases consisted of a mix of searching behaviour and high-speed prey pursuit (<https://vimeo.com/179414575#t=0m33s>). While searching the penguin showed lower flipper beat frequencies (133 ± 48 BPM, $n=809$) paired with greater flipper amplitudes ($53^\circ\pm14^\circ$) when compared to prey pursuit (BPM: 162 ± 44 , $n=113$, Welch's t-test: $t_{232}=-13.37$, $p<0.001$; amplitude: $45^\circ\pm7^\circ$, $t_{152}=6.39$, $p<0.001$). Flipper beat frequency increased slightly but consistently towards the end of the bottom phase (BPM: $R^2=0.017$, $F_{1,484}=8.164$, $p=0.004$, Figure 4d), most likely as a

299 result of the penguin often starting its ascent back to the surface not long after successful prey
300 captures (<https://vimeo.com/179414575#t=1m45s>).

301 **Surface breathing & underwater exhalation**

302 Frame counts of the video footage during 12 random selected surface periods between dives
303 showed that the penguin lifted its head out of the water to breathe for only brief moments
304 (average duration: 0.77 ± 0.22 s, $n=193$); for the majority of the time at the surface the bird kept
305 its head under water (1.53 ± 1.19 s, $n=182$) (<https://vimeo.com/179414575#t=2m25s>). Duration
306 of breathing intervals increased with ongoing duration of the surface period (Pearson
307 correlation: $\rho=0.45$, $F_{1,191}=47.4$, $p<0.001$) indicating increased respiration activity in preparation
308 for the next dive (Figure 5).

309 During the dive, exhalation regularly occurred at the onset of phases with increased acceleration
310 (i.e. prey pursuit). Such exhalations were brief but performed with substantial force; air was
311 jetted from the nostrils as a fine gas spurt (<https://vimeo.com/179418254>). During the passive
312 phase of the ascent, the penguin frequently exhaled as indicated by a stream of large bubbles
313 released from the nostrils. The bird released substantial amounts of air on the last few meters
314 immediately prior to reaching the surface (<https://vimeo.com/179414575#t=2m18s>). While
315 some of this air may have been released from the plumage (c.f. Davenport et al. 2011) bubbles
316 seem principally to originate from the frontal head region; there was no visible major gas release
317 from the penguin's back region.

318 **Discussion**

319 The high-quality video footage provided a substantial amount of new insights into the foraging
320 behaviour of ~~Y~~yellow-eyed penguins and their benthic habitat. While it is impossible to draw far-
321 ranging conclusions from only a single deployment, it nevertheless highlights that high-
322 definition cameras provide a new tool facilitating the examination of various aspects of the
323 foraging ecology of marine predators through direct observation. It can be particularly useful to

324 verify and calibrate behaviours measured with other types of devices such as TDR and
325 accelerometers.

326 **Device effects**

327 Attaching external recording devices to diving animals always comes at the cost of
328 compromising their streamlined body shape (e.g. Ludynia et al., 2012), a problem that can be
329 mitigated via device shape, size and attachment position (Bannasch, Wilson & Culik, 1994). At
330 the surface there were no indications that the penguin was negatively affected by the device; the
331 bird did not exhibit balancing problems which externally attached devices can cause in smaller
332 species (Chiaradia et al., 2005), nor did it peck at the device frequently which suggests aberrant
333 behaviour (Wilson & Wilson, 1989). Moreover, the number of successful prey captures further
334 suggests that the bird's foraging capabilities were not drastically affected by the video logger.
335 With the exception of two unsuccessful blue cod encounters, all events classified as prey pursuit
336 were merely accelerations that did not end in any obvious prey encounter. The bird was one of
337 the few breeders that raised two chicks to fledging in an otherwise poor breeding season.

338 **Predator-prey interactions & prey species importance**

339 In line with previous descriptions of ~~Y~~yellow-eyed penguins as primarily benthic foragers
340 (Mattern et al., 2007), the penguin's prey pursuit and captures recorded during the camera
341 operation indeed all occurred at the sea floor. Swimming very close to the seafloor could serve
342 several purposes. It could be a strategy to flush out benthic prey that blends in with the
343 substrate, but it could also mean the penguin has a greater chance to see its prey from the side,
344 and thus reduce the effect of prey camouflage. Opalfish, for example, are very well camouflaged
345 and very difficult to make out from above (Roberts, Stewart & Struthers, 2015). This species
346 seems to principally rely on its camouflage as means of predator avoidance since none of the
347 opalfish captures involved a chase. In contrast, during both successful blue cod encounters,
348 extended high-speed chases ensued before the fish was ultimately captured. Blue cod and
349 opalfish differ significantly in their anatomy with the small, slender opalfish presumably lacking
350 the physical prowess for prolonged swimming when compared to muscular blue cod (Roberts,

351 Stewart & Struthers, 2015). When facing an air breathing predator, the latter strategy is likely
352 advantageous as the predator's increased energy requirements for pursuit make escape a more
353 likely outcome for the prey. The penguin's hasty ascent and subsequent failure to consume a
354 blue cod it captured after a 22-second-long chase demonstrates the efficacy of this evasion
355 strategy.

356 Both opalfish and blue cod have previously been found to be among the most important prey
357 items in the ~~Y~~yellow-eyed penguin's diet (van Heezik, 1990a; Moore & Wakelin, 1997). While
358 both fish species have comparable energetic values ($\sim 20 \text{ kJ g}^{-1}$, Browne et al., 2011), the body
359 mass of opalfish is considerably lower when compared to blue cod (van Heezik, 1990a,b). It is
360 possible that the energy gain from catching blue cod justifies the expenditure to catch it, while
361 the easier-to-catch opalfish might need to be caught in larger quantities. However, recent studies
362 suggest that blue cod might be suboptimal prey for chick-rearing ~~Y~~yellow-eyed penguins due to
363 their size (Browne et al., 2011; Mattern et al., 2013) so that the penguin's ability to locate prey
364 such as opalfish might be a decisive factor with regards to reproductive success.

365 **Benthic environment**

366 Judging from the total time the bird spent over a benthic environment dominated by coarse sand
367 and sediment ripples (65.9% of total bottom time) as well as almost exclusive encounters of
368 opalfish over such habitat (Figures 1 & 3), it can be assumed that the penguin focussed
369 principally on this species. Blue cod encounters were associated with the presence of horse
370 mussels. These large bivalves protrude from the seafloor and provide hard substrate for other
371 epibenthic taxa, thereby increasing local benthic biodiversity (Cummings et al., 1998). Benthic
372 habitat with increased benthic biodiversity is generally more attractive to a variety of benthic
373 fish species, most likely due to enhanced feeding conditions (Cranfield et al., 2001). Our video
374 data also suggests that the fish use the bivalves and associated cavities as shelter to avoid
375 capture (<https://vimeo.com/179414777>).

376 The majority of prey pursuits occurred in areas that featured anthozoans, principally sea
377 anemones (Figures 1 & 3). Anemones are known to play an important role as refugia and feeding

378 habitats for small fish (Elliott, 1992) and could therefore be another indicator for locally
379 increased biodiversity. Brittle stars on the other hand, although equally abundant, seemed to be
380 of lesser relevance with regards to prey encounters. So, it appears that examining the
381 composition of the benthic habitat alone might enable assessment of which prey types penguins
382 are foraging for, though more data are required before conclusions can be drawn. However, this
383 already hints at the potential for wide-ranging habitat analysis of at-sea movements in benthic
384 predators, provided that spatial distribution of the different benthic habitats can be obtained.
385 While in our specific case, no such habitat maps exist, planned further deployments of video
386 loggers are expected to provide the necessary environmental information.

387 Deploying video loggers on penguins could enable detailed mapping of the benthic habitat
388 within the species home ranges. Yellow-eyed penguins are known to have preferred individual
389 foraging areas often with little overlap between birds (Moore, 1999). Moreover, the birds tend to
390 often dive along the seafloor when swimming towards their foraging grounds (Mattern et al.
391 2007) so that camera logger data in combination with GPS information can be used to establish
392 spatial biodiversity indices and benthic habitat maps.

393 The outer ranges of the marine habitat of ~~Y~~yellow-eyed penguins from the Otago Peninsula is
394 subject to bottom fisheries which have a profound effect on benthic ecosystems (e.g. Hinz et al.,
395 2009; Queirós et al., 2006; Schratzberger and Jennings, 2002). Yellow-eyed penguins have been
396 found to forage in the wake of trawl fisheries, potentially to the detriment of their reproductive
397 success (Mattern et al., 2013). Changes in sediment structure and epibenthic biodiversity as a
398 result of bottom trawl disturbance likely negatively affect the penguins' foraging success
399 (Browne et al., 2011). Camera loggers can help to determine how much of the penguins' foraging
400 habitat has been compromised by fishing activities and what the consequences are for this
401 species' foraging behaviour and success.

402 Beyond investigations of behaviour in a wider environmental context, our study also shows the
403 potential application of camera loggers for the investigation of physiological aspects of marine
404 animals.

Flipper movements

Our observations of flipper movements, i.e. strong flipper movements at the beginning of a dive that decrease with depth, and cessation of flipper movements during ascent, align with findings reported in other penguins. Using accelerometers, Sato et al. (2002) found that King penguins showed vigorous flipper beating at the beginning of a dive to counter positive buoyancy. With increasing depth, air volume in the penguin's body becomes compressed, reducing its buoyancy so that fewer flipper beats are required. That this also applies to flipper amplitude (Figure 45) was not detectable by using body acceleration as the only measure. A more elaborate system of sensors and magnets attached to flippers was used on Magellanic penguins which allowed the recording of both flipper amplitudes and beat frequencies (Wilson & Liebsch, 2003). However, the system is known to be prone to failure, rendering the use of back-mounted wide-angle cameras a much more reliable alternative. Flipper beat frequencies and amplitudes are directly related to energy expenditure (Kooyman & Ponganis, 1998; Sato et al., 2011). They provide the means for the quantification of energy budgets (Wilson & Liebsch, 2003) and subsequently can be used to assess individual fitness in relation to foraging success and subsequent reproductive performance (Kooyman & Ponganis, 1998).

We provide evidence that the ascent phase in penguins is largely passive, as has been suggested using both accelerometers and magnets (Sato et al., 2002; Wilson & Liebsch, 2003). Sato et al. (2002) concluded that during ascent penguins benefit from expanding air volume in their body which increases their buoyancy as they get closer to the surface. Penguins also actively slow down their ascent and it was argued that this could be achieved by increasing the attack angles of their flippers to increase drag (Sato et al., 2002). Judging from body movements apparent in the video data during the ascent phases we suggest that the ~~Y~~yellow-eyed penguin indeed adjusted flipper attack angles while ascending, although this seems to be more for steering. Based on the video footage it appears that the bird uses controlled exhalation towards the end of the ascent to control speed (<https://vimeo.com/179414575#t=2m18s>).

Respiration

The video data provide new insights into the respiration of yellow-eyed penguins. To date it was unclear whether penguins exhale regularly while diving. Various studies estimated diving air volume via a penguin's buoyancy calculated from its ascent speeds at the end of dives (Sato et al., 2002, 2011). However, the accuracy of this approach is compromised if the penguins were to exhale prior to their final ascent (Ponganis, St Leger & Scadeng, 2015). The video data clearly showed that the penguin generally exhaled when accelerating during prey pursuit so that models estimating diving air volume via the proxy buoyancy must take acceleration into account. The fact that the penguin exhaled when accelerating probably serves the purpose of reducing blood CO₂ and mobilizing O₂ from oxygen stores for prey pursuit. Such pursuits must be costly in terms of oxygen consumption as is evident from the observed consecutive prey encounters during one single dive, which resulted in the penguin letting go of the second fish after a rapid ascent to the surface (<https://vimeo.com/179414724#t=3m07s>). Unlike seals that have been found to exhale when ascending from deep dives, most likely to reduce the drop in blood oxygen (Hooker et al., 2005), the penguin principally exhaled during the second half of the ascent possibly indicating adjustment of buoyancy and ascent speed (but see also Davenport et al. 2011). Reoxygenation during the surface period in penguins is highly optimized (Wilson et al., 2003). Inhalation events at the surface are brief so that the bird can frequently lower its head into the water, presumably in an effort to look out for potential predators (e.g. sharks, sea lions; Seddon et al., 2013). Extensive exhalation prior to resurfacing also prevents pulmonary barotrauma and facilitates immediate inhalation once back at the surface.

Conclusions

The deployment of a full HD video logger on a yellow-eyed penguin resulted in a versatile visual data set that provided a variety of information well beyond what was initially intended. Enhanced video quality allows detailed analysis of the benthic environment as well as prey encounter rates and prey composition. In combination with GPS data, the potential for a

comprehensive survey of benthic ecosystems is substantial highlighting the multi-disciplinary potential of such data.

A large field of view achieved through wide-angle lenses furthermore allows detailed analysis of flipper movements, which to date could only be achieved through elaborate modelling of accelerometer data (Sato et al., 2002, 2011) or use of complicated magnetic logger setups (Wilson & Liebsch, 2003). Neither of these setups provided information about exhalation, which appears to play a much more important role during diving than previously thought. When comparing video data recorded here with videos from previously published studies (e.g. Watanabe and Takahashi, 2013, <https://vimeo.com/268905870>) it becomes clear that greater visual fidelity of full HD cameras comes along with a much wider range of quantifiable data. This creates a new opportunity for a more holistic approach to study the diving behaviour of marine animals that integrates behaviour, physiology and their environment.

Depending on which behaviours are quantified, the manual analysis of video data can be quite time-consuming. For example, flipper beats and angles require a frame-by-frame analysis; an average dive duration of 3 minutes translated to 5400 frames per dive. However, the higher the resolution and quality of the video footage, the greater the potential to develop machine learning algorithms (such as Google Cloud Video Intelligence; <https://cloud.google.com/video-intelligence/>) that may be used to automate the analysis process. For more basic analyses such as prey composition and encounter rates, but also determination of environmental parameters, there already exist software solutions that offer an enhanced workflow, for example the video annotation software BORIS (<http://www.boris.unito.it/>).

Obviously, there are still limitations to the use of camera loggers. Restrictions arise from the battery life as well as the memory to store high definition video data. In our case, 15 minutes of footage resulted in video file sizes of 1.5 gigabytes. Moreover, the deployment with the camera set-up we used requires a certain amount of predictability, particularly knowledge about how soon after departure the bird is likely to engage in behaviours that are of interest (e.g. prey pursuit). For all these reasons, the technology currently available is best suited for short-term

484 deployments on central place foragers. Although video data recorded on animals performing
485 long-term foraging trips (e.g. Magellanic penguins, Boersma & Rebstock, 2009) might still
486 deliver valuable data, this has to be weighed against the fact that external devices inevitably
487 have an effect on the animal's foraging ability (Bannasch, Wilson & Culik, 1994; Ludynia et al.,
488 2012). This could be alleviated by incorporating further mechanisms to control camera
489 recording (e.g. duty-cycling of recording function, pressure control). While the use of animal-
490 borne cameras for scientific research is still in its early day, the enormous potential of this
491 technology will doubtlessly result in devices incorporating more elaborate functionality in the
492 future.

493 **Acknowledgments**

494 We would like to thank Horst Mattern, Melanie Young and Jim Watts for help in the field, and
495 Leon Berard for first preliminary evaluation of the video data. Special thanks are due to Bruce
496 McKinlay (Department of Conservation) for supporting this project and facilitating the
497 permitting process of a novel bio-logging method.

References

- Bannasch R., Wilson RP., Culik BM. 1994. Hydrodynamic aspects of design and attachment of a back-mounted device in penguins. *The Journal of Experimental Biology* 194:83–96.
- BirdLife International. 2012. Megadyptes antipodes. In: *The IUCN Red List of Threatened Species*. e.T22697800A40186242.
- Boersma PD., Rebstock GA. 2009. Foraging distance affects reproductive success in Magellanic penguins. *Marine Ecology Progress Series* 375:263–275. DOI: 10.3354/meps07753.
- Browne T., Lalas C., Mattern T., Van Heezik Y. 2011. Chick starvation in yellow-eyed penguins: Evidence for poor diet quality and selective provisioning of chicks from conventional diet analysis and stable isotopes. *Austral Ecology* 36:99–108. DOI: 10.1111/j.1442-9993.2010.02125.x.
- Chiaradia A., Ropert-coudert Y., Healy M., Knott N. 2005. Finding the balance : the effect of the position of external devices on little penguins. *Polar Bioscience* 18:46–53.
- Cranfield HJ., Carbines G., Michael KP., Dunn A., Stotter DR., Smith DJ. 2001. Promising signs of regeneration of blue cod and oyster habitat changed by dredging in Foveaux Strait, southern New Zealand. *New Zealand Journal of Marine and Freshwater Research* 35:897–908. DOI: 10.1080/00288330.2001.9517052.
- Cummings VJ., Thrush SF., Hewitt JE., Turner SJ. 1998. The influence of the pinnid bivalve *Atrina zelandica* (Gray) on benthic macroinvertebrate communities in soft-sediment habitats. *Journal of Experimental Marine Biology and Ecology* 228:227–240. DOI: 10.1016/S0022-0981(98)00028-8.
- Davenport J., Hughes RN., Shorten M., Larsen PS. 2011. Drag reduction by air release promotes fast ascent in jumping emperor penguins-a novel hypothesis. *Marine Ecology Progress Series* 430:171–182. DOI: 10.3354/meps08868.
- Ellenberg U., Mattern T. 2012. *Yellow-eyed penguin - review of population information*.

Wellington, New Zealand. DOI: <https://doi.org/10.13140/RG.2.2.21606.83523>.

Elliott J. 1992. The role of sea anemones as refuges and feeding habitats for the temperate fish *Oxylebius pictus*. *Environmental Biology of Fishes* 35:381–400. DOI: 10.1007/BF00004991.

Gómez-Laich A., Yoda K., Zavalaga C., Quintana F. 2015. Selfies of Imperial Cormorants (*Phalacrocorax atriceps*): What Is Happening Underwater? *PLOS ONE* 10:1–18. DOI: 10.1371/journal.pone.0136980.

Handley JM., Pistorius P. 2015. Kleptoparasitism in foraging gentoo penguins *Pygoscelis papua*. *Polar Biology* 39:391–395. DOI: 10.1007/s00300-015-1772-2.

van Heezik Y. 1990a. Seasonal, geographical, and age-related variations in the diet of the Yellow-eyed Penguin (*Megadyptes antipodes*). *New Zealand Journal of Zoology* 17:201–212.

van Heezik Y. 1990b. Diets of yellow-eyed, Fiordland crested, and little blue penguins breeding sympatrically on Codfish Island, New Zealand. *New Zealand Journal of Zoology* 17:543–548.

van Heezik Y., Davis LS. 1990. Effects of food variability on growth rates, fledging sizes and reproductive success in the Yellow-eyed Penguin *Megadyptes antipodes*. *Ibis* 132:354–365.

Hinz H., Prieto V., Kaiser MJ. 2009. Trawl disturbance on benthic communities: Chronic effects and experimental predictions. *Ecological Applications* 19:761–773. DOI: 10.1890/08-0351.1.

Hooker SK., Miller PJO., Johnson MP., Cox OP., Boyd IL. 2005. Ascent exhalations of Antarctic fur seals: a behavioural adaptation for breath-hold diving? *Proceedings of the Royal Society of London B: Biological Sciences* 272:355–363. DOI: 10.1098/rspb.2004.2964.

Kane SA., Zamani M. 2014. Falcons pursue prey using visual motion cues: new perspectives from animal-borne cameras. *Journal of Experimental Biology* 217:225–234. DOI: 10.1242/jeb.092403.

Kooyman GL., Ponganis PJ. 1998. The physiological basis of diving to depth: birds and mammals. *Annual review of physiology* 60:19–32. DOI: 10.1146/annurev.physiol.60.1.19.

548 Ludynia K., Dehnhard N., Poisbleau M., Demongin L., Masello JF., Quillfeldt P. 2012. Evaluating
 549 the Impact of Handling and Logger Attachment on Foraging Parameters and Physiology in
 550 Southern Rockhopper Penguins. *PLoS ONE* 7:e50429. DOI: 10.1371/journal.pone.0050429.
 551 Mattern T., Ellenberg U., Houston DMDM., Davis LSLs. 2007. Consistent foraging routes and
 552 benthic foraging behaviour in yellow-eyed penguins. *Marine Ecology Progress Series*
 553 343:295–306. DOI: 10.3354/meps06954.
 554 Mattern T., Ellenberg U., Houston DMDM., Lamare M., Davis LSLs., Van Heezik Y., Seddon PJPJ.
 555 2013. Straight line foraging in yellow-eyed penguins: new insights into cascading fisheries
 556 effects and orientation capabilities of marine predators. *PloS ONE* 8:e84381. DOI:
 557 10.1371/journal.pone.0084381.
 558 Moll RJ., Millspaugh JJ., Beringer J., Sartwell J., He Z. 2007. A new “view” of ecology and
 559 conservation through animal-borne video systems. *Trends in Ecology & Evolution* 22:660–
 560 668. DOI: <http://dx.doi.org/10.1016/j.tree.2007.09.007>.
 561 Moore PJ., Wakelin MD., Douglas ME., McKinlay B., Nelson D., Murphy B. 1995. *Yellow-eyed*
 562 *penguin foraging study, south-eastern New Zealand, 1991-1993*. Wellington, N.Z.:
 563 Department of Conservation. Available at [http://www.doc.govt.nz/Documents/science-](http://www.doc.govt.nz/Documents/science-and-technical/sr83a.pdf)
 564 [and-technical/sr83a.pdf](http://www.doc.govt.nz/Documents/science-and-technical/sr83a.pdf).
 565 Moore PJ. 1999. Foraging range of the Yellow-eyed penguin *Megadyptes antipodes*. *Marine*
 566 *Ornithology* 27:49–58.
 567 Moore PJ., Wakelin MD. 1997. Diet of the Yellow-eyed penguin *Megadyptes antipodes*, South
 568 Island, New Zealand, 1991-1993. *Marine Ornithology* 25:17–29.
 569 Moreau M., Siebert S., Buerkert A., Schlecht E. 2009. Use of a tri-axial accelerometer for
 570 automated recording and classification of goats’ grazing behaviour. *Applied Animal*
 571 *Behaviour Science* 119:158–170. DOI: <http://dx.doi.org/10.1016/j.applanim.2009.04.008>.
 572 Noda T., Kawabata Y., Arai N., Mitamura H., Watanabe S. 2014. Animal-mounted

gyroscope/accelerometer/magnetometer: in situ measurement of the movement
performance of fast-start behaviour in fish. *J Exp Mar Biol Ecol* 451. DOI:
10.1016/j.jembe.2013.10.031.

Ponganis PJ, Van Dam RP, Marshall G, Knowler T, Levenson DH. 2000. Sub-ice foraging behavior
of emperor penguins. *Journal of Experimental Biology* 203:3275–3278.

Ponganis PJ, St Leger J, Scadeng M. 2015. Penguin lungs and air sacs: implications for
baroprotection, oxygen stores and buoyancy. *Journal of Experimental Biology* 218:720–730.
DOI: 10.1242/jeb.113647.

Queirós AM, Hiddink JG, Kaiser MJ, Hinz H. 2006. Effects of chronic bottom trawling
disturbance on benthic biomass, production and size spectra in different habitats. *Journal of
Experimental Marine Biology and Ecology* 335:91–103. DOI: 10.1016/j.jembe.2006.03.001.

R Core Team. 2014. R: A Language and Environment for Statistical Computing.

Roberts CD, Stewart AL, Struthers CD. 2015. *The Fishes of New Zealand*. Wellington, New
Zealand: Te Papa Press.

Ruxton GD. 2006. The unequal variance t-test is an underused alternative to Student's t-test and
the Mann–Whitney U test. *Behavioral Ecology* 17:688–690. DOI: 10.1093/beheco/ark016.

Sakamoto KQ, Takahashi A, Iwata T, Trathan PN. 2009. From the Eye of the Albatrosses: A Bird-
Borne Camera Shows an Association between Albatrosses and a Killer Whale in the
Southern Ocean. *PLoS ONE* 4:e7322. DOI: 10.1371/journal.pone.0007322.

Sato K, Naito Y, Kato A, Niizuma Y, Watanuki Y, Charrassin JB, Bost C-A, Handrich Y, Le Maho
Y. 2002. Buoyancy and maximal diving depth in penguins. *Journal of Experimental Biology*
205:1189–1197.

Sato K, Shiomi K, Marshall G, Kooyman GL, Ponganis PJ. 2011. Stroke rates and diving air
volumes of emperor penguins: implications for dive performance. *Journal of Experimental
Biology* 214:2854–2863. DOI: 10.1242/jeb.055723.

598 Schratzberger M., Jennings S. 2002. Impacts of chronic trawling disturbance on meiofaunal
599 communities. *Marine Biology* 141:991–1000. DOI: 10.1007/s00227-002-0895-5.

600 Seddon PJ., Ellenberg U., van Heezik Y. 2013. Yellow-eyed penguin (*Megadyptes antipodes*). In:
601 Garcia Borboroglu P, Boersma PD eds. *Penguins: Natural History and Conservation*. Seattle &
602 London: University of Washington Press, 91–110.

603 Sutton GJ., Hoskins AJ., Arnould JPY. 2015. Benefits of Group Foraging Depend on Prey Type in a
604 Small Marine Predator, the Little Penguin. *PLoS ONE* 10:e0144297. DOI:
605 10.1371/journal.pone.0144297.

606 Takahashi A., Kokubun N., Mori Y., Shin H-C. 2008. Krill-feeding behaviour of gentoo penguins as
607 shown by animal-borne camera loggers. *Polar Biology* 31:1291–1294.

608 Thiebot J-B., Ito K., Raclot T., Poupart T., Kato A., Ropert-Coudert Y., Takahashi A. 2016. On the
609 significance of Antarctic jellyfish as food for Adélie penguins, as revealed by video loggers.
610 *Marine Biology* 163:1–8. DOI: 10.1007/s00227-016-2890-2.

611 Thiebot J-B., Arnould JP., Gómez-Laich A., Ito K., Kato A., Mattern T., Mitamura H., Noda T.,
612 Poupart T., Quintana F., Raclot T., Ropert-Coudert Y., Sala JE, Seddon PJ., Sutton GJ., Yoda K.,
613 Takahashi A. 2017. Jellyfish and other gelata as food for four penguin species - insights
614 from predator-borne videos. *Frontiers in Ecology and the Environment* 15:437–441. DOI:
615 10.1002/fee.1529.

616 Valletta JJ., Torney C., Kings M., Thornton A., Madden J. 2017. Applications of machine learning in
617 animal behaviour studies. *Animal Behaviour* 124:203–220. DOI:
618 10.1016/j.anbehav.2016.12.005.

619 Volpov BL., Rosen DAS., Hoskins AJ., Lourie HJ., Dorville N., Baylis AMM., Wheatley KE., Marshall
620 G., Abernathy K., Semmens J., Hindell MA., Arnould JPY. 2016. Dive characteristics can
621 predict foraging success in Australian fur seals (*Arctocephalus pusillus doriferus*) as
622 validated by animal-borne video. *Biology Open* 5:262 LP-271.

623 Watanabe YY., Takahashi A. 2013. Linking animal-borne video to accelerometers reveals prey
624 capture variability. *Proceedings of the National Academy of Sciences* 110:2199–2204. DOI:
625 10.1073/pnas.1216244110.

626 Watanuki Y., Daunt F., Takahashi A., Newell M., Wanless S., Sato K., Miyazaki N. 2008.
627 Microhabitat use and prey capture of a bottom-feeding top predator, the European shag,
628 shown by camera loggers. *Marine Ecology Progress Series* 356:283–293. DOI:
629 10.3354/meps07266.

630 Wilmers CC., Nickel B., Bryce CM., Smith JA., Wheat RE., Yovovich V. 2015. The golden age of bio-
631 logging: how animal-borne sensors are advancing the frontiers of ecology. *Ecology*
632 96:1741–1753. DOI: 10.1890/14-1401.1.

633 Wilson RP., Pütz K., Peters G., Culik BM., Scolaro JA., Charrassin J-B., Ropert-Coudert Y. 1997.
634 Long-term attachment of transmitting and recording devices to penguins and other
635 seabirds. 25:101–106.

636 Wilson RP., Simeone A., Luna-Jorquera G., Steinfurth A., Jackson S., Fahlman A. 2003. Patterns of
637 respiration in diving penguins: is the last gasp an inspired tactic? *The Journal of*
638 *experimental biology* 206:1751–1763. DOI: 10.1242/jeb.00341.

639 Wilson R., Liebsch N. 2003. Up-beat motion in swinging limbs: new insights into assessing
640 movement in free-living aquatic vertebrates. *Marine Biology* 142:537–547. DOI:
641 10.1007/s00227-002-0964-9.

642 Wilson RP., Wilson MT. 1989. A peck activity record for birds fitted with devices. *Journal of Field*
643 *Ornithology* 60:104–108.

644