

# Rhizobacterial communities of five co-occurring desert halophytes

Yan Li<sup>1,2,3</sup>, Yan Kong<sup>4,5</sup>, Dexiong Teng<sup>2</sup>, Xueni Zhang<sup>1,2</sup>, Xuemin He<sup>1,2</sup>, Yang Zhang<sup>6</sup>, Guanghui Lv<sup>Corresp. 1,2</sup>

<sup>1</sup> Key Laboratory of Oasis Ecology of Education Ministry, Xinjiang University, Urumqi, Xinjiang, China

<sup>2</sup> Institute of Arid Ecology and Environment, Xinjiang University, Urumqi, Xinjiang, China

<sup>3</sup> Ecology Post-doctoral Research Station, Xinjiang University, Urumqi, Xinjiang, China

<sup>4</sup> School of Life Science and Biotechnology, Shanghai Jiaotong University, Shanghai, China

<sup>5</sup> SJTU-Yale Joint Center for Biostatistics, Shanghai Jiaotong University, Shanghai, China

<sup>6</sup> College of Resource and Environment Sciences, Xinjiang University, Urumqi, Xinjiang, China

Corresponding Author: Guanghui Lv

Email address: ler@xju.edu.cn

**Background:** Recently, researches have begun to investigate the microbial communities associated with halophytes. Both rhizobacterial community composition and the environmental drivers of community assembly have been addressed. However, few studies have explored the structure of rhizobacterial communities associated with halophytic plants that are co-occurring in arid, salinized areas. **Methods:** Five halophytes were selected for study: these co-occurred in saline soils in the Ebinur Lake Nature Reserve, located at the western margin of the Gurbantunggut Desert of Northwestern China. Halophyte-associated bacterial communities were sampled, and the bacterial 16S rDNA V3-V4 region amplified and sequenced using the Illumina Miseq platform. The bacterial community diversity and structure were compared between the rhizosphere and bulk soils, as well as among the rhizosphere samples. The effects of plant species identity and soil properties on the bacterial communities were also analyzed. **Results:** Significant differences were observed between the rhizosphere and bulk soil bacterial communities. Diversity was higher in the rhizosphere than in the bulk soils. Abundant taxonomic groups (from phylum to genus) in the rhizosphere were much more diverse than in bulk soils. Proteobacteria, Firmicutes, Actinobacteria, Bacteroidetes and Planctomycetes were the most abundant phyla in the rhizosphere, while Proteobacteria and Firmicutes were common in bulk soils. Overall, the bacterial community composition were not significantly differentiated between the bulk soils of the five plants, but community diversity and structure differed significantly in the rhizosphere. The diversity of *Halostachys caspica*, *Halocnemum strobilaceum* and *Kalidium foliatum* associated communities was lower than that of *Limonium gmelinii* and *Lycium ruthenicum* communities. Furthermore, the composition of *Halostachys caspica* and *Halocnemum strobilaceum* communities was very

different from that of *Limonium gmelinii* and *Lycium ruthenicum* communities. The diversity and community structure were influenced by soil EC, pH and nutrient content (TOC, SOM, TON and AP); of these, the effects of EC on bacterial community composition were less important than those of soil nutrients. **DISCUSSION:** Halophytic plant species played an important role in shaping associated rhizosphere bacterial communities. When salinity levels were constant, soil nutrients emerged as key factors structuring bacterial communities, while EC played only a minor role. Pairwise differences among the rhizobacterial communities associated with different plant species were not significant, despite some evidence of differentiation. Further studies involving more halophyte species, and individuals per species, are necessary to elucidate plant species identity effects on the rhizosphere for co-occurring halophytes.

# Rhizobacterial communities of five co-occurring desert halophytes

Yan LI<sup>1,2,3</sup>, Yan Kong<sup>4,5</sup>, Dexiong Teng<sup>2</sup>, Xueni Zhang<sup>1,2</sup>, Xuemin HE<sup>1,2</sup>,  
Yang Zhang<sup>6</sup>, Guanghui Lv<sup>1,2,\*</sup>

1 Key Laboratory of Oasis Ecology, Education Ministry, Urumqi, China

2 Institute of Arid Ecology and Environment, Xinjiang University, Urumqi, China

3 Ecology Post-doctoral Research Station, Xinjiang University, Urumqi, China

4 School of Life Science and Biotechnology, Shanghai Jiaotong University, Shanghai, China

5 SJTU-Yale Joint Center for Biostatistics, Shanghai Jiaotong University, Shanghai, China

6 College of Resource and Environment Sciences, Xinjiang University, Xinjiang, China

\* Corresponding author, email: [ler@xju.edu.cn](mailto:ler@xju.edu.cn)

# ABSTRACT

**Background:** Recently, researches have begun to investigate the microbial communities associated with halophytes. Both rhizobacterial community composition and the environmental drivers of community assembly have been addressed. However, few studies have explored the structure of rhizobacterial communities associated with halophytic plants that are co-occurring in arid, salinized areas.

**Methods:** Five halophytes were selected for study: these co-occurred in saline soils in the Ebinur Lake Nature Reserve, located at the western margin of the Gurbantunggut Desert of Northwestern China. Halophyte-associated bacterial communities were sampled, and the bacterial 16S rDNA V3–V4 region amplified and sequenced using the Illumina Miseq platform. The bacterial community diversity and structure were compared between the rhizosphere and bulk soils, as well as among the rhizosphere samples. The effects of plant species identity and soil properties on the bacterial communities were also analyzed.

**Results:** Significant differences were observed between the rhizosphere and bulk soil bacterial communities. Diversity was higher in the rhizosphere than in the bulk soils. Abundant taxonomic groups (from phylum to genus) in the rhizosphere were much more diverse than in bulk soils. Proteobacteria, Firmicutes, Actinobacteria, Bacteroidetes and Planctomycetes were the most abundant phyla in the rhizosphere, while Proteobacteria and Firmicutes were common in bulk soils. Overall, the bacterial community composition were not significantly differentiated between the bulk soils of the five plants, but community diversity and structure differed significantly in the rhizosphere. The diversity of *Halostachys caspica*, *Halocnemum strobilaceum* and *Kalidium foliatum* associated communities was lower than that of *Limonium gmelinii* and *Lycium ruthenicum* communities. Furthermore, the composition of *Halostachys caspica* and *Halocnemum strobilaceum* communities was very different from that of *Limonium gmelinii* and *Lycium ruthenicum* communities. The diversity and community structure were influenced by soil EC, pH and nutrient content (TOC, SOM, TON and AP); of these, the effects of EC on bacterial community composition were less important than those of soil nutrients.

**DISCUSSION:** Halophytic plant species played an important role in shaping associated rhizosphere bacterial communities. When salinity levels were constant, soil nutrients emerged as key factors structuring bacterial communities, while EC played only a minor role. Pairwise differences among the rhizobacterial communities associated with different plant species were not significant, despite some evidence of differentiation. Further studies involving more halophyte species, and individuals per species, are necessary to elucidate plant species identity effects on the rhizosphere for co-occurring halophytes.

# INTRODUCTION

Salinization is a serious land degradation problem, as high salinity limits plant growth. Salts may accumulate in soils as a result of natural processes, such as mineral weathering, dust collection and precipitation, or artificial processes, such as irrigation (Oosterbaan 1988); both may lead to saline soils that make it difficult for plants to absorb moisture from the soil. Halophytes are salt-tolerant plants that can grow in areas with salt (NaCl) concentrations higher than 400 mM (Flowers 2004; English & Colmer 2011). In saline soil environments, halophytes play an important role in carbon sequestration, nutrient mineralization, nutrient cycling and improvement of the micro-environment (Cao et al. 2014; Chaudhary et al. 2015), and may have great potential to preserve ecosystems.

Salinity tolerance in halophytes is not solely due to physiological mechanisms and their genetic regulation (Vasquez et al. 2005), but also to complex ecological processes within the plant rhizosphere and phyllosphere; microorganisms inhabiting the roots and leaves of halophytes may significantly contribute to their salinity tolerance (Ruppel et al. 2013). Many microorganisms have plant growth-promoting activities and confer salt tolerance on halophytic plants (Nabti et al. 2007; Sgroy et al. 2009; Jha et al. 2012; Mapelli et al. 2013). Recent studies of halophyte-associated microbial communities have addressed the ecological and environmental drivers underlying community assembly and recruitment (Jha et al. 2012; Borruso et al. 2014; Marasco et al. 2016; Chaudhary et al. 2017; Tian & Zhang 2017). These studies have revealed that bacterial communities in the halophyte rhizosphere are distinctly different from those of non-halophytic plants, containing a larger proportion of halophilic bacteria (Al-Mailem et al. 2010). Many halophilic bacteria have been identified or isolated from halophyte roots, soils and desert habitats, including species belonging to the following genera: *Alkalimonas*, *Bacillus*, *Brachybacterium*, *Brevibacterium*, *Cronobacter*, *Halobacillus*, *Halomonas*, *Marinococcus*, *Methylibium*, *Nesterenkonia*, *Oceanobacillus*, *Staphylococcus*, *Stenotrophomonas*, *Virgibacillus* and *Zhihengliuella* (Sgroy et al. 2009; Siddikee et al. 2010; Tang et al. 2011; Shi et al. 2012; Zhou et al. 2012; Ramadoss et al. 2013; Borsodi et al. 2015; Zhao et al. 2016).

The diversity and composition of the rhizosphere bacterial community depends not only on the plant species, but also on various soil parameters (Tian & Gao 2014; Rodriguez-Blanco et al. 2015; Pii et al. 2016; Song et al. 2017). Different plant species, or even genotypes within species, tend to assemble distinct rhizobacterial communities (Chaudhary et al. 2015). For example, in the rhizosphere of *Aster tripolium*, Actinobacteria, Firmicutes and Proteobacteria are the most abundant bacterial phyla, and *Bacillus* the dominant genus (Szymanska et al. 2016b). In contrast, Acidimicrobiales, Myxococcales and Sphingomonadales are common in the rhizosphere of *Halimione portulacoides* and *Sarcocornia perennis* (Oliveira et al. 2014). The *Puccinellia limosa* rhizosphere is dominated by *Halomonas* and *Nesterenkonia* species (Borsodi et al. 2015). Similar rhizobacterial communities may be found in different environments when the same plant species is present (Smalla et al. 2001; Berg & Smalla 2009). However, in certain environments, such as hypersaline soils, plant species identity plays only a minor role compared to soil salinity in shaping microbial community structure (Borruso et al. 2014); this result has been confirmed by studies of desert soils (Li et al. 2013). Overall, both plant species identity and soil type can be important, depending on abiotic and biotic conditions (Berg & Smalla 2009).

While many studies have investigated the effects of plant species identity and various soil properties, as well as other factors (i.e. temperature, geographical distance), on structuring microbial communities in the rhizosphere, few studies have characterized the rhizobacterial communities associated with halophytic plants found in arid, saline environments. In a Mediterranean salt marsh in Southeastern Spain, a study of eight

halophytes (*Asteriscus maritimus*, *Arthrocnemum macrostachyum*, *Frankenia corymbosa*, *Halimione portulacoides*, *Limonium cossonianum*, *Limonium caesium*, *Lygeum spartum* and *Suaeda vera* Forsskal) examined the soil microbiological and biochemical properties of the rhizosphere, revealing that soil microbial activity and microbial-related soil properties, such as aggregate stability, were determined by the plant species. However, the community composition of the microbes was not examined (Caravaca et al. 2005). Another study recently compared the microbial communities associated with three dominant halophytes (*Aeluropus*, *Salicornia* and *Suaeda*) in a coastal region of India (Chaudhary et al. 2015). To date, in halophytic species occurring in arid desert environments, little is known regarding rhizobacterial community assembly and the relative contributions of rhizosphere effects versus salinity to this process.

In this study, the bacterial communities associated with five halophytes (*Halocnemum strobilaceum*, *Halostachys caspica*, *Limonium gmelinii*, *Lycium ruthenicum* and *Kalidium foliatum*) growing in arid, saline environments were characterized. The diversity and structure of rhizosphere bacterial communities was investigated using an Illumina MiSeq sequencing approach. The study goals were to: 1) compare the bacterial communities of the rhizosphere with those found in bulk soil samples (i.e. non-rhizosphere samples), in order to understand the effects of plant species identity on bacterial communities; 2) compare the rhizosphere community composition of the five halophytic species so as to look for similarities across species; and, 3) evaluate the relative contributions of plant species identity and soil salinity in structuring rhizosphere bacterial communities in arid, saline habitats.

## MATERIALS & METHODS

### Study area and sample collection

Soil samples were collected from the Ebinur Lake Wetland, Xinjiang, China (44.595°N, 83.552°E) in July 2017, following previously established protocols (Chaudhary et al. 2015; Edwards et al. 2015). The Ebinur Lake Nature Reserve is located at the western margin of the Gurbantunggut Desert in Xinjiang, China. Conditions are windy in the Reserve, which has a typical dry, continental climate, with an annual average precipitation of 105 mm and evaporation of 1315 mm. Soils in the Reserve are highly salinized and alkalized, with an average electrical conductivity (EC) of 5.41 mS/cm and pH of 8.77 in surface soils 0–10 cm deep; the mean soil water content is 7.19% (Zhang et al. 2014). As such, many halophytic species grow in this region. In this study, five co-occurring halophytic plants species, four shrubs (*Halocnemum strobilaceum*, *Halostachys caspica*, *Lycium ruthenicum* and *Kalidium foliatum*,) and one perennial herb (*Limonium gmelinii*), were selected for study.

Thirty samples were collected in total: 15 from the rhizosphere and 15 paired bulk soil samples. Each rhizosphere sample came from a different plant; sampled individuals were distributed within a 1 km radius of the GPS coordinates provided above (Fig S1). Three healthy individuals were randomly selected from each species to be sampled. Plant roots were dug up using a shovel, in order to collect all roots to a depth of approximately 35–45 cm. Excess soil was manually shaken from the roots, but any soil still attached after shaking (a layer ~1 mm thick) was retained for study. Each root sample was immediately placed into a sterile flask with 30 ml of sterile Phosphate Buffered Saline (PBS) solution (137 mmol/L NaCl, 2.7 mmol/L KCl, 8.5 mmol/L Na<sub>2</sub>HPO<sub>4</sub>, 1.5 mmol/L KH<sub>2</sub>PO<sub>4</sub>, pH 7.3). Bulk soil samples were collected from sites 30–40 cm away from the roots of a given plant; soils were collected to a depth of approximately 40 cm. About 100 g of fresh soil from each sample was stored in a sterile plastic bag; these were immediately transported back to the lab on ice. In the lab, the flasks containing root samples were stirred vigorously with sterile forceps to clean all the soil from the root surfaces. This soil was then poured into a 50 ml sterile Falcon tube ready for DNA extraction.

# **Soil chemical analysis**

Soil samples were dried in a hot air oven at 105°C for 48 hours to determine the soil water content (SWC). After drying, samples were ground and sieved through 2 mm mesh. The electrical conductivity (EC) and pH were measured in a 1:2.5 (w:v) soil to water mix. The total organic carbon (TOC) and soil organic matter (SOM) were estimated using a spectrophotometer, after oxidizing soil samples with K<sub>2</sub>Cr<sub>2</sub>O<sub>4</sub> (Yang 1987). The total nitrogen (TON) was determined using the Kjeldahl method (Honda 1962). To measure available phosphorus (AP), samples were digested with HClO<sub>4</sub>-H<sub>2</sub>SO<sub>4</sub>, and then the Mo-Sb colorimetric method was used for quantification (Agrochemistry Committee of the Chinese Soil Society 1983).

# **DNA extraction, amplification and sequencing**

Rhizosphere samples were concentrated by pipetting 1 mL of the PBS/soil mix into a 2 mL sterile tube and centrifuging for 1 min at 10,000 g. The supernatant was discarded leaving only the soil fraction behind. About 0.2 g of soil (wet weight) from each bulk sample was transferred to a 2 mL sterile tube. Then, the total genomic DNA was extracted using an E.Z.N.A<sup>TM</sup> Mag-Bind Soil DNA Kit (OMEGA) following the manufacturer's instructions. DNA samples were inspected on a 1.0% agarose gel and quantified using a Nanodrop 2000 spectrophotometer (Nanodrop Technologies, Wilmington DE). The bacterial 16S rDNA V3–V4 region was amplified and sequenced for analysis. PCR products were visualized using electrophoresis on 1.5% agarose gels and purified using VAHTS<sup>TM</sup> DNA Clean Beads (Vazyme, Nanjing, China). Finally, about 10 ng of DNA from each sample was sequenced on the Illumina MiSeq platform by Sangon Technology Co., Ltd. (Shanghai, China). The sequence data has been submitted to the NCBI Sequence Read Archive database under accession number SRP129060.

# **Sequence preprocessing and OTU assignment**

Quality control of raw sequencing data was conducted following Schmieder & Edwards (Schmieder & Edwards 2011). Ambiguous bases with Phred quality score < 20 at the end of a read and fragments containing Ns were trimmed. Reads with length < 200 nucleotides (nt) were removed, and the remaining reads truncated to 400–450 nt sequences. Chimeric sequences were identified with UCHIME (Edgar et al. 2011) and discarded (109–1397 chimeras per sample). The filtered sequences were then clustered into OTUs at a 97% similarity level. A representative sequence from each OTU was selected for both taxonomic annotation using the Ribosomal Database Project (RDP) classifier (Wang et al. 2007) and also to BLAST against the Silva and NCBI databases (Quast et al. 2013). OTUs with an RDP classification threshold below 0.8 or with identity and coverage lower than 90% were marked as unclassified. Singletons and sequences aligning to the mitochondria or chloroplast were removed (4–1450 reads per sample). Finally, the number of sequences in each sample was normalized by random resampling to the smallest sample size (n=35000) prior to calculation of species diversity indices (Shannon, Simpson, Chao1, ACE, and Good's coverage) using Mothur ver 1.30.1 (Schloss et al. 2009). OTU richness was calculated using the *vegan* package ver. 2.1-10 (Dixon 2003) in R ver. 3.2. Rarefaction analysis was implemented in Mothur 1.30.1 and a rarefaction curve produced in R. Using R, a species accumulation curve was also constructed (with the *vegan* package) and diagrams depicting bacterial community structure (composition and relative abundance) at multiple taxonomic ranks (phylum, class, order, family and genus) were generated.

# **Statistical analysis**

One-way ANOVAs were used to test for differences in alpha diversity indices (OTU richness, Shannon diversity index and Good's coverage index) as well as soil physicochemical properties. A principal coordinates analysis

(PCoA) was performed on weighted UniFrac distances (using the *vegan* package) to compare community composition among samples. A UPGMA tree depicting clustering relationships among samples was produced based on Bray-Curtis beta diversity distance metrics. ANOSIM (999 permutations) and Adonis statistics, available in the *vegan* package, were calculated to evaluate differences in rhizosphere community composition among the five study species. Venn diagrams illustrating similarities/differences in OTU composition between samples were produced with the *Venn-Diagram* package, ver. 1.6.16. Furthermore, STAMP 2.1.3 (Parks et al. 2014) and LEfSe 1.1.0 (Segata et al. 2011) were implemented to identify differentially abundant groups among samples. Pearson's correlation coefficients were calculated between the community diversity and richness indices and each soil property, and also among soil properties. Correspondence analysis (CA) was performed using the *vegan* package to investigate links between community structure, soil properties and plant species.

## RESULTS

### Soil properties

The average soil water content (SWC) of bulk soils was  $16.40 \pm 4.57$  %. The electrical conductance (EC) was  $6.30 \pm 1.21$  mS/cm and pH  $8.14 \pm 0.27$ . The mean total organic carbon (TOC), soil organic matter (SOM), total nitrogen (TON) and available phosphorus (AP) were  $8.05 \pm 4.15$  g/kg,  $13.87 \pm 7.15$  g/kg,  $0.48 \pm 0.23$  g/kg and  $0.82 \pm 0.14$  g/kg, respectively. *Halocnemum strobilaceum* and *Halostachys caspica* soils had higher EC and lower TOC, SOM and TON compared to those of other species ( $P < 0.05$ ). The AP content in *Halocnemum strobilaceum* soils was significantly lower than for other species ( $P < 0.05$ ) (Table 1).

### Bacterial community diversity

In total, 1.83 Gb of raw sequence data was obtained from all samples; after quality controls, a total of 1.18 Gb of clean sequence reads were available for further analysis. Good's coverage for all samples was higher than 0.98 (Table S1). Rarefaction curves stabilized as the number of sequences increased (Fig. S2), suggesting that bacterial communities were reasonably well-characterized. Species accumulation curves nearly reached a plateau, where the number of OTUs did not increase with sample size, indicating that the sample size was sufficient for data analysis (Fig. S3). After removal of chimeras, plant sequences and singletons, a total of 1315341 reads were obtained from soil samples; these were grouped into 8087 OTUs. OTU richness was higher in rhizosphere versus bulk soil samples ( $P < 0.01$ ; Fig. 1). The number of OTUs identified in the rhizosphere of each species ranged from a minimum of 2342 (for *Halocnemum strobilaceum*) to a maximum of 4602 (*Limonium gmelinii*); within species, the number of OTUs present in all three replicates ranged from 317 to 729. In bulk soil samples, the number of OTUs detected for each species ranged from 1108 to 3688, and between 94 and 220 OTUs were common to all three replicates within species (Table S1).

Bacterial community diversity was higher in the rhizosphere versus bulk soil samples (ANOVA  $P < 0.01$ ) (Fig. 1). Rhizobacterial diversity did not differ among *Halocnemum strobilaceum*, *Halostachys caspica* and *Kalidium foliatum*, but was lower in these species than in *Limonium gmelinii* and *Lycium ruthenicum* ( $P < 0.05$ ). Diversity in *Lycium ruthenicum* bulk soil samples was higher than in the other four plant species ( $P < 0.05$ ). There were 1005, 1001, 677, 589 and 510 OTUs exclusive to the rhizobacterial communities associated with *Halostachys caspica*, *Limonium gmelinii*, *Lycium ruthenicum*, *Kalidium foliatum*, and *Halocnemum strobilaceum*, respectively; of these, 135, 187, 242, 88 and 63 OTUs, respectively, were found in all within-species replicates. Comparing across species, 242 OTUs were identified in all bulk soil samples and 647 OTUs in all rhizosphere samples; only 31 and 87 of these OTUs, respectively, were found in all three individuals of each species (Fig. 2). Abundant OTUs (i.e., those accounting for  $> 0.1\%$  of sequences) common to the



rhizospheres of all halophytes belonged to 16 genera, including *Acinetobacter*, *Aliifodinibius*, *Citrobacter*, *Deferrisoma*, *Exiguobacterium*, *Gracilimonas*, *Halomonas*, *Marinobacter*, *Pseudomonas*, *Thiopfundum* and others. Considering bulk soils, the most abundant OTUs came from four genera: *Acinetobacter*, *Citrobacter*, *Exiguobacterium* and *Pseudomonas* (Fig. S4).

### Bacterial community structure

A total of 36 phyla, 61 classes, 201 families and 617 genera were identified over all samples. In the bulk soil samples, Proteobacteria and Firmicutes were the dominant phyla. Gammaproteobacteria and Bacilli were the dominant classes, and *Acinetobacter*, *Bacillus*, *Citrobacter*, *Exiguobacterium* and *Pseudomonas* were the most abundant genera (Fig. 3). In the rhizosphere samples, Acidobacteria, Actinobacteria, Bacteroidetes, Chloroflexi, Firmicutes, Planctomycetes, Proteobacteria and Verrucomicrobia were the most abundant phyla. Actinobacteria, Bacilli, Cytophagia, Flavobacteriia, Planctomycetia, Sphingobacteriia, Alphaproteobacteria, Deltaproteobacteria and Gammaproteobacteria were the most abundant classes. The most common genera included *Acinetobacter*, *Aliifodinibius*, *Citrobacter*, *Deferrisoma*, *Exiguobacterium*, *Geminicoccus*, *Gp10*, *Gracilimonas*, *Halomonas*, *Marinobacter*, *Pseudomonas* and *Thiopfundum* (Fig. 3).

### Differences between rhizosphere and bulk soil samples

The most abundant bacterial groups in bulk soil communities were less common in rhizosphere communities, whereas some low abundance groups in the former were more common in the latter. At the phylum level, the Firmicutes and Proteobacteria was relatively less abundant in the rhizosphere, especially in the case of Firmicutes ( $P < 0.001$ ), whereas the Acidobacteria, Actinobacteria, Bacteroidetes, Chloroflexi, Planctomycetes and Verrucomicrobia were more abundant. A similar pattern was also observed at lower taxonomic ranks; however, overall, there were more abundant groups in the rhizosphere. Comparing bulk soil, the following genera were lower in abundance: *Acinetobacter*, *Citrobacter*, *Exiguobacterium*, *Halomonas* and *Pseudomonas*. Meanwhile, *Aciditerrimonas*, *Aliifodinibius*, *Deferrisoma*, *Fodinicurvata*, *Geminicoccus*, *Gp10*, *Gracilimonas*, *Marinobacter*, *Nitriliruptor*, *Planococcus* and *Thiopfundum* were more abundant in the rhizosphere samples ( $P < 0.01$ ) (Fig. 3).

Community composition differed significantly between rhizosphere and bulk soil samples (ANOSIM,  $R = 0.961$ ,  $P = 0.001$ ). A PCoA analysis illustrates these differences in community structure (Fig. 4a). Rhizosphere and bulk soil samples are separated on the first PCoA axis, which explained 82.33% of the variance. Bulk soil samples clustered together, indicating a high degree of similarity among their bacterial communities. However, rhizosphere samples were less tightly aggregated than bulk soil samples. Consistent with the PCoA results, the UPGMA tree also distinguished rhizosphere communities from bulk soil communities (Fig. 4b).

### Community structure differences among the five halophytes

Significant variation in community structure was observed among the five species' rhizospheres ( $R^2 = 0.703$ ,  $P = 0.001$ ). However, pairwise differences between each species pair, though relatively large, were not significant ( $P > 0.05$ ) (Table 2). The rhizobacterial community of *Lycium ruthenicum* was most similar to that of *Limonium gmelinii*. Meanwhile, the *Halocnemum strobilaceum* rhizobacterial community was most dissimilar to those of other species. As determined by a LEfSe analysis, rhizobacterial community composition differed among the five halophytes (Fig. 5). The genera *Gimesia* and *Pelagibius* were significantly more abundant in the *Lycium ruthenicum* rhizosphere; *Albidovulum*, *Bauldia*, *Deferrisoma*, *Geminicoccus*, *Gp10* and *Thiohalomonas* were more common in the *Limonium gmelinii* rhizosphere, *Citrobacter*, *Exiguobacterium* and *Pseudomonas* in the *Halostachys caspica* rhizosphere, *Gracilimonas*, *Jiangella*, *Marinimicrobium*, *Planococcus* and *Pontibacter* in

the *Kalidium foliatum* rhizosphere, and *Fodinicurvata*, *Halomonas*, *Mesorhizobium* and *Salegentibacter* in the *Halocnemum strobilaceum* rhizosphere.

In contrast, an Adonis analysis found no differences in community composition among bulk soil samples from different species ( $R^2 = 0.300$ ,  $P = 0.413$ ). However, LEfSe analyses found differences in composition in bulk soil communities for four out of the five species (Fig. 6). The bacterial community associated with *Lycium ruthenicum* was relatively distinct from that of other species, mainly due to differences in the abundance of the following families: Acidimicrobiaceae, Anaerolineaceae, Alteromonadaceae, Bacillaceae, Chromatiaceae, Demequinaceae, Planctomycetaceae, Puniceicoccaceae, Rhodobiaceae and Sprospiraceae. . Meanwhile, *Kalidium foliatum* communities were indistinguishable from those of the other four species.

### Correlations between bacterial diversity, community structure and soil properties

Relationships between microbial community diversity and structure, and soil bio-chemical properties were assessed with Pearson correlation coefficients and canonical correspondence analysis (CCA). The soil TOC, TON and AP were all strongly positively correlated, while EC was negatively correlated with all other soil variables except pH. Bacterial community diversity and richness were positively correlated with soil TON and AP, but negatively correlated with EC (Table 3) The first canonical axis (CCA1) was negatively correlated with EC, while the second canonical axis (CCA2) was positively correlated with SWC, but negatively correlated with EC. The SWC, TOC, SOM, TON and AP were more important in determining the bacterial community composition (as represented by longer arrows) than pH and EC, indicating that EC played only a minor role in the shaping of community structure. Rhizosphere communities from *Lycium ruthenicum* and *Limonium gmelinii* samples were positively associated with higher SWC, TON and AP. Meanwhile, *Halostachys caspica* rhizosphere communities were positively correlated with EC, and *Halocnemum strobilaceum* communities negatively correlated (Fig. 7a). Considering the bulk soil samples, CCA1 was positively correlated with EC and negatively correlated with the TOC, TON and pH. The second axis (CCA2) was positively correlated with AP and SWC. Apart from *Lycium ruthenicum*, the other species' samples were negatively correlated with SOM, TON and AP, but positively correlated with EC. The TON, AP and pH had a stronger influence on community structure than SOM or EC (Fig. 7b).

## DISCUSSION

### Bacterial community structure in saline soils as compared to other environments

Soil salinity has important effects on the distribution of plant communities, their composition and diversity (Xi et al. 2016), and may also affect soil bacterial diversity and community structure (Fang et al. 2016; Pavloudi et al. 2016). Plant communities in salinized habitats are dominated by halophytes, and the abundance and diversity of associated microbial communities, in saline or hypersaline terrestrial environments, is usually low (Takekawa et al. 2006; Jiang et al. 2007; Foti et al. 2008). In this study, bacterial communities were very low in richness and diversity compared to documented communities in forests, grasslands and agricultural areas (Rampelotto et al. 2013), maize crop soils (Garcia-Salamanca et al. 2013) and even other saline soils (Canfora et al. 2014); however, metrics were consistent with those found in other “extreme” hypersaline soils in semiarid Mediterranean regions (Canfora et al. 2015). The soils surrounding *Halocnemum strobilaceum* had the lowest diversity and richness, perhaps as a result of the relatively high soil EC and low nutrient availability. Soils associated with *Lycium ruthenicum* were the most diverse of all studied plant species. Considering that nutrient availability in *Lycium ruthenicum*-associated soils was only mediocre, other factors (i.e., plant community

composition) likely affected the soil bacterial community (Ravit et al. 2003; Cao et al. 2014). Indeed, based on field observations, the plant community surrounding focal *Lycium ruthenicum* individuals was more diverse and had higher percent cover than for other species.

Bulk soil samples had similar, relatively simple bacterial communities. Only four genera, *Acinetobacter*, *Citrobacter*, *Exiguobacterium* and *Pseudomonas*, were common, which differs from the bacterial communities of saline soils in Inner Mongolia (Borruso et al. 2014) and the Shandong Peninsula coast (Tian & Zhang 2017). Differences in community composition may be due to distinct climatic and/or soil properties among geographic regions (Ben-David et al. 2011). However, the dominance of Bacilli (Firmicutes) and  $\gamma$ -proteobacteria in the study region is consistent with previous studies (Tang et al. 2011; Borsodi et al. 2013), confirming that these two taxa are important in saline or hypersaline environments. A low level of structural differentiation was also found when comparing communities associated with different plant species, as determined by a LEfSe analysis. The highest differentiation was observed between *Lycium ruthenicum* and the other species; in contrast, *Kalidium foliatum* communities were indistinguishable from those of other species. This suggests that soil bacterial communities are highly similar in arid environments, at least on a small geographic scale. Microhabitat similarities may produce such similar bacterial communities, as low plant cover and a lack of plant litter are common to arid environments.

# **Bacterial community composition in the rhizosphere versus bulk soils**

Rhizosphere effects may be an important driving force shaping microbial communities and leading to compositional differences between the rhizosphere and the bulk soils (Morgan & Whipps 2001). In this study, bacterial communities differed between the rhizosphere and bulk soils, with higher community diversity and richness in the rhizosphere, consistent with previous studies (Avis et al. 2008; Borruso et al. 2014; Edwards et al. 2015; Chaudhary et al. 2017; Yang et al. 2017). Higher diversity may be a result of root exudates, which can raise nutrient concentrations (Li et al. 2014). While soil properties were not measured exhaustively here, the soil TOC and TON contents estimated in the rhizosphere of *Limonium gmelinii* and *Lycium ruthenicum* were approximately 5-7 times higher than in the bulk soils. However, it should be noted that bulk soils may have greater microbial richness than the rhizosphere in some cases (Shange et al. 2012; Carbonetto et al. 2014; Gomes et al. 2014; Tian & Zhang 2017). These divergent results suggest that the soil type and plant species identity have complex effects on bacterial communities, with the strength of their effects depending on abiotic and biotic conditions (Berg & Smalla 2009).

In addition to diversity differences, the composition of bacterial communities also differed between the rhizosphere and bulk soil samples, as revealed by PCoA and cluster analysis. Compared to bulk soils, the rhizosphere communities had more groups with relative abundance > 1%, such as the following phyla: Actinobacteria, Bacteroidetes, Firmicutes, Planctomycetes and Proteobacteria. Dominance by a greater number of groups has also been reported in other saline ecosystems, both marine and terrestrial (Tang et al. 2011; Marasco et al. 2013; Oliveira et al. 2014; Soussi et al. 2016). At the genus level, many genera were abundant in the rhizosphere, for example *Acinetobacter*, *Bacillus*, *Citrobacter*, *Deferrisoma*, *Exiguobacterium*, *Haliea*, *Halomonas*, *Marinimicrobium*, *Marinobacter*, *Methylohalomonas*, *Microbulbifer*, *Planococcus*, *Pseudomonas* and *Thiopfundum* (Fig. 3, Fig. S5). Meanwhile, the richness of groups common in bulk soils (i.e.  $\gamma$ -proteobacteria and Firmicutes) was reduced in the rhizosphere; for example, the abundance of Firmicutes was about 80% lower. The low abundance of Firmicutes in the rhizosphere has been reported many times; see one such case study in barley, where Firmicutes is almost excluded from the rhizosphere (Bulgarelli et al. 2015).

Although  $\gamma$ -proteobacteria were less abundant in the rhizosphere, as observed by a decrease in the number of occurrences of *Acinetobacter*, *Citrobacter* and *Pseudomonas* species,  $\gamma$ -proteobacteria remained the most abundant class, as found in many plant-associated bacterial communities (Mukhtar et al. 2017).

Shifts in bacterial community composition in bulk soils versus the rhizosphere may be the consequence of active selection by plants (Kowalchuk et al. 2002). As many endophytes and bacteria colonizing root surfaces have beneficial effects, such as nitrogen fixation, phytohormone production, nutrient supply and pathogen suppression (Rosenblueth & Martinez-Romero 2006; Hardoim et al. 2008), they typically promote plant growth and can alleviate salt stress in halophytes (Ali et al. 2015). Some *Microbulbifer* and *Planococcus* species have the ability to degrade complex hydrocarbons (See-Too et al. 2017). Meanwhile, *Bacillus*, *Exiguobacterium*, *Halomonas*, *Planococcus* and *Pseudomonas* can generate 1-aminocyclopropane-1- carboxylic acid (ACC) deaminase to convert the ethylene precursor ACC into ammonia and  $\alpha$ -ketobutyrate; this has the effect of lowering the ethylene concentration within plant tissues, reducing its constraining effect on root elongation and general plant growth (Siddikee et al. 2010). Moreover, some *Bacillus*, *Halomonas* and *Pseudomonas* species in can produce indole-3-acetic acid (IAA) to confer salt tolerance (Tiwari et al. 2011). *Halomonas* species are able to tolerate a wide set of abiotic stresses, and promote plant growth via IAA production, phosphate solubilization, nitrogen fixation and more (Mapelli et al. 2013). In rhizobacterial communities, *Bacillus* and *Pseudomonas* species are critically important for plant growth; these species have multiple functional activities, including phosphate solubilization, phytopathogen inhibition (Prashar et al. 2014) and auxin production. Why these species are less abundant in the rhizosphere (versus bulk soils) remains unclear. It could be caused by competition among rhizobacteria, but this hypothesis needs verification. Moreover, the effects of many other common genera in the rhizosphere (whether beneficial, harmful or neutral), as well their inter-relationships (mutualistic or competitive) with plant growth promoting species, need further investigation.

### **Differences among halophytes in rhizobacterial communities and relationship to soil properties**

A number of studies have demonstrated that microbial community composition is plant species specific (Andreote et al. 2009; Poli et al. 2016), leading to unique rhizobacterial communities among species. Here, five halophyte-associated rhizobacterial communities varied both in diversity (ANOVA  $P < 0.01$ ) and structure (Adonis  $R^2 = 0.703$ ,  $P = 0.001$ ). Rhizobacterial diversity was lower in *Halostachys caspica*, *Halocnemum strobilaceum* and *Kalidium foliatum* than in *Limonium gmelinii* and *Lycium ruthenicum* ( $P < 0.05$ ). While community composition also seemed to vary, differences were not significant (Table 2). *Halomonas* was the dominant genera in *Halocnemum strobilaceum* communities (32.4% of total abundance), a significantly higher proportion than in the other plant-associated communities; this trend is consistent with previous studies (Al-Mailem et al. 2010; Marasco et al. 2016), including of other halophytes (Borsodi et al. 2015). In the *Halostachys caspica* rhizosphere, *Exiguobacterium* was the most common genus, followed by *Citrobacter*, *Acinetobacter* and *Pseudomonas*. Meanwhile, in the *Kalidium foliatum* rhizosphere, *Halomonas*, *Exiguobacterium* and *Gracilimonas* were most abundant. The rhizobacterial communities of *Limonium gmelinii* and *Lycium ruthenicum* were highly similar (Adonis  $R^2 = 0.324$ ,  $P = 0.108$ ), perhaps as a result of similar soil properties. The relatively high abundance of *Exiguobacterium* in the rhizosphere is unique to this study, suggesting that it is potentially important in the study region, but this requires further investigation.

Furthermore, among replicates for a given plant species, only a small proportion of OTUs occurred in all three. Interestingly, these shared OTUs were usually highly abundant overall, whereas OTUs that were found only in one or two individuals were mostly rare OTUs (with a relative abundance less than five). Thus, bacterial

communities may have individual specificity, potentially influenced by a plant's genotype, age or health (Zhu et al. 2013; Belimov et al. 2015). The OTUs found across all five plant species belong to 16 genera in total, and some, such as *Deferrisoma*, *Exiguobacterium*, *Geminicoccus*, *Gracilimonas* and *Marinimicrobium*, were also included in the taxa identified in the LEfSe. As successful colonizers of halophyte plants in saline-arid lands, these may be good plant growth promotion rhizobacteria (PGPR) candidates.

Soil properties are also important determinants of soil microbial communities (Szymanska et al. 2016a; Rathore et al. 2017). In particular, soil salinity is considered a primary environmental factor, directly or indirectly driving the composition and diversity of prokaryotic communities (Marasco et al. 2016; Zhong et al. 2016). In this study, soil EC had a positive influences on the bulk soil bacterial communities, and other soil factors also shaped bacterial community composition. Soil nutrients (TON, TOC and AP) were most important for *Lycium ruthenicum*. Similarly, a recent study found that variation in prokaryotic community structure was significantly correlated with the TON and  $PO_4^{3-}$  concentration (Zhong et al. 2016). In the rhizosphere, EC was negatively correlated with rhizobacterial community diversity and structure, though its influence on community structure was only minor as revealed by the CCA. In contrast, the effects of soil nutrients, such as TOC, SOM, TON and AP, were much stronger than those of EC or soil pH. The low diversity and compositional variation seen in *Halostachys caspica* and *Halocnemum strobilaceum* rhizobacterial communities were negatively correlated with soil TOC, SOM, TON and AP. These results imply that, when salinity levels are similar among habitats, the influence of salinity on the rhizosphere microbial community structure is relatively weak. In these cases, rhizobacterial community structure is more strongly influenced by the host plant identity and soil nutrient content, whereas salinity has a minor effect.

Overall, variation in rhizobacterial community structure among the five study plant species was significant, though pairwise differences did not reach significance despite the large number of OTUs that were unique to each species. This may be explained by the relatively low total abundance of rare OTUs, as well as the low proportion of shared OTUs in a given plant species. While abundant OTUs were shared across all five plant species (and accounted for a large proportion of total richness in each), they differed in relative abundance. This suggests that, in similar saline habitats, plants tend to select similar bacterial species to colonize roots, perhaps as a consequence of adaptation to salinity stress; these species may act to promote plant growth or alleviate salt stress. Coevolution between microorganisms and associated plant species has been addressed in leguminous *Rhizobium* species (Wang et al. 2018). Community differences (especially in the richness of abundant groups) may be closely related to differences in root exudates or plant litter between plant species (Chaudhary et al. 2015), as supported by the results here. However, the effects of other factors, such as ion concentration, and interactions between microbiomes within a community (Chen et al. 2013; Poosakkannu et al. 2017) cannot be excluded.

## CONCLUSIONS

The present study investigated the composition and diversity of rhizobacterial communities in five co-occurring halophytic species growing in salinized, arid desert soils within the Ebinur Lake Wetland Reserve in Northwestern China. Significant differences were found between the rhizosphere and bulk soil communities, both in diversity and bacterial composition. Diversity was higher in the rhizosphere than in the bulk soils. Abundant taxonomic groups (from phylum to genus) in the rhizosphere were much more diverse than in bulk soils. Actinobacteria, Bacteroidetes, Firmicutes, Planctomycetes and Proteobacteria were the most abundant

phyla in the rhizosphere, while Firmicutes and Proteobacteria were common in bulk soils. Comparing among species, significant differences in rhizobacterial diversity and identity were observed. The diversity of *Halostachys caspica*, *Halocnemum strobilaceum* and *Kalidium foliatum* associated communities was lower than that of *Limonium gmelinii* and *Lycium ruthenicum* communities. Furthermore, the composition of *Halostachys caspica* and *Halocnemum strobilaceum* communities was very different from that of *Limonium gmelinii* and *Lycium ruthenicum* communities. Thus, plant species identity can have important effects on root-associated bacterial communities. Diversity was positively correlated with soil nutrients, including TOC, SOM, TON and AP, but negatively correlated with EC, though the effects of EC were much lower than those of soil nutrient content. In conclusion, halophytic plant species played an important role in shaping associated rhizosphere bacterial communities. Furthermore, when salinity levels were constant, soil nutrients emerged as key factors structuring bacterial communities. These results provide insight into the nature of halophyte microbial communities in arid regions, as well as the factors shaping these communities. However, pairwise differences among the five rhizobacterial communities were not significant, despite some evidence for differentiation among plant species. Further studies involving more halophyte species and individuals per species are necessary to elucidate plant species identity effects on the rhizosphere for co-occurring halophytes.

# ACKNOWLEDGMENTS

We would like to thank LetPub, Elizabeth Tokarz at the Yale University and Dr. Emily Drummond at the University of British Columbia for their assistance with English language and grammatical editing of the manuscript. We would also like to thank Dr. Rodrigo Taketani and the anonymous reviewers for their comments and suggestions that improved the quality of our manuscript.

# ADDITIONAL INFORMATION AND DECLARATIONS

**Fig. S1** Collection locations of soil samples used in the present study.

**Fig. S2** Rarefaction curves for bacterial OTUs, clustering at 97% sequence similarity. Error bars indicate standard errors.

**Fig. S3** Species accumulation curves for soil samples. The horizontal bars within boxes represent the medians, while the tops and bottoms of the boxes represent 75th and 25th quartiles, respectively. Plus signs (+) indicate outliers.

**Fig. S4** Phylogeny and taxonomic classification of abundant OTUs shared among either rhizosphere or bulk soil communities.

**Fig. S5** Relative abundance of common phyla and genera in bulk soil and rhizosphere bacterial communities.

**Table S1** Alpha diversity indices of bacterial communities in rhizosphere and bulk soil samples.

# TABLE CAPTIONS

**Table 1** Soil characteristics of bulk soil samples from five halophytes.

**Table 2** Adonis analysis of bacterial community composition for bulk soil and rhizosphere samples.

**Table 3** Pearson correlation coefficients among soil chemical properties, and between soil properties and

community diversity.

# FIGURE LEGENDS

**Fig. 1** Multiple comparisons of OTU richness (a) and  $\alpha$ -diversity (Shannon's index) (b) between rhizospheric and bulk soil bacterial communities, and among plant species within soil type. Within boxes, horizontal bars indicate medians, while the tops and bottoms of boxes illustrate 75th and 25th quartiles, respectively. Small circles represent outliers in bulk samples. G, B, J, S, Z indicate bulk soil samples from *Lycium ruthenicum*, *Limonium gmelinii*, *Halocnemum strobilaceum*, *Halostachys caspica*, and *Kalidium foliatum*, respectively, while GQ, BX, JM, SM and ZZ denote the corresponding rhizosphere samples from each species. Significance levels (p values) are provided for among-species comparisons within soil type (bulk versus rhizosphere), as well as between soil types.

**Fig. 2** Venn diagram showing the OTUs shared among different samples. a: rhizosphere samples, b: bulk soil samples. Information on OTU numbers is provided as follows: the total number of OTUs detected across all three replicates for each species (the number of OTUs shared among the three replicates).

**Fig. 3** Diagrams depicting community composition and relative abundance at different taxonomic levels. a: phylum, b: class, and c: genus.

**Fig. 4** Principal coordinate analysis (PCoA) (a) and UPGMA clustering diagram (b) of soil samples. G, B, J, S and Z represent bulk soil samples from *Lycium ruthenicum*, *Limonium gmelinii*, *Halocnemum strobilaceum*, *Halostachys caspica* and *Kalidium foliatum*, respectively, while GQ, BX, JM, SM and ZZ represent rhizosphere samples associated with these species.

**Fig. 5** LEfSe analysis at multiple taxonomic levels comparing rhizobacterial community composition for five focal plant species. (a) Cladogram illustrating the taxonomic groups explaining the most variation among rhizobacterial communities. Each ring represents a taxonomic level, with phylum (p), class (c), order (o), family (f) and genus (g) emanating from the center to the periphery. Each circle is a taxonomic unit found in the dataset, with circles or nodes shown in colors (other than yellow) indicating where a taxon was significantly more abundant. (b) Histogram of the LDA scores computed for groups with differential abundance among the rhizobacterial communities of the five plant species.

**Fig. 6** LEfSe analysis of bacterial community composition for bulk soil samples from four plant species. (a) Cladogram illustrating the taxonomic groups that explain the most variation among the bacterial communities. (b) Histogram of the LDA scores computed for groups with differential abundance among the bacterial communities of the four plant species. *Kalidium foliatum*-associated communities did not form a separate group (from other plant species' associated communities) and, therefore, these are absent from the cladogram and histogram.

**Fig. 7** Canonical correspondence analysis (CCA) of the effect of soil properties on bacterial community structure in the rhizosphere (a) and bulk soil samples (b). The soil properties are indicated with arrows, and include soil pH, electrical conductivity (EC), total organic carbon (TOC), soil organic matter (SOM), total nitrogen (TON), phosphorous (AP) and soil water content (SWC). The percentage of variation explained by each axis is provided. GQ, BX, JM, SM and ZZ represent rhizosphere samples associated with *Lycium ruthenicum*, *Limonium gmelinii*, *Halocnemum strobilaceum*, *Halostachys caspica* and *Kalidium foliatum*, respectively. G, B, J, S and Z represent bulk soil samples from *Lycium ruthenicum*, *Limonium gmelinii*, *Halocnemum strobilaceum*, *Halostachys caspica* and *Kalidium foliatum*, respectively.



# REFERENCES

- Agrochemistry Committee of the Chinese Soil Society. 1983. Conventional analytical methods for soil agrochemistry. *Beijing: Science Press*.
- Al-Mailem DM, Sorkhoh NA, Marafie M, Al-Awadhi H, Eliyas M, and Radwan SS. 2010. Oil phytoremediation potential of hypersaline coasts of the Arabian Gulf using rhizosphere technology. *Bioresource Technology* 101:5786-5792. 10.1016/j.biortech.2010.02.082
- Ali Z, Ullah N, Naseem S, Inam-Ul-Haq M, and Jacobsen HJ. 2015. Soil bacteria conferred a positive relationship and improved salt stress tolerance in transgenic pea (*Pisum sativum* L.) harboring Na<sup>+</sup>/H<sup>+</sup> antiporter. *Turkish Journal of Botany* 39:962-+. 10.3906/bot-1505-50
- Andreote FD, Rossetto PB, Mendes R, Avila LA, Labate CA, Pizzirani-Kleiner AA, Azevedo JL, and Araujo WL. 2009. Bacterial community in the rhizosphere and rhizoplane of wild type and transgenic eucalyptus. *World Journal of Microbiology & Biotechnology* 25:1065-1073. 10.1007/s11274-009-9990-9
- Avis TJ, Gravel V, Antoun H, and Tweddell RJ. 2008. Multifaceted beneficial effects of rhizosphere microorganisms on plant health and productivity. *Soil Biology & Biochemistry* 40:1733-1740. 10.1016/j.soilbio.2008.02.013
- Belimov AA, Puhalsky IV, Safronova VI, Shaposhnikov AI, Vishnyakova MA, Semenova EV, Zinovkina NY, Makarova NM, Wenzel W, and Tikhonovich IA. 2015. Role of plant genotype and soil conditions in symbiotic plant-microbe interactions for adaptation of plants to Cadmium-polluted soils. *Water Air and Soil Pollution* 226. Artn 26410.1007/S11270-015-2537-9
- Ben-David EA, Zaady E, Sher Y, and Nejdat A. 2011. Assessment of the spatial distribution of soil microbial communities in patchy arid and semi-arid landscapes of the Negev Desert using combined PLFA and DGGE analyses. *Fems Microbiology Ecology* 76:492-503. 10.1111/j.1574-6941.2011.01075.x
- Berg G, and Smalla K. 2009. Plant species and soil type cooperatively shape the structure and function of microbial communities in the rhizosphere. *Fems Microbiology Ecology* 68:1-13. 10.1111/j.1574-6941.2009.00654.x
- Borruso L, Bacci G, Mengoni A, De Philippis R, and Brusetti L. 2014. Rhizosphere effect and salinity competing to shape microbial communities in *Phragmites australis* (Cav.) Trin. ex-Steud. *Fems Microbiology Letters* 359:193-200. 10.1111/1574-6968.12565
- Borsodi AK, Barany A, Krett G, Marialigeti K, and Szili-Kovacs T. 2015. Diversity and ecological tolerance of bacteria isolated from the rhizosphere of halophyton plants living nearby Kiskunsag soda ponds, Hungary. *Acta Microbiologica Et Immunologica Hungarica* 62:183-197. 10.1556/030.62.2015.2.8
- Borsodi AK, Felfoldi T, Mathe I, Bogнар V, Knab M, Krett G, Jurecska L, Toth EM, and Marialigeti K. 2013. Phylogenetic diversity of bacterial and archaeal communities inhabiting the saline Lake Red located in Sovata, Romania. *Extremophiles* 17:87-98. 10.1007/s00792-012-0496-2
- Bulgarelli D, Garrido-Oter R, Munch PC, Weiman A, Droge J, Pan Y, McHardy AC, and Schulze-Lefert P. 2015. Structure and function of the bacterial root microbiota in wild and domesticated barley. *Cell Host & Microbe* 17:392-403. 10.1016/j.chom.2015.01.011
- Canfora L, Bacci G, Pinzari F, Lo Papa G, Dazzi C, and Benedetti A. 2014. Salinity and bacterial diversity: to what extent does the concentration of salt affect the bacterial community in a saline soil? *Plos One* 9:e106662. 10.1371/journal.pone.0106662

- Canfora L, Lo Papa G, Antisari LV, Bazan G, Dazzi C, and Benedetti A. 2015. Spatial microbial community structure and biodiversity analysis in "extreme" hypersaline soils of a semiarid Mediterranean area. *Applied Soil Ecology* 93:120-129. 10.1016/j.apsoil.2015.04.014
- Cao D, Shi FC, Koike T, Lu ZH, and Sun JK. 2014. Halophyte plant communities affecting enzyme activity and microbes in saline soils of the Yellow River delta in China. *Clean-Soil Air Water* 42:1433-1440. 10.1002/clen.201300007
- Caravaca F, Aiguacil MM, Torres P, and Roldan A. 2005. Plant type mediates rhizospheric microbial activities and soil aggregation in a semiarid Mediterranean salt marsh. *Geoderma* 124:375-382. 10.1016/j.geoderma.2004.05.010
- Carbonetto B, Rascovan N, Alvarez R, Mentaberry A, and Vazquez MP. 2014. Structure, Composition and metagenomic profile of soil microbiomes associated to agricultural land use and tillage systems in Argentine pampas. *Plos One* 9. ARTN e9994910.1371/journal.pone.0099949
- Chaudhary DR, Gautam RK, Yousuf B, Mishra A, and Jha B. 2015. Nutrients, microbial community structure and functional gene abundance of rhizosphere and bulk soils of halophytes. *Applied Soil Ecology* 91:16-26. 10.1016/j.apsoil.2015.02.003
- Chaudhary DR, Rathore AP, Kumar R, and Jha B. 2017. Spatial and halophyte-associated microbial communities in intertidal coastal region of India. *International Journal of Phytoremediation* 19:478-489. 10.1080/15226514.2016.1244168
- Chen YL, Chen BD, Hu YJ, Li T, Zhang X, Hao ZP, and Wang YS. 2013. Direct and indirect influence of arbuscular mycorrhizal fungi on abundance and community structure of ammonia oxidizing bacteria and archaea in soil microcosms. *Pedobiologia* 56:205-212. 10.1016/j.pedobi.2013.07.003
- Dixon P. 2003. VEGAN, a package of R functions for community ecology. *Journal of Vegetation Science* 14:927-930. DOI 10.1111/j.1654-1103.2003.tb02228.x
- Edgar RC, Haas BJ, Clemente JC, Quince C, and Knight R. 2011. UCHIME improves sensitivity and speed of chimera detection. *Bioinformatics* 27:2194-2200. 10.1093/bioinformatics/btr381
- Edwards J, Johnson C, Santos-Medellin C, Lurie E, Podishetty NK, Bhatnagar S, Eisen JA, and Sundaresan V. 2015. Structure, variation, and assembly of the root-associated microbiomes of rice. *Proceedings of the National Academy of Sciences of the United States of America* 112:E911-E920. 10.1073/pnas.1414592112
- English JP, and Colmer TD. 2011. Salinity and waterlogging tolerances in three stem-succulent halophytes (Tecticornia species) from the margins of ephemeral salt lakes. *Plant and Soil* 348:379-396. 10.1007/s11104-011-0924-6
- Fang TT, Pan RS, Jiang J, He F, and Wang H. 2016. Effect of salinity on community structure and naphthalene dioxygenase gene diversity of a halophilic bacterial consortium. *Frontiers of Environmental Science & Engineering* 10. ARTN 1610.1007/s11783-016-0888-0
- Flowers TJ. 2004. Improving crop salt tolerance. *Journal of Experimental Botany* 55:307-319. 10.1093/jxb/erh003
- Foti MJ, Sorokin DY, Zacharova EE, Pimenov NV, Kuenen JG, and Muyzer G. 2008. Bacterial diversity and activity along a salinity gradient in soda lakes of the Kulunda Steppe (Altai, Russia). *Extremophiles* 12:133-145. 10.1007/s00792-007-0117-7
- Garcia-Salamanca A, Molina-Henares MA, van Dillewijn P, Solano J, Pizarro-Tobias P, Roca A, Duque E,

- and Ramos JL. 2013. Bacterial diversity in the rhizosphere of maize and the surrounding carbonate-rich bulk soil. *Microbial Biotechnology* 6:36-44. 10.1111/j.1751-7915.2012.00358.x
- Gomes NCM, Cleary DFR, Pires ACC, Almeida A, Cunha A, Mendonca-Hagler LCS, and Smalla K. 2014. Assessing variation in bacterial, composition between the rhizospheres of two mangrove tree species. *Estuarine Coastal and Shelf Science* 139:40-45. 10.1016/j.ecss.2013.12.022
- Hardoim PR, van Overbeek LS, and van Elsas JD. 2008. Properties of bacterial endophytes and their proposed role in plant growth. *Trends in Microbiology* 16:463-471. 10.1016/j.tim.2008.07.008
- Honda C. 1962. Rapid procedure for determination of nitrogen in soil by kjeldahl method. *J Sci Soil Manure* 33:195-200.
- Jha B, Gontia I, and Hartmann A. 2012. The roots of the halophyte *Salicornia brachiata* are a source of new halotolerant diazotrophic bacteria with plant growth-promoting potential. *Plant and Soil* 356:265-277. 10.1007/s11104-011-0877-9
- Jiang HC, Dong HL, Yu BS, Liu XQ, Li YL, Ji SS, and Zhang CLL. 2007. Microbial response to salinity change in Lake Chaka, a hypersaline lake on Tibetan plateau. *Environmental Microbiology* 9:2603-2621. 10.1111/j.1462-2920.2007.01377.x
- Kowalchuk GA, Buma DS, de Boer W, Klinkhamer PGL, and van Veen JA. 2002. Effects of above-ground plant species composition and diversity on the diversity of soil-borne microorganisms. *Antonie Van Leeuwenhoek International Journal of General and Molecular Microbiology* 81:509-520. Doi 10.1023/A:1020565523615
- Li K, Liu RY, Zhang HX, and Yun JL. 2013. The diversity and abundance of bacteria and oxygenic phototrophs in saline biological desert crusts in Xinjiang, Northwest China. *Microbial Ecology* 66:40-48. 10.1007/s00248-012-0164-1
- Li XZ, Rui JP, Mao YJ, Yannarell A, and Mackie R. 2014. Dynamics of the bacterial community structure in the rhizosphere of a maize cultivar. *Soil Biology & Biochemistry* 68:392-401. 10.1016/j.soilbio.2013.10.017
- Mapelli F, Marasco R, Rolli E, Barbato M, Cherif H, Guesmi A, Ouzari I, Daffonchio D, and Borin S. 2013. Potential for plant growth promotion of rhizobacteria associated with salicornia growing in Tunisian hypersaline soils. *Biomed Research International*. Artn 24807810.1155/2013/248078
- Marasco R, Mapelli F, Rolli E, Mosqueira MJ, Fusi M, Bariselli P, Reddy M, Cherif A, Tsiamis G, Borin S, and Daffonchio D. 2016. *Salicornia strobilacea* (synonym of *Halocnemum strobilaceum*) grown under different tidal regimes selects rhizosphere bacteria capable of promoting plant growth. *Frontiers in Microbiology* 7. ARTN 128610.3389/fmicb.2016.01286
- Marasco R, Rolli E, Fusi M, Cherif A, Abou-Hadid A, El-Bahairy U, Borin S, Sorlini C, and Daffonchio D. 2013. Plant growth promotion potential is equally represented in diverse grapevine root-associated bacterial communities from different biopedoclimatic environments. *Biomed Research International*. Artn 49109110.1155/2013/491091
- Morgan JAW, and Whipps JM. 2001. Methodological approaches to the study of rhizosphere carbon flow and microbial population dynamics. In: Pinton A, Varanini Z, Nannipieri P (eds) *The rhizosphere: biochemistry and organic substances at the soil-plant interface Marcel Dekker, New York*:373-409.
- Mukhtar S, Ishaq A, Hassan S, Mehnad S, Mirza MS, and Malik KA. 2017. Comparison of microbial communities associated with halophyte (*Salsola stocksii*) and non-halophyte (*Triticum aestivum*)

- 633 using culture-independent approaches. *Polish Journal of Microbiology* 66:353-364.
- 634 Nabti E, Sahnoune M, Adirad S, Van Dommelen A, Ghoul M, Schmid M, and Hartmann A. 2007. A halophilic  
635 and osmotolerant *Azospirillum brasilense* strain from Algerian soil restores wheat growth under saline  
636 conditions. *Engineering in Life Sciences* 7:354-360. 10.1002/elsc.200720201
- 637 Oliveira V, Gomes NCM, Cleary DFR, Almeida A, Silva AMS, Simoes MMQ, Silva H, and Cunha A. 2014.  
638 Halophyte plant colonization as a driver of the composition of bacterial communities in salt marshes  
639 chronically exposed to oil hydrocarbons. *Fems Microbiology Ecology* 90:647-662. 10.1111/1574-  
640 6941.12425
- 641 Oosterbaan RJ. 1988. Effectiveness and social/environmental impacts of irrigation projects: a critical review.  
642 In ILRI Annual Report 1988, international institute for land reclamation and improvement,.  
643 Wageningen, The Netherlands:18-34.
- 644 Parks DH, Tyson GW, Hugenholtz P, and Beiko RG. 2014. STAMP: statistical analysis of taxonomic and  
645 functional profiles. *Bioinformatics* 30:3123-3124. 10.1093/bioinformatics/btu494
- 646 Pavloudi C, Oulas A, Vasileiadou K, Sarropoulou E, Kotoulas G, and Arvanitidis C. 2016. Salinity is the  
647 major factor influencing the sediment bacterial communities in a Mediterranean lagoonal complex  
648 (Amvrakikos Gulf, Ionian Sea). *Marine Genomics* 28:71-81. 10.1016/j.margen.2016.01.005
- 649 Pii Y, Borruso L, Brusetti L, Crecchio C, Cesco S, and Mimmo T. 2016. The interaction between iron  
650 nutrition, plant species and soil type shapes the rhizosphere microbiome. *Plant Physiology and*  
651 *Biochemistry* 99:39-48. 10.1016/j.plaphy.2015.12.002
- 652 Poli A, Lazzari A, Prigione V, Voyron S, Spadaro D, and Varese GC. 2016. Influence of plant genotype on the  
653 cultivable fungi associated to tomato rhizosphere and roots in different soils. *Fungal Biology* 120:862-  
654 872. 10.1016/j.funbio.2016.03.008
- 655 Poosakkannu A, Nissinen R, and Kytoviita MM. 2017. Native arbuscular mycorrhizal symbiosis alters foliar  
656 bacterial community composition. *Mycorrhiza* 27:801-810. 10.1007/s00572-017-0796-6
- 657 Prashar P, Kapoor N, and Sachdeva S. 2014. Rhizosphere: its structure, bacterial diversity and significance.  
658 *Reviews in Environmental Science and Bio-Technology* 13:63-77. 10.1007/s11157-013-9317-z
- 659 Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, Peplies J, and Glockner FO. 2013. The SILVA  
660 ribosomal RNA gene database project: improved data processing and web-based tools. *Nucleic Acids*  
661 *Research* 41:D590-D596. 10.1093/nar/gks1219
- 662 Ramadoss D, Lakkineni VK, Bose P, Ali S, and Annapurna K. 2013. Mitigation of salt stress in wheat  
663 seedlings by halotolerant bacteria isolated from saline habitats. *Springerplus* 2. Artn 610.1186/2193-  
664 1801-2-6
- 665 Rampelotto PH, Ferreira AD, Barboza ADM, and Roesch LFW. 2013. Changes in diversity, abundance, and  
666 structure of soil bacterial communities in Brazilian Savanna under different land use systems.  
667 *Microbial Ecology* 66:593-607. 10.1007/s00248-013-0235-y
- 668 Rathore AP, Chaudhary DR, and Jha B. 2017. Seasonal patterns of microbial community structure and enzyme  
669 activities in coastal saline soils of perennial halophytes. *Land Degradation & Development* 28:1779-  
670 1790. 10.1002/ldr.2710
- 671 Ravit B, Ehrenfeld JG, and Haggblom MM. 2003. A comparison of sediment microbial communities  
672 associated with *Phragmites australis* and *Spartina alterniflora* in two brackish wetlands of New Jersey.  
673 *Estuaries* 26:465-474. Doi 10.1007/Bf02823723

- Rodriguez-Blanco A, Sicardi M, and Frioni L. 2015. Plant genotype and nitrogen fertilization effects on abundance and diversity of diazotrophic bacteria associated with maize (*Zea mays* L.). *Biology and Fertility of Soils* 51:391-402. 10.1007/s00374-014-0986-8
- Rosenblueth M, and Martinez-Romero E. 2006. Bacterial endophytes and their interactions with hosts. *Molecular Plant-Microbe Interactions* 19:827-837. 10.1094/Mpmi-19-0827
- Ruppel S, Franken p, and Witzel K. 2013. Properties of the halophyte microbiome and their implications for plant salt tolerance. *Functional Plant Biology* 40:940–951.
- Schloss PD, Westcott SL, Ryabin T, Hall JR, Hartmann M, Hollister EB, Lesniewski RA, Oakley BB, Parks DH, Robinson CJ, Sahl JW, Stres B, Thallinger GG, Van Horn DJ, and Weber CF. 2009. Introducing mothur: open-source, platform-independent, community-supported software for describing and comparing microbial communities. *Applied and Environmental Microbiology* 75:7537-7541. 10.1128/Aem.01541-09
- Schmieder R, and Edwards R. 2011. Quality control and preprocessing of metagenomic datasets. *Bioinformatics* 27:863-864. 10.1093/bioinformatics/btr026
- See-Too WS, Chua KO, Lim YL, Chen JW, Convey P, Mohidin TBM, Yin WF, and Chan KG. 2017. Complete genome sequence of *Planococcus donghaensis* JH1(T), a pectin-degrading bacterium. *Journal of Biotechnology* 252:11-14. 10.1016/j.jbiotec.2017.05.005
- Segata N, Izard J, Waldron L, Gevers D, Miropolsky L, Garrett WS, and Huttenhower C. 2011. Metagenomic biomarker discovery and explanation. *Genome Biology* 12. ARTN R6010.1186/gb-2011-12-6-r60
- Sgroy V, Cassan F, Masciarelli O, Del Papa MF, Lagares A, and Luna V. 2009. Isolation and characterization of endophytic plant growth-promoting (PGPB) or stress homeostasis-regulating (PSHB) bacteria associated to the halophyte *Prosopis strombulifera*. *Applied Microbiology and Biotechnology* 85:371-381. 10.1007/s00253-009-2116-3
- Shange RS, Ankumah RO, Ibekwe AM, Zabawa R, and Dowd SE. 2012. Distinct Soil Bacterial Communities Revealed under a Diversely Managed Agroecosystem. *Plos One* 7. ARTN e4033810.1371/journal.pone.0040338
- Shi SJ, O'Callaghan M, Jones EE, Richardson AE, Walter C, Stewart A, and Condron L. 2012. Investigation of organic anions in tree root exudates and rhizosphere microbial communities using in situ and destructive sampling techniques. *Plant and Soil* 359:149-163. 10.1007/s11104-012-1198-3
- Siddikee MA, Chauhan PS, Anandham R, Han GH, and Sa T. 2010. Isolation, characterization, and use for plant growth promotion under salt stress, of ACC deaminase-producing halotolerant bacteria derived from coastal soil. *Journal of Microbiology and Biotechnology* 20:1577-1584. 10.4014/jmb.1007.07011
- Smalla K, Wieland G, Buchner A, Zock A, Parzy J, Kaiser S, Roskot N, Heuer H, and Berg G. 2001. Bulk and rhizosphere soil bacterial communities studied by denaturing gradient gel electrophoresis: Plant-dependent enrichment and seasonal shifts revealed. *Applied and Environmental Microbiology* 67:4742-4751. Doi 10.1128/Aem.67.10.4742-4751.2001
- Song LY, Yang S, Liu HJ, and Xu J. 2017. Geographic and environmental sources of variation in bacterial community composition in a large-scale municipal landfill site in China. *Applied Microbiology and Biotechnology* 101:761-769. 10.1007/s00253-016-7917-6
- Soussi A, Ferjani R, Marasco R, Guesmi A, Cherif H, Rolli E, Mapelli F, Ouzari HI, Daffonchio D, and Cherif

- 715 A. 2016. Plant-associated microbiomes in arid lands: diversity, ecology and biotechnological  
716 potential. *Plant and Soil* 405:357-370. 10.1007/s11104-015-2650-y
- 717 Szymanska S, Plociniczak T, Piotrowska-Seget Z, and Hryniewicz K. 2016a. Endophytic and rhizosphere  
718 bacteria associated with the roots of the halophyte *Salicornia europaea* L. - community structure and  
719 metabolic potential. *Microbiological Research* 192:37-51. 10.1016/j.micres.2016.05.012
- 720 Szymanska S, Plociniczak T, Piotrowska-Seget Z, Zloch M, Ruppel S, and Hryniewicz K. 2016b. Metabolic  
721 potential and community structure of endophytic and rhizosphere bacteria associated with the roots of  
722 the halophyte *Aster tripolium* L. *Microbiological Research* 182:68-79. 10.1016/j.micres.2015.09.007
- 723 Takekawa JY, Miles AK, Schoellhamer DH, Athearn ND, Saiki MK, Duffy WD, Kleinschmidt S,  
724 Shellenbarger GG, and Jannusch CA. 2006. Trophic structure and avian communities across a salinity  
725 gradient in evaporation ponds of the San Francisco Bay estuary. *Hydrobiologia* 567:307-327.  
726 10.1007/s10750-006-0061-z
- 727 Tang J, Zheng AP, Bromfield ESP, Zhu J, Li SC, Wang SQ, Deng QM, and Li P. 2011. 16S rRNA gene  
728 sequence analysis of halophilic and halotolerant bacteria isolated from a hypersaline pond in Sichuan,  
729 China. *Annals of Microbiology* 61:375-381. 10.1007/s13213-010-0137-x
- 730 Tian XY, and Zhang CS. 2017. Illumina-based analysis of endophytic and rhizosphere bacterial diversity of  
731 the coastal halophyte *Messerschmidia sibirica*. *Frontiers in Microbiology* 8. Artn  
732 228810.3389/Fmicb.2017.02288
- 733 Tian YQ, and Gao LH. 2014. Bacterial diversity in the rhizosphere of cucumbers grown in soils covering a  
734 wide range of cucumber cropping histories and environmental conditions. *Microbial Ecology* 68:794-  
735 806. 10.1007/s00248-014-0461-y
- 736 Tiwari S, Singh P, Tiwari R, Meena KK, Yandigeri M, Singh DP, and Arora DK. 2011. Salt-tolerant  
737 rhizobacteria-mediated induced tolerance in wheat (*Triticum aestivum*) and chemical diversity in  
738 rhizosphere enhance plant growth. *Biology and Fertility of Soils* 47:907-916. 10.1007/s00374-011-  
739 0598-5
- 740 Vasquez EA, Glenn EP, Brown JJ, Guntenspergen GR, and Nelson SG. 2005. Salt tolerance underlies the  
741 cryptic invasion of North American salt marshes by an introduced haplotype of the common reed  
742 *Phragmites australis* (Poaceae). *Marine Ecology Progress Series* 298:1-8. DOI 10.3354/meps298001
- 743 Wang Q, Garrity GM, Tiedje JM, and Cole JR. 2007. Naive Bayesian classifier for rapid assignment of rRNA  
744 sequences into the new bacterial taxonomy. *Applied and Environmental Microbiology* 73:5261-5267.  
745 10.1128/Aem.00062-07
- 746 Wang XL, Cui WJ, Feng XY, Zhong ZM, Li Y, Chen WX, Chen WF, Shao XM, and Tian CF. 2018. Rhizobia  
747 inhabiting nodules and rhizosphere soils of alfalfa: A strong selection of facultative microsymbionts.  
748 *Soil Biology & Biochemistry* 116:340-350. 10.1016/j.soilbio.2017.10.033
- 749 Xi HY, Feng Q, Zhang L, Si JH, Chang ZQ, Yu TF, and Guo R. 2016. Effects of water and salinity on plant  
750 species composition and community succession in Ejina Desert Oasis, northwest China.  
751 *Environmental Earth Sciences* 75. Artn 13810.1007/S12665-015-4823-7
- 752 Yang GM. 1987. Photometric determination of organic matter in soil. *Soils and Fertilizers*:43-44.
- 753 Yang Y, Wang N, Guo XY, Zhang Y, and Ye BP. 2017. Comparative analysis of bacterial community  
754 structure in the rhizosphere of maize by high-throughput pyrosequencing. *Plos One* 12. ARTN  
755 e017842510.1371/journal.pone.0178425

756 Zhang XN, Yang XD, and Lv GH. 2014. Diversity patterns and response mechanisms of desert plants to the  
757 soil environment along soil water and salinity gradients. *Acta Ecologica Sinica* 36:3206-3215.

758 Zhao S, Zhou N, Zhao ZY, Zhang K, Wu GH, and Tian CY. 2016. Isolation of endophytic plant growth-  
759 promoting bacteria associated with the halophyte *Salicornia europaea* and evaluation of their  
760 promoting activity under salt stress. *Current Microbiology* 73:574-581. 10.1007/s00284-016-1096-7

761 Zhong ZP, Liu Y, Miao LL, Wang F, Chu LM, Wang JL, and Liu ZP. 2016. Prokaryotic community structure  
762 driven by salinity and ionic concentrations in plateau lakes of the Tibetan Plateau. *Appl Environ*  
763 *Microbiol* 82:1846-1858. 10.1128/AEM.03332-15

764 Zhou ML, Chen WM, Chen HY, and Wei GH. 2012. Draft genome sequence of *Mesorhizobium alhagi*  
765 CCNWXJ12-2(Tau), a novel salt-resistant species isolated from the desert of Northwestern China.  
766 *Journal of Bacteriology* 194:1261-1262. 10.1128/Jb.06635-11


767 Zhu YB, Tian JQ, Shi FY, Su L, Liu KK, Xiang MC, and Liu XZ. 2013. Rhizosphere bacterial communities  
768 associated with healthy and Heterodera glycines-infected soybean roots. *European Journal of Soil*  
769 *Biology* 58:32-37. 10.1016/j.ejsobi.2013.05.001

770

# Figure 1

Multiple comparisons of OTU richness (a) and  $\alpha$ -diversity (Shannon's index) (b) between rhizospheric and bulk soil bacterial communities, and among plant species within soil type.

Within boxes, horizontal bars indicate medians, while the tops and bottoms of boxes illustrate 75th and 25th quartiles, respectively. Small circles represent outliers in bulk samples. G, B, J, S, Z indicate bulk soil samples from *Lycium ruthenicum*, *Limonium gmelinii*, *Halocnemum strobilaceum*, *Halostachys caspica*, and *Kalidium foliatum*, respectively, while GQ, BX, JM, SM and ZZ denote the corresponding rhizosphere samples from each species. Significance levels (p values) are provided for among-species comparisons within soil type (bulk versus rhizosphere), as well as between soil types.






# Figure 2


Venn diagram showing the OTUs shared among different samples.

a: rhizosphere samples, b: bulk soil samples. Information on OTU numbers is provided as follows: the total number of OTUs detected across all three replicates for each species (the number of OTUs shared among the three replicates).

**a**




**b**



# Figure 3

Diagrams depicting community composition and relative abundance at different taxonomic level.


a: phylum, b: class, c: genus. com/", 2.34.



# Figure 4

Principal coordinate analysis (PCoA) (a) and UPGMA clustering diagram (b) of soil samples

G, B, J, S and Z represent bulk soil samples from *Lycium ruthenicum*, *Limonium gmelinii*, *Halocnemum strobilaceum*, *Halostachys caspica* and *Kalidium foliatum*, respectively, while GQ, BX, JM, SM and ZZ represent rhizosphere samples associated with these species.




# Figure 5


LEfSe analysis at multiple taxonomic levels comparing rhizobacterial community composition for five focal plant species.

(a) Cladogram illustrating the taxonomic groups explaining the most variation among rhizobacterial communities. Each ring represents a taxonomic level, with phylum (p), class (c), order (o), family (f) and genus (g) emanating from the center to the periphery. Each circle is a taxonomic unit found in the dataset, with circles or nodes shown in colors (other than yellow) indicating where a taxon was significantly more abundant. (b) Histogram of the LDA scores computed for groups with differential abundance among the rhizobacterial communities of the five plant species.

a



b






# Figure 6

LEfSe analysis of bacterial community composition for bulk soil samples from four plant species.


(a) Cladogram illustrating the taxonomic groups that explain the most variation among the bacterial communities. (b) Histogram of the LDA scores computed for groups with differential abundance among the bacterial communities of the four plant species. *Kalidium foliatum*-associated communities did not form a separate group (from other plant species' associated communities) and, therefore, these are absent from the cladogram and histogram.



# Figure 7

Canonical correspondence analysis (CCA) of the effect of soil properties on bacterial community structure in the rhizosphere (a) and bulk soil samples (b)

The soil properties are indicated with arrows, and include soil pH, electrical conductivity (EC), total organic carbon (TOC), soil organic matter (SOM), total nitrogen (TON), phosphorous (AP) and soil water content (SWC). The percentage of variation explained by each axis is provided. GQ, BX, JM, SM and ZZ represent rhizosphere samples associated with *Lycium ruthenicum*, *Limonium gmelinii*, *Halocnemum strobilaceum*, *Halostachys caspica* and *Kalidium foliatum*, respectively. G, B, J, S and Z represent bulk soil samples from *Lycium ruthenicum*, *Limonium gmelinii*, *Halocnemum strobilaceum*, *Halostachys caspica* and *Kalidium foliatum*, respectively.





**Table 1**(on next page)

Soil characteristics of bulk soil samples from five halophytes.

**Table 1** Soil characteristics of bulk soil samples from five halophytes.

	TOC (g/kg)	SOM (g/kg)	TON (g/kg)	AP (g/kg)	pH	EC (mS/cm)	SWC (%)
<i>Lycium ruthenicum</i>	9.14 ± 3.43a	15.75 ± 5.92a	0.58 ± 0.24a	0.89 ± 0.15a	8.23 ± 0.37	5.56 ± 1.26b	19.73 ± 2.18a
<i>Limonium gmelinii</i>	10.78 ± 1.60a	18.59 ± 2.77a	0.60 ± 0.10a	0.80 ± 0.08a	8.33 ± 0.24	6.61 ± 0.91ab	17.02 ± 3.51a
<i>Kalidium foliatum</i>	11.27 ± 5.66a	19.43 ± 9.76a	0.64 ± 0.25a	0.92 ± 0.08a	8.02 ± 0.25	5.65 ± 0.53b	16.45 ± 6.11a
<i>Halostachys caspica</i>	5.53 ± 0.95b	9.53 ± 1.63b	0.34 ± 0.03ab	0.82 ± 0.06a	8.05 ± 0.16	6.78 ± 1.42ab	10.42 ± 2.57b
<i>Halocnemum strobilaceum</i>	3.15 ± 1.09b	5.43 ± 1.88b	0.21 ± 0.02b	0.62 ± 0.06b	8.05 ± 0.30	7.14 ± 1.46a	17.71 ± 3.16a
mean	8.05 ± 4.15	13.87 ± 7.15	0.48 ± 0.23	0.82 ± 0.14	8.14 ± 0.27	6.30 ± 1.21	16.40 ± 4.57

Values are given as means ( ± standard error) (n = 3).

Different letters indicate significant differences among five halophytes at  $P < 0.05$  level.

## **Table 2**(on next page)

Adonis analysis of bacterial community composition for bulk soil and rhizosphere samples.

**Table 2** Adonis analysis of bacterial community composition for bulk soil and rhizosphere samples.

	<i>Lycium ruthenicum</i>	<i>Limonium gmelinii</i>	<i>Halocnemum strobilaceum</i>	<i>Halostachys caspica</i>	<i>Kalidium foliatum</i>
<i>Lycium ruthenicum</i>		0.323(0.3)	0.399(0.199)	0.319(0.289)	0.367(0.089)
<i>Limonium gmelinii</i>	0.324(0.108)		0.145(0.486)	0.042(0.894)	0.086(0.615)
<i>Halocnemum strobilaceum</i>	0.697(0.126)	0.718 (0.098)		0.096(0.605)	0.175(0.499)
<i>Halostachys caspica</i>	0.528(0.104)	0.513(0.11)	0.750(0.104)		0.016(0.904)
<i>Kalidium foliatum</i>	0.457(0.097)	0.504(0.102)	0.521(0.098)	0.574(0.114)	

data was shown in format of  $R^2$  ( $P$  value); values below diagonal are for rhizosphere samples, and above diagonal for bulk samples.

**Table 3**(on next page)

Pearson correlation coefficients among soil chemical properties, and between soil properties and community diversity.

**Table 3** Pearson correlation coefficients among soil chemical properties, and between soil properties and community diversity.

	TOC	TON	AP	SWC	pH	EC	OTU richness	Shannon index
TOC							0.446	0.484
TON	0.970***						0.549*	0.620*
AP	0.728**	0.761***					0.559*	0.609*
SWC	0.479*	0.476*	0.137				0.485	0.435
pH	0.420	0.467	0.151	0.301			0.504	0.459
EC	-0.211	-0.235	-0.126	-0.206	0.235		-0.093	-0.054

\* indicates significance level,  $*P < 0.05$ ,  $**P < 0.01$ ,  $***P < 0.001$ .

EC: electrical conductivity, TOC: total organic carbon, SOM: soil organic matter, TON: total nitrogen, AP: available phosphorous, SWC: soil water content.