

European population trends and current conservation status of an endangered steppe-bird species: the Dupont's lark *Chersophilus duponti* (#27939)

1

First submission

Editor guidance

Please submit by **21 May 2018** for the benefit of the authors (and your \$200 publishing discount).



Structure and Criteria

Please read the 'Structure and Criteria' page for general guidance.



Custom checks

Make sure you include the custom checks shown below, in your review.



Raw data check

Review the raw data. Download from the [materials page](#).



Image check

Check that figures and images have not been inappropriately manipulated.

Privacy reminder: If uploading an annotated PDF, remove identifiable information to remain anonymous.

Files

Download and review all files from the [materials page](#).

3 Figure file(s)

12 Table file(s)

1 Raw data file(s)

! Custom checks

Vertebrate animal usage checks



Have you checked the authors [ethical approval statement](#)?



Were the experiments necessary and ethical?



Have you checked our [animal research policies](#)?



Structure your review

The review form is divided into 5 sections.
Please consider these when composing your review:

1. BASIC REPORTING
2. EXPERIMENTAL DESIGN
3. VALIDITY OF THE FINDINGS
4. General comments
5. Confidential notes to the editor






 You can also annotate this PDF and upload it as part of your review

When ready [submit online](#).





Editorial Criteria

Use these criteria points to structure your review. The full detailed editorial criteria is on your [guidance page](#).





BASIC REPORTING

-  Clear, unambiguous, professional English language used throughout.
-  Intro & background to show context. Literature well referenced & relevant.
-  Structure conforms to [PeerJ standards](#), discipline norm, or improved for clarity.
-  Figures are relevant, high quality, well labelled & described.
-  Raw data supplied (see [PeerJ policy](#)).

EXPERIMENTAL DESIGN

-  Original primary research within [Scope of the journal](#).
-  Research question well defined, relevant & meaningful. It is stated how the research fills an identified knowledge gap.
-  Rigorous investigation performed to a high technical & ethical standard.
-  Methods described with sufficient detail & information to replicate.

VALIDITY OF THE FINDINGS

-  Impact and novelty not assessed. Negative/inconclusive results accepted. *Meaningful* replication encouraged where rationale & benefit to literature is clearly stated.
-  Conclusions are well stated, linked to original research question & limited to supporting results.
-  Data is robust, statistically sound, & controlled.
-  Speculation is welcome, but should be identified as such.



The best reviewers use these techniques

Tip

Support criticisms with evidence from the text or from other sources

Example

Smith et al (J of Methodology, 2005, V3, pp 123) have shown that the analysis you use in Lines 241-250 is not the most appropriate for this situation. Please explain why you used this method.

Give specific suggestions on how to improve the manuscript

Your introduction needs more detail. I suggest that you improve the description at lines 57- 86 to provide more justification for your study (specifically, you should expand upon the knowledge gap being filled).

Comment on language and grammar issues

The English language should be improved to ensure that an international audience can clearly understand your text. Some examples where the language could be improved include lines 23, 77, 121, 128 - the current phrasing makes comprehension difficult.

Organize by importance of the issues, and number your points

- 1. Your most important issue*
- 2. The next most important item*
- 3. ...*
- 4. The least important points*

Please provide constructive criticism, and avoid personal opinions

I thank you for providing the raw data, however your supplemental files need more descriptive metadata identifiers to be useful to future readers. Although your results are compelling, the data analysis should be improved in the following ways: AA, BB, CC

Comment on strengths (as well as weaknesses) of the manuscript

I commend the authors for their extensive data set, compiled over many years of detailed fieldwork. In addition, the manuscript is clearly written in professional, unambiguous language. If there is a weakness, it is in the statistical analysis (as I have noted above) which should be improved upon before Acceptance.

European population trends and current conservation status of an endangered steppe-bird species: the Dupont's lark *Chersophilus duponti*

Julia Gómez-Catasús ^{Corresp., 1}, Cristian Pérez-Granados ^{1,2}, Adrián Barrero ¹, Gerard Bota ³, David Giralt ³, Germán M. López-Iborra ², David Serrano ⁴, Juan Traba ¹

¹ Terrestrial Ecology Group (TEG-UAM), Department of Ecology, Universidad Autónoma de Madrid, Madrid, Spain

² Multidisciplinary Institute for Environmental Studies "Ramón Margalef", Department of Ecology, Universidad de Alicante, Alicante, Spain

³ Biodiversity and Animal Conservation Lab, Forest Sciences Center of Catalonia (CTFC), Solsona, Catalonia, Spain

⁴ Department of Conservation Biology, Estación Biológica de Doñana (EBD-CSIC), Sevilla, Spain

Corresponding Author: Julia Gómez-Catasús

Email address: julia.gomez@uam.es

Background

Steppe-birds face drastic population declines throughout Europe. The Dupont's lark *Chersophilus duponti* is an endangered steppe-bird species whose European distribution is restricted to Spain. This scarce passerine bird could be considered an 'umbrella species', since its population trends may reveal the conservation status of shrub-steppes. However, Dupont's lark population trends of the whole Spanish (and therefore European) population are unknown, so an updated and rigorous assessment is needed.

In this work, we evaluated Dupont's lark population trends in Europe employing the most recent and largest compiled database until date (92 populations and 12 years). In addition, we assessed the species threat category according to current applicable criteria (approved in March 2017) in the Spanish Catalogue of Threatened Species (SCTS), which have never been applied to the Dupont's lark nor to any other Spanish species. Finally, we compared the resulting threat categories with current conservation status at European, national and regional level.

Methods

We fitted Switching Linear Trend models (software TRIM - *Trends and Indices for Monitoring data*) to evaluate population trends at national and regional scale (i.e. per Autonomous Community) during the period 2004 - 2015. In addition, the finite multiplicative annual rate (lambda) obtained from the TRIM analysis was employed to estimate the percentage of population size change in a 10-year period. In accordance, a threat category was assigned following A1 and A2 criteria applicable in the SCTS.

Results

Trends showed an overall 3.9% annual decline rate for the Spanish population (moderate decline, following TRIM). Regional analyses showed a high inter-regional variability. Andalusia and Castile-Leon showed a steep decline, experiencing over 5% annual declining change rate. Trends were classified as uncertain in Aragon, Castile-La Mancha, Catalonia, Community of Valencia, Navarre and Region of Murcia, due to variability in trends between years and populations, and to the high proportion of missing values.

On the other hand, we forecasted a 32.8% average decline during the next 10 years. Attending to these

results the species should be listed as 'Vulnerable' at national scale (SCTS). At the regional level, the conservation status of the species is of particular concern in Andalusia and Castile-Leon, where the species qualifies to be listed as 'Endangered'.

Discussion

This work highlights the worrying conservation status of the European Dupont's lark population, facing a 3.9% annual declining rate. Under this scenario, the urgent implementation of a wide-range conservation plan is vital to ensure the conservation of this steppe-bird species. Besides, the legal responsibility of administrations to law enforcement in matter of nature protection and cataloguing endangered species, is crucial to reverse declining population trends of this and other endangered taxa.

1 **European population trends and current conservation status of an**
2 **endangered steppe-bird species: the Dupont's lark *Chersophilus duponti***

3 Julia Gómez-Catasús¹; Cristian Pérez-Granados^{1,2}; Adrián Barrero¹; Gerard Bota³; David Giralt³;

4 Germán M. López-Iborra²; David Serrano⁴; Juan Traba¹

5 ¹ Terrestrial Ecology Group (TEG-UAM), Department of Ecology, Universidad Autónoma de
6 Madrid. C/ Darwin 2, 23049 Madrid, Spain.

7 ² Multidisciplinary Institute for Environmental Studies "Ramón Margalef", Department of
8 Ecology, Universidad de Alicante, PO Box 99, 03080 Alicante, Spain.

9 ³ Biodiversity and Animal Conservation Lab. Forest Sciences Center of Catalonia (CTFC). Crta.
10 de Sant Llorenç de Morunys a Port del Comte km 2, 25280 Solsona, Catalonia, Spain.

11 ⁴ Department of Conservation Biology, Estación Biológica de Doñana (EBD-CSIC). Avda
12 Américo Vespucio 26, 41092 Sevilla, Spain.

13 Corresponding Author:

14 Julia Gómez-Catasús¹

15 E-mail address: julia.gomez@uam.es

17 **Abstract**

18 **Background**

19 Steppe-birds face drastic population declines throughout Europe. The Dupont's lark
20 *Chersophilus duponti* is an endangered steppe-bird species whose European distribution is
21 restricted to Spain. This scarce passerine bird could be considered an 'umbrella species', since its
22 population trends may reveal the conservation status of shrub-steppes. However, Dupont's lark
23 population trends of the whole Spanish (and therefore European) population are unknown, so an
24 updated and rigorous assessment is needed.

25 In this work, we evaluated Dupont's lark population trends in Europe employing the most recent
26 and largest compiled database until date (92 populations and 12 years). In addition, we assessed
27 the species threat category according to current applicable criteria (approved in March 2017) in
28 the Spanish Catalogue of Threatened Species (SCTS), which have never been applied to the
29 Dupont's lark nor to any other Spanish species. Finally, we compared the resulting threat
30 categories with current conservation status at European, national and regional level.

31 **Methods**

32 We fitted Switching Linear Trend models (software TRIM - *Trends and Indices for Monitoring*
33 *data*) to evaluate population trends at national and regional scale (i.e. per Autonomous
34 Community) during the period 2004 – 2015. In addition, the finite multiplicative annual rate (λ)
35 obtained from the TRIM analysis was employed to estimate the percentage of population size
36 change in a 10-year period. In accordance, a threat category was assigned following A1 and A2
37 criteria applicable in the SCTS.

38 **Results**

39 Trends showed an overall 3.9% annual decline rate for the Spanish population (moderate decline,
40 following TRIM). Regional analyses showed a high inter-regional variability. Andalusia and
41 Castile-Leon showed a steep decline, experiencing over 5% annual declining change rate. Trends
42 were classified as uncertain in Aragon, Castile-La Mancha, Catalonia, Community of Valencia,
43 Navarre and Region of Murcia, due to variability in trends between years and populations, and to
44 the high proportion of missing values. On the other hand, we forecasted a 32.8% average decline
45 during the next 10 years. Attending to these results the species should be listed as ‘Vulnerable’ at
46 national scale (SCTS). At the regional level, the conservation status of the species is of particular
47 concern in Andalusia and Castile-Leon, where the species qualifies to be listed as ‘Endangered’.

48 **Discussion**


49 Our results highlight the worrying conservation status of the European Dupont’s lark population,
50 facing a 3.9% annual declining rate. Under this scenario, the urgent implementation of a wide-
51 range conservation plan is vital to ensure the conservation of this steppe-bird species. Besides,
52 the legal responsibility of administrations to law enforcement in matter of nature protection and
53 cataloguing endangered species, is crucial to reverse declining population trends of this and other
54 endangered taxa.

55 **Introduction**

56 Steppes and pseudo-steppes are two of the most important habitats for preservation of bird
57 diversity since 55% of European bird species listed in the IUCN Red List are highly dependent
58 on these habitats (Burfield 2005). Moreover, 83% of steppe-bird species show an unfavorable
59 conservation status in Europe (Burfield and Van Bommel 2004, Burfield 2005). This is a
60 consequence of the accelerated process of land use changes faced by steppe-like habitats, with
61 dramatic consequences on steppe-bird populations across Europe (Benton *et al.* 2003, Burfield

62 and Van Bommel 2004, Santos and Suárez 2005). Main habitat-related threats, and therefore
63 drivers of steppe-bird populations decline are: (i) changes on land use (afforestation, new crops,
64 infrastructure development, mining, rubbish dumps; Burfield 2005; Laiolo and Tella 2006a,
65 Gómez-Catasús *et al.* 2016, 2018); (ii) agricultural intensification (landscape homogenization,
66 irrigation, increase of agrochemicals; Donald *et al.* 2001, Benton *et al.* 2003, Brotons *et al.* 2004,
67 Burfield 2005); and (iii) land abandonment and changes on agriculture and livestock
68 management (Madroño *et al.* 2004, Burfield 2005).

69 Spain is the stronghold for steppe-birds in Western Europe, harbouring a major proportion of
70 their total European breeding population (Burfield 2005). However, most of Spanish steppe-bird
71 populations declined during the 1990 – 2000 period (Burfield 2005) and afterwards (BirdLife
72 International 2015). A species of a particular conservation concern is the Dupont's lark
73 *Chersophilus duponti* (Vieillot, 1820), identified amongst the 65 priority bird species inhabiting
74 steppes (Burfield and Van Bommel 2004) and one of the scarcest passerine birds in Europe. The
75 species is classified as 'Near Threatened' in the IUCN Red List (BirdLife International 2017)
76 and as 'Vulnerable' in both the European Red List of Birds (BirdLife International 2015) and in
77 the Spanish Catalogue of Threatened Species (Royal Decree 139/2011, 4th February). Its
78 European geographic range is restricted to Spain spreading over 1,480 km² (Suárez 2010), and its
79 population has been estimated at 1,300-2,400 breeding pairs (Garza *et al.* 2003, Tella *et al.* 2005,
80 Suárez 2010). The European population of Dupont's lark qualifies to be considered as an
81 Evolutionary Significant Unit (*sensu* Moritz 1994, Casacci *et al.* 2014), since they are isolated
82 and genetically and morphologically differentiated from the African ones (García *et al.* 2008,
83 Suárez 2010).


84 The species inhabits flat (<10-15% of slope) shrub-steppes, avoiding dry pastures and cereal
85 fields (Garza *et al.* 2005, Seoane *et al.* 2006, Pérez-Granados *et al.* 2017a). Habitat
86 fragmentation and land-use changes, common issues in steppe ecosystems, have been
87 documented as the main threats for the species (Tella *et al.* 2005, Íñigo *et al.* 2008, Garza and
 Traba 2016, Pérez-Granados *et al.* 2016, Gómez-Catasús *et al.* 2018). Therefore, the Dupont's
89 lark could be considered an 'umbrella species' (Frankel and Soulé 1981), since its population
90 trends may reveal the conservation status of shrub-steppes.

91 Dupont's lark population trends have been previously addressed globally (Suárez 2010) or in a
92 sample of populations (Tella *et al.* 2005, Pérez-Granados and López-Iborra 2013, 2014). Despite
93 results of all these studies show declining population trends, none of them derived population
94 change estimates using appropriate statistical methods. Moreover, current trends of the whole
95 Spanish (and European) population are unknown, so an updated and rigorous assessment is
96 needed. This updated information would allow assessing the conservation status of the species
97 according to a formal set of criteria at two spatial scales: national and regional (i.e. per
98 Autonomous Community where the species is present). The importance of both spatial scales
99 relies on the competence of the Spanish Autonomous Communities in nature protection and,
100 specifically, in listing and cataloguing endangered species (Law 42/2007, 13th December). The
101 Spanish Ministry of Agriculture and Fisheries, Food and Environment has the jurisdiction to list
102 the species at national scale in the Spanish Catalogue of Threatened Species (SCTS, Law
103 42/2007, 13th December) and to elaborate the National Conservation Strategy of endangered
104 species. On the other hand, each Autonomous Community is legally bound to list species in its
105 Regional Catalogue of Threatened Species (RCTS), at least with the same category than at the
106 national level. Besides, they have the competence to elaborate and implement both Conservation

107 and Recovery Plans for those species classified as ‘Vulnerable’ and ‘Endangered’, respectively.

108 Thus, regional population trends are crucial to assess whether species conservation status is
109 especially worrying in specific regions and if the category of threat should be increased in the
110 pertinent Catalogues.

111 Species included in the SCTS were listed in 2011 (Royal Decree 139/2011), but listing criteria
112 applicable in the SCTS were modified in March 2017 (Royal Decree 139/2011, 4th February;
113 Resolution 6th March 2017), to accommodate IUCN ones (IUCN 2012). However, conservation
114 status of catalogued species in the SCTS has not been reviewed after this modification. To our
115 knowledge, new criteria have never been applied to the Dupont’s lark nor to any other Spanish
116 species and, therefore, an assessment of the category of threat assigned under the new criteria is
117 needed.



118 In this work, we aimed to evaluate Dupont's lark population trends during the 2004 - 2015 period
119 at both national and regional scale  per region or also called Autonomous Communities),
120 using the largest database ever compiled. We also carried out a comprehensive assessment of the
121 conservation status of the Dupont’s lark according to quantitative threshold criteria of reduction
122 in population size (A1 and A2 criteria, see below) under current applicable criteria in the SCTS
123 (Resolution 6th March 2017). Finally, we also aimed to assess whether the current threat
124 category of the species at European (European Red List of Birds), national (SCTS) and regional
125 levels (RCTS) agrees with Dupont’s lark populations trends.

126 **Materials and Methods**

127 *Data collection*

128 The ethics committee of Animal Experimentation of the Autonomous University of Madrid as an
129 Organ Enabled by the Community of Madrid (Resolution 24th September 2013) for the

130 evaluation of projects based on the provisions of Royal Decree 53/2013, 1st February, has
131 provided full approval for this purely observational research (CEI 80-1468-A229).

132 We compiled data for 92 Dupont's lark populations during the 2004-2015 period. We considered
133 a population as those **habitat patches** separated by less than 1 km (termed subpopulation in
 Suárez 2010). **Our dataset accounted for 41.6% of the 221 populations surveyed during the II**
135 National Survey (2004-2006; Suárez 2010) and includes a temporal range between one and 12 

136 years (mean \pm SD = 5.36 ± 2.77 years). **Besides**, all the Autonomous communities where the
137 species occurs are included in the dataset (Fig. 1) (Suárez 2010).

138 The Dupont's lark population size is difficult to quantify due to the extremely shy and elusive
139 behavior of the species and the concentration of singing activity mainly before dawn. Therefore,
140 surveys of the species rely on auditory contacts. Bird censuses were carried out during the
141 breeding season (March-June depending on phenological differences; Garza *et al.* 2010)
142 approximately 1 hour before dawn, when singing activity peaks. Birds were counted by linear
143 transects (500 m inner belt width; Garza *et al.* 2010) or by territory mapping (Bibby *et al.* 2000),
144 since they produced similar population size estimates (Pérez-Granados and López-Iborra 2017).

145 A slightly different census method, consisting in a network of point counts, was performed in
146 Catalonia and Region of Murcia populations (comprising less than 5% of all populations).
147 Anyway, counting method remained constant throughout the study period within each region,
148 making inter-annual data comparable. Linear transects were designated to cover the whole
149 population (Suarez 2010), and were walked at constant speed georeferencing singing males with
150 a GPS and noting all males singing simultaneously. Transects were walked once under the linear
151 transect method and 2-4 times under the mapping method. Number of territories per population
152 was estimated by mapping all records and taking into account clusters of registrations and birds

153 heard simultaneously (Garza *et al.* 2010, Pérez-Granados and López-Iborra 2017). Population
154 size estimates refer to the minimum number of territories (mapping method), or minimum
155 number of males (line transect method) per population.

156 *Trend analysis*

157 Changes on population estimates were evaluated using the software TRIM (*Trends and Indices*
158 *for Monitoring data*. TRIM v. 3.54. Pannekoek and Van Strien 2006a). TRIM fits log-linear
159 models and was employed because: i) it allows to analyze time series with absence of data in
160 some years, a common issue in long-time series; and ii) it takes into account overdispersion and
161 serial correlation of data (Pannekoek and Van Strien 2005). TRIM calculates indices that
162 represent the effect of change between years, which indicates relative variation of the total
163 population size. From these indices, a mean annual change rate is estimated and a trend category
164 is assigned (Pannekoek and Van Strien 2006a). At the first time-point, the index value is 1 and is
165 taken as point-reference for quantifying the relative temporal trends in the subsequent years. This
166 technique has been broadly employed for the analysis of temporal series in bird populations (e.g.
167 Paradis *et al.* 2002, Wretenberg *et al.* 2007, Delgado *et al.* 2009).

168 We fitted Switching Linear Trend models to evaluate both national and regional Dupont's lark
169 trends during the period 2004 - 2015. TRIM employs a stepwise selection of change-points in
170 trends using Wald-tests for the significance of change-points. When the difference between
171 parameters before and after a change-point does not differ from zero (default significance
172 threshold: 0.2), the corresponding change-point is removed from the model attending to the
173 parsimony principle (Pannekoek and Van Strien 2005). The best-fit models were selected
174 according to Goodness-of-fit tests (Likelihood ratio test and Chi-squared) and Akaike
175 information criterion (AIC). A model with a significance value higher than 0.05 indicates that

176 data fit a Poisson distribution and, therefore, the model could be accepted. Indices, overall slope
177 and Wald tests remain reliable in case of lack-of-fit (Pannekoek and Van Strien 2005). In case of
178 overdispersion or serial correlation (default TRIM threshold: >3.0 and >0.4 respectively;
179 Pannekoek and Van Strien 2006b) Wald-test for the significance of slope was employed
180 (Pannekoek and Van Strien 2005). While the whole set of 92 populations was used to analyse
181 national trends, regional subsets were subsequently extracted to analyse regional trends (see
182 Table 1 for sample size in each region).

183 *Threat category*

184 We evaluated Dupont's lark category of threat according to A1 (population size reduction over
185 the last 10 years or three generations, whichever the longer) and A2 (population size reduction
186 within the next 10 years or three generations, whichever the longer) criteria applicable in the
187 SCTS. We used recent trends to forecast future population trends of the species, since its
188 geographic range reduction (Traba *et al.* 2016) and the lack of conservation measures (Tella *et*
189 *al.* 2005, Suárez 2010, Pérez-Granados and López-Iborra 2014) predict similar population trends
190 in the next years.

191 The finite multiplicative annual rate (λ) was obtained from the TRIM analysis. This value was
192 employed to estimate the percentage of population size change in a 10-year period following the
193 equation below:

$$194 \quad \text{Percentage of change in a 10-year period (\%)} = (\lambda^{10} - 1) \cdot 100$$

195 We assigned a threat category according to population size reduction estimated over the last 10
196 years (A1 criterion; 'Endangered' $\geq 70\%$ 'Vulnerable' $\geq 50\%$) and forecasted in the next 10
197 years (A2 criterion; 'Endangered' $\geq 50\%$ 'Vulnerable' $\geq 30\%$) at both national and regional

198 scale. Lastly, categories were compared with the current threat categories for the Dupont's lark
199 in the European Red List of Birds, the SCTS and the RCTS.

200 **Results**

201 *Spanish (European) population trend*

202 The best Switching Linear Trend model for all Dupont's lark populations did not fit to a log-
203 linear distribution (Chi-square, $\chi^2 = 684.92$, $df = 389$, $p < 0.001$; Likelihood Ratio, $LR = 722.30$,
204 $df = 389$, $p < 0.001$; $AIC = -55.70$). Overdispersion and serial correlation values were relatively
205 low (1.70 and 0.34, respectively), but 55.8% of counts were missing values. The stepwise
206 procedure revealed six significant change-points in trends (Fig. 2; Table S1). Population size-
207 index experienced an overall 41.3% decline (95% CI, -50.2 to -32.5) from 2004 to 2015.

208 ~~Besides~~, the extinction of 26 populations (hereafter local extinction events), which represents
209 28% of the set of study populations, was registered in this period (Table S2). The overall slope
210 parameter showed a 3.9% annual decrease (95% CI, -4.9 to -2.8%), which corresponds to a
211 moderate decline according to TRIM criteria (Pannekoek and Van Strien 2006a).

212 *Regional population trends*

213 Regional trends showed high variability between regions (Table 1; Fig. 3). Switching Linear
214 Trend models for Aragon (AR), Navarre (NA) and Region of Murcia (RM) populations fitted to
215 a log-linear distribution (χ^2 and LR p-values > 0.05), while goodness-of-fit tests for models of
216 Andalusia (AN) and Community of Valencia (CV) were near to acceptance values (χ^2 and LR p-
217 values > 0.01 ; Table 1). However, Castile-La Mancha (CM) and Castile-Leon (CL) models did
218 not fit to a log-linear distribution (χ^2 and LR p-values < 0.01 ; Table 1). Overdispersion and serial
219 correlation values were of less concern for all models except for Catalonia (CA; Table 1), so we
220 relied on Wald-tests for best-model selection. Proportion of missing values was higher than 50%

221 for AR, CM, CA and NA models, and sample sizes were small for all regions (i.e. less than 15
222 populations) except for CM and CL (Table 1). Significant change-points in slope were
223 incorporated in all models except for AR, CA and NA (Fig. 3; Table S3-S10), these three
224 showing a constant slope throughout the study period. Trend analyses showed mean overall
225 decreases in AN (70.0%), CL (50.8%), CM (59.0%), CV (26.8%) and NA (11.8%) during the
226 2004 – 2015 period (Table 2). However, mean overall trends were positive in AR (17.1%), CA
227 (48.2%) and RM (37.4%) populations (Table 2). Average annual change rates showed a steep
228 decline for AN and CL populations, higher than 5% per year (Table 1; Fig. 3). Population trends
229 of AR, CA, CM, CV, NA and RM were classified as uncertain (Table 1; Fig. 3). Local extinction
230 events were registered mainly in CL (9), AN (6), and CM (6) (Table S2). Frequency of local
231 extinction events were higher in CA (100%, only one population under study which was
232 ultimately recolonized in 2015), AN (50%), CV (37.5%), NA (33.3%), CL (31%) and CM
233 (23.1%).

234 *Threat category*

235 According to estimated mean annual rate of change (-3.9%), Dupont's lark population size in
236 Spain has been reduced on average by 32.8% over the last 10 years and will be reduced by the
237 same percentage in the next 10 years (Table 2). This reduction in population size does not entail
238 the classification of the Dupont's lark at any category of threat in Spain according to A1 criterion
239 (Table 2). However, the Dupont's lark should be classified as 'Vulnerable' in the SCTS
240 according to A2 criterion (Table 2).

241 Regional analyses showed that the species should classify as 'Vulnerable' in AN and CL
242 according to past population trends (A1 criterion) and no category of threat is assigned in the rest
243 of the Regional Catalogues (Table 2). Nevertheless, the species should classify at least as


244 'Vulnerable' in all the Regional Catalogues according to forecasted population declines (A2
245 criterion) and Spanish legislation (Table 2). Specifically, the species should be upgraded to
246 'Endangered' in AN and CL in agreement with A2 criterion (Table 2).

247 **Discussion**

248 Our results evidence the worrying trends of the Spanish Dupont's larks population, the only
249 bastion of this endangered steppe-bird in Europe. The species exhibited an estimated annual
250 decline rate of 3.9% during the last decade, which agrees qualitatively with previous work on
251 Dupont's lark population trends at particular areas of its Spanish distribution (Tella *et al.* 2005,
252 Pérez-Granados and López-Iborra 2013). Our results are also in concordance with declining
253 trends described for most of steppe-bird species in the Iberian Peninsula during the last decades
254 (Burfield 2005, BirdLife International 2015). Habitat loss and alteration (in terms of either
255 availability or quality) through agricultural intensification, abandonment of traditional extensive
256 livestock and other land use changes (e.g. ploughing and afforestation promoted by the Common
257 Agricultural Policy, tree crops, irrigated lands, infrastructure development), are some of the
258 anthropic activities known to impact on shrub-steppes (Santos and Suárez 2005), and have been
259 repeatedly cited as the main causes of Dupont's lark negative population trends (Tella *et al.*
260 2005, Íñigo *et al.* 2008, Garza and Traba 2016, Pérez-Granados *et al.* 2017b, Gómez-Catasús *et*
261 *al.* 2016, 2018).

262 In this study, we compiled the most exhaustive and updated database for Dupont's lark
263 population trends. We considered that our sampling coverage is representative of the Iberian
264 (European) distribution, leading to reliable results for the population trend analysis. Most regions
265 were significantly represented in this sample, ranging between 43% of the total regional
266 population for CL, 48% for CM and 100% for AN, CA, CV, NA and RM. However, we only

267 were able to compile data on 10 populations for AR (10.5% of the 95 populations surveyed in
268 2004-2006; Suárez 2010), the region that concentrates the majority of the Spanish Dupont's lark
269 population (Suárez 2010). Thus, overall trends results (3.9% annual decline rate) may be
270 somewhat biased due to absence of data in some important populations. Therefore, future
271 population trend analyses incorporating a higher proportion of the regional populations in AR are
272 needed. Accordingly, priority should be given to standardize long-term monitoring, particularly
273 in those large populations in Aragon.

274 One additional precaution is related to the lack of fit in models, probably due to slight
275 overdispersion in data (i.e. variance greater than the mean). This could be due to unknown
276 factors not incorporated into the models, which could influence on trends (Quinn and Keough
277 2002, Crawley 2007). For instance, interannual variability in population trends encompassed by
278 the significant change-points (Table S1; Table S3-S10) could be explained by natural
279 stochasticity, either demographic or environmental (Lande 1987), as well as density-dependent
280 interactions (Bjørnstad and Grenfell 2001). Demographic stochasticity may be an important
281 driver of the observed oscillations between years, since Dupont's lark seems to fit to a
282 metapopulation structure with local extinction events and colonization processes (e.g. Alfés
283 population in CA; Bota *et al.* 2016).  This produces high variability in TRIM yearly indices (i.e.
284 overdispersion), and therefore hinders to obtain generalized population trends over time. On the
285 other hand, interannual variability may be also associated to environmental stochasticity and
286 fluctuations on abiotic factors such as climate (Delgado *et al.* 2009) due to its effects on food
287 availability (Wiens 1989, Lemoine *et al.* 2007), reproductive success (Bolger *et al.* 2005, Van de
288 Pol *et al.* 2010) or annual survival (Robinson *et al.* 2007), among others. Future research should
289 focus on disentangling the mechanisms underlying variability on trends in order to incorporate

290 new covariates in models and improve their Goodness-of-fit. Anyway, the lack of fit would not
291 invalidate indices, overall slope and Wald tests (Pannekoek and Van Strien 2005), and
292 consequently main results about Dupont's lark population trends remain reliable.

293 We found large differences between regions in population trends; drastic declining trends
294 (annual declining rate higher than 5%) occurred in AN and CL, while trends were classified as
295 uncertain in the other regions (AR, CM, CA, CV, NA and RM). Uncertainty in trends may be
296 due to two typical handicaps in long-term databases: (i) high variability between years and
297 populations (within a region) that produces large Confidence Intervals (i.e. overdispersion); and
298 (ii) high proportion of missing values (Atkinson *et al.* 2006). As we stated above, overdispersion
299 was low except for CA, which could be explained by the extinction-recolonization process
300 undergone by the single population in this region (Bota *et al.* 2016). ~~Besides~~, the percentage of
301 missing values (Table 1) exceeded the recommended threshold of 20-50% for TRIM analyses
302 (Pannekoek and Van Strien 2005). These two analytical constraints have negligible effects at
303 national scale but less reliable estimates are expected to be obtained with small-size samples (i.e.
304 regional analysis; Atkinson *et al.* 2006). This probably explains uncertain population trends for
305 AR, CA, CV, NA and RM. Consequently, results for some regional trends should be treated with
306 caution, especially when a low proportion of populations were included in the regional analyses
307 (e.g. AR; see above).

308 The comprehensive assessment of the conservation status of the Dupont's lark yielded a higher
309 category of threat according to A2 criterion (future population trends) than A1 criterion (past
310 population trends). The fulfillment of one criterion is enough to classify the species at the highest
311 category of threat. Thus, according to A2 criterion, the Dupont's lark is correctly listed as
312 'Vulnerable' in the European Red List of Birds, in the SCTS and in the Regional Catalogues of

313 CM, CV and RM. Of particular concern, however, are Dupont's lark populations in AN and CL,
314 where the species qualifies to be listed as 'Endangered'. However, CL has not yet elaborated a
315 RCTS, while the species is currently listed as 'Vulnerable' in AN. In the other regions (AR, CA
316 and NA), the species should be classified as 'Vulnerable' according to the category of threat
317 assigned in the SCTS (Law 42/2007, 13th December). If the same assessment would have been
318 carried out using previous applicable criteria in the SCTS (before March 2017; Dirección
319 General para la Conservación de la Naturaleza 2004), the cataloguing scenario would have
320 changed drastically. Under the old criteria the Dupont's lark should have been listed as
321 'Endangered' (A2 criterion; population size reduction of $\geq 40\%$ within the next 20 years),
322 evidencing the effects that listing criteria modification may have on the management and
323 conservation of threatened species.

324 In this study, we assessed the conservation status of the Dupont's lark according to A criteria,
325 since we had no reliable data for including other criteria in our analyses. Therefore, a similar
326 comprehensive assessment should be carried out considering the remaining listing SCTS criteria
327 (reduction in area of occupancy and/or population viability analysis; Resolution 6th March 2017)
328 to elucidate whether or not the species should be classified as 'Endangered', ensuring proper
329 listing of the species at both European and national level. For instance, consensus among experts
330 (D criteria; Resolution 6th March 2017) upon the need of its reclassification as "Endangered"
331 exists (Tella *et al.* 2005, Pérez-Granados and López-Iborra 2014, Garza and Traba 2016). Future
332 research should focus on accurately estimating the reduction in area of occupancy. Besides, a
333 population viability analysis could be carried out to assess the risk of extinction in the coming
334 years, although estimating reliable demographic parameters for the whole population of this
335 secretive species is challenging.

336 **Conclusions**

337 Despite methodological constraints due to slight overdispersion, missing data, and low
338 proportion of populations incorporated for AR, we believe that our results are conclusive. The
339 European Dupont's lark population faces a 3.9% annual declining rate, entailing an expected
340 average population decline of 32.8% within the next 10 years. The pressures faced by the species
341 have not ceased during the last years (Tella *et al.* 2005, Íñigo *et al.* 2008, Garza and Traba 2016),
342 and may be expected to increase in the future due to strong fragmentation and high vulnerability
343 to stochastic factors (Laiolo and Tella 2006b, Vögeli *et al.* 2010, Méndez *et al.* 2011, Gómez-
344 Catasús *et al.* 2018). Under this scenario, the implementation of a wide-range conservation plan
345 within the Iberian distribution is vital to ensure the conservation of the species. According to
346 Spanish legislation the elaboration of a Conservation Plan is mandatory for those species
347 classified as 'Vulnerable', as the Dupont's lark since 2004 (Orden MAM/2784/2004), and this is
348 within the competence of the Autonomous Communities. In addition, Autonomous Communities
349 are legally obligated to comply with current legislation in cataloguing endangered species (Law
350 42/2007, 13th December). Therefore, the species should be classified as 'Endangered' in
351 Andalusia and Castile-Leon, and as 'Vulnerable' in Aragon, Catalonia and Navarra. In this
352 context, the legal responsibility of administrations to law enforcement is crucial to reverse
353 declining population trends of this and other endangered taxa.

354 **Acknowledgements**

355 The authors wish to thank all the people that disinterestedly provided their data on Dupont's lark
356 populations in Spain.

357 **References**

358 Atkinson, P. W., Austin, G. E., Rehfisch, M. M., Baker, H., Cranswick, P., Kershaw, M.,

- 359 Robinson, J., Lagnston, R. H. W., Stroud, D. A., Van Turnhout, C. and Maclean, I. M.
360 (2006). Identifying declines in waterbirds: The effects of missing data, population
361 variability and count period on the interpretation of long-term survey data. *Biological*
362 *Conservation*, 130: 549–559.
- 363 Benton, T. G., Vickery, J. A. and Wilson, J. D. (2003). Farmland biodiversity: is habitat
364 heterogeneity the key? *Trends of Ecology and Evolution*, 18: 182–188.
- 365 Bibby, C. J., Burgess, N. D., Hill, D. A. and Mustoe, S. (2000). *Bird census techniques*. London,
366 U.K.: Academic Press.
- 367 BirdLife International. (2015). European Red List of Birds. Luxembourg: Office for Official
368 Publications of the European Communities.
- 369 BirdLife International. (2017). *Chersophilus duponti*. The IUCN Red List of Threatened Species.
370 Available at: <http://dx.doi.org/10.2305/IUCN.UK.2017-1.RLTS.T22717380A111109346.en>
371 (Accessed 1 July 2017).
- 372 Bjørnstad, O.N. and Grenfell, B.T. (2001). Noisy clockwork: time series analysis of population
373 fluctuations in animals. *Science*, 293: 638–643.
- 374 Bolger, D. T., Patten, M. A. and Bostock, D. C. (2005). Avian reproductive failure in response to
375 an extreme climatic event. *Oecologia*, 142: 398–406.
- 376 Bota, D., Giralt, D. and Guixé, D. (2016). *La alondra ricotí en Catalunya: evolución histórica de*
377 *una población en el límite del área de distribución*. II Workshop Grupo de Expertos en la
378 Alondra ricotí. UAM-Madrid.
- 379 Brotons, L., Mañosa, S. and Estrada, J. (2004). Modelling the effects of irrigation schemes on the
380 distribution of steppe birds in Mediterranean farmland. *Biodiversity and Conservation*, 13:
381 1039–1058.
- 382 Burfield, I. J. (2005). The conservation status of steppic birds in Europe. In G. Bota, M. B.
383 Morales, S. Mañosa, and J. Camprodon (Eds.), *Ecology and conservation of steppe-land*
384 *birds* (pp. 69–102). Lynx Edicions, Barcelona, Spain.
- 385 Burfield, I. and Van Bommel, F. (2004). *Birds in Europe: population estimates, trends and*
386 *conservation status*. BirdLife International. Cambridge, UK.
- 387 Casacci, L. P., Barbero, F. and Balletto, E. (2014). The “Evolutionarily Significant Unit” concept
388 and its applicability in biological conservation. *Italian Journal of Zoology*, 81: 182–193.
- 389 Crawley, M. J. (2007). *The R Book*. John Wiley & Sons, Hoboken, New Jersey, USA.
- 390 Delgado, M. P., Morales, M. B., Traba, J. and De la Morena, E. L. G. (2009). Determining the
391 effects of habitat management and climate on the population trends of a declining steppe
392 bird. *Ibis*, 151: 440–451.
- 393 Dirección General para la Conservación de la Naturaleza. (2004). *Criterios orientadores para la*
394 *inclusión de taxones y poblaciones en catálogos de especies amenazadas*. Ministerio de

- 395 Medio Ambiente, Madrid, Spain.
- 396 Donald, P. F., Green, R. E. and Heath, M. F. (2001). Agricultural intensification and the collapse
397 of Europe's farmland bird populations. *Proceedings of the Royal Society of London B*, 268:
398 25–29.
- 399 Frankel, O. H. and Soulé, M. E. (1981). *Conservation and Evolution*. Cambridge University
400 Press, Cambridge, United Kingdom.
- 401 García, J. T., Suárez, F., Garza, V., Calero-Riestra, M., Hernández, J. and Pérez-Tris, J. (2008).
402 Genetic and phenotypic variation among geographically isolated populations of the globally
403 threatened Dupont's lark *Chersophilus duponti*. *Molecular phylogenetics and evolution*, 46:
404 237–251.
- 405 Garza, V., Suárez, F. and Carriles, E. (2010). El censo actual: diseño y métodos de muestreo y
406 estima de las poblaciones. In F. Suárez (Ed.), *La alondra ricotí (Chersophilus duponti)* (pp.
407 175–194). Dirección General para la Biodiversidad, Ministerio de Medio Ambiente y
408 Medio Rural y Marino, Madrid, Spain.
- 409 Garza, V., Suárez, F., Herranz, J., Traba, J., De la Morena, E. L. G., Morales, M. B., González,
410 R. and Castañeda, M. (2005). Home range, territoriality and habitat selection by the
411 Dupont's lark *Chersophilus duponti* during the breeding and postbreeding periods. *Ardeola*,
412 52: 133–146.
- 413 Garza, V. and Traba, J. (2016). Retos para la conservación de una especie amenazada. Alondra
414 ricotí, el fantasma del páramo. *Quercus*, 359: 24–33.
- 415 Garza, V., Traba, J. and Suárez, F. (2003). Is the European population of Dupont's Lark
416 *Chersophilus duponti* adequately estimated? *Bird Study*, 50: 309–311.
- 417 Gómez-Catasús, J., Barrero, A., Garza, V. and Traba, J. (2016) Alondra ricotí – *Chersophilus*
418 *duponti*. In A. Salvador and M.B. Morales (eds.), *Enciclopedia virtual de vertebrados*
419 *Espanoles*. Madrid: Museo Nacional de Ciencias Naturales. Retrieved from: [http://](http://www.vertebradosibericos.org)
420 www.vertebradosibericos.org
- 421 Gómez-Catasús, J. Garza, V. and Traba, J. (2018) Wind farms affect the occurrence, abundance
422 and population trends of small passerine birds: The case of the Dupont's lark. *Journal of*
423 *Applied Ecology*. DOI: 10.1111/1365-2664.13107
- 424 Íñigo, A., Garza, V., Tella, J. L., Laiolo, P., Suárez, F. and Barov, B. (2008). *Action Plan for the*
425 *Dupont's Lark Chersophilus duponti in the European Union*. Madrid, Spain.
- 426 IUCN. (2012). *IUCN Red List Categories and Criteria: Version 3.1. Second edition*. IUCN,
427 Gland, Switzerland and Cambridge, UK.
- 428 Laiolo, P. and Tella, J. L. (2006a). Fate of unproductive and unattractive habitats: recent changes
429 in Iberian steppes and their effects on endangered avifauna. *Environmental Conservation*,
430 33: 223–232.
- 431 Laiolo, P. and Tella, J. L. (2006b). Landscape bioacoustics allows detection of the effects of

- 432 habitat patchiness on population structure. *Ecology*, 87: 1203–1214.
- 433 Lande, R. (1987). Extinction thresholds in demographic models of territorial populations. *The*
434 *American Naturalist*, 130: 624-635.
- 435 Lemoine, N., Bauer, H.G., Peintinger, M. and Böhning-Gaese, K. (2007). Effects of climate and
436 land-use changes on species abundance in a central European bird community.
437 *Conservation Biology*, 21: 495–503.
- 438 Madroño, A., González, C. and Atienza, J. C. (2004). *Libro rojo de las aves de España*.
439 Dirección General para la Biodiversidad- SEO/Birdlife, Madrid, Spain.
- 440 Méndez, M., Tella, J. L. and Godoy, J. A. (2011). Restricted gene flow and genetic drift in
441 recently fragmented populations of an endangered steppe bird. *Biological Conservation*,
442 144: 2615–2622.
- 443 Moritz, C. (1994). Defining “Evolutionarily Significant Units” for conservation. *Trends in*
444 *Ecology and Evolution*, 9: 373–375.
- 445 Pannekoek, J. and Van Strien, A. (2005). *TRIM 3 Manual (Trends and Indices for Monitoring*
446 *data)*. Statistics Netherlands, Voorburg, Holland.
- 447 Pannekoek, J. and Van Strien, A. (2006a). TRIM version 3.54 (Trends & Indices for Monitoring
448 data). Statistics Netherlands, Voorburg, Holland.
- 449 Pannekoek, J. and Van Strien, A. (2006b). Frequently Asked Questions Updated 2006 (TRIM
450 version 3.51). Statistics Netherlands, Voorburg, Holland. Available at: [http://www.bird-](http://www.bird-research.jp/1_shiryo/trim/Q&A_ENG.pdf)
451 [research.jp/1_shiryo/trim/Q&A_ENG.pdf](http://www.bird-research.jp/1_shiryo/trim/Q&A_ENG.pdf) (Accessed 1 July 2017).
- 452 Paradis, E., Baillie, S. R., Sutherland, W. J. and Gregory, R. D. (2002). Exploring density-
453 dependent relationships in demographic parameters in population of birds at a large spatial
454 scale. *Oikos*, 97: 293–307.
- 455 Pérez-Granados, C. and López-Iborra, G. M. (2013). Census of Breeding Birds and Population
456 Trends of the Dupont’s Lark *Chersophilus duponti* in Eastern Spain. *Ardeola*, 60: 143–150.
- 457 Pérez-Granados, C. and López-Iborra, G. M. (2014). ¿Por qué la alondra ricotí debe catalogarse
458 como “En peligro de extinción”? *Quercus*, 337: 18–25.
- 459 Pérez-Granados, C. and López-Iborra, G. M. (2017). Assesment of Counting Methods Used for
460 Estimating the Number of Territorial Males in the Endangered Dupont’s Lark. *Ardeola*, 64:
461 5–14.
- 462 Pérez-Granados, C., Osiejuk, T., and López-Iborra, G. M. (2016). Habitat fragmentation effects
463 and variations in repertoire size and degree of song sharing among close Dupont’s Lark
464 *Chersophilus duponti* populations. *Journal of Ornithology*, 157: 471-482.
- 465 Pérez-Granados, C., Lopez-Iborra, G. M., and Seoane, J. (2017a). A multi-scale analysis of
466 habitat selection in peripheral populations of the endangered Dupont’s Lark *Chersophilus*
467 *duponti*. *Bird Conservation International*, 27: 398-413.

- 468 Pérez-Granados, C., López-Iborra, G. M., Garza, V. and Traba, J. (2017b). Breeding biology of
469 the endangered Dupont's Lark *Chersophilus duponti* in two separate Spanish shrub-steppes.
470 *Bird Study*, 64: 328-338.
- 471 Quinn, G. P. and Keough, M. J. (2002). *Experimental design and data analysis for biologists*.
472 Cambridge University Press, Cambridge, UK.
- 473 Robinson, R. A., Baillie, S. R. and Crick, H. Q. (2007). Weather-dependent survival:
474 implications of climate change for passerine population processes. *Ibis*, 149: 357-364.
- 475 Santos, T. and Suárez, F. (2005). Biogeography and population trends of iberian steppe bird. In
476 G. Bota, M. B. Morales, S. Mañosa, and J. Camprodon (Eds.), *Ecology and conservation of*
477 *steppe-land birds* (pp. 69–102). Lynx Edicions, Barcelona, Spain.
- 478 Seoane, J., Justribo, J. H., García, F., Retamar, J., Rabadan, C. and Atienza, J. C. (2006).
479 Habitat-suitability modelling to assess the effects of land-use changes on Dupont's lark
480 *Chersophilus duponti*: a case study in the Layna Important Bird Area. *Biological*
481 *Conservation*, 128: 241–252.
- 482 Suárez, F. (2010). *La alondra ricotí (Chersophilus duponti)*. Dirección General para la
483 Biodiversidad, Ministerio de Medio Ambiente y Medio Rural y Marino Medio Rural y
484 Marino, Madrid, Spain.
- 485 Tella, J. L., Vögeli, M., Serrano, D. Y. and Carrete, M. (2005). Status of the threatened Dupont's
486 lark in Spain: overestimation, decline and extinction of local populations. *Oryx*, 39: 1–5.
- 487 Traba, J., Gómez-Catasús, J., García-Antón, A., Barrero, A. and Garza, V. (2016). Bases
488 científicas para la elaboración de la estrategia nacional de conservación de la alondra ricotí.
489 Informe inédito. Fundación Biodiversidad.
- 490 Van De Pol, M., Ens, B. J., Heg, D., Brouwer, L., Krol, J., Maier, M., Exo, K., Oosterbeek, K.,
491 Lok, T., Eising, C. M. and Koffijberg, K. (2010). Do changes in the frequency, magnitude
492 and timing of extreme climatic events threaten the population viability of coastal birds?
493 *Journal of Applied Ecology*, 47: 720-730.
- 494 Vögeli, M., Serrano, D., Pacios, F. and Tella, J. L. (2010). The relative importance of patch
495 habitat quality and landscape attributes on a declining steppe-bird metapopulation.
496 *Biological Conservation*, 143: 1057–1067.
- 497 Wiens, J.A. (1989). *The Ecology of Bird Communities. Vol 2. Processes and Variations*.
498 Cambridge University Press, Cambridge, UK.
- 499 Wretenberg, J., Lindström, A., Svensson, S. and Pärt, T. (2007). Linking agricultural policies to
500 population trends of Swedish farmland birds in different agricultural regions. *Journal of*
501 *Applied Ecology*, 44: 933–941.

Figure 1(on next page)

Dupont's lark distribution in Spain according to Suárez, 2010 (light grey) and Dupont's lark populations included in this study (black).

The names of the Autonomous Communities where the species is present, are shown. The arrow refers to an isolated region belonging to the Community of Valencia. AN: Andalusia. AR: Aragon. CA: Catalonia. CL: Castile-Leon. CM: Castile-La Mancha. CV: Community of Valencia. NA: Navarre. RM: Region of Murcia.




Figure 2 (on next page)

Population size indices estimated by the Switching Linear Trend model for 92 Dupont's lark populations during the 2004 - 2015 period.

Time-points incorporated in the model as significant change-points on population trends are marked with asterisk (*).




Figure 3(on next page)

Population size indices predicted by Switching Linear Trend models during the 2004 - 2015 period for each Autonomous Community.

Time-points incorporated in models as significant change-points on population trends are marked with asterisk (*). AN: Andalusia. AR: Aragon. CA: Catalonia. CL: Castile-Leon. CM: Castile-La Mancha. CV: Community of Valencia. NA: Navarre. RM: Region of Murcia.




Table 1 (on next page)

Results of regional Switching Linear Trend models through the time series 2004-2015.

AN: Andalusia. AR: Aragon. CA: Catalonia. CL: Castile-Leon. CM: Castile-La Mancha. CV: Community of Valencia. NA: Navarre. RM: Region of Murcia.

	AN	AR	CA	CL	CM	CV	NA	RM
Number of populations	12	10	1	29	26	8	3	3
Local extinction events	6	0	1	9	6	3	1	0
Missing values (%)	38.2	81.6	58.3	49.1	63.1	44.8	63.9	47.2
Annual change rate (%)	-10.9	+1.5	+3.6	-8.4	+1.5	-2.5	-1.1	+2.6
95% Confidence Interval	[-16.2; -5.7]	[-2.3; +5.2]	[-33.5; +40.8]	[-10.0; -6.7]	[-2.1; +5.1]	[-5.7; +0.7]	[-7.9; +5.6]	[-2.2; +7.5]
TRIM Trend ^a	Steep decline	Uncertain	Uncertain	Steep decline	Uncertain	Uncertain	Uncertain	Uncertain
Wald-test change rate	-	-	0.04	-	-	-	-	-
p-value	-	-	> 0.05	-	-	-	-	-
Goodness-of-fit test								
Chi-squared (χ^2)	98.98	11.56	-	187.13	152.34	63.00	2.00	4.98
p-value χ^2	0.0158*	> 0.05	-	< 0.01	< 0.01	0.0152*	> 0.05	> 0.05
Likelihood Ratio (LR)	100.81	11.85	-	211.67	139.36	63.53	2.24	5.44
p-value LR	0.0115*	> 0.05	-	< 0.01	< 0.01	0.0136*	> 0.05	> 0.05
AIC	-41.19	-10.15	-	-74.33	-24.64	-18.47	-3.76	-18.56
Overdispersion	1.39	1.01	6.67	1.29	1.69	1.43	0.66	0.23
Serial correlation	0.09	-0.18	-0.06	0.39	0.20	0.30	-	0.06

- 1 P-values of accepted models are marked in bold
- 2 P-values of models near to acceptance threshold are marked with asterisk (*)
- 3 ^a Trend classification attending to TRIM criteria (Pannekoek and Van Strien 2006b)

Table 2 (on next page)

Assessment of Dupont's lark threat category.

Overall and average annual change rate obtained from trend analysis, and current threat category at National and Regional Catalogues of Endangered Species are shown. In addition, population size change in a 10-year period and corresponding threat category attending to A1 and A2 criteria applicable in the SCTS (Resolution 6th March 2017) are provided. The 95% Confidence Intervals are shown in brackets. Threat categories: Sensitive to Habitat Alteration (SHA), Vulnerable (VU) and Endangered (EN). AN: Andalusia. AR: Aragon. CA: Catalonia. CL: Castile-Leon. CM: Castile-La Mancha. CV: Community of Valencia. NA: Navarre. RM: Region of Murcia. SP: Spain.

	Overall change rate (%) from 2004 to 2015	Average annual change rate (%)	Current category of threat	Change rate for 10 years (%)	Category of threat – A1 criterion	Category of threat – A2 criterion
AN	-70.0 [-87.3; -52.7]	-10.9 [-16.2; -5.7]	VU ^a	-68.5 [-82.9; -44.4]	VU [EN; None]	EN [EN; VU]
AR	+17.1 [-30.2; +64.5]	+1.5 [-2.3; +5.2]	SHA ^b	+16.1 [-20.8; +66.0]	None [None; None]	VU* [VU*; VU*]
CA	+48.2 [-536.3; +632.6]	+3.6 [-33.5; +40.8]	-	+42.4 [-98.3; +2.9·10 ³]	None [EN; None]	VU* [EN; VU*]
CL	-50.8 [-60.8; -40.8]	-8.4 [-10.0; -6.7]	-	-58.4 [-65.1; -50.0]	VU [VU; VU]	EN [EN; EN]
CM	-59.0 [-78.9; -39.1]	+1.5 [-2.1; +5.1]	VU ^c	+16.1 [-19.1; +64.4]	None [None; None]	VU* [VU*; VU*]
CV	-26.8 [-55.3; +1.7]	-2.5 [-5.7; +0.7]	VU ^d	-22.4 [-44.4; +7.2]	None [None; None]	VU* [VU; VU*]
NA	-11.8 [-77.2; +53.6]	-1.1 [-7.9; +5.6]	SHA ^e	-10.5 [-56.1; +72.4]	None [VU; None]	VU* [EN; VU*]
RM	+37.4 [-53.6; +128.4]	+2.6 [-2.2; +7.5]	VU ^f	+29.3 [-19.9; +106.1]	None [None; None]	VU* [VU*; VU*]
SP	-41.3 [-50.2; -32.5]	-3.9 [-4.9; -2.8]	VU	-32.8 [-39.5; -24.7]	None [None; None]	VU [VU; none]

1 ^a Decree 23/2012 of 14 February 2012

2 ^b Decree 49/1995 of 28 March 1995

3 ^c Decree 33/1998 of 5 May 1998

4 ^d Decree 32/2004 of 27 February 2004

5 ^e Decree 563/1995 of 27 November 1995

6 ^f Law 7/1995 of 21 April 1995

7 * Minimum category of threat in accordance to the category of threat in the SCTS (Law 42/2007, 13th
8 December)