

Phylogenetic congruence of lichenised fungi and algae is affected by spatial scale and phylogenetic diversity

The role of species' interactions in structuring biological communities remains unclear. Mutualistic symbioses, involving close positive interactions between two distinct organismal lineages, provide an excellent means to explore the roles of both evolutionary and ecological processes in determining how positive interactions affect community structure. In this study, we investigate patterns of co-diversification between fungi and algae for a range of New Zealand lichens at the community, genus, and species levels and explore explanations for possible patterns related to spatial scale and pattern, phylogenetic diversity of the taxa considered, and the levelsampling replication. We assembled six, independent datasets to compare patterns in phylogenetic congruence with varied spatial extent of sampling, phylogenetic diversity and level of specimen replication. For each dataset, we used the DNA sequences from the ITS regions of both the fungal and algal genomes from lichen specimens to produce genetic distance matrices. Phylogenetic congruence between fungi and algae was quantified using distance-based redundancy analysis and we used geographic distance matrices in Moran's eigenvector mapping and variance partitioning to evaluate the effects of spatial variation on the quantification of phylogenetic congruence. Phylogenetic congruence was highly significant for all datasets and a large proportion of variance in both algal and fungal genetic distances was explained by partner genetic variation. Spatial variables, primarily at large and intermediate scales, were also important for explaining genetic diversity patterns in all but one very small-scale dataset. Interestingly, spatial structuring was stronger for fungal than algal genetic variation. As the spatial extent of the samples increased,so too did the proportion of explained variation that was shared between the spatialvariables and the partners' genetic variation. Different lichen taxa showed some variation in their phylogenetic congruence and spatial genetic patterns and where greater sample replication was used, the amount of variation explained by partner genetic variation increased. Our results suggest that the phylogenetic congruencepattern, at least at small spatial scales, is

likely due to reciprocal co-adaptation or co-dispersal. However, the detection of these patterns varies among different lichen taxa, across spatial scales and with different levels of sample replication. This work provides insight into the complexities faced in determining how evolutionary and ecological processes may interact to generate diversity in symbiotic association patterns at the population and community levels. Further, it highlights the critical importance of considering sample replication, taxonomic diversity and spatial scale in designing studies of co-diversification.

1 **Phylogenetic congruence of lichenised fungi and algae is affected by spatial scale and phylogenetic** 2 **diversity**

3 Hannah L. Buckley*, Arash Rafat, Johnathon D. Ridden, Robert H. Cruickshank, Hayley J. Ridgway,
4 Adrian M. Paterson

5 Department of Ecology, PO Box 85084, Lincoln University, Lincoln, Canterbury 7467, New Zealand

6 * Author for correspondence: Hannah L. Buckley (Hannah.Buckley@lincoln.ac.nz)

7 INTRODUCTION

8 Ecologists still do not have a full understanding of how species' interactions affect the structure of
9 biological communities. We have a long history of theoretical and empirical work on the roles of
10 competition and predation (Chase & Leibold 2003; Tilman 1982), but a much poorer understanding of
11 how positive interactions, such as facilitation and mutualisms, drive community phenomena, such as
12 species' diversity (Gross 2008; Stachowicz 2001). Over the last few decades, there has been an increase
13 in interest in the role of positive interactions, with many empirical studies showing that it is the balance
14 between positive and negative interactions that is important in structuring communities (e.g., Bertness &
15 Callaway 1994; Elias et al. (2008; LaJeunesse 2002; Thrall et al. 2007; Waterman et al. 2011).


16 Because mutualistic symbioses involve very close positive interactions between two distinct
17 organismal lineages, they provide an excellent opportunity to specifically explore how positive
18 interactions influence community structure and to evaluate the relative importance of evolutionary and
19 ecological processes in the way that positive interactions affect community structure. Tightly interacting
20 taxa, such as host-parasite systems and certain plants and their insect pollinators, typically show high
21 degrees of phylogenetic congruence between hosts and associates, where the phylogeny of one taxon
22 closely tracks the phylogeny of the partner taxon (Cuthill & Charleston ; Light & Hafner 2008; Quek et al.
23 2004; Subbotin et al. 2004), largely due to co-evolutionary processes. What is not as clear is whether
24 most obligate species-level symbiotic relationships, such as ~~lichens~~ and corals, also have measurable seen in
25 levels of codivergence. Typically, these diffuse symbiotic and mutualistic interactions are complicated by
26 variation in the specificity of the partners or where more than one symbiotic partner is involved, such as
27 vascular plants and their root-inhabiting mutualists (Hollants et al. 2013; Lanterbecq et al. 2010; Walker et
28 al. 2014). Further, it is much less clear that the causal processes involved in these coevolving symbiotic
29 relationships will produce a pattern of codivergence given the increased opportunity for host switching.

30 Lichens are a classic example of a mutualistic symbiosis. Lichen thalli are the result of an
31 association between a fungus (the mycobiont) and a photobiont, which is usually a green alga, but may
32 also be a cyanobacterium (Nash 2008). Around 3-4% of lichens are tripartite involving a symbiosis of

both a green alga and a cyanobacterium (Henskens et al. 2012). In all lichens, the photobiont provides photosynthate to the mycobiont, which in turn provides habitat, water and nutrients to the photobiont (Honegger 1991; Nash 2008). The symbionts of lichens are relatively poorly known because they are very often difficult to culture and identify, particularly many of the photobiont partners (Ahmadjian 1993; del Campo et al. 2013; Grube & Muggia 2010; Honegger 1991). However, in the last two decades a great deal of molecular work has been done to address this, showing variable patterns in partner identity and other patterns of association (DePriest 2004). This variable, and seemingly diffuse, mutualism provides a complex model system for addressing questions regarding the role of evolutionary processes in forming and driving ecological patterns in species interactions and how this affects community structure.

The first step in understanding how a mutualistic symbiosis might affect community structure is to determine whether or not there is specificity in the symbiosis. In the case of lichens, we know that although there is no evidence for very tight co-evolutionary relationships at the species level, phylogenetic patterns have been observed where particular fungal taxa preferentially partner with particular algal taxa (Fernández-Mendoza et al. 2011; Yahr et al. 2004) resulting in a correlation in the respective genetic distances of each partner. For example, Widmer et al. (2012) found similar genetic structures for the lichen symbiosis between the mycobiont, *Lobaria pulmonaria*, and its photobiont, *Dictyochoropsis reticulata* within Europe. Conversely, it seems that some lichenised algal taxa are capable of partnering with a range of fungal taxa (Beck 1999); thus, the specificity (*sensu* Piercey-Normore & DePriest 2001) of the symbiosis appears to be driven by fungal selectivity (*sensu* Piercey-Normore & DePriest 2001). For example, Nelsen & Gargas (2008) observed a lack of phylogenetic congruence between the algae and fungi in specimens from the lichen genus *Lepraria*. Thus, it appears that there is variability in association patterns among different lichen taxa (Fahselt 2008).

Several mechanisms are proposed to underpin the patterns in phylogenetic congruence observed for lichens. First, co-evolutionary processes, whereby one partner adapts to take better advantage of the symbiosis, may lead to a reciprocal adaptive evolutionary change in the other partner, although there is little evidence for this in the literature (Yahr et al. 2006). Second, many lichens asexually reproduce, either by fragmentation or specialised structures (Walser 2004), so that the resulting offspring lichens contain clones of their parents, otherwise known as ‘vertical transmission’ (Dal Grande et al. 2012; Werth & Scheidegger 2011). Sexual reproduction in green algal photobionts (apart from those in the Trentepholiales) is thought to be extremely rare within lichen thalli (Friedl & Büdel 2008; Sanders 2005), despite evidence of recombination within these taxa (Kroken & Taylor 2000). If symbiont co-dispersal is coupled with genetic drift, a pattern of co-diversification is likely to emerge. However, ‘horizontal transmission’ of photobionts into newly forming thalli is thought to occur, such as in the form of escaped zoospores (Beck et al. 1998), and population genetics studies show evidence of algal switching (Dal Grande et al. 2012; Kroken & Taylor 2000; Nelsen & Gargas 2008; Piercey-Normore & DePriest 2001).

Further, most green algal photobionts commonly occur in a free-living state, although for some taxa, such as *Trebouxia*, much about their life cycles and availability in the environment is unknown (Sanders 2005). A free-living state is likely to decrease the importance of vertical transmission and decrease the congruence of phylogenetic patterns. Third, spatial structure in fungal and algal distributions could drive patterns in phylogenetic congruence. Spatial structure in either fungi or algae could arise through dispersal limitation, habitat fragmentation, or niche differentiation, such as variation in habitat preferences. For example, (Werth et al. 2007) showed that the mycobiont of the cyanobacterial lichen, *Lobaria pulmonaria*, varied genetically over spatial scales of less than a few kilometres in a pasture-woodland landscape and suggested that this could be caused by dispersal limitation among habitat patches. Peksa & Škaloud (2011) showed that the spatial distribution patterns of ~~the~~ *Asterochloris* in Europe and North America, the green algal photobiont for two different lichen genera, were driven by substrate type and relative exposure to rain and sun. If spatial structure in fungal and algal distributions resulted in limited availability of one or both partners relative to the other, this would lead to a congruent phylogenetic pattern.  example, Marini et al. (2011) found that communities of the 822 species of epiphytic lichens within Italy with different photobiont types (Chlorococcoid green algae, Cyanobacteria or *Trentepohlia*) showed different biogeographic patterns across climatically different areas. Thus, if spatial structure in genetic variation results from differential distributions of algal or fungal ecotypes, this could result in phylogenetic congruence for specimens compared across environmental gradients. For example, the patterns in variable algal selectivity that Vargas Castillo & Beck (2012) observed within the genus *Caloplaca* in the Atacama Desert in Northern Chile were related to changing habitat conditions along an altitudinal gradient.

Although much recent research has shown that lichenised fungi specialise on particular algae regardless of the availability of other species, most work has been conducted at the within-species and within-genus levels, and much less often at higher levels of phylogenetic diversity. Such patterns and their explanations, like most ecological phenomena, are likely to be scale-dependent and related to both small scale processes, such as the dispersal of lichen propagules, as well as larger scale biogeographic processes and climatic variation. In addition, these patterns are likely to depend on the amount of phylogenetic diversity contained within the dataset considered. We expect that if co-evolutionary processes play a role, then phylogenetic congruence should be stronger when considering higher levels of phylogenetic diversity because they are the accumulation of a longer period of evolutionary change. To examine these patterns and effects, we tested patterns of association for a range of New Zealand lichens at the community, genus, and species levels. We assembled six independent datasets that varied in the spatial extent of sampling, phylogenetic diversity and the level of specimen replication so that we could compare patterns in phylogenetic congruence across these variables. We have taken a novel approach to the analysis of phylogenetic congruence that uses Moran's eigenvector mapping, distance-based

redundancy analysis and variance partitioning, which allows us to evaluate the effects of sampling on the quantification of phylogenetic congruence. We interpret the patterns in the light of the relative importance of the mechanisms driving variation in co-diversification patterns.

METHODS

Lichen specimen collection

Lichen thallus samples were collected from many mapped locations around both the North and South Islands of New Zealand (under New Zealand Department of Conservation low-impact research and collection permit, CA-31641-FLO). Samples were taken either from the ground, or from trees and structures like fence posts and stored in paper envelopes. Each sample was identified to the lowest taxonomic level possible and assigned a specific sample code. Six, non-overlapping sample sets were collected (Table 1): (1) Multiple lichen species collected on mountain beech (*Nothofagus solandri* var. *cliffortioides*) trees within less than 1 km² of Flock Hill Station (Buckley 2011), (2) *Usnea* spp. specimens from Flock Hill Station, (3) *Usnea* spp. specimens from Craigieburn Forest Park, (4) *Usnea* spp. specimens from sites around New Zealand, and (5) *Ramalina* spp. specimens from around New Zealand (Figure 1).

Molecular analysis

Total DNA was extracted from surface sterilised lichens using Plant DNA Mini Kit (Bioline, London, UK) following the manufacturer's instructions. Photobiont and mycobiont ITS rRNA were amplified using specific algal (nr-SSU-1780-5' Algal and ITS4) and fungal (nr-SSU-1780-5' Fungal and ITS) primers described by (Piercey-Normore & DePriest 2001). PCR amplification was performed in a 25 µl reaction volume. Each 25 µl reaction contained 1× GoTaq® Green Master Mix, 5 pmol of each primer, 10 µg purified bovine serum albumin (BSA; New England BioLabs, Ipswich, MA, USA) and 1 µl of the extracted DNA (25-30 µg/µl). The thermal cycle for the algal reaction was as follows: initial denaturation at 94°C for 2 min, then cycles of denaturation at 94°C for 30 s, annealing at 50°C for 45 s, extension at 72°C for 2 min, then a final extension at 72°C for 7 min. The thermal cycle for fungal rRNA amplification was the same as the algal one except the annealing time was increased to 54°C.

Data analysis

For each of the six datasets, algal and fungal DNA sequences were aligned separately using 'Prankster' (Löytynoja & Goldman 2005) with the default parameters. These alignments were used to calculate genetic distance matrices from the raw distances using uncorrected p-distances (Paradis 2006) implemented by the 'dist.dna' function in the 'ape' package in R (R Core Team 2013). We repeated the analyses using a genetic distance matrix calculated using the TN93 substitution model (Tamura & Nei

1993). The results (not shown) were congruent with those obtained using p-distances, so we present the p-distance results only.

To relate fungal and algal genetic distances to each other and to describe their variation in space, distance matrices were used in a combined analysis using Moran's eigenvector mapping and variance partitioning, following (Borcard et al. 2011, pp. 258). This analysis was used to describe and partition the variation in algal genetic distances between (a) the fungal genetic distance matrix and (b) a matrix of spatial variables. The spatial variables were derived using the Moran's eigenvector maps (MEMs) procedure (Borcard et al. 2011) implemented using the 'pcnm' function in the R package 'vegan', which uses a principal coordinates analysis to represent different scales of spatial variation for the given set of sample locations (Borcard et al. 2011). MEM analysis produces one fewer spatial variables than there are sample points, describing all possible spatial variation in the data from broad scale variation to very fine scale variation. Only the most important subset of these spatial variables (MEMs) was included in a distance-based redundancy analysis (db-RDA) analysis, which relates multivariate data (algal genetic distance matrix) to explanatory matrices (fungal genetic distance matrix and spatial variables). The selected MEMs were those that were significantly related to the algal distance matrix in a distance-based RDA and forward selection procedure using the 'capscale' and 'ordistep' functions in the 'vegan' package in R (Oksanen et al. 2013). Variance partitioning calculations were conducted following procedures outlined in (Borcard et al. 2011). The 'capscale' function performs a redundancy analysis that seeks the series of linear combinations of the explanatory factors that best describe variation in the response matrix, constrained by the two explanatory matrices (Borcard et al. 2011). The variance partitioning procedure computes R^2 canonical values analogous to the adjusted R^2 values produced in multiple regression (Peres-Neto et al. 2006). The analysis indicates how much total variation in the response matrix (e.g., algal genetic distance) is explained by each of the explanatory matrices alone, as well as the component of shared variation, e.g., spatially structured variation in fungal genetic distances. This analysis was also performed using the fungal genetic distance matrix as the dependent matrix and the algal genetic distance matrix as the explanatory matrix to allow comparison of the degree of spatial correlation in each of the matrices.

To test the significance of the phylogenetic congruence between fungi and algae for each of the six datasets, we used the Procrustes approach to co-phylogeny (PACo, Balbuena et al. 2012). This procedure performs a principal coordinates analysis on the algal genetic distance matrix followed by a Procrustes rotation of the fungal genetic distance matrix, while retaining the information that algae and fungi are paired in particular lichen specimens (Balbuena et al. 2012). A sum of squares is calculated from the individual residuals for each specimen that represents the lack of fit of the fungal genetic distance matrix onto the principal coordinate analysis result for the algal genetic distance matrix (Balbuena et al. 2012). The algal-fungal pairing matrix, i.e., which alga is paired with which fungus, is then randomised

10,000 times and the sums of square values recalculated. The observed sum of squares value is then compared to the distribution of values from the randomisations to determine the probability of obtaining the observed result under random expectation (Balbuena et al. 2012). The magnitude of the residual for each lichen specimen shows its relative lack of fit to a co-diversification pattern. Therefore, for three datasets for which we had additional information on specimen traits, we compared individual residuals among specimens to determine which ones contributed most to the observed association pattern. These three datasets (and trait information) were the Flock Hill community (growth form), Flock Hill *Usnea* and New Zealand *Usnea* datasets (apothecia present or absent). Raw genetic and geographic distance matrices are provided as supplemental files.

RESULTS

The six datasets captured a wide range of geographic extent and, unsurprisingly, the datasets with greatest numbers of different lichen morphotypes contained the greatest fungal genetic diversities (Table 1). Algal genetic diversity had only a weak, positive correlation with fungal genetic diversity (Pearson's $r = 0.39$; $n = 6$).

The db-RDA showed that spatial variables were important for explaining genetic diversity patterns in all but the Flock Hill community dataset (Table 2, Significant MEMs). Large-scale variables, e.g. low numbered MEMs, were important for all spatially-structured datasets. Genetic variation in *Ramalina* algae and fungi was significantly related to several large-scale MEMs, showing that spatial pattern in relatedness varied at the larger scales within this spatial extent, such as between the North and South Islands (Table 2). Some intermediate scale MEMs were also important, illustrating additional, more complex, spatial patterns. Similarly, for the *Usnea* datasets, large-scale MEMs were of greatest importance, along with some intermediate-scale, but no very fine-scale, MEMs.

Variance partitioning showed that a large proportion of variance in both the algal and fungal genetic variation was explained by genetic variation in the partner (Figure 2). In general, the proportion of explained variation was high, at 75% or more (Table 2, Figure 2). Interestingly, across all datasets for which spatial variation was important, the fungal genetic variation was better explained by spatial variables than was the algal genetic variation for the same lichens (Figure 2). As the spatial extent of the samples increased, so too did the proportion of explained variation that was shared between the spatial variables and the partners' genetic variation (Figure 2).

PACo analysis showed that the co-diversification for all datasets, regardless of whether the dependent matrix was the algae or the fungi, was significant at the $\alpha = 0.05$ level (Table 1), indicating that it is very unlikely that the correlations in the fungal and algal genetic distances have arisen by chance. When comparing the individual residuals from the PACo analysis, for the Flock Hill community dataset, the

Usnea and *Ramalina* specimens, which made up all but one of the foliose specimens, had relatively good fit to the co-diversification pattern, compared to most other taxa (Figure 3). For the Flock Hill *Usnea* and New Zealand *Usnea* datasets, the presence of apothecia did not appear to be related to the relative specimen contribution to the lack of phylogenetic congruence (Figure 3).

DISCUSSION

We show that there is a strong and significant pattern of phylogenetic congruence for the algae and fungi found in New Zealand lichens, both across multiple taxa within a community and within two widespread genera sampled at spatial extents from less than 1 km², to the whole country (*c.* 260,000 km²). For instance, at the small spatial scales of the Flock Hill study site, there was surprisingly high phylogenetic congruence, as demonstrated by the high variance explained by partner genetic variation. Such congruence suggests that fungi and algae are not randomly distributed across the lichen symbiosis for the taxa considered. Our results show that this non-random pattern is affected by (1) spatial scale, (2) taxon considered, (3) phylogenetic diversity and (4) level of sample replication.

The amount of genetic variation explained by spatially-structured partner genetic variation increased with increasing spatial extent (Figure 2) suggesting that a large amount of phylogenetic congruence is likely to be due to the distributions of fungi and algae at larger, rather than smaller, spatial scales. In addition, the spatial structuring of *Usnea* and *Ramalina* fungal genetic distances was more prominent than for algae, suggesting that drivers of fungal distributions were more important in determining these congruence patterns than drivers of algal distributions. It appears that the algae are more dependent on the distribution of the fungi than the fungi are on the algae. This is in contrast to the findings of (Marini et al. 2011), who suggested that the richness of lichens within Italy was dependent on the distribution of different algal types. However, this study encompassed much larger phylogenetic diversity, i.e., kingdom-level diversity in photobionts, than is considered here. Thus, the strong spatial signal in our results show that at large spatial scales, at least some of the co-diversification pattern appears likely to be due to processes other than co-evolution or vertical transmission. If factors such as variation in habitat preferences or photobiont availability were important, we would expect to see an increase in the proportions of genetic variance being explained by spatially-structured variation in the genetic distances of the partners. This is indeed the result we observe across the four *Usnea* datasets. One other paper that mentions the effects of spatial scale on phylogenetic congruence patterns is a study of *Cladonia* lichens across six discrete rosemary scrub sites in three regions in Florida (Yahr et al. 2004). Their findings illustrated that, across Florida, photobiont genetic variation was not significantly spatially structured, despite different fungi occurring at different sites. These contrasting findings suggest that different patterns of phylogenetic congruence may occur in different lichen taxa.

The variation in the patterns from the PACo for individual lichen specimens also suggest that the taxon considered may affect the observed phylogenetic congruence pattern. The fruticose taxa had lower residual values in the PACo analysis showing that they contributed relatively more to the phylogenetic congruence pattern than did crustose and foliose taxa. However, these lichen specimens were primarily *Usnea* and *Ramalina* suggesting that this variation may have a phylogenetic basis, rather than being due to the life form. There was little pattern observed when we related the presence of apothecia to their contribution to the co-diversification pattern for the *Usnea*-only datasets (Figure 3) suggesting that, at least at this scale and level of replication, ability for the fungal component to sexually reproduce had little to do with the observed pattern. Residual values are not comparable among datasets, so it is not possible to evaluate these differences among spatial scales. Overall, these results are consistent with the variable patterns in lichen co-diversification in the literature (Ruprecht et al. 2012; Vargas Castillo & Beck 2012).

Our study encompassed a range of phylogenetic diversities for both fungi and algae. Specifically, when fungal diversity was very high, algal diversity was also relatively high, despite the very small spatial scale (Table 1, Flock Hill community dataset). For *Usnea*, when fungal genetic diversity was low, algal diversity was low or high depending on whether the spatial scale was small or large, respectively (Table 1). However, if we consider only the two datasets at the smallest spatial scale, despite the variation in genetic diversity, the pattern of phylogenetic congruence did not vary; the Flock Hill datasets both show high levels of phylogenetic congruence despite having the largest difference in genetic diversity (Figure 2). This is consistent with arguments suggesting that coevolution is not an important driver of the lichen symbiosis (Yahr et al. 2006), because if coevolution was important, then we would expect to see an increase in the degree of phylogenetic congruence with increasing genetic diversity.

By contrasting the result from the NZ *Usnea* dataset and the NZ *Usnea* replicated dataset, we can consider the effects of within-site replication on the phylogenetic congruence signal. This result shows that where greater replication was used, the amount of variation explained by partner genetic variation increased. This highlights the importance of considering sampling design in studies of phylogenetic congruence.

This work leads us to more questions regarding variation in phylogenetic congruence patterns and its causes, which are likely to be scale-dependent. We need better understanding of the factors influencing association patterns including algal availability and niche differentiation, dispersal (metapopulation dynamics), and reproductive traits. In particular, we need to understand the effects of lichen reproductive modes better as many lichens reproduce both sexually and asexually. When the fungal component sexually reproduces, the symbiosis must re-form which gives the fungus an opportunity to change symbiotic partners. With asexual reproduction, the fungus and alga disperse together, and so, a predominance of asexual reproduction may be one of the reasons we see such a strong co-diversification

signal in these taxa. However, the results of this study suggest that the spatial distributions of the fungi and algae may also be important in determining the nature of the symbiosis, particularly at larger spatial scales, as has been observed in some parasite lineages (e.g., du Toit et al. 2013). Thus, we need to do more work on algal and fungal availability to determine if habitat preferences and/or dispersal limitation led to some of the spatial patterns that we see. In addition, we need better understanding of the availability of free living algae to lichens and what their microhabitat preferences are.

This study is consistent with previous research showing that, despite the diffuse nature of the lichen mutualistic symbiosis, there is strong selectivity within the association. We show that despite spatial structuring in algal, and particularly fungal, distributions at large spatial scales, the phylogenetic congruence pattern, at least at small spatial scales, is due to reciprocal co-adaptation or, more likely, to co-dispersal. However, the influence of these processes is likely to differ among different lichen taxa. This work gives us insight into some of the complexities we face in determining how evolutionary and ecological processes may interact to generate diversity in symbiotic association patterns at the population and community levels.

ACKNOWLEDGEMENTS

The authors would like to thank: Elizabeth Bargh, Jennifer Bannister, Alison Knight, Mike Bowie, John Marris, Dan Blachon, Brad Case and Sam Case for specimen collection; Ursula Brandes, Hamish Maule, Ben Myles and Natalie Scott for field assistance; Richard Hill for land access permission; Ben Myles, David Galloway, Jennifer Bannister, and Alison Knight for lichen identification; Dalin Brown, Seelan Baskarathevan and Chantal Probst for assistance with molecular analysis; Norma Merrick for DNA sequencing; Brad Case for GIS work; the Lincoln University Spatial Ecology and Molecular Ecology Groups for discussion.

REFERENCES

- Ahmadjian V. 1993. *The lichen symbiosis*. New York: Wiley.
- Balbuena JA, Míguez-Lozano R, and Blasco-Costa I. 2012. PACo: A novel Procrustes application to cophylogenetic analysis. *PLOS ONE* 8:e61048.
- Beck A. 1999. Photobiont Inventory of a Lichen Community Growing on Heavy-Metal-Rich Rock. *The Lichenologist* 31:501-510.
- Beck A, Friedl T, and Rambold G. 1998. Selectivity of photobiont choice in a defined lichen community: inferences from cultural and molecular studies. *New Phytologist* 139:709-720.
- Bertness MD, and Callaway R. 1994. Positive interactions in communities. *Trends in Ecology & Evolution* 9:191-193.
- Borcard D, Gillert R, and Legendre P. 2011. *Numerical Ecology with R*. New York: Springer.
- Buckley HL. 2011. Isolation affects tree-scale epiphytic lichen community structure on New Zealand mountain beech trees. *Journal of Vegetation Science* 22:1062-1071.

- Chase JM, and Leibold MA. 2003. *Ecological Niches: Linking Classical and Contemporary Approaches*. Chicago: The University of Chicago Press.
- Cuthill JH, and Charleston M. Phylogenetic codivergence supports coevolution of mimetic *Heliconius* butterflies. *PLOS ONE* 7:e36464.
- Dal Grande F, Widmer I, Wagner HH, and Scheidegger C. 2012. Vertical and horizontal photobiont transmission within populations of a lichen symbiosis. *Molecular Ecology* 21:3159-3172.
- del Campo EM, Catala S, Gimeno J, del Hoyo A, Martínez-Alberola F, Casano LM, Grube M, and Barreno E. 2013. The genetic structure of the cosmopolitan three-partner lichen *Ramalina farinacea* evidences the concerted diversification of symbionts. *FEMS Microbiology Ecology* 83:310-323.
- DePriest PT. 2004. Early molecular investigations of lichen-forming symbionts: 1986–2001. *Annual Review of Microbiology* 58:273-301.
- du Toit N, van Vuuren BJ, Matthee S, and Matthee CA. 2013. Biogeography and host-related factors trump parasite life history: limited congruence among the genetic structures of specific ectoparasitic lice and their rodent hosts. *Molecular Ecology* 22:5185-5204.
- Elias M, Gompert Z, Jiggins C, and Willmott K. (2008. Mutualistic Interactions Drive Ecological Niche Convergence in a Diverse Butterfly Community. *PLoS Biol* 6:e300.
- Fahselt D. 2008. Individuals and populations of lichens. In: Nash TH, ed. *Lichen Biology*. 2nd Edition ed. Cambridge: Cambridge University Press.
- Fernández-Mendoza F, Domaschke S, García MA, Jordan P, Martín MP, and Printzen C. 2011. Population structure of mycobionts and photobionts of the widespread lichen *Cetraria aculeata*. *Molecular Ecology* 20:1208-1232.
- Friedl T, and Büdel B. 2008. Photobionts. In: Nash TH, ed. *Lichen Biology*. 2nd Edition ed. Cambridge: Cambridge University Press.
- Gross K. 2008. Positive interactions among competitors can produce species-rich communities. *Ecology Letters* 11:929-936.
- Grube M, and Muggia L. 2010. Identifying algal symbionts in lichen symbioses. In: Nimis PL, and Vignes Lebbe R, eds. *Tools for Identifying Biodiversity: Progress and Problems*. Trieste: Edizioni Università di Trieste, 295-299.
- Henskens FL, Green TGA, and Wilkins A. 2012. Cyanolichens can have both cyanobacteria and green algae in a common layer as major contributors to photosynthesis. *Annals of Botany* 110:555-563.
- Hollants J, Leliaert F, Verbruggen H, De Clerck O, and Willems A. 2013. Host specificity and coevolution of Flavobacteriaceae endosymbionts within the siphonous green seaweed *Bryopsis*. *Molecular Phylogenetics and Evolution* 67:608-614.
- Honegger R. 1991. Functional aspects of the lichen symbiosis. *Annual Review of Plant Physiology and Plant Molecular Biology* 42:553-578.
- Kroken S, and Taylor JW. 2000. Phylogenetic Species, Reproductive Mode, and Specificity of the Green Alga *Trebouxia* Forming Lichens with the Fungal Genus *Letharia*. *The Bryologist* 103:645-660.
- LaJeunesse T. 2002. Diversity and community structure of symbiotic dinoflagellates from Caribbean coral reefs. *Marine Biology* 141:387-400.
- Lanterbecq D, Rouse GW, and Eeckhaut I. 2010. Evidence for cospeciation events in the host–symbiont system involving crinoids (Echinodermata) and their obligate associates, the myzostomids (Myzostomida, Annelida). *Molecular Phylogenetics and Evolution* 54:357-371.

- 353 Light JE, and Hafner MS. 2008. Codivergence in Heteromyid Rodents (Rodentia: Heteromyidae)
- 354 and Their Sucking Lice of the Genus *Fahrenholzia* (Phthiraptera: Anoplura). *Systematic*
- 355 *Biology* 57:449-465.
- 356 Löytynoja A, and Goldman N. 2005. An algorithm for progressive multiple alignment of
- 357 sequences with insertions. *Proceedings of the National Academy of Sciences of the United*
- 358 *States of America* 102:10557-10562.
- 359 Marini L, Nascimbene J, and Nimis PL. 2011. Large-scale patterns of epiphytic lichen species
- 360 richness: Photobiont-dependent response to climate and forest structure. *Science of The*
- 361 *Total Environment* 409:4381-4386.
- 362 Nash TH. 2008. Introduction. In: Nash TH, ed. *Lichen Biology*. 2nd Edition ed. Cambridge:
- 363 Cambridge University Press.
- 364 Nelsen MP, and Gargas A. 2008. Dissociation and horizontal transmission of codispersing lichen
- 365 symbionts in the genus *Lepraria* (Lecanorales: Stereocaulaceae). *New Phytologist*
- 366 177:264-275.
- 367 Oksanen J, Blanchet FG, Kindt R, Legendre P, Minchin PR, O'Hara RB, Simpson GL, Solymos
- 368 P, Stevens MHH, and Wagner H. 2013. vegan: Community Ecology Package. R package
- 369 version 2.0-9 ed. <http://CRAN.R-project.org/package=vegan>.
- 370 Paradis E. 2006. *Analysis of phylogenetics and evolution with R*. New York: Springer.
- 371 Peksa O, and Škaloud P. 2011. Do photobionts influence the ecology of lichens? A case study of
- 372 environmental preferences in symbiotic green alga *Asterochloris* (Trebouxiophyceae).
- 373 *Molecular Ecology* 20:3936-3948.
- 374 Peres-Neto PR, Legendre P, Dray S, and Borcard D. 2006. Variation partitioning of species data
- 375 matrices: Estimation and comparison of fractions. *Ecology* 87:2614-2625.
- 376 Piercey-Normore MD, and DePriest PT. 2001. Algal switching among lichen symbioses.
- 377 *American Journal of Botany* 88:1490-1498.
- 378 Quek S-P, Davies SJ, Itino T, and Pierce NE. 2004. Codiversification in ant-plant mutualism:
- 379 stem texture and the evolution of host use in *Crematogaster* (Formicidae: Myrmicinae)
- 380 inhabitants of *Macaranga* (Euphorbiaceae). *Evolution* 58:554-570.
- 381 R Core Team. 2013. R: A language and environment for statistical computing. Vienna, Austria: R
- 382 Foundation for Statistical Computing.
- 383 Ruprecht U, Brunauer G, and Printzen C. 2012. Genetic diversity of photobionts in Antarctic
- 384 lecideoid lichens from an ecological view point. *The Lichenologist* 44:661-678.
- 385 Sanders WB. 2005. Observing microscopic phases of lichen life cycles on transparent substrata
- 386 placed in situ. *The Lichenologist* 37:373-382.
- 387 Stachowicz JJ. 2001. Mutualism, Facilitation, and the Structure of Ecological Communities:
- 388 Positive interactions play a critical, but underappreciated, role in ecological communities
- 389 by reducing physical or biotic stresses in existing habitats and by creating new habitats on
- 390 which many species depend. *BioScience* 51:235-246.
- 391 Subbotin SA, Krall EL, Riley IT, Chizhov VN, Staelens A, De Loose M, and Moens M. 2004.
- 392 Evolution of the gall-forming plant parasitic nematodes (Tylenchida: Anguinidae) and
- 393 their relationships with hosts as inferred from Internal Transcribed Spacer sequences of
- 394 nuclear ribosomal DNA. *Molecular Phylogenetics and Evolution* 30:226-235.
- 395 Tamura K, and Nei M. 1993. Estimation of the number of nucleotide substitutions in the control
- 396 region of mitochondrial DNA in humans and chimpanzees. *Molecular Biology and*
- 397 *Evolution* 10:512-526.
- 398 Thrall PH, Hochberg ME, Burdon JJ, and Bever JD. 2007. Coevolution of symbiotic mutualists
- 399 and parasites in a community context. *Trends in Ecology & Evolution* 22:120-126.
- 400 Tilman D. 1982. *Resource competition and community structure*. Princeton: Princeton University
- 401 Press.

- 402 Vargas Castillo R, and Beck A. 2012. Photobiont selectivity and specificity in Caloplaca species
403 in a fog-induced community in the Atacama Desert, northern Chile. *Fungal Biology*
404 116:665-676.
- 405 Walker DM, Castlebury LA, Rossman AY, and Struwe L. 2014. Host conservatism or host
406 specialization? Patterns of fungal diversification are influenced by host plant specificity in
407 Ophiognomonia (Gnomoniaceae: Diaporthales). *Biological Journal of the Linnean*
408 *Society* 111:1-16.
- 409 Walser J-C. 2004. Molecular evidence for limited dispersal of vegetative propagules in the
410 epiphytic lichen Lobaria pulmonaria. *American Journal of Botany* 91:1273-1276.
- 411 Waterman RJ, Bidartondo MI, Stofberg J, Combs JK, Gebauer G, Savolainen V, Barraclough TG,
412 and Anton P. 2011. The Effects of Above- and Belowground Mutualisms on Orchid
413 Speciation and Coexistence. *The American Naturalist* 177:E54-E68.
- 414 Werth S, Gugerli F, Holderegger R, Wagner HH, Csencsics D, and Scheidegger C. 2007.
415 Landscape-level gene flow in Lobaria pulmonaria, an epiphytic lichen. *Molecular*
416 *Ecology* 16:2807-2815.
- 417 Werth S, and Scheidegger C. 2011. Congruent Genetic Structure in the Lichen-Forming Fungus
418 Lobaria pulmonaria and Its Green-Algal Photobiont. *Molecular Plant-Microbe*
419 *Interactions* 25:220-230.
- 420 Widmer I, Dal Grande F, Excoffier L, Holderegger R, Keller C, Mikryukov VS, and Scheidegger
421 C. 2012. European phylogeography of the epiphytic lichen fungus Lobaria pulmonaria
422 and its green algal symbiont. *Molecular Ecology* 21:5827-5844.
- 423 Yahr R, Vilgalys R, and Depriest PT. 2004. Strong fungal specificity and selectivity for algal
424 symbionts in Florida scrub Cladonia lichens. *Molecular Ecology* 13:3367-3378.
- 425 Yahr R, Vilgalys R, and DePriest PT. 2006. Geographic variation in algal partners of Cladonia
426 subtenuis (Cladoniaceae) highlights the dynamic nature of a lichen symbiosis. *New*
427 *Phytologist* 171:847-860.

Figure 1 (on next page)

Map

Figure 1: Maps showing sample collection locations for (A) the three New Zealand datasets, (B) the Craigieburn *Usnea* dataset and (C) the Flock Hill community and *Usnea* dataset. Note that the Craigieburn samples were collected along a road running along an elevation gradient.




Figure 1: Maps showing sample collection locations for (A) the three New Zealand datasets, (B) the Craigieburn *Usnea* dataset and (C) the Flock Hill community and *Usnea* dataset. Note that the Craigieburn samples were collected along a road running along an elevation gradient.

Figure 2_(on next page)

Variance partitioning graph

Figure 2: Bar charts showing variance partitioning for six independent datasets modelled as algal genetic variance as a function of fungal genetic variance and spatial variation (left) and fungal genetic variance as a function of algal genetic variance and spatial variation (right). The total variation in genetic distance is explained by partner genetic distance (red), independent spatial variation (blue), and spatially-structured variation in partner genetic distances (purple). Unexplained variation is shown in grey. The number of specimens sampled (n) is given for each dataset.




Figure 3_(on next page)

Species' bar charts

Figure 3: Bar charts showing the individual lichen specimen contribution to the Procrustes sums of squares for (A) the New Zealand *Usnea* dataset (n = 111), (B) the Craigieburn *Usnea* dataset (n = 36) and (C) the Flock Hill all specimen dataset (n = 67). Dashed line indicates the median sums of squares value. Bars for (C) are coloured by growth form: crustose (black), foliose (red) and fruticose (green) and for the other two datasets black bars indicate specimens that had apothecia and white bars are those that were asexual. Note that in (C) all but the first from the left of the fruticose specimens were specimens of *Usnea* or *Ramalina*.




Table 1 (on next page)

List of datasets

Table 1: List of the datasets analysed showing the number of specimens sampled, the approximate number of lichen morphotypes, and the number of sites samples. Also given are the maximum distance between two sample points (Spatial extent) and the mean (standard deviation) genetic distance for each matrix. Note that only 35 of the 67 Flock Hill multiple species dataset specimens were mapped and were therefore analysed separately.

Table 1: List of the datasets analysed showing the number of specimens sampled, the approximate number of lichen morphotypes, and the number of sites samples. Also given are the maximum distance between two sample points (Spatial extent) and the mean (standard deviation) genetic distance for each matrix. Note that only 35 of the 67 Flock Hill multiple species dataset specimens were mapped and were therefore analysed separately.

Dataset	Number of specimens	Taxonomic variation	Number of sites	Spatial extent (m)	Algal genetic diversity		Fungal genetic diversity	
					Mean (S.D.)	Range	Mean (S.D.)	Range
NZ <i>Ramalina</i>	21	Few morphotypes (3)	9	581,576	0.08	0.0 – 0.16	0.06	0.0 – 0.11
NZ <i>Usnea</i>	111	Many morphotypes (17)	43	1,251,276	0.09	0.0 – 0.18	0.05	0.0 – 0.10
NZ <i>Usnea</i> replicated	83	Several morphotypes (9)	18	882,910	0.10	0.0 – 0.17	0.05	0.0 – 0.09
Craigieburn <i>Usnea</i>	36	Several morphotypes (6)	1	1,775	0.04	0.0 – 0.15	0.03	0.0 – 0.06
Flock Hill <i>Usnea</i>	66	Few morphotypes (3)	1	1,095	0.03	0.0 – 0.14	0.03	0.0 – 0.08
Flock Hill community mapped	35	Many lichen genera	1	796	0.06	0.0 – 0.22	0.13	0.0 – 0.33
Flock Hill community	67	Many lichen genera	1	1,141	0.08	0.0 – 0.49	0.15	0.0 – 0.38

Table 2_(on next page)

Results table

Table 2: Results from db-RDA with variance partitioning and Procrustes approach to co-phylogeny giving the P -alue from a test randomising the association matrix for fungi and algae for the six independent datasets and thefull Flock Hill community dataset. Note that only 35 of the 67 Flock Hill multiple species dataset specimens were mapped and were therefore analysed separately from the full dataset. Variance partitioning divides the total variance up into portions explained by the partner genetic distance matrix (Partner), the purely spatial portion (Space), the spatially structured variation in the partner matrix (Shared) and unexplained variation (Unexpl.). The significant MEMs are given in order of their significance. For each dataset, there are $n-1$ MEMs in the total set and smaller MEM numbers represent larger-scale spatial pattern.

Table 2: Results from db-RDA with variance partitioning and Procrustes approach to co-phylogeny giving the P -value from a test randomising the association matrix for fungi and algae for the six independent datasets and the full Flock Hill community dataset. Note that only 35 of the 67 Flock Hill multiple species dataset specimens were mapped and were therefore analysed separately from the full dataset. Variance partitioning divides the total variance up into portions explained by the partner genetic distance matrix (Partner), the purely spatial portion (Space), the spatially structured variation in the partner matrix (Shared) and unexplained variation (Unexpl.). The significant MEMs are given in order of their significance. For each dataset, there are $n-1$ MEMs in the total set and smaller MEM numbers represent larger-scale spatial pattern.

Dataset	Significant MEMs	Partner	Shared	Space	Unexpl.	P
Dependent matrix: Algae						
NZ <i>Ramalina</i>	1, 5, 3, 2, 7	0.21	0.65	0.09	0.06	0.016
NZ <i>Usnea</i>	1, 4, 6, 12, 17, 7, 3, 9, 2, 5, 45, 8, 63, 38, 41, 28, 54	0.39	0.41	0.08	0.12	< 0.001
NZ <i>Usnea</i> replicated	4, 5, 1, 3	0.89	0.07	0.00	0.04	< 0.001
Craigieburn <i>Usnea</i>	2, 9	0.87	0.00	0.01	0.12	0.021
Flock Hill <i>Usnea</i>	16	0.62	0.01	0.01	0.35	0.030
Flock Hill community mapped	none	0.84	0.03	0.02	0.12	< 0.001
Flock Hill community	-	0.71	-	-	0.29	< 0.001
Dependent matrix: Fungi						

NZ <i>Ramalina</i>	1, 3, 5, 7, 9, 10, 2,	0.32	0.62	0.02	0.03	0.025
NZ <i>Usnea</i>	1, 4, 12, 3, 8, 6, 9, 45, 17, 54, 13, 38, 41, 16, 5, 56, 7	0.45	0.23	0.12	0.2	< 0.001
NZ <i>Usnea</i> replicated	4, 3, 31, 1, 17, 5, 9	0.72	0.01	0.02	0.25	< 0.001
Craigieburn <i>Usnea</i>	7, 12, 13, 15, 16, 3	0.55	0.02	0.3	0.14	0.024
Flock Hill <i>Usnea</i>	1	0.62	0.00	0.08	0.30	0.022
Flock Hill community mapped	none	0.73	0.02	0.03	0.22	< 0.001
Flock Hill community	-	0.6	-	-	0.40	< 0.001