

1 **Bite marks on the frill of a juvenile *Centrosaurus* from the Late Cretaceous Dinosaur**
2 **Provincial Park Formation**

3
4 **David W. E. Hone¹**

5 **Darren H. Tanke²**

6 **Caleb M. Brown²**

7
8 1. School of Biological and Chemical Sciences, Queen Mary, University of London,
9 London, UK

10 2. Royal Tyrrell Museum, Drumheller, Alberta, Canada

11
12 **Corresponding author:** David Hone, d.hone@qmul.ac.uk

13

14 **Abstract:**

15 Bite marks on bones can provide critical information about interactions between carnivores
16 and animals they consumed (or attempted to) in the fossil record. Data from such interactions
17 is somewhat sparse but is hampered by a lack of records in the scientific literature. Here we
18 present a rare instance of feeding traces on the frill of a juvenile ceratopsian dinosaur from
19 the late Campanian Dinosaur Park Formation of Alberta. It is difficult to determine the likely
20 tracemaker(s) but the strongest candidate is a small-bodied theropod such as a dromaeosaur or
21 juvenile tyrannosaur. This marks the first documented case of carnivore consumption of a
22 juvenile ceratopsid, but may represent scavenging as opposed to predation.

23

24 **Keywords:** ceratopsian, carnivore-consumed, ecology, tooth-marked

25

26 **Introduction:**

27 Bite marks on the bones of fossils can provide important information as to the
28 palaeoecology of ancient ecosystems and as indicators of trophic interactions between
29 animals. In the case of the non-avian dinosaurs (hereafter simply ‘dinosaurs’), bite marks
30 (that are healing, healed and peri- or post-mortem) can allow inferences about both inter- and
31 intraspecific interactions in various clades. This includes inferences about cannibalism (Bell
32 & Currie, 2010; Hone & Tanke, 2015), scavenging (Hone & Watabe, 2010), intraspecific
33 combat (Tanke & Currie, 1998), interspecific combat (Happ, 2008), prey preferences
34 (Jacobsen, 1998), and attempted predation (De Palma et al., 2013). However, there are major
35 problems with the use of bite mark data which has limited its potential.

36 Although tooth-marks are not uncommon for dinosaurs, they are considerably more
37 common in tyrannosaur-dominated faunas (Fiorillo, 1991) and can be regularly seen in some
38 formations such as Dinosaur Provincial Park (authors pers. obs.). Even so, relatively few

39 marks have been described in details to date, which limits comparisons or large-scale
40 assessments of patterns across multiple traces (though see e.g. Jacobsen, 1998).

41 Identification of both parties associated with bite marks (i.e. both the carnivore and
42 the consumed sensu Hone & Tanke, 2015) is often difficult, limiting the available
43 information. Bitten specimens are often fragmentary, and as bite marks are commonly found
44 on isolated elements, these are often not diagnostic to genera or species. Similarly, bites
45 marks are often difficult to attribute to tracemakers (e.g. see Hone & Chure, in press),
46 although specimens that include shed teeth of a feeding carnivore (e.g. Currie & Jacobsen,
47 1995; Maxwell & Ostrom, 1995; Hone et al., 2010), or where there are single credible
48 candidates for the tracemaker (e.g. Bell & Currie, 2010) support identifications.

49 Finally, there are often difficulties in interpreting bite mark data and the actions of
50 the tracemakers (Chure, Fiorillo, & Jacobsen, 2000; Robinson, Jasinski & Sullivan, 2015). It
51 is difficult to separate out scavenging events from those associated with late stage carcass
52 consumption of a prey item without supporting taphonomic data (e.g. see Hone & Watabe,
53 2010). Bites may have been made by multiple tracemakers, or at different times, and traces
54 can potentially be altered through erosion or transport which restricts interpretations.

55 Collectively then, this makes interpretations difficult, although it also means that
56 every recorded bite event may be valuable as it is only through the collection and assessment
57 of large datasets that confident interpretations may be made. In this context, unusual or rare
58 marks may be especially important for determining the range of possible interactions and
59 events based on theropod bites.

60 Here we describe a number of small marks on a partial frill of a juvenile ceratopsian
61 (referred to *Centrosaurus apertus*). Bite marks on ceratopsians are known (e.g., Erickson et
62 al., 1996; Jacobsen, 1998; Happ, 2008, Fowler et al., 2006) but are restricted to larger bodied
63 animals making this the first description of bites on such a young animal. Determining the

64 tracemaker is not possible given the range of possible candidates but this may represent an
65 example of a small-bodied carnivore (i.e., Dromaeosauridae, Troodontidae or juvenile
66 Tyrannosauridae) feeding on the young of a much larger-bodied taxon.

67

68 **Materials and Methods:**

69 The present specimen (TMP 2014.012.0036) represents a fragment of the squamosal
70 of a subadult centrosaurine ceratopsid, from the lower Dinosaur Park Formation (Campanian)
71 of Southern Alberta. It was found by DHT and collected under ~~and~~-a Park Research and
72 Collection Permit (No. 14-095) from Alberta Tourism, Parks and Recreation, as well as a
73 Permit to Excavate Palaeontological Resources (No. 14-018) from Alberta Culture and
74 Tourism and the Royal Tyrrell Museum of Palaeontology, both issued to CMB, and is
75 accessioned at the Royal Tyrrell Museum of Palaeontology, Drumheller.

76 The fossil was collected from the surface of a multi-taxic bonebed in the core area of
77 Dinosaur Provincial Park (UTM, 12U: 464,462 E; 5,621,335 N, WGS 84). Stratigraphically,
78 the specimen is from the lower Dinosaur Park Formation (~5 m above the contact with the
79 underlying Oldman Formation), and falls between the radiometrically dateable Jackson
80 Coulee (min. 76.32 Ma) and Plateau (75.60 +/- 0.02 Ma) bentonites (Dave Eberth, pers.
81 comm., 2017). This confidently places the specimen within the *Corythosaurus-Centrosaurus*
82 zone (Ryan et al., 2012; Mallon et al., 2013), and as result, is here referred to *Centrosaurus*
83 *apertus* as this is ~~this is~~ the only centrosaurine ceratopsid species known to occur in this well
84 sampled (>20 diagnostic skulls, and ~20 bonebeds) interval (Eberth and Getty, 2005; Brown,
85 2013).

86

87 **Description:**

Specimen TMP 2014.012.0036 is identified as a fragment of squamosal of a small centrosaurine ceratopsid dinosaur (Fig 1). This is subtriangular in shape and approximately 8 cm per side and just over 1 cm thick. It represents the posterior corner of the lateral margin of the squamosal and is from a position just ventral to the suture with the parietal (Fig 2). It was broken in several places prior to fossilisation, but part of the original lateral margin remains intact and shows the scalloped edge of the frill.

Four independent lines of evidence suggest this element derived from juvenile/subadult animal. Firstly, despite limited wear to the element, the majority of the surface is unweathered and shows the distinctly striated long grained bone texture of juvenile centrosaurine frill elements (Sampson, Ryan & Tanke, 1997; Brown, Russell & Ryan, 2009; Tumarkin-Deratzian, 2010). Secondly, the preserved lateral margin of the element is straight, and bears no evidence of the imbrication of the loci undulations that develop ontogenetically (Sampson, Ryan & Tanke, 1997). Thirdly, the partially preserved epioossification loci, is without fused epioossification seen in many (but not ubiquitously preserved) adults (Sampson, Ryan & Tanke, 1997; Horner and Goodwin 2008). Finally, the cross-sectional ~~the~~ thickness of the element (<10 mm) and the overall small size of the one preserved episquamosal loci (see Fig 1) indicate a small absolute size of the entire squamosal. Taken together, this suggest the animal was below osteologically adult maturity (cf Hone, Farke, & Wedel, 2016), and falls into the juvenile age class established by Sampson, Ryan & Tanke (1997).

The absolute size of the animal in life is difficult to estimate from the limited remains, but comparison with a sample of 24 more complete juvenile/subadult squamosals derived from monodominant centrosaurine bonebeds (*Centrosaurs apertus*, *Coronosaurus brinkmani*, *Pachyrhinosaurus lakustai*), suggest the complete squamosal would have had a marginal length of approximately 204 mm, and a maximum length of approximately 293 mm. For comparison, osteologically mature *C. apertus* specimens have squamosals ranging in

113 marginal length of 258-373 mm (mean = 322 mm), total length of 288-481 mm (mean = 401
114 mm), for skulls ranging in basal skull length of 660-868 mm (mean = 779 mm). The suggests
115 the tooth-marked squamosal represents an individual with linear skull measures around two-
116 thirds to three-quarters (64-73%) the size of the average ontogenetically adult *Centrosaurus*
117 *apertus* skull, and approximately one-half (48-61%) the size of the largest *Centrosaurus*
118 *apertus* skull. Although this may not sound small in comparison, due to the cubic scaling of
119 mass relative linear measures, this equates to an animal less than one-third (~29%), and less
120 than one-seventh (~13%), the mass of the average and largest adult, respectively. This also
121 likely represents an underestimate due to potential negative allometry of the skull relative to
122 the body.

123 The specimen as preserved has a light coloured and dark coloured side, presumably
124 the former being somewhat bleached by exposure to the sun and rain prior to discovery. The
125 texture on the surface (fine striations) is similar on both sides, suggesting this is a genuine
126 feature and not the result of erosion or exposure. It is not possible to confidently determine
127 which surface is internal and which is external, so as a result the lighter coloured side is
128 referred to as 'Side A', with the darker side as 'Side B'. A number of features and marks are
129 seen on the specimen that are described below and are numbered as in Figure 3. Part of the
130 lateral margin of the element is broken (which is common in isolated parts of certatopsian
131 frills), but one aspect of this retains a natural edge.

132

133 Side A (Figure 3A):

134

135 1. A groove on the surface of the bone, which has a counterpart (i) on side B.

136 2. A thin score that cuts through the cortex. This is filled in with fossil sediment and so it is
137 hard to see margins well and it means the depth of this cannot be measured. It is long and
138 especially narrow being 18 mm by 1mm at the widest, and mostly c. 0.5 mm wide.
139 3. A small oval mark (6.5 by 3 mm) near the margin of the bone. This is uneven and slightly
140 'Z' shaped.
141 4-6. A series of marks that resemble cracks. As with trace 2 there is some infill of the marks
142 so the margins are not entirely clear. Number 5 is rather irregular and 4 in particular matches
143 other very small cracks in general form.
144 7. A slight mark on the edge of the bone, near the broken margin. It is small and oval in shape
145 and parallel to the frill margin. The mark is 5 mm long by 1.5 mm wide.
146 8. A small but deep mark on the margin that is associated with some damage to the frill
147 margin. The mark is 5 mm long, 1.7 mm deep, and as it is at the broken margin, the width
148 cannot be determined.

149

150 Side B (Figure 3B):

151 i. A long groove that has some slight damage to one edge of it. This ~~is~~ runs parallel to mark 1
152 on side A.
153 ii. Two shallow scores, one is broad and the second very thin that departs the former at a
154 shallow angle. The thin side branch does not cut across the fibers of the bone cleanly. The
155 larger trace is 18 mm long and up to 1.25 mm wide.
156 iii. A short and proportionally deep penetration of the bone, which appears to be broken at the
157 margins. The mark is 11.5 mm long, up to 4 mm wide, and ~~is~~ 3 mm deep (it is deeper
158 proximally and becomes more shallow towards the margin). There is a little wear internally
159 as it is smooth in places including the margins.

iv. A comparatively broad mark that is up to 11.75 mm long, 2.25 mm wide, and ~~is~~ approximately 1 mm deep. The trace is slightly curved along its length.

v. This is a small and narrow score mark that is 17 mm long and 1 mm wide, and closely associated with mark iv. The depth cannot be measured accurately, but is estimated to be under 0.5 mm. This is subparallel to ii and iii.

vi. A triangular mark that lies at the margin of the piece. The mark is 7 mm long, as preserved, and 1.8 mm deep. This lies close to mark iii.

167

168 Discussion

The specimen here shows a mixture of mark types which are considered to be the result of a combination of effects. The element was found as an isolated piece and not from ~~of~~ one of the ceratopsian bonebeds that are common in Dinosaur Provincial Park. Given the isolated nature of the fragment (removed from the rest of the skeleton, and the abraded nature of the breaks), it is likely to have undergone some transport and erosion (and perhaps chemical alteration) given that it was not associated with any other parts of a young *Centrosaurus*. This also means that its exact taphonomic history is unknown and thus caution is required when interpreting the limited data.

Breaks to ceratopsian frills are common and thus there is little to take from the separation of the element from the rest of the skull, or the broken margin. Although these are major breaks to this small bone, there is some wear at the edges (suggesting transport and perhaps chemical wear) and the breaks are not clearly associated with possible bites. On side A in particular there are a series of cracks (4-6) on the surface that align with the natural striations on the bone (see Fig 3A) and the larger manifestations of the long-grained bone texture associated with immature frills. Although they are subparallel to each other which is a very common feature of theropod bite marks e.g. Currie & Jacobsen, 1995; Chure, Fiorillo, &

185 Jacobsen, 2000; Hone & Watabe, 2010), they also align very well with the general orientation
186 of fibers and smaller cracks on the (dorsal) surface. Mark 7 is an odd shape that does not
187 resemble a bite mark and as it is close to the break of the frill margin, it is suggested that this
188 may be part of an impact that lead to this damage, possibly through trampling or transport.
189 Although different in form, the marks at point ii are likely also cracks resulting from the same
190 stress as these also primarily align with the natural form of the bone and the cracks seen on
191 the dorsal surface.

192 Traces 1 and i are considered the remains of vascular grooves. They are both broad
193 and shallow and very smooth making them quite unlike typical bite marks.

194 Mark 3 is less clearly defined than other traces on the bone and the shallow and
195 rounded nature of this make it likely to be part of another vascular groove as with marks 1
196 and i. Similarly, marks iv and v are subparallel which is a common feature of bite marks
197 however they are also rather irregular in shape and do not track each other closely as would
198 be expected for adjacent teeth in a jaw. These traces are also smooth and worn, and broad and
199 shallow which is unlike most bite marks, though their identity is unclear – they may be more
200 vascular pathways, or eroded damage, or perhaps both.

201 Traces ii, iv and v are difficult to interpret and may be considered bite marks but this
202 is uncertain. Mark ii is slightly tear-drop shaped and does not follow the grain of the bone as
203 with the above traces so is not part of a crack associated with long grain bone texture. It is
204 however relatively shallow and smooth unlike typical bite marks, although and perhaps
205 altered through erosion. This may therefore be the result of a small impact during transport.
206 Marks iv and v lie sub-parallel to one another as is common for dinosaurian bite marks, but
207 trace iv has a somewhat sinusoidal pattern and does not closely track v as would be expected
208 if they were delivered by consecutive teeth in the jaws. As with trace ii they are somewhat
209 broad and smooth which is not normal for bite marks.

210 Marks 8 and vi are relatively deep into the cortex and come at the margins of the
211 piece and thus could potentially represent bites that penetrate the cortex and thus may have in
212 part led to the breaking off of the piece. These traces are therefore tentatively assigned as bite
213 marks, but may well be the result of damage from transport and erosion.

214 This leaves two traces on the specimen that are confidently interpreted as bite marks,
215 trace 2 on the side A and iii on side B. Mark 2 is a narrow trace which does correspond in
216 general form to other bite marks seen on bones from the Dinosaur Park Formation (though
217 these are typically considerably larger – DWEH pers obs). This is a long and thin ‘diamond’
218 shape tapering to points at each end, although there is also some damage to the margins of
219 this where the bone splintered as the mark was inflicted or perhaps through later erosion. It
220 corresponds to a drag mark (sensu Hone & Watabe, 2010) where the tooth does not break
221 through the cortex of the bone.

222 Mark iii is ~~isn~~ close in morphology to a bite and drag (sensu Hone & Watabe, 2010)
223 where the tooth penetrates deep into the bone and then is pulled back. This corresponded with
224 the orientation of the bite which is from medial to distal on the frill being deeper more
225 proximally, and is more shallow towards the frill margin. The trace does have an unusual
226 morphology which is slightly expanded laterally underneath the bite (not sure this is
227 explained well here – hard to describe). This may due to an effect such as resting in shallow
228 water where particles might swirl inside the cavity causing erosion to expand this without
229 damaging more of the surface bone.

230

231 *Tracemaker identity:*

232 The marks here do not correspond well to those of non-dinosaurian carnivores known
233 from Dinosaur Provincial Park and thus can be ruled out. There are lizards, crocodiles,
234 champsosaurs, and mammals known which could potentially have bitten on dinosaur bone.

带格式的: 字体颜色: 红色

235 However, extant crocodiles tend to splinter bones when biting and also leave sub-circular
236 punctures not seen here (e.g. see Naju & Blumenschine, 2006; Drumheller and Brochu, 2014;
237 Botfalvai, Prondvai & Ősi, 2014) and large lizards tend to leave curved traces because the
238 head sweeps in an arc during feeding (D'Amore & Blumenschine, 2009). There are not
239 currently bite marks assigned to champsosaurs, but they might be expected to feed in similar
240 ways to either or even both of these techniques (based on their gross anatomy and
241 phylogenetic ancestry) which would not match the traces seen here, and they are piscivorous.
242 The marks also do not correspond with inferred traces from mammals known from the
243 underlying Oldman Formation of Alberta (Longrich and Ryan, 2010).

244 With these ruled out, the most likely candidates are therefore the non-avian theropods.
245 Three clades of toothed, carnivorous, forms are known from these beds: tyrannosaurs,
246 dromaeosaurs, troodontids as well as the genus *Richardoestesia* which is of uncertain
247 affinities (Currie, 2005). Although at adult size, the tyrannosaurs are very large, bite marks
248 from smaller individuals remain a possibility.

249 Mark 2 is a good match for the very thin and blade-like teeth of dromaeosaurs and
250 troodontids which would leave proportionally thin traces. Indeed, these marks are a good
251 match in general form for bite marks left by dromaeosaurs in the formation which can be
252 positively identified because of a shed tooth (Currie & Jacobsen, 1995). Long and straight
253 bites from tyrannosaurs are typically left as a result of scrape feeding where the premaxillary
254 teeth are drawn across the cortex (Hone & Watabe, 2010) and usually leave multiple
255 subparallel traces that are broad because of the D-shaped nature of the teeth and these are
256 therefore rather unlike mark 2.

257 The morphology of trace iii however, is very different from that of 2, being much
258 more broad and deep. As noted above, this shape may have been exaggerated by later
259 erosion, but this would still be different to the relatively thin and well-defined mark 2.

260 Although slightly elongate, this is closest to a puncture mark (sensu Hone & Watabe, 2010)
261 and would be a good match for a tyrannosaur tooth (premaxillary or maxillary / dentary).
262 Similarly, the traces 3, 8, and vi, if they are bites, would more closely match tyrannosaurs
263 given their general broad and deep nature. At least some deep puncture wounds that may be
264 attributed to larger dromaeosaurs are known (Gignac et al., 2010) and such traces do seem to
265 be relatively rare. Even when a dromaeosaur tooth was punctured into a pterosaur bone with
266 enough force to remove the tooth this was not driven deep into the bone and there were no
267 other associated punctures (Currie and Jacobsen, 1995).

268 The mixture of trace morphology, coupled with the likely erosion of at least some
269 marks makes the identity of the tracemaker difficult to determine. It may have been a
270 dromaeosaurid or young tyrannosaur, or possibly both. Although we are not aware of any bite
271 marks on dinosaur fossils that can be attributed to multiple species this is something which
272 might be predicted – modern carcasses may be fed on by multiple species through
273 kleptoparasitism (Höner et al., 2002) or simply feeding on carrion after the original predator
274 has moved on (Lanszki et al., 2015).

275

276 *Interpretation:*

277 In all cases (2, 3, 8, iii, vi) the traces are well separated from one another and not a
278 series of punctures or sub-parallel marks that are typical of theropod bite traces. Marks may
279 be inconsistent in this regard thanks to the different lengths of theropod teeth in the jaws and
280 possible absences etc. such that a bite may only result in one or two teeth engaging with the
281 bone. In the case of traces 8 and vi which abut the broken margins, these may represent a bite
282 on the now missing part of the frill where only a single tooth contacted the squamosal. Single
283 traces made by theropod teeth are certainly known in a number of cases (e.g. some traces in
284 Erickson & Olson, 1996; Tanke & Currie, 1998; Gignac et al., 2010; Hone & Tanke, 2015;)

285 and so despite the unusual arrangement of these traces, we are confident that several of these
286 do represent bite marks.

287 Superposition of the two sides of the squamosal piece (Fig 4) shows that marks 3, iii,
288 and vi are close to one another and 3 and iii even partially overlap. However, iii lies at a very
289 different angle to the other marks and this is hard to reconcile as being associated with them.
290 In contrast, traces 3 and vi are in a similar location and have a similar orientation suggesting
291 they may be the result of a single bite engaging both sides of the frill.

292 No major muscle groups or soft tissues such as fat deposits are likely associated with
293 the squamosal of ceratopsian dinosaurs. As such, feeding on this part of the skull was likely a
294 result of late stage carcass consumption (see Hone & Rauhut, 2010 and references therein)
295 whereby feeding only occurred as a result of the more nutritious aspects of the carcass having
296 been exploited (Fig 5). The small size of the animal may imply that the carcass was exploited
297 quickly – indeed, large theropods like tyrannosaurs were apparently capable of processing
298 and consuming most or all of a juvenile dinosaur (Chin et al., 1998). As a result, although
299 juvenile dinosaurs were likely common components of dinosaurian faunas, they were at least
300 in part rare in the fossil record as a result of destruction by theropod feeding (Hone & Rauhut,
301 2010). As a result, despite the apparent preferences for feeding on juvenile dinosaurs, most
302 described bite marks are on the bones of adults which may have resisted being consumed and
303 destroyed (even by large tyrannosaurs) and thus feeding traces on a juvenile dinosaur remain
304 unusual. Perhaps the size and shape of ceratopsian crania, even in juveniles, made them
305 difficult to process or required an excess of handling effort for a relatively low reward.

306

307 **Conclusions:**

308 Bite marks remain an important source of information on trophic interactions between carnivores
309 and consumed species. Such traces attributes to tyrannosaurs are more common than for other
310 theropod dinosaurs but even so few have been described in detail despite the information that may
311 be available to help interpret their ecology and behaviour. This first evidence of likely scavenging on

带格式的: 无间隔, 缩进: 首行缩进: 0 厘米, 行距: 单倍行距

312 a non-adult animal adds to the known diversity of animals apparently fed on by Late Cretaceous
313 tyrannosaurs.
314

315 **Acknowledgements:**

316 We thank Marie-Hélène Trudel-Aubry for her artwork as used in figure 5. We thank Brandon
317 Strilisky for his help as collections manager and David Eberth for preliminary updated
318 radiometric dates for the specimen.
319

320 **References:**

- 321 Bell PR, Currie PJ. 2010. A tyrannosaur jaw bitten by a confamilial: scavenging or fatal
322 agonism? *Lethaia* 43:278–281.
- 323 Botfalvai G, Prondvai E, Ősi A. 2014. Inferred bite marks on a Late Cretaceous (Santonian)
324 bothremydid turtle and a hylaeochampsid crocodilian from Hungary. *Cretaceous*
325 *Research*, 50:304-317.
- 326 Brown CM, Russell AP, Ryan MJ. 2009. Pattern and transition of surficial bone texture of the
327 centrosaurine frill and their ontogenetic and taxonomic implications. *Journal of*
328 *Vertebrate Paleontology* 29:132-141.
- 329 Brown CM. 2013. Advances in quantitative methods in dinosaur palaeobiology: a case study
330 in horned dinosaur evolution. PhD thesis, University of Toronto.
- 331 Chin K, Tokaryk TT, Erickson GM, Calk LC. 1998. A king-sized theropod coprolite. *Nature*
332 393:680-682.
- 333 Chure DJ, Fiorillo AR, Jacobsen R. 2000. Prey bone utilization by predatory dinosaurs in the
334 Late Jurassic of North America, with comments on prey bone use by dinosaurs
335 throughout the Mesozoic. *Gaia* 15:227–232.
- 336 Currie PJ. 2005. Theropods, including birds. In: Currie PJ, Koppelhus EB, eds. *Dinosaur*
337 *Provincial Park*. Bloomington: Indiana University Press, 367-397.

338 Currie PJ, Jacobsen AR. 1995. An azhdarchid pterosaur eaten by a velociraptorine theropod.
339 *Canadian Journal of Earth Sciences* 32:922–925.

340 D'Amore DC, Blumenshine RJ. 2009. Komodo monitor (*Varanus komodoensis*) feeding
341 behavior and dental function reflected through tooth marks on bone surfaces, and the
342 application to ziphodont paleobiology. *Paleobiology* 35:525-552.

343 DePalma RA, Burnham DA, Martin LD, Rothschild BM, Larson PL. 2013. Physical evidence
344 of predatory behavior in *Tyrannosaurus rex*. *Proceedings of the National Academy of*
345 *Sciences* 110:12560-12564.

346 Drumheller SK, Brochu CA. 2014. A diagnosis of Alligator mississippiensis bite marks with
347 comparisons to existing crocodylian datasets. *Ichnos* 21:131-146.

348 Eberth DA, Getty MA. 2005. Ceratopsian bonebeds: occurrence, origins, and significance. In:
349 Currie PJ, Koppelhus EB eds. *Dinosaur Provincial Park: a spectacular ancient*
350 *ecosystem revealed*. Bloomington: Indiana University Press, 501-536.

351 Erickson GM, van Kirk SD, Su J, Levenston ME, Caler WE, Carter DR. 1996. Bite–force
352 estimation for *Tyrannosaurus rex* from bone–marks. *Nature* 382:706–708.

353 Erickson GM, Olson KH. 1996. Bite marks attributable to *Tyrannosaurus rex*: preliminary
354 description and implications. *Journal of Vertebrate Paleontology* 16:175-178.

355 Fiorillo AR. 1991. Prey bone utilisation by predatory dinosaurs. *Palaeogeography,*
356 *Palaeoclimatology, Palaeoecology* 88:157–166.

357 Fowler DW, Sullivan RM. 2006. A ceratopsid pelvis with toothmarks from the Upper
358 Cretaceous Kirtland Formation, New Mexico: evidence of Late Campanian
359 tyrannosaurid feeding behaviour. *New Mexico Museum of Natural History and Science*
360 *Bulletin* 35:127–130.

361 Gignac PM, Makovicky PJ, Erickson GM, Walsh RP, 2010. A description of *Deinonychus*
 362 *antirrhopus* bite marks and estimates of bite force using tooth indentation simulations.
 363 *Journal of Vertebrate Paleontology* 30:1169-1177.

364 Gilmore CW. 1914. A new ceratopsian dinosaur from the Upper Cretaceous of Montana, with
 365 note on *Hypacrosaurus*. Smithsonian Miscellaneous Collections 63:1-10.

366 Happ J. 2008. An analysis of predator-prey behavior in a head-to-head encounter between
 367 *Tyrannosaurus rex* and *Triceratops*. In: Larson P, Carpenter K. eds. *Tyrannosaurus rex*
 368 *the Tyrant King*. Bloomington: Indiana University Press, 355-370

369 Hone DWE, Chure DJ. In press. Difficulties in assigning trace makers from theropodan bite
 370 marks: an example from a young diplodocoid sauropod. *Lethaia*.

371 Hone DWE, Rauhut OWM. 2010. Feeding behaviour and bone utilisation by theropod
 372 dinosaurs. *Lethaia* 43:232-244.

373 Hone DWE, Tanke DH. 2015. Pre-and postmortem tyrannosaurid bite marks on the remains
 374 of *Daspletosaurus* (Tyrannosaurinae: Theropoda) from Dinosaur Provincial Park,
 375 Alberta, Canada. *PeerJ* 3:p.e885.

376 Hone DWE, Watabe M. 2010. New information on the feeding behaviour of tyrannosaurs.
 377 *Acta Palaeontologica Polonica* 55:627-634.

378 Hone DWE, Choiniere J, Sullivan C, Xu X, Pittman M, Tan Q. 2010. New evidence for a
 379 tropic relationship between the dinosaurs *Velociraptor* and *Protoceratops*.
 380 *Palaeogeography, Palaeoclimatology, Palaeoecology* 291:488-492.

381 Hone DWE, Farke AA, Wedel MJ. 2016. Ontogeny and the fossil record: what if anything is
 382 an adult dinosaur? *Biology Letters* 12:20150947.

383 Höner OP, Wachter B, East ML, Hofer H. 2002. The response of spotted hyaenas to long -
 384 term changes in prey populations: functional response and interspecific kleptoparasitism.
 385 *Journal of Animal Ecology* 71:236-246.

386 Horner JR, Goodwin MB. 2008. Ontogeny of cranial epi-ossifications in *Triceratops*. *Journal*
 387 *of Vertebrate Paleontology* 28:134-144.

388 Jacobsen AR. 1998. Feeding behavior of carnivorous dinosaurs as determined by tooth marks
 389 on dinosaur bones. *Historical Biology* 13:17–26.

390 Lanszki J, Kurys A, Heltai M, Csányi S, Ács K. 2015. Diet composition of the golden jackal
 391 in an area of intensive big game management. *Annales Zoologici Fennici* 52:243-255.

392 Longrich NR, Ryan MJ. 2010. Mammalian tooth marks on the bones of dinosaurs and other
 393 Late Cretaceous vertebrates. *Palaeontology* 53:703-709.

394 Mallon JC, Evans DC, Ryan MJ, Anderson JS. 2013. Megaherbivorous dinosaur turnover in
 395 the Dinosaur Park Formation (upper Campanian) of Alberta, Canada. *Palaeogeography,*
 396 *Palaeoclimatology, Palaeoecology* 350:124-138.

397 Maxwell WD, Ostrom JH. 1995. Taphonomy and paleobiological implications of
 398 *Tenontosaurus-Deinonychus* associations. *Journal of Vertebrate Paleontology* 15:707-
 399 712.

400 Naju JK, Blumenschine RJ. 2006: A diagnosis of crocodile feeding traces on larger mammal
 401 bone, with fossil examples from the Plio-Pleistocene Olduvai Basin, Tanzania. *Journal*
 402 *of Human Evolution* 50:142–162.

403 Robinson RF, Jasinski SE, Sullivan RM, 2015 Theropod bite marks on dinosaur bones:
 404 indications of a scavenger, predator or both?; and their taphonomic implications. New
 405 *Mexico Museum of Natural History and Science Bulletin* 68:275-282.

406 Ryan MJ, Evans DC, Currie PJ, Brown CM, Brinkman D. 2012. New leptoceratopsids from
 407 the Upper Cretaceous of Alberta, Canada. *Cretaceous Research* 35:69-80.

408 Sampson SD, Ryan MJ, Tanke DH. 1997. Craniofacial ontogeny in centrosaurine dinosaurs
 409 (Ornithischia: Ceratopsidae): taxonomic and behavioral implications. *Zoological Journal*
 410 *of the Linnean Society* 121:293-337.

411 Tanke DH, Currie P. 1998. Head-biting behavior in theropod dinosaurs: paleopathological
412 evidence. *Gaia* 15:167–184.

413 Tumarkin-Deratzian AR. 2010. Histological evaluation of ontogenetic bone surface texture
414 changes in the frill of *Centrosaurus apertus*. In: Ryan MJ, Chinnery-Allgeier BJ,
415 Eberth DA, eds. *New Perspectives on Horned Dinosaurs, the Royal Tyrrell Museum*
416 *Ceratopsian Symposium*. Bloomington: Indiana University Press, 251-263.

417

418

419

420

421 Fig 1. Photographs of TMP 2014.012.0036 showing side A and side B. Scale bar is 50 mm
422 long.

423

424 Fig 2. Reconstructed skull of a juvenile *Centrosaurus apertus* of approximately similar
425 ontogenetic status to that of TMP 2014.012.0036 (A) in right lateral view, next to that of
426 an adult (B). The two skulls are to scale with one another. The squamosal is highlighted
427 in medium grey and the approximate outline of the specimen preserved here is in dark
428 grey. Reconstruction of the juvenile skull based largely on USNM 7951 (Gilmore, 1914),
429 with additions from TMP 1982.016.0011 and 1996.175.0064, adult based on YPM 2015.

430

431 Fig 3. Interpretative drawing of TMP 2014.012.0036 showing side A and side B. Numbers
432 relate to various areas of interest as described in the text. Pale grey areas mark areas of
433 wear to the bone, dark grey areas represent major features, and black areas are those that
434 penetrate deep into the cortex. The thicker lines on the margins represent the natural
435 margin of the element (see also figure 2). Scale bar is 50 mm long.

436

437 Fig 4. Interpretative drawing of TMP2014.012.0036 flipped such that the bite marks from the
438 dorsal and ventral sides both appear. Dark grey areas represent major features, and black
439 areas are those that penetrate deep into the cortex. The thicker line on the margins
440 represent the natural margin of the element (see also figure 2). Scale bar is 50 mm long.

441

442 Fig 5. Although the identity of the tracemaker of the marks on the *Centrosaurus* frill
443 fragment is uncertain, here we present a speculative reconstruction of scavenging by a
444 juvenile *Gorgosaurus*. Artwork by Marie-Hélène Trudel-Aubry and used with their
445 permission.

446

447