

Table 1. Comparison of morphological variables between males (n = 113) and females (n = 15) of *Hyla eximia*. SVL = snout-vent length, HL = head length, HW = head width, AL = arm length, FL = forearm length, FML = femur length, TL = tibia length, BM = body mass; SSDi = Sexual Size Dimorphism index. Means for each variable reported $\pm$ standard error; values in parentheses show variable ranges.

Variables	Males	Females	SSDi	Mann-Whitney <i>U</i>	<i>P</i>
SVL	29.9 $\pm$ 0.39 (26.9 - 32.7)	33.5 $\pm$ 0.49 (29.0 - 37.5)	1.09	-4.656	0.003
HL	9.93 $\pm$ 0.67 (8.8 - 11.6)	10.4 $\pm$ 0.95 (8.8 - 12.8)	1.04	-2.022	0.043
HW	9.99 $\pm$ 0.68 (8.9 - 12.0)	10.9 $\pm$ 0.65 (9.9 - 12.1)	1.09	-3.937	0.008
AL	6.63 $\pm$ 0.68 (4.9 - 7.9)	7.14 $\pm$ 0.41 (6.4 - 8.0)	1.07	-3.033	0.002
FL	6.60 $\pm$ 0.65 (5.1 - 7.9)	7.02 $\pm$ 0.31 (6.5 - 7.7)	1.06	-2.374	0.017
FML	15.34 $\pm$ 0.76 (14.0 - 17.4)	16.20 $\pm$ 1.16 (14.0 - 18.0)	1.05	-2.866	0.004
TL	15.59 $\pm$ 0.94 (13.5 - 17.9)	16.95 $\pm$ 1.13 (15.5 - 19.0)	1.08	-3.852	0.001
BM	2.5 $\pm$ 0.39 (1.4 - 3.5)	3.4 $\pm$ 0.74 (2.0 - 4.4)		-4.676	0.001