

A taxonomic reassessment of *Piramys auffenbergi*, a neglected turtle from the late Miocene of Piram Island, Gujarat, India (#30116)

1

First submission

Editor guidance

Please submit by **17 Sep 2018** for the benefit of the authors (and your \$200 publishing discount).



Structure and Criteria

Please read the 'Structure and Criteria' page for general guidance.



Raw data check

Review the raw data. Download from the [materials page](#).



Image check

Check that figures and images have not been inappropriately manipulated.

Privacy reminder: If uploading an annotated PDF, remove identifiable information to remain anonymous.

Files

Download and review all files from the [materials page](#).

4 Figure file(s)

2 Other file(s)



Structure your review

The review form is divided into 5 sections. Please consider these when composing your review:

1. BASIC REPORTING
2. EXPERIMENTAL DESIGN
3. VALIDITY OF THE FINDINGS
4. General comments
5. Confidential notes to the editor

 You can also annotate this PDF and upload it as part of your review

When ready [submit online](#).





Editorial Criteria

Use these criteria points to structure your review. The full detailed editorial criteria is on your [guidance page](#).

BASIC REPORTING

-  Clear, unambiguous, professional English language used throughout.
-  Intro & background to show context. Literature well referenced & relevant.
-  Structure conforms to [PeerJ standards](#), discipline norm, or improved for clarity.
-  Figures are relevant, high quality, well labelled & described.
-  Raw data supplied (see [PeerJ policy](#)).

EXPERIMENTAL DESIGN

-  Original primary research within [Scope of the journal](#).
-  Research question well defined, relevant & meaningful. It is stated how the research fills an identified knowledge gap.
-  Rigorous investigation performed to a high technical & ethical standard.
-  Methods described with sufficient detail & information to replicate.

VALIDITY OF THE FINDINGS

-  Impact and novelty not assessed. Negative/inconclusive results accepted. *Meaningful* replication encouraged where rationale & benefit to literature is clearly stated.
-  Data is robust, statistically sound, & controlled.
-  Speculation is welcome, but should be identified as such.
-  Conclusions are well stated, linked to original research question & limited to supporting results.

Standout reviewing tips

3



The best reviewers use these techniques

Tip

Support criticisms with evidence from the text or from other sources

Example

Smith et al (J of Methodology, 2005, V3, pp 123) have shown that the analysis you use in Lines 241-250 is not the most appropriate for this situation. Please explain why you used this method.

Give specific suggestions on how to improve the manuscript

Your introduction needs more detail. I suggest that you improve the description at lines 57- 86 to provide more justification for your study (specifically, you should expand upon the knowledge gap being filled).

Comment on language and grammar issues

The English language should be improved to ensure that an international audience can clearly understand your text. Some examples where the language could be improved include lines 23, 77, 121, 128 – the current phrasing makes comprehension difficult.

Organize by importance of the issues, and number your points

1. Your most important issue
2. The next most important item
3. ...
4. The least important points

Please provide constructive criticism, and avoid personal opinions

I thank you for providing the raw data, however your supplemental files need more descriptive metadata identifiers to be useful to future readers. Although your results are compelling, the data analysis should be improved in the following ways: AA, BB, CC

Comment on strengths (as well as weaknesses) of the manuscript

I commend the authors for their extensive data set, compiled over many years of detailed fieldwork. In addition, the manuscript is clearly written in professional, unambiguous language. If there is a weakness, it is in the statistical analysis (as I have noted above) which should be improved upon before Acceptance.

A taxonomic reassessment of *Piramys auffenbergi*, a neglected turtle from the late Miocene of Piram Island, Gujarat, India

Gabriel S Ferreira^{Corresp., 1, 2}, Saswati Bandyopadhyay³, Walter G Joyce^{Corresp. 4}

¹ Faculdade de Filosofia, Ciências e Letras de Ribeirão Preto, Universidade de São Paulo, Ribeirão Preto, Brazil

² Fachbereich Geowissenschaften, Eberhard-Karls-Universität Tübingen, Tübingen, Germany

³ Geological Studies Unit, Indian Statistical Institute, Kolkata, India

⁴ Departement für Geowissenschaften, University of Fribourg, Freiburg, Switzerland

Corresponding Authors: Gabriel S Ferreira, Walter G Joyce

Email address: gsferreira@usp.br, walter.joyce@unifr.ch

Background. *Piramys auffenbergi* was described as an emydine turtle based on a well-preserved skull retrieved from late Miocene deposits exposed on Piram Island, India. The description and figures provided in the original publication are vague and do not support assignment to Emydinae. This taxon has mostly been ignored by subsequent authors.

Material & Methods. We reexamine the holotype specimen and provide an extensive description and diagnosis for *Piramys auffenbergi* and included this taxon in a global character-taxon matrix for Pleurodira.

Results. The presence of a processus trochlearis pterygoidei and the absence of nasals conclusively show pleurodiran affinities. Inclusion of *Piramys auffenbergi* in a phylogenetic analysis retrieves this taxon within Stereogenyini and as sister to the Asian taxa *Shweboemys pilgrimi* and *Brontochelys gaffneyi*.

Discussion. Our reexamination of the holotype of *Piramys auffenbergi* confidently rejects the original assessment of this taxon as an emydine testudinoid and conclusively shows affinities with the pleurodiran clade Stereogenyini instead. Even though most taxa from this lineage are thought to be coastal turtles, all Asian stereogenyines were collected from continental deposits, suggesting a more diverse paleoecology for the group.

A taxonomic reassessment of *Piramys auffenbergi*, a neglected turtle from the late Miocene of Piram Island, Gujarat, India

Gabriel S. Ferreira^{1,2}, Saswati Bandyopadhyay³, Walter G. Joyce⁴

¹Faculdade de Filosofia, Ciências e Letras de Ribeirão Preto, Universidade de São Paulo, Ribeirão Preto, Brazil

²Fachbereich Geowissenschaften, Eberhard Karls Universität Tübingen, Tübingen, Germany

³Geological Studies Unit, Indian Statistical Institute, Kolkata, India

⁴Departement für Geowissenschaften, Universität Freiburg, Freiburg, Switzerland

* Walter G. Joyce, walter.joyce@unifr.ch, walter.g.joyce@gmail.com

ABSTRACT

Background. *Piramys auffenbergi* was described as an emydine turtle based on a well-preserved skull retrieved from late Miocene deposits exposed on Piram Island, India. The description and figures provided in the original publication are vague and do not support assignment to Emydinae. This taxon has mostly been ignored by subsequent authors.

Material & Methods. We reexamine the holotype specimen and provide an extensive description and diagnosis for *Piramys auffenbergi* and included this taxon in a global character-taxon matrix for Pleurodira.

Results. The presence of a processus trochlearis pterygoidei and the absence of nasals conclusively show pleurodiran affinities. Inclusion of *Piramys auffenbergi* in a phylogenetic

analysis retrieves this taxon within Stereogenyini and as sister to the Asian taxa *Shweboemys pilgrimi* and *Brontochelys gaffneyi*.

Discussion. Our reexamination of the holotype of *Piramys auffenbergi* confidently rejects the original assessment of this taxon as an emydine testudinoid and conclusively shows affinities with the pleurodiran clade Stereogenyini instead. Even though most taxa from this lineage are thought to be coastal turtles, all Asian stereogenyines were collected from continental deposits, suggesting a more diverse paleoecology for the group.

INTRODUCTION

Piramys auffenbergi was described by Prasad (1974) based on a single skull (GSI 18133) retrieved from Neogene sediments exposed on the small island of Piram, Gujarat State, India (Fig. 1). Prasad (1974) identified the taxon as an emydine, but the diagnosis and the description are general and do not reveal any emydine affinities. Today, only cryptodiran cheloniids, geoemydids (formerly part of Emydidae), testudinids, and trionychids occur in India (TTWG, 2017), but stereogenyine pleurodires inhabited the area as recently as the Plio/Pleistocene (Gaffney et al., 2011).

Here we provide an extensive description and diagnosis of *Piramys auffenbergi* based on a reexamination of the holotype specimen. Phylogenetic analysis concludes that this taxon is a representative of Stereogenyini, a clade that includes other South Asian forms, namely the Plio/Pleistocene *Shweboemys pilgrimi* Swinton, 1939 from Myanmar, and the late Oligocene *Brontochelys gaffneyi* (Wood, 1970) from Pakistan. We also provide an updated account of the paleoecology and biogeography of the group, concluding that, although they were most likely

adapted (or at least highly tolerant) to salty waters, stereogenyines were still restricted geographically and not as widespread as modern sea turtles.

Institutional abbreviations.— GSI, Geological Survey of India, Kolkata, India.

SYSTEMATIC PALEONTOLOGY

PLEURODIRA Cope, 1865

PELOMEDUSOIDES Broin, 1988

PODOCNEMIDIDAE Cope, 1868

STEREOGENYINI Gaffney et al., 2011

STEREOGENYITA Gaffney et al., 2011

Piramys auffenbergi Prasad, 1974

Holotype.— GSI 18133, a partial skull (Prasad, 1974, figs. 2–4, pl. 3.1; Figure 2).

Type locality and horizon.— Piram Island (Fig. 1), Gulf of Khambhat (formerly Gulf of Cambay), Gujarat, India; conglomerate beds, middle Siwaliks (Prasad, 1974), Dhok Pathan age, late Miocene (Nanda, Sehgal & Chauhan, 2017).

Referred material.— No specimens have been referred to date.

Diagnosis.—*Piramys auffenbergi* can be diagnosed as a representative of Pleurodira by the presence of the processus trochlearis pterygoidei and of Pelomedusoides by the absence of nasal bones. *Piramys auffenbergi* can be diagnosed as a representative of Stereogenyini by the small entrance to the antrum postoticum, the condylus mandibularis projecting ventrally from the cavum tympani, and the posteriorly broad skull. *Piramys auffenbergi* is distinguished from the other members of Stereogenyini by the following combination of characters: a pinched snout (not present in *Latentemys plowdeni* Gaffney et al., 2011, *Bairdemys venezuelensis* [Wood &

Díaz de Gamero, 1971], and *B. sanchezi* Gaffney et al., 2008), a deep interorbital groove on the prefrontal and frontal (otherwise present in *Cordichelys antiqua* [Andrews, 1903]), foramen nervi trigemini located above the level of the sulcus palatinopterygoideus (in contrast to the more ventrally displaced foramen nervi trigemini in *Shweboemys pilgrimi* and *Brontochelys gaffneyi*), the lower temporal emargination rising above the ventral level of the orbit (shallower in *C. antiqua* and unknown for other Stereogenyita), and the foramen stapedio-temporale placed anteriorly (autapomorphic character).

DESCRIPTION

Preservations—The skull was likely buried intact but suffered much weathering damage more recently. The bone is nevertheless of good quality and sutures are well preserved. The palate and ventral aspects of the basicranium are still obscured by matrix, but the dermatocranium and the dorsal parts of the basicranium are exposed (Fig. 2). The margins of the external nares and the median portions of the labial margins are damaged, but the remaining margins likely approximate their original outlines. The temporal roofing, by contrast, is massively eroded and the extent of the upper temporal emarginations can therefore not be assessed anymore. Much of the postorbitals, parietals, quadratojugals, squamosals, and the supraoccipital are missing. Faint scute sulci are apparent on the postorbitals and parietals (Fig. 2A).

Prefrontal.—The prefrontals are relatively small, paired elements that form the anterodorsal aspects of the orbit and the anterior third of the narrow interorbital space. The anterior margins are damaged, but the prefrontals likely formed the dorsal roof of the external nares, **as nasals were likely absent**. The prefrontals contact the frontals posteriorly along a straight, transverse

suture, the maxillae ventrally along a straight, horizontal suture, and one another along a straight, median suture. The descending branch of the prefrontal contacts the frontal posteriorly and the maxilla ventrally, but potential, more-distal contacts with the vomer and palatine are obscured by matrix (Fig. 2C). The prefrontals jointly form a deep, median groove on the dorsal skull surface (Fig. 2A).

Frontal.—The frontals are the only bones that are undamaged. The dorsal plate of the frontals forms the posterior two thirds of the interorbital space and posterodorsal margins of the anterolaterally facing orbits. The frontals contact the prefrontals anteriorly along a straight, transverse suture, the postorbitals laterally along parasagittal sutures of uneven length, the parietals posteriorly along straight, transverse sutures, and one another along a straight, median suture. Within the orbits, the frontals contact the prefrontals anteriorly and the postorbitals laterally. The median groove formed by the prefrontals continues posteriorly onto the frontals (Fig. 2A), but fades to oblivion towards the posterior margins of these elements.

Parietal.—Symmetric damage to the dorsal plate of the parietal creates the illusion that the parietals are relatively small elements that define the margins of well-developed upper temporal emarginations, but all margins show signs of damage and the extent of this emargination cannot be assessed with any confidence. It is therefore only clear that the dorsal plates of the parietals contact the frontals anteriorly along straight, transverse sutures and the postorbitals along oblique, anterolateral sutures. Together with the postorbitals, the parietals form a pair of sulci that converge towards the posterior and that likely trace the outline of a large internarial scute (scute ii of Ferreira et al., 2015) that covered the prefrontals, frontals, and the medial portions of the postorbitals and parietals. The descending process of the parietals contacts the pterygoid anterior to the foramen nervi trigemini, forms the dorsal margin of this foramen, and contacts the

prootic and supraoccipital posterior to it (Fig. 2A). The foramen nervi trigemini is located above the level of the sulcus palatinoptyergoideus and is well separated from the foramen stapedio-temporale by the prootic (Fig. 2C). More anterior contacts, if present, are obscured by matrix.

Postorbital.—The posterior margins of the postorbitals show signs of damage and it is therefore not possible to assess if they originally contributed to the upper temporal emarginations, although the thickness of their damaged margins makes this implausible. However, as the intact posterior margins converge, it is unlikely that they were significantly larger than preserved. The dorsal plates of the postorbitals at least contact the frontals medially, the parietals posteromedially, the jugals ventrally, and the quadratojugals posterolaterally. The descending process of the postorbital forms much of the posterior wall of the orbit, but its distal contacts are obscured by matrix. We are not able to discern a dorsal pocket within the upper limits of the posterior wall of the orbit.

Jugal.—The jugals are relatively large elements that form the posteroventral aspects of the orbit. Although the posteroventral aspects are damaged on both sides, enough is preserved to indicate that a well-developed lower temporal emargination is developed that rises to the level of the lower third of the orbit and that is anterodorsally framed by the jugals (Fig. 2B). The jugals contact the postorbitals dorsally along horizontal sutures, the quadratojugal posteriorly along a transverse suture, and the maxillae anteroventrally along a slightly oblique suture. The jugals contact the maxilla anteriorly and the postorbitals laterally on the surface of the skull, but matrix obscures possible contacts within the orbit.

Quadratojugal.—Only a fragment of the quadratojugal remains on the right side of the skull (Fig. 2B). It is therefore unclear if the quadratojugal contributes to the upper or lower temporal emarginations, although a contribution to the lower temporal emargination appears highly likely.

Only a short, anteromedial contact is preserved with the postorbital, an elongate anterior contact with the jugal, and remnants of the posterior contact of the quadrate anterior to the cavum tympani.

Squamosal.—A fragment of the right squamosal is available at the back of the skull (Fig. 2A). This fragment contacts the quadrate anterolaterally and the opisthotic anteromedially within the upper temporal fossa. A likely contribution of the squamosal to the region posterior to the cavum tympani cannot be discerned from that of the quadrate. However, the massive nature of the damaged quadrates suggests that the squamosal did not contribute to the antrum postoticum.

Premaxilla.—The anterior tip of the skull is damaged, but the median, cleft like notch appears to be genuine, as much of the left labial margin is intact (Fig. 2D). We are unable to discern the contacts, or even the presence, of premaxillae.

Maxilla.—The maxillae can only be observed in anterior and lateral views, as the palate is hidden by matrix. On the skull surface, the high maxillae contact the jugals below the orbit along transverse sutures and the prefrontals dorsally along horizontal suture. These contacts can partially be traced within the orbits, but matrix obscures their full extent. It is unclear if an anterior contact is developed with the premaxillae. Likely contacts with the palatines are obscured by matrix. The maxillae otherwise form the labial margins of the jaw, the anteroventral margins of the orbit, and the lateral margin of the broad external nares.

Prootic.—Only the left prootic can be observed in dorsal view and its sutures are obscured by localized damage to the surface of the skull. The left prootic clearly contacts the descending process of the parietal anteromedially and the quadrate laterally. Contacts are certainly present with the supraoccipital posteromedially and the opisthotic posteriorly, but their orientation and size cannot be estimated with confidence. The left prootic partially roofs the foramen nervi

trigemini and together with the left quadrate forms the foramen stapedio-temporale, which is situated at the anterior margin of the otic capsule and oriented anteriorly (Fig. 2A).

Opisthotic.—Only the external aspects of the opisthotics can be observed. Within the upper temporal fossa, the opisthotics contact the supraoccipital anteromedially, the quadrate anterolaterally, the squamosal posterolaterally, and the exoccipital posteromedially.

Quadrate.—Much of the quadrates **remains** on both sides of the skull, but most surfaces are damaged, making it difficult to discern its original shapes and contacts. At the very least, the quadrate contacts the quadratojugal anterodorsally and the squamosal posteriorly along the skull surface. Within the upper temporal fossa, the quadrate contacts the squamosal posteriorly, the opisthotic posteromedially, and the prootic anteromedially. Although damage to the middle ear is significant, it is apparent that the quadrates fully enclosed the incisura columella auris and Eustachian tube and that the quadrates fully formed the highly reduced antrum postoticum. However, matrix obscures the presence of a possible precolumellar fossa. In lateral view (Fig. 2C), the mandibular condyle of the quadrate projects ventrally, beyond the ventral outline of the cavum tympani.

Supraoccipital.—The supraoccipital is heavily eroded and some of its contacts are obscured by damage. It is therefore not possible to assess its possible contribution to the dorsal skull roof or the length of the supraoccipital crest. Within the upper temporal fossa, the supraoccipital certainly contacts the parietal anteromedially, the prootic anterolaterally, the opisthotic posterolaterally, and the exoccipital posteriorly. In posterior view, the supraoccipital roofs the foramen magnum and contacts the exoccipitals ventrally.

Exoccipital.—The exoccipitals are both present, but their ventral portions are obscured by matrix. These bones form the lateral margins of the foramen magnum, contact the supraoccipital

dorsally, the opisthotic anterolaterally, but their likely ventral contacts with the basioccipital are obscured. Although the occipital condyle is heavily damaged, the sulcus left by the contact between the exoccipitals laterally and the basioccipital ventrally are still visible (Fig. 2E).

MATERIAL & METHODS

Matrix & Phylogenetic Analysis

We integrated the holotype of *Piramys auffenbergi* into the phylogenetic matrix of Ferreira et al. (2018) to assess its relationships to other pleurodires. The matrix was modified through the addition of a new character (ch. 245: PM, median notch on upper jaw; see Supplementary Files) and by updating the scorings of some taxa (see Supplementary Files for full list of changes). As in the original analysis (Ferreira et al., 2018), twelve characters (14, 18, 19, 71, 95, 96, 99, 101, 119, 129, 174, 175) were interpreted as forming morphoclines and were ordered in the analysis. The resulting matrix was analyzed in TNT v. 1.5 (Goloboff & Catalano, 2016) using a traditional search with 2000 replicates of Wagner trees, random seed set to 0, branch-swapping algorithm Tree Bisection Reconnection (TBR), hold = 0, and collapsing zero-length branches according to rule '1'. The most parsimonious trees (MPTs) were subject to a second round of TBR and a strict consensus was obtained from the resulting MPTs. Consistency (CI) and Retention (RI) indexes, Bremer support, and resampling values (bootstrap and jackknife, using 1000 resamples for calculating absolute and difference of frequencies; Goloboff et al., 2003) were retrieved using implemented functions in TNT.

RESULTS

The search yielded 270 MPTs with 1134 steps (CI=0.288; RI=0.748). The strict consensus (see Supplementary Files) differs slightly from the results of Ferreira et al. (2018) by yielding a polytomy in the clade that includes the extant *Erymnochelys madagascariensis* (Grandidier 1867) and *Peltocephalus dumerilianus* (Schweigger 1812). *Piramys auffenbergi* is retrieved with relatively high support (Bremer support = 2; see Supplementary Files) inside Stereogenyini in a polytomy with *Stereogenys cromeri* Andrews, 1901 and *Shweboemys pilgrimi* (Fig. 3).

DISCUSSION & CONCLUSION

Alpha taxonomy and phylogenetic relationships

Prasad (1974) documented in the original description and figures of the holotype of *Piramys auffenbergi* the presence of large nasals, elongate prefrontals, small frontals that do not contribute to the orbits, and deep upper temporal emarginations. Prasad (1974) unfortunately did not explain why he felt *Piramys auffenbergi* to be an emydine, but we note that the presence of nasals is neither consistent with an assignment to Emydinae in particular, nor Testudinoidea more generally, as testudinoids universally lack nasal bones. Our re-examination of the type specimen allows us to better evaluate the cranial morphology and taxonomic affinities of this taxon and to present a more detailed list of diagnostic characters.

The most apparent differences in our interpretation of the skull is that nasals are absent, that the prefrontals are situated at the front of the orbits, that the frontals broadly contribute to the orbits, that the margin of the upper temporal emargination is not preserved, and that a trochlear process is present on the pterygoid. The presence of a trochlear process of the pterygoid and the absence of nasals clearly hint at pleurodiran and pelomedusoids affinities for *Piramys auffenbergi*, respectively (Gaffney, Tong & Meylan, 2006). The trochlear process is almost

completely embedded in matrix, but its lateral tip can be seen anterior to the otic capsule (Fig. 2C).

The posteriorly broadened skull, small anterior opening of the antrum postoticum, and a mandibular condyle projecting ventrally from the cavum tympani in lateral view (Fig. 2) suggest stereogenyine affinities (Gaffney et al., 2011; Ferreira et al., 2015). The results of our phylogenetic analysis (Fig. 3) support this initial hypothesis, even though *Piramys auffenbergi* is scored as “unknown” for several characters that support the monophyly of Stereogenyini. The majority of characters that resolve relationships within Stereogenyini, e.g., presence of a secondary palate with a midline cleft (Gaffney et al., 2011; Ferreira et al., 2015), refer to features in the ventral region of the skull. Unfortunately, this region is covered by matrix in GSI 18133, hampering a more detailed account of its taxonomic affinities. Nevertheless, the pinched snout (ch. 41, state 1), premaxillae protruding anteriorly beyond the dorsal edge of the apertura narium externa (ch. 43, state 1), the small opening of the antrum postoticum (ch. 86, state 2), the ventral projection of the mandibular condyle (ch. 90, state 1), and the median notch in the upper jaw (ch. 246, state 2) supports *Piramys auffenbergi* as a Stereogenyini with close affinities to *Stereogenys cromeri* and *Shweboemys pilgrimi* (Fig. 3).

Comparisons of *Piramys auffenbergi* with other Asian stereogenyines reveal osteological and chronological distinctness. The Bugti Hills in Pakistan, the type locality of *Brontochelys gaffneyi*, are now considered to be late Oligocene in age (Welcome et al., 2001; see discussion in the next section) whereas the Irrawaddy Beds in Myanmar, the type locality of *Shweboemys pilgrimi*, are thought to be Pliocene/Pleistocene age (Thein, Nu & Pyone, 2012). The sediments exposed on Piram Island are dated as late Miocene (Nanda, Sehgal & Chauhan, 2017) and *Piramys auffenbergi* therefore is temporally distinct. *Piramys auffenbergi* differs from

Brontochelys gaffneyi by the presence of a pinched snout, anterodorsally facing orbits, and a median notch in the upper jaw. On the other hand, *Piramys auffenbergi* differs from *Shweboemys pilgrimi* by the presence of an interorbital space that is narrower than the diameter of the orbits and the straight prefrontal-frontal suture. Furthermore, *Piramys auffenbergi* can be distinguished from both other Asian stereogenyines by the presence of a narrow interorbital groove, the more posterodorsally located foramen nervi trigemini, and the anterior, instead of dorsal, opening of the foramen stapedio-temporale. We are therefore able to confirm the validity of *Piramys auffenbergi*, though not as a testudinoid cryptodire, but rather as a podocnemidid pleurodire.

Paleoecology and biogeography

Members of Stereogenyini are currently thought to be durophagous, because they have broad and flat triturating surfaces combined with a secondary palate (Ferreira et al., 2015), features that are prevalent among extant turtles with durophagous dietary preferences (Foth, Rabi & Joyce, 2017). The palate is not exposed in the holotype of *Piramys auffenbergi* and we are therefore not able to assess the diet of this taxon with confidence. However, the high maxillae, anterodorsally oriented orbits, and posteriorly broadened skull with space for well-developed adductor muscles (Fig. 3) are features consistent with this life style. Considering the placement of *Piramys auffenbergi* within Stereogenyini we therefore predict the presence of durophagous features in this taxon as well.

Members of Stereogenyini are furthermore **though** to have had a marine, or at least coastal, lifestyle based on limb morphology (Weems & Knight, 2013), shell morphology (Pérez-García, 2017), and depositional environments (Ferreira et al., 2015). Although most stereogenyine taxa have indeed been recovered from sediments that represent coastal or open sea depositional

environments (Ferreira et al., 2015), we here note that all known Asian stereogenyines were collected from continental deposits, at least as indicated by the prevalence of terrestrial mammals in combination with fresh water aquatic fish and reptiles. In particular, the type of *Brontochelys gaffneyi* was collected from late Oligocene (not Miocene as reported Wood, 1970; Gaffney et al., 2011) continental sediments exposed in the region of Dera Bugti, Pakistan (Marivaux et al., 1999; Welcome et al., 2001), *Shweboemys pilgrimi* from the continental Plio/Pleistocene Irrawaddy Beds of Myanmar (Thein, Nu & Pyone, 2012), and, as demonstrated herein, *Piramys auffenbergi* from the continental late Miocene middle Siwaliks of India (Nanda, Sehgal & Chauhan, 2017). Given that all Asian taxa also form a clade (Fig. 3), it appears that the more continental clade Stereogenyita is nested within the more marine adapted clade Stereogenyini (Fig. 3), indicating a certain amount of ecological fluidity within the group. However, as deltaic and marine sediments often interfinger finely and as estuarine and coastal turtles are expected to be found in numerous coastal facies, we suggest that the available data is not sufficient to establish a rigorous pattern for the moment. Nevertheless, the fact that our phylogenetic analysis retrieves two geographically separated clades, the Afro-Asian Stereogenyita and the South American Bairdemydita, suggests that the even more salt tolerant forms were not highly marine, as greater amounts of dispersal and less endemism were otherwise to be expected.

Despite the broad geographic distribution of stereogenyines across North and South America, Africa, and India, they did not achieve the degree of cosmopolitanism seen in modern sea turtles (*Chelonoidae*; TTWG 2017). Instead, it appears that they were restricted to equatorial and subtropical regions of the northern hemisphere (Fig. 4). This hints at the possibility that their distribution was limited by cooler temperatures outside of the tropics.

ACKNOWLEDGEMENTS

We would like to thank all officials and supporting staff at the Curatorial Division of the Geological Survey of India and of the Indian Museum for generously providing access to collections under their care, in particular Senior Geologist A. Bhattacharya, Senior Geologist M. R. Moulick, and Senior Geologist A. K. Mondal. This contribution was funded by grants from the Swiss National Science Foundation to WGJ (SNF 20021_153502/2) and the Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP) to GSF (2014/25379-5 and 2016/03934-2).

REFERENCES

- Andrews CW. 1901.** Preliminary note on some recently discovered extinct vertebrates from Egypt (Part II). *Geological Magazine* **8**:436–444.
- Andrews CW. 1903.** On some pleurodiran chelonians from the Eocene of the Fayum, Egypt. *Annals and Magazine of Natural History Series* **7**:115–122.
- Broin F. 1988.** Les tortues et le Gondwana. Examen des rapports entre le fractionnement du Gondwana et la dispersion géographique des tortues pleurodires à partir du Crétacé. *Studia Palaeocheloniologica* **2**:103–142.
- Cope ED. 1865.** Third contribution to the herpetology of tropical America. *Proceedings of the Academy of Natural Sciences of Philadelphia* **1865**:185–198.
- Cope ED. 1868.** On the origin of genera. *Proceedings of the Academy of Natural Sciences of Philadelphia* **1868**:242–300.
- Dacqué E. 1912.** Die fossilen Schildkröten Aegyptens. *Geologische und Palaeontologische Abhandlungen* **14**:275–337.

- 321 **Ferreira GS, Bronzati M, Langer MC, Sterli J. 2018.** Phylogeny, biogeography and
- 322 diversification patterns of side-necked turtles (Testudines: Pleurodira). *Royal Society Open*
- 323 *Science* **5**:171773 DOI 10.1098/rsos.171773.
- 324 **Ferreira GS, Rincón AD, Solórzano A, Langer MC. 2015.** The last marine pelomedusoids
- 325 (*Testudines: Pleurodira*): a new species of *Bairdemys* and the paleoecology of *Stereogenyina*.
- 326 *PeerJ* **3**:e1063 DOI 10.7717/peerj.1063.
- 327 **Foth C, Rabi M, Joyce WG. 2017.** Skull shape variation in recent and fossil Testudinata and its
- 328 relation to habitat and feeding ecology. *Acta Zoologica* **98**:310–325 DOI 10.1111/azo.12181.
- 329 **Gaffney ES, Meylan PA, Wood RC, Simons E, Campos DA. 2011.** Evolution of the side-
- 330 necked turtles: the Family Podocnemididae. *Bulletin of the American Museum of Natural*
- 331 *History* **350**:1–237 DOI 10.1206/350.1.
- 332 **Gaffney ES, Scheyer TM, Johnson KG, Bocquentin J, Aguilera OA. 2008.** Two new species
- 333 of the side necked turtle genus, *Bairdemys* (Pleurodira, Podocnemididae), from the Miocene
- 334 of Venezuela. *Paläontologische Zeitschrift* **82**:209–229 DOI 10.1007/BF02988411.
- 335 **Gaffney ES, Tong H, Meylan PA. 2006.** Evolution of the side-necked turtles: the families
- 336 Bothremydidae, Euraxemydidae, and Araripemydidae. *Bulletin of the American Museum of*
- 337 *Natural History* **300**:1–700 DOI 10.1206/0003-0090(2006)300[1:EOTSTT]2.0.CO;2.
- 338 **Gaffney ES, Wood RC. 2002.** *Bairdemys*, a new side-necked turtle (Pelomedusoides:
- 339 Podocnemididae) from the Miocene of the Caribbean. *American Museum Novitates* **3359**:1–
- 340 28 DOI 10.1206/0003-0082(2002)359<0001:BANSNT>2.0.CO;2.
- 341 **Goloboff PA, Catalano SA. 2016.** TNT version 1.5, including a full implementation of
- 342 phylogenetic morphometrics. *Cladistics* **32**:221–238 DOI 10.1111/cla.12160.

- Goloboff PA, Farris JS, Källersjö M, Oxelman B, Ramírez MJ, Szumik CA. 2003.**
Improvements to resampling measures of group support. *Cladistics* **19**:324–332 DOI
10.1111/j.1096-0031.2003.tb00376.x
- Grandidier A. 1867.** Liste des reptiles nouveaux decouverts, en 1866, sur la côte sud-ouest de
Madagascar. *Revue et Magasin de Zoologie Pure et Appliquée* **19**:232–234.
- Marivaux L, Vianey-Liaud M, Welcomme J-L. 1999.** Première découverte de Cricetidae
(Rodentia, Mammalia) Oligocènes dans le synclinal Sud de Gandoï (Bugti Hills,
Balouchistan, Pakistan). *Comptes Rendus de l'Académie des Sciences* **329**:839–844.
- Nanda AC, Sehgal RK, Chauhan PR. 2017.** Siwalik-age faunas from the Himalayan Foreland
Basin of South Asia. *Journal of Asian Earth Sciences* **162**:54–68 DOI
10.1016/j.jseaes.2017.10.035.
- Pérez-García A. 2017.** New information and establishment of a new genus for the Egyptian
Paleogene turtle ‘*Stereogenys*’ *libyca* (Podocnemididae, Erymnochelyinae). *Historical
Biology* 1–11 DOI 10.1080/08912963.2017.1374383.
- Prasad KN. 1974.** The vertebrate fauna from Piram Island, Gujarat, India. *Memoires of the
Geological Survey of India, New Series* **41**:1–23.
- Schweigger AF. 1812.** Prodromus Monographiae Cheloniorum. *Konigsberger Archiv für
Naturwissenschaft und Mathematik* **1**:406–458.
- Swinton WE. 1939.** A new fresh-water tortoise from Burma. *Records of the Geological Survey
of India* **74**:548–551.
- Thein ZMM, Nu TT, Pyone WW. 2012.** Lithofacies and depositional environment of the
Irrawaddy Formation in Myogintha Area, Pauk Township, Magway region. *Universities
Research Journal* **5**:69–82.

- TTWG [Turtle Taxonomy Working Group]. 2017. Turtles of the world: annotated checklist and atlas of taxonomy, synonymy, distribution, and conservation status (8th ed). *Chelonian Research Monographs* 7:1–292.
- Weems RE, Knight JL. 2013. A new species of *Bairdemys* (Pelomedusoides: Podocnemididae) from the Oligocene (Early Chattian) Chandler Bridge Formation of South Carolina, USA, and its paleobiogeographic implications for the genus. In: Brinkman D, Holroyd P, Gardner J, eds. *Morphology and evolution of turtles, Vertebrate paleobiology and paleoanthropology series*. Netherlands: Dordrecht, 289–303.
- Welcome J-L, Benammi M, Crochet J-Y, Marivaux L, Métais G, Antoine P-O, Baloch I. 2001. Himalayan forelands: paleontological evidence for Oligocene detrital deposits in the Bugti Hills (Balochistan, Pakistan). *Geological Magazine* 138:397–405.
- Wood RC. 1970. A review of the fossil Pelomedusidae (Testudines, Pleurodira) of Asia. *Breviora* 357:1–24.
- Wood RC, Díaz de Gamero ML. 1971. *Podocnemis venezuelensis*, a new fossil pelomedusid (Testudines, Pleurodira) from the Pliocene of Venezuela and a review of the history of *Podocnemis* in South America. *Breviora* 376:1–23.

Figure 1: The type locality of *Piramys auffenbergi*.

(A) A simplified map of India. Gujarat State is highlighted in purple, all other states in various shades of grey. (B) Detailed view of the Gulf of Khambhat. Piram Island is highlighted in purple.

Figure 2: GSI 18133, *Piramys auffenbergi*, holotype, late Miocene of Piram Island, Gujarat, India.

Photographs and illustrations of specimen in (A) dorsal, (B) right lateral, (C) left lateral, (D) anterior, and (E) posterior views. Abbreviations: *fnt*, foramen nervi trigemini; *fr*, frontal; *fst*, foramen stapedio-temporale; *ju*, jugal; *mx*, maxilla; *op*, opisthotic; *pa*, parietal; *pf*, prefrontal; *po*, postorbital; *pr*, prootic; *ptp*, processus trochlearis pterygoidei; *qj*, quadratojugal; *qu*, quadrate; *so*, supraoccipital; *sq*, squamosal.

Figure 3: Phylogenetic hypothesis of Stereogenyina.

A time-calibrated cladogram depicting a portion of the strict consensus topology of 270 MPTs with 1134 steps retrieved from the phylogenetic analysis using ordered characters. Dark lines highlight the known temporal distribution of a taxon and the colored continent symbols highlight their known spatial distribution. The full tree is provided in the Supplementary File S2.

Figure 4: Geographic distribution of stereogenyines.

Section of world map indicating localities of extinct taxa assigned to Stereogenyini. Red circles indicate Stereogenyita, green circles Bairdemydita, and blue circles non-Stereogenyita and non-Bairdemydita stereogenyines. 1, *Bairdemys venezuelensis*; 2, *B. thalassica* Ferreira et al., 2015; 3, *B. sanchezi*; 4, *B. winklerae* Gaffney et al., 2008; 5, *B. hartsteini* Gaffney & Wood, 2002; 6, "*B.*" *healeyorum* Weems & Knight, 2013; 7, *Mogharemys blanckenhorni* (Dacqu , 1912); 8, *Cordichelys antiqua*; 9, *Latentemys plowdeni*; 10, *Stereogenys cromeri*; 11, *Lemurchelys diasphax* Gaffney et al., 2011; 12, *Brontochelys gaffneyi*; 13, *Shweboemys pilgrimi*; 14, *Piramys auffenbergi*. Abbreviations: EG, Egypt, MM, Myanmar, PK, Pakistan, PR, Puerto Rico, US, United States of America, VE, Venezuela.

Supplemental File S1: Character taxon matrix used in phylogenetic analysis, including full character list and character state definitions.

Supplemental File S2: List of changes to Ferreira et al. (2018)'s matrix, strict consensus tree retrieved from the phylogenetic analysis, bootstrap, jackknife and Bremer support values, and list of common synapomorphies for Stereogenyina.

Figure 1

The type locality of *Piramys auffenbergi*.

(A) A simplified map of India. Gujarat State is highlighted in purple, all other states in various shades of grey. (B) Detailed view of the Gulf of Khambhat. Piram Island is highlighted in purple.




Figure 2

GSI 18133, *Piramys auffenbergi*, holotype, late Miocene of Piram Island, Gujarat, India.

Photographs and illustrations of specimen in (A) dorsal, (B) right lateral, (C) left lateral, (D) anterior, and (E) posterior views. Abbreviations: *fnt*, foramen nervi trigemini; *fr*, frontal; *fst*, foramen stapedio-temporale; *ju*, jugal; *mx*, maxilla; *op*, opisthotic; *pa*, parietal; *pf*, prefrontal; *po*, postorbital; *pr*, prootic; *ptp*, processus trochlearis pterygoidei; *qj*, quadratojugal; *qu*, quadrate; *so*, supraoccipital; *sq*, squamosal.




Figure 3

Phylogenetic hypothesis of Stereogenyina.

A time-calibrated cladogram depicting a portion of the strict consensus topology of 270 MPTs with 1134 steps retrieved from the phylogenetic analysis using ordered characters. Dark lines highlight the known temporal distribution of a taxon and the colored continent symbols highlight their known spatial distribution. The full tree is provided in the Supplementary File S2.




Figure 4

Geographic distribution of stereogenyines.

Section of world map indicating localities of extinct taxa assigned to Stereogenyini. Red circles indicate Stereogenyita, green circles Bairdemydita, and blue circles non-Stereogenyita and non-Bairdemydita stereogenyines. 1, *Bairdemyys venezuelensis*; 2, *B. thalassica* Ferreira et al., 2015; 3, *B. sanchezi*; 4, *B. winklerae* Gaffney et al., 2008; 5, *B. hartsteini* Gaffney & Wood, 2002; 6, "*B.*" *healeyorum* Weems & Knight, 2013; 7, *Mogharemys blanckenhorni* (Dacqu  , 1912); 8, *Cordichelys antiqua*; 9, *Latentemys plowdeni*; 10, *Stereogenys cromeri*; 11, *Lemurchelys diasphax* Gaffney et al., 2011; 12, *Brontochelys gaffneyi*; 13, *Shweboemys pilgrimi*; 14, *Piramys auffenbergi*. Abbreviations: EG, Egypt, MM, Myanmar, PK, Pakistan, PR, Puerto Rico, US, United States of America, VE, Venezuela.

