

# Anti-predator defences of a bombardier beetle: is bombing essential for successful escape from frogs?

Shinji Sugiura <sup>Corresp. 1</sup>

<sup>1</sup> Graduate School of Agricultural Science, Kobe University, Kobe, Japan

Corresponding Author: Shinji Sugiura  
Email address: ssugiura@people.kobe-u.ac.jp

Some animals, such as the bombardier beetles (Coleoptera: Carabidae: Brachinini), have evolved chemical defences against predators. When attacked, bombardier beetles can discharge noxious chemicals at temperatures of approximately 100°C from the tip of their abdomens, “bombing” their attackers. Although many studies to date have investigated how bombardier beetles discharge defensive chemicals against predators, relatively little research has examined how predators modify their attacks on bombardier beetles to avoid being bombed. In this study, I observed the black-spotted pond frog *Pelophylax nigromaculatus* (Anura: Ranidae) attacking the bombardier beetle *Pheropsophus jessoensis* under laboratory conditions. In Japan, *Pe. nigromaculatus* is a generalist predator in grasslands where the bombardier beetle frequently occurs. Almost all the frogs (92.9%) observed rejected live bombardier beetles; 67.9% stopped their attacks once their tongues touched the beetles, and 25.0% spat out the beetles immediately after taking the beetles into their mouths. No beetle bombed a frog before being taken into a frog’s mouth. All beetles taken into mouths bombed the frogs. Only 7.1% of the frogs swallowed live bombardier beetles after being bombed in the mouth. When dead beetles were provided instead, 85.7% of the frogs rejected the dead beetles, 71.4% stopped their attacks after their tongues touched the beetles, and 14.3% spat out the beetles. Only 14.3% of the frogs swallowed the dead beetles. The results suggest that the frogs tended to stop their predatory attack before receiving a bombing response from the beetles. Therefore, bombing was not essential for the beetles to successfully defend against the frogs. Using its tongue, *Pe. nigromaculatus* may be able to rapidly detect a deterrent chemical or physical characteristics of its potential prey *Ph. jessoensis* and thus avoid injury by stopping its predatory attack before the beetle bombs it.

# **Anti-predator defences of a bombardier beetle: is bombing essential for successful escape from frogs?**

Shinji Sugiura

Graduate School of Agricultural Science, Kobe University, Kobe, Japan

Corresponding author: S. Sugiura, [ssugiura@people.kobe-u.ac.jp](mailto:ssugiura@people.kobe-u.ac.jp), [sugiura.shinji@gmail.com](mailto:sugiura.shinji@gmail.com)

Running title: Bombardier beetle defences against frogs

# ABSTRACT

Some animals, such as the bombardier beetles (Coleoptera: Carabidae: Brachinini), have evolved chemical defences against predators. When attacked, bombardier beetles can discharge noxious chemicals at temperatures of approximately 100°C from the tip of their abdomens, “bombing” their attackers. Although many studies to date have investigated how bombardier beetles discharge defensive chemicals against predators, relatively little research has examined how predators modify their attacks on bombardier beetles to avoid being bombed. In this study, I observed the black-spotted pond frog *Pelophylax nigromaculatus* (Anura: Ranidae) attacking the bombardier beetle *Pheropsophus jessoensis* under laboratory conditions. In Japan, *Pe. nigromaculatus* is a generalist predator in grasslands where the bombardier beetle frequently occurs. Almost all the frogs (92.9%) observed rejected live bombardier beetles; 67.9% stopped their attacks once their tongues touched the beetles, and 25.0% spat out the beetles immediately after taking the beetles into their mouths. No beetle bombed a frog before being taken into a frog’s mouth. All beetles taken into mouths bombed the frogs. Only 7.1% of the frogs swallowed live bombardier beetles after being bombed in the mouth. When dead beetles were provided instead, 85.7% of the frogs rejected the dead beetles, 71.4% stopped their attacks after their tongues touched the beetles, and 14.3% spat out the beetles. Only 14.3% of the frogs swallowed the dead beetles. The results suggest that the frogs tended to stop their predatory attack before receiving a bombing response from the beetles. Therefore, bombing was not essential for the beetles to successfully defend against the frogs. Using its tongue, *Pe. nigromaculatus* may be able to rapidly detect a deterrent chemical or physical characteristics of its potential prey *Ph. jessoensis* and thus avoid injury by stopping its predatory attack before the beetle bombs it.

# INTRODUCTION

36

37 Physical and chemical defences have evolved in many organisms to protect against natural  
 38 enemies (Edmunds, 1974; Eisner, Eisner & Siegler, 2005). For example, some plant and animal  
 39 species have developed physical deterrents such as thorns and spines (Edmunds, 1974; Cooper &  
 40 Owen-Smith, 1986; Eisner, 2003; Sugiura & Yamazaki, 2014; Sugiura, 2016; Ito, Taniguchi &  
 41 Billen, 2016), while other species produce defensive chemicals, including toxic substances, to  
 42 prevent themselves from being eaten (Eisner, 2003; Eisner, Eisner & Siegler, 2005; Derby, 2007;  
 43 Mithöfer & Boland, 2012). Organisms whose defence mechanisms can cause severe injury to  
 44 their natural enemies have also evolved warning signals, such as conspicuous body colouration  
 45 or particular sounds (Lev-Yadun, 2001; Ruxton, Sherratt & Speed, 2004; Inbar & Lev-Yadun,  
 46 2005; Bonacci et al., 2008; Lev-Yadun, 2009; Bura et al., 2016; Sugiura & Takanashi, 2018). In  
 47 response, predators have evolved specific abilities to avoid such well-defended prey by  
 48 recognising warning colouration or detecting chemical signals (Edmunds, 1974; Endler, 1991;  
 49 Ruxton, Sherratt & Speed, 2004; Skelhorn & Rowe, 2006; Williams et al., 2010).

50 Adult bombardier beetles (Coleoptera: Carabidae: Brachinini) bomb, i.e. discharge noxious  
 51 chemicals from the tip of their abdomens at temperatures of approximately 100°C, when they are  
 52 disturbed or attacked (Aneshansley et al., 1969; Dean, 1979; Eisner, 2003; Eisner, Eisner &  
 53 Siegler, 2005; Arndt et. al., 2015). Such ejection of hot chemicals is only known in the  
 54 coleopteran family Carabidae. Previous studies have investigated how bombardier beetles  
 55 successfully defend against predators (Eisner, 1958; Eisner & Meinwald, 1966; Eisner & Dean,  
 56 1976; Dean, 1980a; Eisner, Eisner & Aneshansley, 2005; Eisner et al., 2006). Bombardier  
 57 beetles can aim their abdominal discharge in virtually any direction, spraying various parts of  
 58 their own bodies (e.g., legs and dorsal surface) with the toxic chemicals (Eisner & Aneshansley,  
 59 1999). Dean (1980b) reported that predators displayed intense responses to the unheated

chemical discharges of bombardier beetles in experiments. This suggests that the cooled chemicals coating the beetles' body surfaces function as the primary defence against predators. Successful defence mediated by chemicals on the body surfaces of beetles may reduce the costs of spraying (bombing). Further research is needed to clarify the relative importance of chemical toxicity and heat for overall successful anti-predatory defence.

Frogs and toads are important predators of carabid beetles (Laroche, 1974a,b). However, bombardier beetles have rarely been found in the gut contents and faeces of frogs and toads (Laroche, 1974a,b; Sarashina, Yoshihisa & Yoshida, 2011; except Mori, 2008), suggesting that bombing prevents toads and frogs from swallowing and ingesting beetles (Eisner & Meinwald, 1966; Dean, 1980a; Eisner, 2003; Sugiura & Sato, 2018). Still, only a few studies have investigated the factors that cause anuran predators to stop preying on bombardier beetles (Dean, 1980b). Elucidating these ecological factors would contribute to a better understanding of the evolution of anti-predatory defences in insects.

This study aims to investigate the responses of the black-spotted pond frog *Pelophylax nigromaculatus* (Hallowell) (Anura: Ranidae) to the defensive behaviour of the bombardier beetle *Pheropsophus jessoensis* (Morawitz). *Pheropsophus jessoensis* is a bombardier beetle found in East Asia (Ueno et al., 1985; Jung et al., 2012); the beetle is a common inhabitant of farmlands, grasslands, and forest edges in Japan (Habu & Sadanaga, 1965; Ueno et al., 1985; Yahiro et al., 1992; Ishitani & Yano, 1994; Fujisawa et al., 2012; Ohwaki et al., 2015; Sugiura & Sato, 2018). Adult *Ph. jessoensis* eject toxic chemicals (1,4-benzoquinone and 2-methyl-1,4-benzoquinone) at a temperature of approximately 100°C from their rear ends in response to predator attacks (Video S1; Kanehisa and Murase, 1977; Kanehisa, 1996). *Pelophylax nigromaculatus* is a true frog inhabiting wetlands and farmlands of East Asia (Liu et al., 2010; Tsuji et al., 2011; Komaki et al., 2015), being one of the most abundant frog species of

traditional agricultural landscapes including farmlands, grasslands, and forest edges (Hirai, 2002; Honma, Oku & Nishida, 2006; Tsuji et al., 2011; Matsushashi & Okuyama, 2015). Using its tongue, *Pe. nigromaculatus* readily catches and swallows smaller prey (Video S2; Honma, Oku & Nishida, 2006). *Pelophylax nigromaculatus* is a generalist predator that has been reported to prey on carabid beetles (Maeda & Matsui, 1999; Hirai & Matsui, 1999; Sano & Shinohara, 2011; Sarashina, Yoshihisa & Yoshida, 2011). As *Ph. jessoensis* and *Pe. nigromaculatus* co-occur in the same grassland habitats, this frog species is a potential predator of adult *Ph. jessoensis*. In early June 2016, I offered an adult *Ph. jessoensis* to an adult *Pe. nigromaculatus* under laboratory conditions. The frog attacked the beetle, but stopped the attack immediately after its tongue touched the beetle. No bombing sounds were heard, suggesting that the frog ceased its attack before the beetle bombed. Therefore, I hypothesised that bombing is not essential when *Ph. jessoensis* seeks to avoid being swallowed by *Pe. nigromaculatus*. To test this hypothesis, I observed *Pe. nigromaculatus* attacking *Ph. jessoensis* under laboratory conditions using a digital video camera. Acceptance or rejection of prey was carefully investigated using slow-motion videos. Furthermore, both dead and live beetles were used to test whether bombing is essential for successful defence against predatory attacks by *Pe. nigromaculatus*. Finally, I discuss the importance of primary and secondary defences in terms of overall anti-predation defence.

## MATERIALS AND METHODS

### Sampling

Approximately 100 adult *Ph. jessoensis* were collected from grasslands and forest edges in Kato (34°54'N, 135°02'E, 120 m above sea level), Hyogo Prefecture, central Japan, from May to

August in 2016, 2017, and 2018 (cf. Sugiura & Sato, 2018). Body weight was measured to the nearest 0.1 mg using an electronic balance (PA64JP, Ohaus, Tokyo, Japan). Study individuals were maintained separately in plastic cases (diameter: 85 mm; height: 25 mm) with wet tissue paper in the laboratory at 25°C. Dead larvae of *Spodoptera litura* (Fabricius) (Lepidoptera: Noctuidae) were provided as food (Sugiura & Sato, 2018). Beetles were not used repeatedly in different feeding experiments. All experiments were conducted  $22.6 \pm 4.0$  (means  $\pm$  standard errors; range: 4–69) days after the beetles were collected.

Approximately 100 individuals of *Pe. nigromaculatus* were collected from wetlands and forest edges in Takarazuka-shi (34°53'N, 135°17'E, 230 m above sea level), Sanda-shi (34°57'N, 135°11'E, 180 m above sea level), and Sayo-cho (35°02'N, 134°20'E, 180 m above sea level), Hyogo Prefecture, central Japan, from May to August in 2016, 2017, and 2018. The distances between these sites and the sampling site of *Ph. jessoensis* ranged from 15.6 to 65.4 km.

Although *Pe. nigromaculatus* has recently been classified as near threatened (NT) in the Japanese Red Data List (Ministry of the Environment of Japan, 2017), this species was abundant at the collection sites. Both juveniles and adults were collected. Body weight was measured to the closest 0.01 g using an electronic balance (EK-120A, A&D, Tokyo). Small and large frogs were maintained separately in small (120 × 85 × 130 mm, length × width × height) and large plastic cages (120 × 185 × 130 mm, length × width × height), respectively, in the laboratory at 25°C. Live larvae of *S. litura*, *Tenebrio molitor* Linnaeus (Coleoptera: Tenebrionidae), and *Zophobas atratus* Fabricius (Coleoptera: Tenebrionidae) were provided as food. Frogs were starved for 24 h before the feeding experiments to standardise their hunger level (cf. Honma, Oku & Nishida, 2006). As with the beetles, individual frogs were not used repeatedly. The experiments were conducted  $18.6 \pm 2.5$  (means  $\pm$  standard errors; range: 4–66) days after the frogs were collected. The frogs were released after the experiments were completed.

# **Feeding experiments**

Feeding experiments were all conducted at 25°C. To start, a frog was placed in a transparent plastic container (120 × 85 × 130 mm, length × width × height). Then, a transparent glass petri dish (45 mm in diameter, 15 mm in height) containing a live bombardier beetle was placed outside the plastic container where the frog could see it. Frogs that did not try to attack the beetle (34.1%) were not used for the feeding experiments. However, frogs that ignored *Ph. jessoensis* did not respond to other prey (i.e. *T. molitor* larvae). If a frog displayed attacking behaviour (i.e. opening the mouth and shooting out the tongue to capture prey; Video S2), a live beetle was then placed in the container with the frog. The resulting behaviours were recorded on video using a digital camera (iPhone 6 plus, Apple) at 240 frames per second. If the frog did not swallow the beetle, palatable prey (a *T. molitor* larva) was offered to the frog several minutes after beetle rejection to determine whether the frog was hungry. If a frog swallowed the beetle, I observed whether it vomited the beetle within 330 min of swallowing it (cf. Sugiura & Sato, 2018). Vomited beetles were checked to see whether they were still alive. Frogs that did not vomit after swallowing were considered to have digested the beetle. Frog faeces were examined after the experiment to confirm whether the beetles were digested. In total, 28 frogs and 28 live bombardier beetles were used in the experiments. The means ± standard errors of the frog and live beetle body weights were 10.23 ± 1.39 g ( $n = 28$ ) and 213.0 ± 10.0 mg ( $n = 28$ ), respectively.

A second set of frogs were presented with dead adult beetles to test whether the bombing response is essential for deterring a predatory attack. *Pelophylax nigromaculatus* usually does not attack motionless prey. However, in a pilot test, an individual of *Pe. nigromaculatus* attacked



and ingested a dead caterpillar (*S. litura*) when forceps were used to move the caterpillar within the frog's field of view. For this experiment, the bombardier beetles were killed in a freezer at  $-15^{\circ}\text{C}$ . First, a dead beetle was placed in the plastic container ( $120 \times 85 \times 130$  mm, length  $\times$  width  $\times$  height) within the frog's field of view. If the frog did not initially respond to the beetle, forceps were used to move the dead beetle within the frog's field of view again. Frogs that did not attack the dead beetles (29.2%) were not used in these experiments; frogs that ignored *Ph. jessoensis* did not respond to other prey (i.e., *T. molitor* larvae). The predatory behaviours of the frogs were recorded using the same digital video camera. Frogs that did not swallow dead beetles were offered *T. molitor* larvae several minutes after beetle rejection to check whether they were hungry. Twenty-eight frogs and 28 dead beetles were used in this experiment. The means  $\pm$  standard errors of the frog and dead beetle body weights were  $8.65 \pm 1.25$  g ( $n = 28$ ) and  $214.8 \pm 7.6$  mg ( $n = 28$ ), respectively. The mean body weight of frogs that attacked dead beetles did not differ significantly from the mean body weight of frogs that attacked live beetles (*t*-test,  $t = 0.84$ ,  $P = 0.40$ ). The mean body weight of dead beetles did not differ significantly from the mean body weight of live beetles (*t*-test,  $t = -0.14$ ,  $P = 0.89$ ).

Videos of frogs responding to live and dead beetles were played back using QuickTime Player version 10.4 (Apple, Inc.). Frog responses to the bombardier beetles were grouped into four categories (cf. Ito, Taniguchi & Billen, 2016; Matsubara & Sugiura, 2017; Sugiura & Sato, 2018): (1) frogs that touched the beetles with their tongues but did not take the beetles into their mouths; (2) frogs that spat out the beetles after taking them into the mouth; (3) frogs that swallowed beetles but vomited them later; and (4) frogs that swallowed and digested the beetles. I also assessed whether frogs that rejected beetles (1–2) resumed their attacks within 10 seconds.

All experiments were performed in accordance with the Kobe University Animal Experimentation Regulations (Kobe University Animal Care and Use Committee, 27–01, 30–01).

No frogs were seriously injured or killed during the feeding experiments. My study also complies with the current laws of Japan.

## Data analysis

Generalised linear models (GLMs) featuring binomial error distributions and logit links (i.e., logistic regressions) were used to identify factors that contributed to frogs' successful swallowing and digestion of the bombardier beetles. The success or failure (1/0) of frogs' swallowing and digesting beetles was used as the response variable. Frog weight, beetle weight, the frog weight  $\times$  beetle weight interaction, and beetle condition (live or dead) were treated as fixed factors. When the residual deviance was 1.5-fold larger (overdispersion) or smaller (underdispersion) than the residual degrees of freedom, a quasi-binomial error distribution was used rather than a binomial error distribution. Furthermore, fixed factors were subjected to likelihood ratio testing when marginal significance was evident. Thus, the significance of models with and without the factors of interest were compared using the GLMs. All analyses were performed using R ver. 3.3.2 (R Development Core Team 2016).

## RESULTS

In the experiment using live adult bombardier beetles ( $n = 28$ ), 26 frogs (92.9%) rejected the beetles without swallowing them (Fig. 1); 19 frogs (67.9%) stopped attacking the beetles immediately after touching the beetles with their tongues (Fig. 2; Video S3), and 7 frogs (25.0%) spat out the beetles after taking the beetles into their mouths (Fig. 3; Video S4). No beetle bombed a frog before being taken into the frog's mouth. All beetles that were taken into frog

mouths bombed (Video S4). Only two frogs (7.1%) were observed to swallow the bombardier beetles (Table 1) after being bombed in the mouth; one of the frogs successfully digested the beetle, but the other frog vomited the beetle 18 min after swallowing it (Table 1). The vomited beetle was still alive. Of the frogs that took the beetles into their mouths, 88.9% ( $n = 8/9$ ) initially stopped attacking the beetles when their tongues first touched the beetles, but resumed their predatory attack soon thereafter (Fig. 3; Video S4). Frogs that did not swallow beetles consumed other prey (i.e., *T. molitor* larvae) soon thereafter.

When dead beetles were used ( $n = 28$ ), 24 frogs (85.7%) rejected the dead beetles without swallowing them (Fig. 1); 20 frogs (71.4%) stopped attacking the beetles after their tongues touched the dead beetles (Video S5), and 4 frogs (14.3 %) spat out the beetles after taking the beetles into their mouths (Fig. 1). Only 4 frogs (14.3%) swallowed the dead beetles. Similar to the experiment using live beetles, 87.5% of the frogs that took beetles into their mouths ( $n = 7/8$ ) were initially deterred when their tongues first touched the beetles, but continued with their predatory behaviour soon afterwards. The frogs that did not swallow beetles ate other prey (i.e., *T. molitor* larvae) soon thereafter.

The proportion of dead beetles swallowed by frogs (14.3%) was higher than that of live beetles (7.1%). However, the GLM results indicated that the frog swallowing rates of live and dead beetles did not significantly differ (Table 1). Whether beetles were swallowed or not was associated with beetle size, but not frog size (Table 1). Beetles were more likely to be swallowed as beetle size decreased (Fig. 4A). The interaction of frog and beetle weight was not significant (Fig. 4A).

The proportion of dead beetles digested by frogs (14.3%) was higher than the proportion of live beetles swallowed (3.6%). The GLMs indicated that the difference between the digestion rates of live and dead beetles was significant (Table 2). Beetle size affected the digestion rate

(Table 2, Fig. 4B).

# DISUSSION

Here, I found that *Pe. nigromaculatus* frequently rejected *Ph. jessoensis* without attempting to swallow the beetles (Fig. 1). Around 70% of frogs stopped attacking both live and dead beetles immediately after touching the beetles with their tongues. Although the high-speed release of hot noxious chemicals (bombing) protected *Ph. jessoensis* from digestion by the frog *Pe. nigromaculatus* (Figs. 1, 3; Video S4), my findings support the hypothesis that bombing is not essential for *Ph. jessoensis* to successfully evade being swallowed by *Pe. nigromaculatus* (Figs. 1, 3; Video S3). Which factors, then, stopped the frogs from attacking? Three potential reasons can be considered: (1) the frogs recognised the warning colouration of the beetles; (2) the body size of the beetles was too large for the frog to accommodate; and (3) the frogs reflexively avoided the beetles after detecting toxic substances or other deterrent characteristics on the beetles' body surfaces.

The bombardier beetle *Ph. jessoensis* does have a striking yellow and black pattern on its body that could serve as warning colouration (Fig. 1), although this has not been empirically demonstrated. Anuran predators can avoid toxic prey by recognising certain colours or other morphological characteristics and then ignoring those prey (Brower, Brower & Westcott, 1960; Brower & Brower, 1962; Dean, 1980a; Taniguchi et al., 2005; Ito, Taniguchi & Billen, 2016). In fact, 34.1 and 29.2% of frogs did not seek to attack live or dead beetles, respectively, before the feeding experiments commenced, suggesting that *Pe. nigromaculatus* may recognise the body pattern and shape of *Ph. jessoensis* and interpret these as warning signals. However, frogs that ignored *Ph. jessoensis* did not respond to other prey (i.e., *T. molitor* larvae). Therefore, the

experimental conditions used may not be appropriate for analysing foraging by certain frogs. Alternatively, the yellow-and-black pattern of *Ph. jessoensis* may serve as disruptive camouflage; flightless *Ph. jessoensis* walk on soil of forest edges, grasslands, and agricultural fields. Further work is needed to explore the significance of *Ph. jessoensis* colour as a defensive trait.

GLM analysis indicated that beetle size was correlated with beetle-swallowing frequency of frogs (Table 1). *Pelophylax nigromaculatus* has been reported to spit out large prey that they were unable to swallow after taking the prey into their mouths (Honma, 2004; Honma, Oku & Nishida, 2006). However, 67.9% of the frogs in the experiment with live beetles and 71.4% of the frogs in the experiment with dead beetles stopped their predatory attacks before taking the beetles into their mouths (Fig. 1, 2; Video 3, 5). Thus, my results do not provide strong evidence that the frogs could not ingest large beetles. Rather, the amount of chemicals on the beetle body surface may increase with beetle size (see below).

The rapid responses of the frog species *Pe. nigromaculatus* to bombardier beetles (Fig. 2) could be considered a reflex action of the frogs' tongues (cf. Kumai, 1981a, b; Hirakawa, 1989). Frogs are known to use their tongues as a chemical detector (Dean, 1980b; Kumai, 1981a, b; Barlow, 1998) as well as a prey-catching tool (Noel et al., 2017). For example, chemical or electrical stimulation of the tongue can generate reflex responses in *Pe. nigromaculatus* (Kusano & Sato, 1957; Kumai, 1981a, b; Suzuki & Nomura, 1985; Takeuchi, Satou & Ueda, 1986; Hirakawa, 1989). Because *Pe. nigromaculatus* is a generalist predator that can attack a variety of arthropods within its field of view (Hirai & Matsui, 1999; Honma, 2004; Honma, Oku & Nishida, 2006; Sano & Shinohara, 2011; Sarashina, Yoshihisa & Yoshida, 2011), *Pe. nigromaculatus* may have evolved specific responses to toxic prey to avoid being injured by trying to eat them. The results of this study suggest that the tongues of *Pe. nigromaculatus* may be able to rapidly

detect toxic substances or other characteristics on the body surface of the bombardier beetles, and the frogs subsequently avoid the beetles to prevent themselves from being bombed and injured. Previous studies have focused on how frogs and toads use their tongues to catch prey (Ewert, 1970; Nishikawa & Gans, 1996; Monroy & Nishikawa, 2010; Noel et al., 2017). Few reports have explored how frogs and toads use their tongues to detect toxins in/on potential prey (but see Dean, 1980b). Therefore, the tongue responses that I describe in *Pe. nigromaculatus* will likely be evident in other frogs such as the tree frog *Hyla japonica* (Günther) (Taniguchi et al., 2005; Ito, Taniguchi & Billen, 2016; Matsubara & Sugiura, 2017).

## CONCLUSIONS

In one study, the chemicals produced by bombardier beetles' bombing did not stimulate the tongues of toads any less intensely than did the heat from the chemical reaction (Dean, 1980b). Other than this study, the relative importance of the toxic chemicals and heat produced by bombing for the successful escape of bombardier beetles from predators has been largely unexplored. My results support the hypothesis that bombing is not essential when bombardier beetles defend themselves against frog attacks. Furthermore, my findings suggest that (cool) toxic chemicals on the beetles' bodies alone may cause frogs to desist from an attack; thus, chemicals on the body may serve as a primary defence and bombing as a secondary defence. Successful defence by chemicals on the body would reduce the costs associated with bombing, suggesting that beetles may have evolved to use chemicals on the body surface as their primary defence. However, further experiments are required to validate this hypothesis; for example, dead beetles with body surfaces cleaned of chemicals, or palatable prey coated with toxic chemicals, should be offered to frogs.

Many prey animals exhibit multiple anti-predator defences (Edmunds, 1974). Predation pressures imposed by different enemies may encourage prey to diversify defence strategies. Further studies are needed.

## ACKNOWLEDGEMENTS

I am grateful to K. Sakagami, K. Uchida, and A. Ushimaru for providing valuable information about the collection sites. I thank K. Sakagami, W. Higashikawa, and M. Ito for helping to collect the study animals, and S. Matsubara and Y. Maeda for helping to maintain them. T. Takanashi provided valuable advice on this research. M. Bulbert and two anonymous reviewers provided helpful comments on an earlier version of the manuscript.

## ADDITIONAL INFORMATION AND DECLARATIONS

### Data Availability

The following information was supplied regarding data availability:

Raw data is available from the Figshare Digital Repository (<https://figshare.com/s/734fc893bcaadfde3a90>).

### Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/>

## REFERENCES

**Aneshansley DT, Eisner T, Widom JM, Widom B. 1969.** Biochemistry at 100°C: explosive secretory discharge of bombardier beetles (*Brachinus*). *Science* **165**:61–63. DOI

10.1126/science.165.3888.61.

**Arndt EM, Moore W, Lee WK, Ortiz C. 2015.** Mechanistic origins of bombardier beetle (Brachinini) explosion-induced defensive spray pulsation. *Science* **348**:563–567. DOI 10.1126/science.1261166.

**Bonacci T, Aloise G, Brandmayr P, Brandmayr TZ, Capula M. 2008.** Testing the predatory behaviour of *Podarcis sicula* (Reptilia: Lacertidae) towards aposematic and non-aposematic preys. *Amphibia-Reptilia* **29**: 449–453. DOI 10.1163/156853808785111986.

**Barlow LA. 1998.** The biology of amphibian taste. In: Heatwole H, Dawley E, eds. *Amphibian biology, volume 3: sensory perception*. Chipping Norton, Australia: Surrey Beatty & Sons, 743–782.

**Brower JVZ, Brower LP. 1962.** Experimental studies of mimicry. 6. The reaction of toads (*Bufo terrestris*) to honeybees (*Apis mellifera*) and their dronefly mimics (*Eristalis vinetorum*). *The American Naturalist* **96**:297–307. DOI 10.1086/282237.

**Brower LP, Brower JVZ, Westcott PW. 1960.** Experimental studies of mimicry. 5. The reactions of toads (*Bufo terrestris*) to bumblebees (*Bombus americanorum*) and their robberfly mimics (*Mallophora bomboides*), with a discussion of aggressive mimicry. *The American Naturalist* **94**:343–355. DOI 10.1086/282137.

**Bura BL, Kawahara A, Yack J. 2016.** A comparative analysis of sonic defences in Bombycoidea caterpillars. *Scientific Reports* **6**:31469. DOI 10.1038/srep31469.

**Cooper SM, Owen-Smith N. 1986.** Effects of plant spinescence on large mammalian herbivores. *Oecologia* **68**:446–455. DOI 10.1007/BF01036753.

**Dean J. 1979.** Defensive reaction time of bombardier beetles: an investigation of the speed of a chemical defense. *Journal of Chemical Ecology* **5**:691–701. DOI 10.1007/BF00986554.

**Dean J. 1980a.** Encounters between bombardier beetles and two species of toads (*Bufo*



- 348       *americanus*, *B. marinus*): speed of prey-capture does not determine success. *Journal of*
- 349       *Comparative Physiology* **133**:41–50. DOI 10.1007/BF00660180.
- 350   **Dean J. 1980b.** Effects of thermal and chemical components of bombardier beetle chemical
- 351       defense: glossopharyngeal response in two species of toads (*Bufo americanus*, *B. marinus*).
- 352       *Journal of Comparative Physiology* **133**:51–59. DOI 10.1007/BF00660181.
- 353   **Derby CD. 2007.** Escape by inking and secreting: marine molluscs avoid predators through a
- 354       rich array of chemicals and mechanisms. *Biological Bulletin* **213**:274–289. DOI
- 355       10.2307/25066645.
- 356   **Edmunds M. 1974.** *Defense in animals*. Harlow: Longman.
- 357   **Eisner T. 1958.** The protective role of the spray mechanism of the bombardier beetle, *Brachynus*
- 358       *ballistarius* Lec. *Journal of Insect Physiology* **2**:215–220.
- 359   **Eisner T. 2003.** *For love of insects*. Cambridge: The Belknap Press of the Harvard University
- 360       Press.
- 361   **Eisner T, Aneshansley DJ. 1999.** Spray aiming in the bombardier beetle: Photographic
- 362       evidence. *Proceeding of the National Academy of Sciences USA* **96**:9705–9709. DOI
- 363       10.1073/pnas.96.17.9705.
- 364   **Eisner T, Dean J. 1976.** Ploy and counterploy in predator–prey interactions: Orb-weaving
- 365       spiders versus bombardier beetles. *Proceeding of the National Academy of Sciences USA*
- 366       **73**:1365–1367. DOI 10.1073/pnas.96.17.9705.
- 367   **Eisner T, Meinwald J. 1966.** Defensive secretions of arthropods. *Science* **153**:1341–1350. DOI
- 368       10.1126/science.153.3742.1341.
- 369   **Eisner T, Eisner M, Aneshansley DJ. 2005.** Pre-ingestive treatment of bombardier beetles by
- 370       jays: food preparation by “anting” and “sand-wiping”. *Chemoecology* **15**:227–233. DOI
- 371       10.1007/s00049-005-0316-6.

- 372 **Eisner T, Eisner M, Siegler M. 2005.** *Secret weapons: defenses of insects, spiders, scorpions,*  
373 *and other many-legged creatures.* Cambridge: The Belknap Press of the Harvard  
374 University Press.
- 375 **Eisner T, Aneshansley DJ, del Campo ML, Eisner M, Frank JH, Deyrup M. 2006.** Effect of  
376 bombardier beetle spray on a wolf spider: repellency and leg autotomy. *Chemoecology*  
377 **16:**185–189. DOI 10.1007/s00049-006-0346-8.
- 378 **Endler JA. 1991.** Interactions between predators and prey. In: Krebs JR, Davies NB, eds.  
379 *Behavioural ecology : an evolutionary approach.* London, Paris, Berlin, Vienna: Blackwell,  
380 169–196
- 381 **Ewert JP. 1970.** Neural mechanisms of prey-catching and avoidance behaviour in the toad (*Bufo*  
382 *bufo* L.). *Brain, Behavior and Evolution* **3:**36–56. DOI 10.1159/000125462.
- 383 **Fujisawa T, Lee CM, Ishii M. 2012.** Species diversity of ground beetle assemblages in the  
384 distinctive landscapes of the Yodo River flowing through northern Osaka Prefecture,  
385 central Japan. *Japanese Journal of Environmental Entomology and Zoology* **23:**89–100.  
386 DOI 10.11257/jjeez.23.89.
- 387 **Habu A, Sadanaga K. 1965.** Illustrations for identification of larvae of the Carabidae found in  
388 cultivated fields and paddy-fields (III). *Bulletin of the National Institute of Agricultural*  
389 *Sciences, Series C: Plant Pathology and Entomology* **19:**81–216 (in Japanese).
- 390 **Hirai T. 2002.** Ontogenetic change in the diet of the pond frog, *Rana nigromaculata*. *Ecological*  
391 *Research* **17:**639–644. DOI 10.1046/j.1440-1703.2002.00521.x.
- 392 **Hirai T, Matsui M. 1999.** Feeding habits of the pond frog, *Rana nigromaculata*, inhabiting rice  
393 fields in Kyoto, Japan. *Copeia* **1999:**940–947. DOI 10.2307/1447969.
- 394 **Hirakawa T. 1989.** A Study of averaged waves of reflex discharges in the frog hypoglossal  
395 nerve elicited by electrical stimulation of glossopharyngeal afferent. *Journal of the Kyushu*

396 *Dental Society* **43**:693–709. (in Japanese with English summary). DOI 10.2504/kds.43.693.

397 **Honma A. 2004.** MOMO VideoArchives: momo040902cj01b. [http://movspec.mus-](http://movspec.mus-nh.city.osaka.jp/ethol/showdetail.php?movieid=momo040902cj01b)

398 [nh.city.osaka.jp/ethol/showdetail.php?movieid=momo040902cj01b](http://movspec.mus-nh.city.osaka.jp/ethol/showdetail.php?movieid=momo040902cj01b). Accessed: 28 January

399 2018.

400 **Honma A, Oku S, Nishida T. 2006.** Adaptive significance of death feigning posture as a

401 specialized inducible defence against gape-limited predators. *Proceedings of the Royal*

402 *Society B: Biological Science* **273**:1631–1636. DOI 10.1098/rspb.2006.3501.

403 **Inbar M, Lev-Yadun S. 2005.** Conspicuous and aposematic spines in the animal kingdom.

404 *Naturwissenschaften* **92**:170–172. DOI 10.1007/s00114-005-0608-2.

405 **Ishitani M, Yano K. 1994.** Species composition and seasonal activities of ground beetles

406 (Coleoptera) in a fig orchard. *Japanese Journal of Entomology* **62**:201–210.

407 **Ito F, Taniguchi K, Billen J. 2016.** Defensive function of petiole spines in queens and workers

408 of the formicine ant *Polyrhachis lamellidens* (Hymenoptera: Formicidae) against an ant

409 predator, the Japanese tree frog *Hyla japonica*. *Asian Myrmecology* **8**:1–6. DOI

410 10.20362/am.008014.

411 **Jung JK, Kim ST, Lee SY, Park CK, Lee EH, Lee JH. 2012.** Ground beetle (Coleoptera:

412 Carabidae) assemblage in the urban landscape, Korea. *Journal of Ecology and Field Biology*

413 **35**:79–89. DOI 10.5141/JEFB.2012.012.

414 **Kanehisa K. 1996.** Secretion of defensive substance by Carabidae and Brachinidae. *Bulletin of*

415 *the Research Institute for Bioresources, Okayama University* **4**:9–23 (in Japanese with

416 English summary).

417 **Kanehisa K, Murase M. 1977.** Comparative study of the pygidial defensive systems of carabid

418 beetles. *Applied Entomology and Zoology* **12**:225–235. DOI 10.1303/aez.12.225.

419 **Komaki S, Igawa T, Lin SM, Tojo K, Min MS, Sumida M. 2015.** Robust molecular

phylogeny and palaeodistribution modelling resolve a complex evolutionary history:  
glacial cycling drove recurrent mtDNA introgression among *Pelophylax* frogs in East Asia.  
*Journal of Biogeography* **42**:2159–2171. DOI 10.1111/jbi.12584.

**Kumai T. 1981a.** Reflex response of the hypoglossal nerve induced by gustatory stimulation of  
the frog tongue. *Brain Research* **208**:432–435. DOI 10.1016/0006-8993(81)90572-2.

**Kumai T. 1981b.** Reflex response of the hypoglossal nerve induced by chemical stimulation of  
the tongue and electrical stimulation of the glossopharyngeal nerve in the frog. *Japanese  
Journal of Physiology* **31**:625–637. DOI 10.2170/jjphysiol.31.625.

**Kusano K, Sato M. 1957.** Properties of fungiform papillae in frog's tongue. *Japanese Journal of  
Physiology* **157**:324–338. DOI 10.2170/jjphysiol.7.324

**Larochelle A. 1974a.** The American toad as champion carabid beetle collector. *The Pan-Pacifc  
Entomologist* **50**:203–204.

**Larochelle A. 1974b.** Carabid beetles (Coleoptera: Carabidae) as prey of North American frogs.  
*The Great Lakes Entomologist* **7**:147–148.

**Lev-Yadun S. 2001.** Aposematic (warning) coloration associated with thorns in higher plants.  
*Journal of Theoretical Biology* **210**:385–388 DOI 10.1006/jtbi.2001.2315.

**Lev-Yadun S. 2009.** Aposematic (warning) coloration in plants. In: Baluska F, ed. *Plant-  
environment interactions. From sensory plant biology to active plant behavior*. Berlin:  
Springer-Verlag, 167–202 DOI 10.1007/978-3-540-89230-4\_10.

**Liu K, Wang F, Chen W, Tu L, Min MS, Bi K, Fu J. 2010.** Rampant historical mitochondrial  
genome introgression between two species of green pond frogs, *Pelophylax  
nigromaculatus* and *P. plancyi*. *BMC Evolutionary Biology* **10**:201. DOI 10.1186/1471-  
2148-10-201.

**Maeda N, Matsui M. 1999.** *Frogs and toads of Japan: revised edition*. Tokyo: Bun-ichi Sogo

Shuppan.

**Matsubara S, Sugiura S. 2017.** Chemical defence of turnip sawfly larvae against Japanese tree frogs. *Journal of Asia-Pacific Entomology* **20**:225–227. DOI 10.1016/j.aspen.2017.01.001.

**Matsubashi T, Okuyama F. 2015.** *Frogs and toads of Japan*. Tokyo, Japan: Yamakei (in Japanese).

**Monroy JA, Nishikawa K. 2010.** Prey capture in frogs: alternative strategies, biomechanical trade-offs, and hierarchical decision making. *Journal of Experimental Zoology* **315**:61–71. DOI 10.1002/jez.601

**Mori I. 2008.** Predation by introduced bullfrog *Rana catesbeiana* on a breeding male of *Rhacophorus schlegelii* and the other animals. *Bulletin of the Okayama Prefecture Nature Conservation Center* **16**:61–62 (in Japanese).

**Nishikawa KC, Gans C. 1996.** Mechanisms of tongue protraction and narial closure in the marine toad *Bufo marinus*. *Journal of Experimental Biology* **199**:2511–2529.

**Mithöfer A, Boland W. 2012.** Plant defense against herbivores: chemical aspects. *Annual Review of Plant Biology* **63**:431–450. DOI 10.1146/annurev-arplant-042110-103854.

**Noel AC, Guo HY, Mandica M, Hu DL. 2017.** Frogs use a viscoelastic tongue and non-Newtonian saliva to catch prey. *Journal of the Royal Society Interface* **14**: 20160764. DOI 10.1098/rsif.2016.0764.

**Ohwaki A, Kaneko Y, Ikeda H. 2015.** Seasonal variability in the response of ground beetles (Coleoptera: Carabidae) to a forest edge in a heterogeneous agricultural landscape in Japan. *European Journal of Entomology* **112**:135–144. DOI 10.14411/eje.2015.022.

**R Development Core Team. 2016.** *R, a language and environment for statistical computing*. Vienna: R Foundation for Statistical Computing.

**Ruxton GD, Sherratt TN, Speed MP. 2004.** *Avoiding attack: the evolutionary ecology of*

*crypsis, aposematism, and mimicry*. Oxford: Oxford University Press.

**Sano M, Shinohara M. 2011.** Species comparison of frogs food habits during mating seasons in Uenohara, Yamanashi Pref., Japan. *Bulletin of Teikyo University of Science and Technology* **8**:101–111 (in Japanese with English summary).

**Sarashina M, Yoshihisa Y, Yoshida T. 2011.** Stomach contents of invasive Black-spotted Pond frog (*Rana nigromaculata*) in urban landscape of Sapporo City. *Journal of Rakuno Gakuen University* **36**:81–86 (in Japanese with English summary).

**Skelhorn J, Rowe C. 2006.** Taste-rejection by predators and the evolution of unpalatability in prey. *Behavioral Ecology and Sociobiology* **60**:550–555. DOI 10.1007/s00265-006-0199-8.

**Sugiura S. 2016.** Bagworm bags as portable armour against invertebrate predators. *PeerJ* **4**:e1686 DOI 10.7717/peerj.1686.

**Sugiura S, Sato T. 2018.** Successful escape of bombardier beetles from predator digestive systems. *Biology Letters* **14**:20170647. DOI 10.1098/rsbl.2017.0647.

**Sugiura S, Takanashi T. 2018.** Hornworm counterattacks: Defensive strikes and sound production in response to invertebrate attackers. *Biological Journal of the Linnean Society*. DOI 10.1093/biolinnean/blx156.

**Sugiura S, Yamazaki K. 2014.** Caterpillar hair as a physical barrier against invertebrate predators. *Behavioral Ecology* **25**:975–983. DOI 10.1093/beheco/aru080.

**Suzuki H, Nomura H. 1985.** Input-output relation in gustatory linguo-hypoglossal reflex in the frog. *Matsumoto Shigaku: Journal of the Matsumoto Dental College Society* **11**:13–17. (in Japanese with English summary).

**Takeuchi H, Satou M, Ueda K. 1986.** EMG analysis of head muscle during naturally-occurring and electrically- evoked snapping, rejecting and avoiding behavior in the Japanese toad. *Zoological Science* **3**:992.

- 492 **Taniguchi K, Maruyama M, Ichikawa T, Ito F. 2005.** A case of Batesian mimicry between  
493 myrmecophilous staphylinid beetle, *Pella comes*, and its host ant, *Lasius* (*Dendrolasius*)  
494 *spathepus*: an experiment using the Japanese tree frog *Hyla japonica* as a real predator.  
495 *Insectes Sociaux* **52**:320–322. DOI 10.1007/s00040-005-0813-1.
- 496 **Tsuji M, Ushimaru A, Osawa T, Mitsuhashi H. 2011.** Paddy-associated frog declines via  
497 urbanization: A test of the dispersal-dependent-decline hypothesis. *Landscape and Urban*  
498 *Planning* **103**:318–325. DOI 10.1016/j.landurbplan.2011.08.005.
- 499 **Ueno S, Kurosawa Y, Sato M (eds). 1985.** *The Coleoptera of Japan in color vol. II.* Osaka,  
500 Japan: Hoikusha (in Japanese).
- 501 **Williams BL, Hanifin CT, Brodie ED Jr, Brodie ED III. 2010.** Tetrodotoxin affects survival  
502 probability of rough-skinned newts (*Taricha granulosa*) faced with TTX-resistant garter  
503 snake predators (*Thamnophis sirtalis*). *Chemoecology* **20**:285–290. DOI 10.1007/s00049-  
504 010-0057-z.
- 505 **Yahiro K, Fujimoto T, Tokuda M, Yano K. 1992.** Species composition and seasonal  
506 abundance of ground beetles (Coleoptera) in paddy fields. *Japanese Journal of Entomology*  
507 **60**:805–813.

# Figure legend

Figure 1 Behavioural responses of the black-spotted pond frog *Pelophylax nigromaculatus* to live and dead adult individuals of the bombardier beetle *Pheropsophus jessoensis*. ‘Stop attack’: the frogs stopped their attacks after their tongues touched the beetles. ‘Spit out’: the frogs spat out the beetles immediately after taking the beetles into their mouths. ‘Swallow’: the frogs successfully swallowed the beetles. ‘Bombing’: the beetles could be heard bombing. Photo credit: Shinji Sugiura.

Figure 2 Temporal sequence of the frog *Pelophylax nigromaculatus* rejecting a live adult *Pheropsophus jessoensis* without taking the beetle into its mouth. The frog stopped the attack immediately after its tongue touched the beetle. No bombing sounds were heard (see Video S3). Credit: Shinji Sugiura.

Figure 3 Temporal sequence of the frog *Pelophylax nigromaculatus* spitting out a live adult *Pheropsophus jessoensis* after taking the beetle into its mouth. Bombing by the beetle was audible just before the frog spat out the beetle (E–F: 1675–1800 ms; see Video S4). Credit: Shinji Sugiura.

Figure 4. Body size relationships between predator frogs (*Pelophylax nigromaculatus*) and prey beetles (*Pheropsophus jessoensis*). (A) Success or failure of swallowing by frogs. (B) Success or failure of digestion by frogs. Closed circles and triangles indicate the swallow (or digestion) success of live and dead beetles, respectively. Open circles and triangles indicate the swallow (or digestion) failure of live and dead beetles, respectively. Photo credit: Shinji Sugiura.



533 **Supplemental Information**

534

535 Video S1 An adult *Pheropsophus jessoensis* bombing. The beetle discharged toxic chemicals  
536 when the body was stimulated with a pair of forceps. Credit: Shinji Sugiura.

537

538 Video S2. The frog *Pelophylax nigromaculatus* preying on a carabid beetle. The frog opened  
539 its mouth and shot out its tongue to capture prey. Credit: Shinji Sugiura.

540

541 Video S3 The frog *Pelophylax nigromaculatus* rejecting a live *Pheropsophus jessoensis*. The  
542 frog stopped the attack immediately after its tongue touched the beetle. Credit: Shinji Sugiura.

543

544 Video S4 The frog *Pelophylax nigromaculatus* spitting out a live *Pheropsophus jessoensis*. The  
545 frog took the beetle into its mouth but immediately spat out the beetle. Credit: Shinji Sugiura.

546

547 Video S5 The frog *Pelophylax nigromaculatus* rejecting a dead *Pheropsophus jessoensis*. The  
548 frog stopped its attack immediately after its tongue touched the dead beetle. Credit: Shinji

549 Sugiura.

# **Table 1**(on next page)

Table 1. Results of a generalised linear model (GLM) testing potential factors influencing whether the frog *Pelophylax nigromaculatus* successfully swallowed the bombardier beetle *Pheropsophus jessoensis* in feeding experiments.

1

Table 1. Results of a generalised linear model (GLM) testing potential factors influencing whether the frog *Pelophylax nigromaculatus* successfully swallowed the bombardier beetle *Pheropsophus jessoensis* in feeding experiments.

Response variable	Explanatory variable (fixed effect)	Coefficient estimate	SE	<i>t</i> value	<i>P</i> value
Swallowing success <sup>1)</sup>	Intercept	7.601881	4.454207	1.707	0.094
	Frog weight	−0.420817	0.290774	−1.447	0.154
	Beetle weight	−0.060781	0.025084	−2.423	0.019
	Frog weight × Beetle weight	0.002637	0.001457	1.81	0.076 <sup>3)</sup>
	Beetle treatment <sup>2)</sup>	1.387536	0.855616	1.622	0.111

2 <sup>1)</sup> As the residual deviance was 1.5-fold smaller than the residual degrees of freedom, a quasi-binomial error distribution (rather than a  
3 binomial error distribution) was employed. Two and four live and dead beetles, respectively, were swallowed.

4 <sup>2)</sup> Live beetles were used as a reference.

5 <sup>3)</sup> The significance of this factor was checked using the likelihood ratio test ( $P = 0.063$ ).

6

# Table 2 (on next page)

Table 2. Results of a generalised linear model (GLM) testing potential factors influencing whether the frog *Pelophylax nigromaculatus* successfully digested the bombardier beetle *Pheropsophus jessoensis* in feeding experiments.

1

Table 2. Results of a generalised linear model (GLM) testing potential factors influencing whether the frog *Pelophylax nigromaculatus* successfully digested the bombardier beetle *Pheropsophus jessoensis* in feeding experiments.

Response variable	Explanatory variable (fixed effect)	Coefficient estimate	SE	<i>t</i> value	<i>P</i> value
Digestion success <sup>1)</sup>	Intercept	4.471889	4.45551	1.004	0.32
	Frog weight	−0.370908	0.30261	−1.226	0.226
	Beetle weight	−0.04681	0.023848	−1.963	0.055 <sup>3)</sup>
	Frog weight × Beetle weight	0.002329	0.00146	1.596	0.117
	Beetle treatment <sup>2)</sup>	2.011385	1.033049	1.947	0.057 <sup>4)</sup>

2 <sup>1)</sup> As the residual deviance was 1.5-fold smaller than the residual degrees of freedom, a quasi-binomial error distribution (rather than a  
3 binomial error distribution) was employed. One and four live and dead beetles, respectively, were digested.

4 <sup>2)</sup> Live beetles were used as a reference.

5 <sup>3)</sup> The significance of this factor was checked using the likelihood ratio test ( $P = 0.028$ ).


6 <sup>4)</sup> The significance of this factor was checked using the likelihood ratio test ( $P = 0.029$ ).

7

# Figure 1

Behavioural responses of the black-spotted pond frog *Pelophylax nigromaculatus* to live and dead adult individuals of the bombardier beetle *Pheropsophus jessoensis*.


‘Stop attack’: the frogs stopped their attacks after their tongues touched the beetles. ‘Spit out’: the frogs spat out the beetles immediately after taking the beetles into their mouths. ‘Swallow’: the frogs successfully swallowed the beetles. ‘Bombing’: the beetles could be heard bombing. Photo credit: Shinji Sugiura.



# Figure 2

Temporal sequence of the frog *Pelophylax nigromaculatus* rejecting a live adult *Pheropsophus jessoensis* without taking the beetle into its mouth.

The frog stopped the attack immediately after its tongue touched the beetle. No bombing sounds were heard (see Video S3). Credit: Shinji Sugiura.






# Figure 3

Temporal sequence of the frog *Pelophylax nigromaculatus* spitting out a live adult *Pheropsophus jessoensis* after taking the beetle into its mouth.

Bombing by the beetle was audible just before the frog spat out the beetle (1675–1800 ms; see Video S4). Credit: Shinji Sugiura.




# Figure 4

Body size relationships between predator frogs (*Pelophylax nigromaculatus*) and prey beetles (*Pheropsophus jessoensis*).

(A) Success or failure of swallowing by frogs. (B) Success or failure of digestion by frogs. Closed circles and triangles indicate the swallow (or digestion) success of live and dead beetles, respectively. Open circles and triangles indicate the swallow (or digestion) failure of live and dead beetles, respectively. Photo credit: Shinji Sugiura.

A) Success or failure of swallowing by frogs



B) Success or failure of digestion by frogs

