

A needle in a haystack: Mesozoic origin of parasitism in Strepsiptera revealed by first definite Cretaceous primary larva (Insecta)

Hans Pohl ^{Corresp., 1}, Jan Batelka ², Jakub Prokop ², Patrick Müller ³, Margarita I Yavorskaya ¹, Rolf G Beutel ¹

¹ Institut für Zoologie und Evolutionsforschung, Spezielle Zoologie und Entomologie, Friedrich-Schiller Universität Jena, Jena, Germany

² Department of Zoology, Faculty of Science, Charles University Prague, Praha, Czech Republic

³ Unaffiliated, Kāshofen, Germany

Corresponding Author: Hans Pohl
Email address: hans.pohl@uni-jena.de

Twisted winged insects (Strepsiptera) are a highly specialized small order of parasitic insects. Whether parasitism developed at an early or late stage in the evolution of the group was unknown. Here we record and describe the first *definite* Mesozoic strepsipteran primary larva embedded in Burmese amber (~99 million years ago) extending the parasitic life style back by ~50 million years, revealing a Mesozoic origin of parasitism in the group. The extremely small first instar displays all diagnostic characters of strepsipteran immatures of this stage and is nearly identical with those of Mengenillidae, one of the most “ancestral” extant strepsipteran taxa. This demonstrates a remarkable evolutionary stasis over ~100 million years. **The new finding strongly weakens the case of small larvae embedded in Cretaceous amber interpreted as strepsipteran immatures.** They differ in many structural features from extant strepsipteran primary larvae and are very likely parasitic beetle larvae.

1 A needle in a haystack: Mesozoic origin of parasitism in Strepsiptera revealed by first
2 definite Cretaceous primary larva (Insecta)

3 Hans Pohl¹, Jan Batelka², Jakub Prokop², Patrick Müller³, Margarita I. Yavorskaya¹, Rolf G.
 4 Beutel¹

5

6 ¹ Institut für Zoologie und Evolutionsforschung, Friedrich-Schiller-Universität Jena, Germany

7 ² Department of Zoology, Faculty of Science, Charles University Prague, Praha, Czech Republic

8 ³ Käßhofen, Germany

9

10 Corresponding author

11 Hans Pohl

12 Email address: hans.pohl@uni-jena.de

13

Abstract

Twisted winged insects (Strepsiptera) are a highly specialized small order of parasitic insects. Whether parasitism developed at an early or late stage in the evolution of the group was unknown. Here we record and describe the first *definite* Mesozoic strepsipteran primary larva embedded in Burmese amber (~99 million years ago) extending the parasitic life style back by ~50 million years, revealing a Mesozoic origin of parasitism in the group. The extremely small first instar displays all diagnostic characters of strepsipteran immatures of this stage and is nearly identical with those of Mengenillidae, one of the most “ancestral” extant strepsipteran taxa. This demonstrates a remarkable evolutionary stasis over ~100 million years. The new finding strongly weakens the case of small larvae embedded in Cretaceous amber interpreted as strepsipteran immatures. They differ in many structural features from extant strepsipteran primary larvae and are very likely parasitic beetle larvae.

Introduction

Strepsiptera is a highly specialized small order of holometabolous insects (Pohl & Beutel, 2008; 2013). The phylogenetic placement of the group, one of the longest controversies in systematic entomology, was only recently clarified (Niehuis et al., 2012; Peters et al., 2014; Misof et al., 2014; Kjer et al., 2016). The oldest recorded fossils are known from Cretaceous Burmese amber (Grimaldi, Kathirithamby & Schawaroch, 2005; Pohl & Beutel, 2016; Engel et al., 2016), indicating a minimum age of ca. 100 mya. An age of origin of ca. 120 mya was estimated based on molecular data (McMahon, Hayward & Kathirithamby, 2011). However, the confirmed sister group relationship with Coleoptera (Misof et al., 2014) implies that the group originated already in the early Permian or late Carboniferous (McKenna et al., 2015; Toussaint et al., 2017).

Extant strepsipterans are characterized by a conspicuous sexual dimorphism. The winged males are free living but extremely short-lived. With well-developed sensory organs and a highly efficient flight apparatus they are able to find the females within their very short life span of only few hours. Females are wingless and morphologically strongly simplified. They develop as endoparasites of other insects. Only the females of the basal extant Mengenillidae (and probably Bahiixenidae) leave their host and are able to move actively. The females of the majority of the species (Stylopodia) are permanently endoparasitic and only penetrate the host’s abdominal intersegmental membranes with the sclerotized anterior part of their body (Pohl & Beutel,

2008). The primary larvae of Strepsiptera are among the smallest known metazoans with an average size of ca. 230 μm , comparable to unicellular ciliates of the genus *Paramecium* (Pohl, 2002). This size reduction enables the female to produce a huge number of offspring and the minute primary larvae are able to penetrate relatively small insect hosts. The body of the primary larvae is elongated oval to nearly drop shaped. The head is semicircular with well-developed stemmata with cornea lenses. Antennae and labrum are missing. Sternal plates inserted between the coxae with spinulae and fringes of microtrichia on the posterior border of the abdominal sternites produce capillary forces enabling the larvae to stick to and crawl along wet surfaces (Pohl & Beutel, 2008). A unique apomorphic feature within Holometabola is the presence of one pair of long and strongly developed bristle-like cerci. They are inserted on the terminal abdominal segment XI, with strong muscles attached to their base. They enable the minute larvae to jump very efficiently.

The relatively sparse Cretaceous fossil record of adult strepsipteran males reveals that they were already a specialized group of insects in the late Mesozoic. It is very likely that the highly modified antennae with flabellate appendages and numerous specialised dome-shaped chemoreceptors were used for finding females over a relatively large distance as in extant species of the group. Likewise, the uptake of food was apparently very limited at least, as suggested by the strongly simplified mouthparts. Strepsipteran females from the Mesozoic have not yet been discovered yet. Considering the small size of the males (ca. 1.5–3 mm) it was assumed, that they were already endoparasites of other insects (Pohl & Beutel, 2008). However, there was no direct evidence for an endoparasitic life style of the immature stages of Mesozoic strepsipterans. Endoparasitism was only documented since the Eocene by the appearance of Stylopodia, the strepsipteran subgroup with permanently endoparasitic females (Kinzelbach & Pohl, 1994; Henderickx et al., 2013).

Confirmed fossil records of strepsipteran primary larvae are from the Eocene and Miocene, respectively. The oldest fossil from Eocene brown coal (Geisel Valley, Germany) was initially described as a first instar of a scale insect (Coccoidea) (Haupt, 1950). It was later assigned to the extant strepsipteran genus *Stichotrema* (Myrmecolacidae) (Kinzelbach & Lutz, 1985). Finally, it was placed as *Stylopodia incertae sedis* based on a re-examination and a cladistic analysis of characters of primary larvae of all extant families of Strepsiptera (Pohl, 2009). Strepsipteran primary larvae associated with its parent female and its host (Auchenorrhyncha: Delphacidae)

are reported from Dominican amber (15–20 mya (Iturralde-Vinent & MacPhee, 1996; Poinar, 2004)). Other very small “triungulin” larvae assigned to Strepsiptera were described from the Late Cretaceous (Campanian) amber of Manitoba, Canada (Grimaldi, Kathirithamby & Schawaroch, 2005), and a “planidium” from the Upper Cretaceous (Santonian) amber of the Taimyr Peninsula, Siberia (Kathirithamby et al., 2017). The interpretation of the immatures treated in the earlier study was discussed critically, pointing out an entire series of features in conflict with an assignment to Strepsiptera (Beutel et al., 2016). The “planidium” from the Upper Cretaceous amber (Kathirithamby et al., 2017) is most likely a parasitic beetle larva (Batelka et al., accepted) (s.b.).

Kathirithamby et al. (2017) used the term “planidium” for primary larvae of Strepsiptera. However, this is only appropriate for legless larvae of parasitic Diptera or Hymenoptera (Askew, 1971; Stehr, 1991). Triungulinid was introduced by Pierce (1909) for first instars of Strepsiptera based on their similarity with primary larvae of Meloidae or Ripiphoridae (Coleoptera), which were addressed as triungulins. However, as the structural affinities are only superficial and obviously non-homologous, and the first instars differ in important features (e.g. nine abdominal segments in Meloidae and Ripiphoridae versus 11 in primary larvae of Strepsiptera), we prefer the neutral term primary larva for first instars of Strepsiptera. Claws forming a trident with spatulate setae occur in phoretic primary larvae of some genera of Meloidae (Bologna, Turco & Pinto, 2010), but are completely lacking in strepsipteran larvae.

In the present study we describe a minute larva from a piece of Burmese amber, with a habitus and a set of observable features unambiguously confirming a placement in Strepsiptera. The ordinal assignment among Strepsiptera is discussed. Based on the described features the position of other putative strepsipteran larvae (Grimaldi, Kathirithamby & Schawaroch, 2005; Kathirithamby et al., 2017) is critically re-evaluated. The new fossil larva clearly confirms that Mesozoic Strepsiptera were already endoparasites of other insects. The minimum age of endoparasitism is extended back by ca. 50 million years.

102

103 **Material & Methods**

104 **Material**

The piece of Burmite with the strepsipteran larva came from deposits in the Hukawng Valley of Myanmar. The age is estimated as ca. 99 Ma (earliest Cenomanian (Shi et al., 2012)).

The amber with the strepsipteran first instar is presently on long-term loan at the Institute of Zoology, Chinese Academy of Sciences (hereinafter coded as IZAS) (Box 92, Beichen West Road, Chaoyang District, Beijing, 100101, P. R. China) and will be deposited in the Three Gorges Entomological Museum, Chongqing, China after 2027. It is from a mining locality at Noiye Bum (near Tanai Village, 26°21'33.41''N, 96°43'11.88''E) (Cruickshank & Ko, 2003; Grimaldi, Engel & Nascimbene, 2009). The size of the piece of amber is 26 x 22 x 10 mm. Syninclusions are listed in the following: Acari (9 specimens), orthopteran nymph (1), Psocodea (1), Sternorrhyncha (1), apocritan Hymenoptera (2), Berothidae (Neuroptera) (1), Elateridae (1), primary longipedes larvae of Ripiphoridae (46), Polyphaga with unclear affinity (2), "nematoceran" species of Diptera with unclear affinity (2), brachyceran species of Diptera with unclear affinity (2).

As important syninclusions are embedded very close to the strepsipteran larva, it was not possible to isolate the larva and trim the amber piece into a thin plate and mount it on a glass microscope slide. This precludes examining the first instar with oil-immersion lenses and phase or differential interference contrast at a magnification of 1000 x. Therefore, some structures of the minute larva could not be evaluated.

Specimen imaging

The piece of amber was temporarily mounted on coverslips using glycerine. The specimen was observed under two different microscopes: An Axio Zoom.V16 with a PlanNeoFluar Z 1.0x (Carl Zeiss Microscopy GmbH) was used for the overview images and the images were saved as CZI files. For observations and for measurements ZEN 2.3 lite (blue edition) (Carl Zeiss Microscopy GmbH) was used. For higher magnifications, an Olympus IX81 inverted fluorescence microscope with UIS2 objectives, equipped with an ORCA-AG monochromatic 12-bit CCD camera (Hamamatsu) was used. The mirror images were superimposed with Cell^R software (Olympus Soft Imaging Solutions). Sets of photographs were analyzed with Fiji (Schindelin et al., 2012).

Single images were exported with ZEN 2.3 lite or Fiji respectively. Some images were combined with Zerene Stacker (Zerene Systems LLC, Richland, USA). The photographs were processed using Adobe Photoshop® CS6 (Adobe System Incorporated, San Jose, USA) and arranged as plates. Adobe Illustrator® CS6 (Adobe Systems Incorporated, San Jose, USA) was

used for the lettering of the plates. Image stacks of the Olympus IX81 microscope were used for the drawings and description.

Comparative taxonomy and terminology

Data on the morphology of Strepsiptera larvae are taken from (Pohl, 2000; 2002) and the morphological nomenclature used in these studies is applied. Additionally, primary larvae of *Eoxenos laboulbenei* Peyerimhoff, 1919 and *Mengenilla chobauti* Hofeneder, 1910 (both Mengenillidae) were examined (research collection of H.P. at Phyletisches Museum). For comparison primary larvae of *E. laboulbenei* embedded in Canada balsam on glass microscope slides were examined with the same microscope (Olympus IX81) as the fossil.

Results

Preservation

Strongly depressed dorsoventrally, dorsal and ventral side thus difficult to distinguish, especially in abdominal region. Part of left side of head and thorax covered by debris (Fig. 1A).

Morphology

First instar extremely small, total length excluding terminal bristles 197 µm. Wings or wing buds missing. Head semicircular, with recognizable stemmata but lacking antennae. Sides of body subparallel, slightly convex. Dorsum smooth, without recognizable surface structures and largely devoid of setae. Terminal abdominal segment XI with two pairs of strongly developed bristles. Specimen preserved in ventral position

Exposed part of head capsule semicircular in ventral view, with evenly rounded anterior margin and greatest width at hind margin. Visible part distinctly shorter than maximum width. Six large individual stemmata with cornea lenses recognizable posterolaterally on left side in strongly pigmented area, only three visible on right side (Figs. 1A, B, 3A). Labrum not present as separate element. Evenly rounded anterior margin of head capsule apparently forming sharp edge, lacking median emargination (Figs. 1A, B, 3A, B). Antennae not recognizable as prominent structures, largely reduced. Antennal field likely represented by small circular structure adjacent to anteriormost stemma, visible through translucent cuticle (af in Figs. 1B, 3A). Mandibles not visible. Ventral side of head medially covered by strongly modified maxillae

and labium. Maxillae medially fused, forming slightly curved, transverse plate-like structure (Figs. 1B, 3B). Maxillary palps recognizable as circular spots on posterior maxillary margin (mp in Figs. 1B, 3B). Details of labium not visible. Paired anterior tentorial arms visible in posterolateral cephalic region (te in Figs. 1B, 3B). Ecdysial sutures not visible. One seta recognizable on dorsal surface (fs in Fig. 1 B). Cuticle without recognizable surface modifications.

Pro-, meso- and metathorax subequal in length on ventral side. Only inflected lateral margin of nota visible (Fig. 1). Thoracic segments continuously widening from anterior to posterior. Prothorax with one seta on lateral pronotal margin, visible on right side. Conspicuous sternal plate inserted between coxae (prsp in Figs. 1B, 3A). Anterior part of sternal plate broad, posterior part with triangular apex. Spinulae on posterior margin not visible. Meso- and metathorax very similar to prothorax, but sternal plate only visible on the former. Legs only partly visible, short, composed of large, transverse coxa, trochanterofemur, tibia, and one-segmented pad-like tarsus (Fig. 1B). Tarsus without claws. Hind leg with long seta on mesal margin of coxa.

Abdomen composed of eleven segments. Segment II broadest, following segments slightly tapering posteriorly (Fig. 1). Only lateral margins of tergites visible, with two lateral setae on most of them (Figs. 1B, 4B). Sternites I–IX half as long as thoracic segments. One short seta visible on right side of sternite III, inserted close to hind margin. Sternites III–VI with two setae inserted laterally close to hind margin (Fig. 1B). Very fine parallel longitudinal lines on margins of sternites I–VIII may represent spinulae, or alternatively tergal furrows visible through cuticle (Fig. 4A). Tergites IX and X fused, about as long as segments VI–VIII combined, forming large plate-like structure covering terminal segment XI (Fig. 2A). Hind margin of segment XI truncated, with pair of very strongly developed bristles inserted close to midline, and second similar but shorter pair more laterally (Figs. 1B, 2B, 4B).

Diagnosis

Differs from primary larvae of *Eoxenos laboulbenei* by the presence of six stemmata and from primary larvae of *Mengenilla* (both Mengenillidae) and all other known primary larvae of Strepsiptera by the strongly developed second pair of bristles on hind margin of segment XI.

199

200 Discussion

201 The larva can be easily and unambiguously assigned to Strepsiptera, based on several diagnostic
 202 features and also apomorphic character states shared with other first instars of the order. Like
 203 extant strepsipteran primary larvae (Pohl, 2000) the fossil has an elongated oval to drop shaped
 204 body, with a semicircular head with large stemmata with cornea lenses, but lacking a separate
 205 labrum and also antennae as visible prominent structures. One pair of long and strongly
 206 developed bristles inserted on the terminal abdominal segment XI is present (Figs. 1, 5A, B).
 207 Within Strepsiptera, the larva shows close structural affinities with first instars of *Eoxenos*
 208 *laboulbenei*, especially due to the identical equipment with bristles on the last abdominal
 209 segments (Figs 2, 5). A placement close to the root of Strepsiptera *s.l.* or *s.str.* is likely.
 210 However, as only the male adults of extant *Bahiaxenos* and the extinct †*Mengea*, †*Protoxenos*
 211 (both Eocene Baltic amber), †*Cretostylops*, †*Kinzelbachilla*, and †*Phthanoxenos*, (Cretaceous
 212 Burmese amber) are known (Grimaldi, Kathirithamby & Schawaroch, 2005; Pohl, Beutel &
 213 Kinzelbach, 2005; Bravo et al., 2009; Pohl & Beutel, 2016; Engel et al., 2016), a precise
 214 phylogenetic assessment is not possible. It is conceivable that the larva belongs to one of the
 215 three strepsipteran species known from the same fossil site, but a verification is not possible with
 216 the information at hand.

217 A major point demonstrated here is that Cretaceous primary larva of Strepsiptera differ
 218 only in minimal details from extant immatures of basal genera of the order (Mengenillidae). This
 219 documents a high evolutionary stability over ca. 100 million years and clearly suggests that these
 220 extremely miniaturized larvae were already parasitic and produced in high numbers.

221 Large stemmata for identification of a host and the abdominal jumping apparatus are
 222 features clearly linked with parasitism, and the extremely small size suggests a very high number
 223 of offspring like in extant groups (Pohl & Beutel, 2008).

224 The finding of a Cretaceous primary larva nearly identical with those of the extant genus
 225 *Eoxenos* sheds new light on recently described “planidia” assigned to Strepsiptera
 226 (Kathirithamby et al., 2017). Considering the clearly demonstrated evolutionary stability of the
 227 tiny first instars, it appears highly unlikely that at the same time aberrant and unusually large
 228 strepsipteran primary larvae occurred. The “planidia” described by Kathirithamby et al. (2017)
 229 (Kathirithamby et al., 2017) differ in many features from strepsipteran larvae, such as for

instance much larger size (ca. 0.5 mm), anteriorly conical head, absence of large stemmata (“presumably two stemmata”), presence of a membranous cervix, posteriorly widening meso- and metanota, absence of sternal plates, strongly widened femora, lack of setae and spines on the abdominal sternites, only ten abdominal segments, lacking terminal bristles (on segment XI in Strepsiptera), and consequently the lack of a jumping apparatus. It is much more likely that these were simply miniaturized parasitic beetle larvae (Beutel et al., 2016), although their family placement is still open to debate (Batelka et al., accepted).

Considering the large number of primary larvae produced by strepsipteran females of Mengenillidae (ca. 2,500 by a 5 mm large female of *E. laboulbenei* (Silvestri, 1941)), it is surprising at first glance that not more strepsipteran primary larvae have been found in amber. The larva we describe here was only found by chance, like the primary larva from Eocene brown coal (Voigt, 1938). Due to their extremely small size, the larvae are only visible with high magnification (100 x) and can be overlooked very easily. In contrast to primary longipedes larvae of Ripiphoridae (Batelka et al., accepted), extant primary larvae of Strepsiptera show no aggregation behaviour (Pohl, unpublished results). After hatching from the female, the larvae quickly disperse in search of a suitable host. Considering their very small size, their radius of activity is probably limited to a few meters.

Conclusions

Detailed investigations of Burmese amber revealed the first definitive strepsipteran primary larva from the Cretaceous. Diagnostic features are the size of less than 200 µm, an elongated oval body, a semicircular head, and stemmata with cornea lenses in a strongly pigmented area. Apomorphic character states shared with other primary larvae of Strepsiptera are the lack of a separate labrum and prominent antennae, the medially fused maxillae, sternal plates, and the presence of a pair of long and strongly developed bristles inserted on the terminal abdominal segment XI. An evolutionary stability over ca. 100 million years is revealed as the Cretaceous primary larva of Strepsiptera differs only in minimal details from extant Mengenillidae, the sistergroup of the vast majority of the Strepsiptera (ca. 97 % of the species). The recently described “planidium” assigned to Strepsiptera by Kathirithamby et al. (2017) differs in many characters from fossil and extant primary larvae of Strepsiptera and is very likely a parasitic beetle larva.

261

262 Acknowledgments

263 We are grateful to Mr. Weiwei Zhang for providing pieces of amber used in the context of
 264 another publication, and to Assoc. Prof. Ming Bai for arranging this loan. The authors thank
 265 Ondřej Šebesta (Charles University) for his help with examination setup on the Olympus IX81
 266 and Birgit Perner (Leibniz-Institut für Alternsforschung – Fritz-Lippmann-Institut, Jena) for
 267 providing us access to the Axio Zoom.V16. We also thank Benjamin Naumann (Friedrich-
 268 Schiller-Universität Jena) for his help with the Axio Zoom.V16.

269

270 References

- 271 Askew RR 1971. *Parasitic insects*. London: Heinemann Educational Books Ltd.
- 272 Batelka J, Prokop J, Pohl H, Bai M, Zhang W, Beutel RG accepted. Highly specialized
 273 Cretaceous beetle parasitoids (Rhipiphoridae) identified with optimized visualization of
 274 microstructures. *Systematic Entomology*.
- 275 Beutel RG, Zhang WW, Pohl H, Wappler T, Bai M 2016. A miniaturized beetle larva in
 276 Cretaceous Burmese amber: reinterpretation of a fossil “strepsipteran triungulin.” *Insect*
 277 *Systematics & Evolution* 47:83–91.
- 278 Bologna MA, Turco F, Pinto JD 2010. 11.19. Meloidae Gyllenhal, 1810. In: Leschen RAB,
 279 Beutel RG, Lawrence JF eds. *Handbook of Zoology*. New York, 681–693.
- 280 Bravo F, Pohl H, Silva-Neto A, Beutel R 2009. Bahiixenidae, a “living fossil” and a new family
 281 of Strepsiptera (Hexapoda) discovered in Brazil. *Cladistics* 25:614–623.
- 282 Cruickshank RD, Ko K 2003. Geology of an amber locality in the Hukawng Valley, Northern
 283 Myanmar. *Journal of Asian Earth Sciences* 21:441–455.
- 284 Engel MS, Huang D, Breitkreuz LCV, Azar D, Cai C, Alvarado M 2016. A new twisted-wing
 285 parasitoid from mid-Cretaceous amber of Myanmar (Strepsiptera). *Cretaceous Research*
 286 58:160–167.
- 287 Grimaldi DA, Engel MS, Nascimbene PC 2009. Fossiliferous Cretaceous amber from Myanmar
 288 (Burma): Its rediscovery, biotic diversity, and paleontological significance. *American*
 289 *Museum Novitates* 3361:1–71.
- 290 Grimaldi D, Kathirithamby J, Schawaroch V 2005. Strepsiptera and triungula in Cretaceous
 291 amber. *Insect Systematics & Evolution* 36:1–20.

- 292 Haupt H 1950. Die Käfer (Coleoptera) aus der eozänen Braunkohle des Geiseltals. *Geologica.*
293 *Schriftenreihe der geologischen Institute der Universitäten Berlin, Greifswald, Halle,*
294 *Rostock* 6:VII–168.
- 295 Henderickx H, Bosselaers J, Pauwels E, van Hoorebeke L, Boone M 2013. X-ray micro-CT
296 reconstruction reveals eight antennomeres in a new fossil taxon that constitutes a sister clade
297 to *Dundoxenos* and *Triozocera* (Strepsiptera: Corioxenidae). *Palaeontologia Electronica*
298 16:1–16.
- 299 Iturralde-Vinent MA, MacPhee RDE 1996. Age and paleogeographical origin of Dominican
300 amber. *Science* 273:1850–1852.
- 301 Kathirithamby J, Perkovsky EE, Falin ZH, Engel MS 2017. A putative twisted-wing parasitoid
302 planidium (Insecta: Strepsiptera) in Taimyr Upper Cretaceous amber. *Cretaceous Research*
303 69:106–112.
- 304 Kinzelbach R, Lutz H 1985. Stylopoid larva from the Eocene – a spotlight on the phylogeny of the
305 stylopids (Strepsiptera). *Annals of the Entomological Society of America* 78:600–602.
- 306 Kinzelbach R, Pohl H 1994. The fossil Strepsiptera (Insecta: Strepsiptera). *Annals of the*
307 *Entomological Society of America* 87:59–70.
- 308 Kjer KM, Simon C, Yavorskaya M, Beutel RG 2016. Progress, pitfalls and parallel universes: a
309 history of insect phylogenetics. *Journal of the Royal Society Interface* 13:20160363.
- 310 McKenna DD, Wild AL, Kanda K, Bellamy CL, Beutel RG, Caterino MS, Farnum CW, Hawks
311 DC, Ivie MA, Jameson ML, Leschen RAB, Marvaldi AE, McHugh JV, Newton AF,
312 Robertson JA, Thayer MK, Whiting MF, Lawrence JF, Ślipiński A, Maddison DR, Farrell
313 BD 2015. The beetle tree of life reveals that Coleoptera survived end-Permian mass
314 extinction to diversify during the Cretaceous terrestrial revolution. *Systematic Entomology*
315 40:835–880.
- 316 McMahon DP, Hayward A, Kathirithamby J 2011. The first molecular phylogeny of Strepsiptera
317 (Insecta) reveals an early burst of molecular evolution correlated with the transition to
318 endoparasitism. *PLoS ONE* 6:e21206.
- 319 Misof B, Liu S, Meusemann K, Peters RS, Donath A, Mayer C, Frandsen PB, Ware J, Flouri T,
320 Beutel RG, Niehuis O, Petersen M, Izquierdo-Carrasco F, Wappler T, Rust J, Aberer AJ,
321 Aspöck U, Aspöck H, Bartel D, Blanke A, Berger S, Böhm A, Buckley TR, Calcott B, Chen
322 J, Friedrich F, Fukui M, Fujita M, Greve C, Grobe P, Gu S, Huang Y, Jermiin LS, Kawahara

AY, Krogmann L, Kubiak M, Lanfear R, Letsch H, Li Y, Li Z, Li J, Lu H, Machida R, Mashimo Y, Kapli P, McKenna DD, Meng G, Nakagaki Y, Navarrete-Heredia JL, Ott M, Ou Y, Pass G, Podsiadlowski L, Pohl H, Reumont von BM, Schütte K, Sekiya K, Shimizu S, Ślipiński A, Stamatakis A, Song W, Su X, Szucsich NU, Tan M, Tan X, Tang M, Tang J, Timelthaler G, Tomizuka S, Trautwein M, Tong X, Uchifune T, Walz MG, Wiegmann BM, Wilbrandt J, Wipfler B, Wong TKF, Wu Q, Wu G, Xie Y, Yang S, Yang Q, Yeates DK, YOSHIZAWA K, Zhang Q, Zhang R, Zhang W, Zhang Y, Zhao J, Zhou C, Zhou L, Ziesmann T, Zou S, Li Y, Xu X, Zhang Y, Yang H, Wang J, Wang J, Kjer KM, Zhou X 2014. Phylogenomics resolves the timing and pattern of insect evolution. *Science* 346:763–767.

Niehuis O, Hartig G, Grath S, Pohl H, Lehmann J, Tafer H, Donath A, Krauss V, Eisenhardt C, Hertel J, Petersen M, Mayer C, Meusemann K, Peters RS, Stadler PF, Beutel RG, Bornberg-Bauer E, McKenna DD, Misof B 2012. Genomic and morphological evidence converge to resolve the enigma of Strepsiptera. *Current Biology* 22:1309–1313.

Peters RS, Meusemann K, Petersen M, Mayer C, Wilbrandt J, Ziesmann T, Donath A, Kjer KM, Aspöck U, Aspöck H, Aberer A, Stamatakis A, Friedrich F, Hünefeld F, Niehuis O, Beutel RG, Misof B 2014. The evolutionary history of holometabolous insects inferred from transcriptome-based phylogeny and comprehensive morphological data. *BMC Evolutionary Biology* 14:52.

Pierce W 1909. A monographic revision of the twisted winged insects comprising the order Strepsiptera Kirby. *Bulletin of the United States National Museum* 66:1–232.

Pohl H 2000. Die Primärlarven der Fächerflügler – evolutive Trends (Insecta, Strepsiptera). *Kaupia, Darmstädter Beiträge zur Naturgeschichte* 10:1–144.

Pohl H 2002. Phylogeny of the Strepsiptera based on morphological data of the first instar larvae. *Zoologica Scripta* 31:123–134.

Pohl H 2009. The oldest fossil strepsipteran larva (Insecta: Strepsiptera) from the Geisel Valley, Germany (Eocene). *Insect Systematics & Evolution* 40:333–347.

Pohl H, Beutel RG 2008. The evolution of Strepsiptera (Hexapoda). *Zoology* 111:318–338.

Pohl H, Beutel RG 2013. The Strepsiptera-Odyssey: the history of the systematic placement of an enigmatic parasitic insect order. *Entomologia* 1:e4.

353 Pohl H, Beutel RG 2016. †*Kinzelbachilla ellenbergeri* – a new ancestral species, genus and
354 family of Strepsiptera (Insecta). *Systematic Entomology* 41:287–297.

355 Pohl H, Beutel RG, Kinzelbach R 2005. Protoxenidae fam. nov (Insecta, Strepsiptera) from
356 Baltic amber – a “missing link” in strepsipteran phylogeny. *Zoologica Scripta* 34:57–69.

357 Poinar G 2004. Evidence of parasitism by Strepsiptera in Dominican amber. *Biocontrol* 49:239–
358 244.

359 Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch S,
360 Rueden C, Saalfeld S, Schmid B, Tinevez J-Y, White DJ, Hartenstein V, Eliceiri K,
361 Tomancak P, Cardona A 2012. Fiji: an open-source platform for biological-image analysis.
362 *Nature Methods* 9:676–682.

363 Shi G, Grimaldi DA, Harlow GE, Wang J, Wang J, Yang M, Lei W, Li Q, Li X 2012. Age
364 constraint on Burmese amber based on U–Pb dating of zircons. *Cretaceous Research*
365 37:155–163.

366 Silvestri F 1941. Studi sugli “Strepsiptera” (Insecta). I. Ridescrizione e ciclo dell’*Eoxenos*
367 *Laboulbenei* Peyerimhoff. *Bolletino del Laboratorio di Zoologia generale e agraria della*
368 *Facoltà Agraria in Portici* 31:311–341.

369 Stehr F 1991. *Immature insects*. Michigan: Kendall/Hunt.

370 Toussaint EFA, Seidel M, Arriaga-Varela E, Hájek J, Král D, Sekerka L, Short AEZ, Fikáček M
371 2017. The peril of dating beetles. *Systematic Entomology* 42:1–10.

372 Voigt E 1938. Weichteile an fossilen Insekten aus der eozänen Braunkohle des Geiseltales bei
373 Halle (Saale). *Nova Acta Leopoldina, Neue Folge* 6:1–38.

374

375

Figure 1 Strepsiptera primary larva in Burmese amber, ventral view. A. Photomicrograph with an Axio Zoom.V16 with a PlanNeoFluar Z 1.0x. B. Drawing based on photomicrographs with an Olympus IX81 inverted fluorescence microscope with UIS2 objective. Abbreviations: af = antennal field, cb = caudal seta, cx = coxa, fe = femur, fs = frontal seta, lcb = lateral caudal seta, mp = maxillary palp, mssp = mesosternal plate, mt = metanotum, mx = maxilla, prsp = prosternal plate, sbsIX/X = segmental border between abdominal sternites IX/X, sbtVIII/IX = segmental border between abdominal tergites VIII/IX, sI–sIX = abdominal sternites I–XI, st = stemmata, ta = tarsus, te = tentorium, ti = tibia, X = abdominal segment X, XI = abdominal segment XI.

Figure 2 Terminal segments of Strepsiptera primary larva in Burmese amber, drawings based on photomicrographs with an Olympus IX81 inverted fluorescence microscope with UIS2 objective. A. Dorsal view. B. Ventral view. Abbreviations: sbsIX/X = segmental border between abdominal sternites IX/X, sbtVIII/IX = segmental border between abdominal tergites VIII/IX, X = abdominal segment X, XI = abdominal segment XI.

Figure 3 A, B. Strepsiptera primary larva in Burmese amber, head, pro-, and mesothorax, ventral view. C, D. *Eoxenos laboulbenei*, head, pro-, and mesothorax, ventral view. Photomicrographs with an Olympus IX81 inverted fluorescence microscope with UIS2 objective. Abbreviations: af = antennal field, cos = coxal seta, lb = labium, mp = maxillary palp, mssp = mesosternal plate, mx = maxilla, prsp = prosternal plate, st = stemmata, te = tentorium, ti = tibia.

Figure 4 A, B. Strepsiptera primary larva in Burmese amber. A. Meso-, metathorax, and anterior abdominal segments, ventral view. B. Terminal abdominal segments. C, D. *Eoxenos laboulbenei*. C. Abdominal segments III–IX, ventral view. D. Terminal abdominal segments. Photomicrographs with an Olympus IX81 inverted fluorescence microscope with UIS2 objective. Abbreviations: cb = caudal bristle, lcb = lateral caudal bristle, sbsIX/X = segmental border between sternite IX and X, sbsX/XI = segmental border between sternite X and XI, sbtVIII/IX = segmental border between tergite VIII and IX, ti = tibia, X = abdominal segment X, XI = abdominal segment XI.

406 **Figure 5** Primary larva of *Eoxenos laboulbenei*, drawings based on scanning electron
 407 micrographs (modified from Pohl, 2000). A. Ventral view. B. Dorsal view. Abbreviations: af =
 408 antennal field, cb = caudal bristle, cx = coxa, fe = femur, fs = frontal seta, lb = labium, lcb =
 409 lateral caudal bristle, mp = maxillary palp, mx = maxilla, prsp = prosternal plate, sI–sXI =
 410 abdominal sternite I–XI, st = stemma, ta = tarsus, ti = tibia, tI–tX = abdominal tergite I–X.

Figure 1

Strepsiptera primary larva in Burmese amber, ventral view.

A. Photomicrograph with an Axio Zoom.V16 with a PlanNeoFluar Z 1.0x. B. Drawing based on photomicrographs with an Olympus IX81 inverted fluorescence microscope with UIS2 objective. Abbreviations: af = antennal field, cb = caudal seta, cx = coxa, fe = femur, fs = frontal seta, lcb = lateral caudal seta, mp = maxillary palp, mssp = mesosternal plate, mt = metanotum, mx = maxilla, prsp = prosternal plate, sbsIX/X = segmental border between abdominal sternites IX/X, sbtVIII/IX = segmental border between abdominal tergites VIII/IX, sl-sIX = abdominal sternites I-XI, st = stemmata, ta = tarsus, te = tentorium, ti = tibia, X = abdominal segment X, XI = abdominal segment XI.





Figure 2

Terminal segments of Strepsiptera primary larva in Burmese amber, drawings based on photomicrographs with an Olympus IX81 inverted fluorescence microscope with UIS2 objective.

A. Dorsal view. B. Ventral view. Abbreviations: sbsIX/X = segmental border between abdominal sternites IX/X, sbtVIII/IX = segmental border between abdominal tergites VIII/IX, X = abdominal segment X, XI = abdominal segment XI.




Figure 3

Strepsiptera primary larva in Burmese amber in comparison with an extant primary larva.

A, B. Strepsiptera primary larva in Burmese amber, head, pro-, and mesothorax, ventral view. C, D. *Eoxenos laboulbenei*, head, pro-, and mesothorax, ventral view. Photomicrographs with an Olympus IX81 inverted fluorescence microscope with UIS2 objective. Abbreviations: af = antennal field, cos = coxal seta, lb = labium, mp = maxillary palp, mssp = mesosternal plate, mx = maxilla, prsp = prosternal plate, st = stemmata, te = tentorium, ti = tibia.






Figure 4

Strepsiptera primary larva in Burmese amber in comparison with an extant primary larva.

A, B. Strepsiptera primary larva in Burmese amber. A. Meso-, metathorax, and anterior abdominal segments, ventral view. B. Terminal abdominal segments. C, D. *Eoxenos laboulbenei*. C. Abdominal segments III–IX, ventral view. D. Terminal abdominal segments. Photomicrographs with an Olympus IX81 inverted fluorescence microscope with UIS2 objective. Abbreviations: cb = caudal bristle, lcb = lateral caudal bristle, sbsIX/X = segmental border between sternite IX and X, sbsX/XI = segmental border between sternite X and XI, sbsVIII/IX = segmental border between tergite VIII and IX, ti = tibia, X = abdominal segment X, XI = abdominal segment XI.




Figure 5

Primary larva of *Eoxenos laboulbenei*, drawings based on scanning electron micrographs (modified from Pohl, 2000).

A. Ventral view. B. Dorsal view. Abbreviations: af = antennal field, cb = caudal bristle, cx = coxa, fe = femur, fs = frontal seta, lb = labium, lcb = lateral caudal bristle, mp = maxillary palp, mx = maxilla, prsp = prosternal plate, sl-sXI = abdominal sternite I-XI, st = stemma, ta = tarsus, ti = tibia, tl-tX = abdominal tergite I-X.

