

1 **Effects of a six-week weighted-implement throwing program on baseball pitching velocity,**
2 **kinematics, arm stress, and arm range of motion**

3 Joseph A. Marsh¹, Matthew I. Wagshol², Kyle J. Boddy³, Michael E. O'Connell⁴, Sam J. Briend⁵,
4 Kyle E. Lindley⁶, Alex Caravan⁷

5 ^{1, 2, 3, 4, 6, 7} Driveline Baseball, Research & Development, Kent, WA USA

6 ⁵ Driveline Baseball, High Performance, Kent, WA USA

7

8 Corresponding Authors:

9 Kyle J. Boddy, Matthew I. Wagshol, Joseph A. Marsh
10 19612 70th Avenue South, Unit 2-4
11 Kent, WA 98032

12 Email address: kyle@drivelinebaseball.com, mwagshol@drivelinebaseball.com,
13 joe@drivelinebaseball.com

14

15 **Background.** Weighted-baseball training programs are used at the high school, collegiate, and
16 professional levels of baseball. The purpose of this study was to ~~compare-evaluate the effects of~~
17 a six-week training period ~~using-consisting of~~ weighted implements, manual therapy,
18 weightlifting, and other modalities on ~~how they may affect arm stress, arm range of~~
19 ~~motion~~ shoulder external rotation, elbow valgus stress, pitching velocity, and kinematics.

20 **Hypothesis.** ~~A six-week training program using that includes~~ weighted implements may increase
21 ~~pitching velocity along with concomitant~~ There is a gain in pitching velocity with concomitant
22 increases in arm angular velocities, joint kinetics, and shoulder external rotation.

23 **Methods.** Seventeen collegiate and professional baseball pitchers (~~age range 18-23, average:~~
24 ~~19.9~~) training at Driveline Baseball were ~~tested-evaluated~~ with via a combination of an eight-
25 camera motion-capture system, range-of-motion measurements, and radar- and pitch-tracking
26 equipment, ~~both~~ before and after a six-week training period. Each participant received
27 individualized training programs ~~as standard protocol~~, with significant overlap in training
28 methods for all athletes. ~~In addition, it should be noted that individuals in this training program~~
29 ~~used training equipment sold out of Driveline Baseball (Kent, WA), which is owned by one of~~
30 ~~the primary authors of this study, Kyle J. Boddy, and followed prescribed training programs out~~
31 ~~of the aforementioned author's published book Hacking the Kinetic Chain.~~ Twenty-eight
32 biomechanical parameters were computed for each bullpen trial, four arm range-of-motion
33 measurements were taken, and pitching velocities were recorded before and after the training
34 period. Pre- and post-training period data were compared via post-hoc paired *t* tests.

35 **Results.** There was a ~~statistically insignificant~~ no change in pitching velocity across the
36 seventeen subjects (~~p value?~~). ~~Four biomechanical parameters for the holistic group were~~
37 ~~significantly changed after the training period: Among the biomechanical parameters~~
38 ~~computed, four were significantly different after the training period:~~ internal rotational velocity
39 was higher (~~from 4527 to 4759 degrees/second~~), shoulder abduction was lower at ball release
40 (96 to 93°), the shoulder was less externally rotated at ball release (95 to 86°) and shoulder
41 adduction torque was higher (~~from 103 to 138 N-m~~). Among the arm ~~arm laxity~~ range of motion
42 measurements, four were significantly different after the training period: the shoulder internal
43 rotation range of motion and total range of motion for both the dominant and non-dominant
44 arm.

45 When the group was divided into those who gained pitching velocity and those who did not,
46 the group that gained pitching velocity showed no significant increase in shoulder external
47 rotation, or elbow valgus stress.

48 **Conclusions.** ~~Following a six-week weighted implement program, pitchers did not show a~~
49 ~~statistically-significant change in velocity, joint kinetics, or shoulder and elbow range of motion.~~
50 ~~When comparing pitchers who gained velocity versus pitchers who did not, no statistically~~
51 ~~significant changes were seen in joint kinetics and shoulder range of motion, as a result of~~
52 ~~positive developments in kinematics and range of motion as demonstrated in the study.~~

Commented [MH1]: Was this an addition based on reviewer comments? I would suggest putting this at the end of the document as a "Disclosures" or "Conflicts" section.

Formatted: Font: Italic

Commented [MH2]: Avoid this.

Commented [MH3]: Keep consistent with above.


Commented [MH4]: Add SD or SE metrics.

Formatted: Font: Not Bold

Commented [MH5]: Confirm. Above you say shoulder internal rom and total rom was significant?

Formatted: Font: Not Bold

Formatted: Font: Not Bold



54 **Introduction**

55 Studies on underweight and overweight baseballs have shown a positive training effect on the
56 throwing velocity of regulation-weight baseballs (DeRenne et al., 2005; DeRenne et al., 1990;
57 Egstrom et al., 1960; DeRenne, 1985). Additionally, studies have also shown no negative effects
58 of throwing underweight and overweight implements on pitching control or injury risk
59 (DeRenne et al., 2005; DeRenne et al., 1994).

60 A recent biomechanical study shows that pitching slightly underweight and overweight
61 baseballs can produce variations in kinematics (specifically arm, trunk, pelvis, and shoulder
62 velocities) without increased arm kinetics ~~<CAN SOMEONE INCLUDE THESE>~~ (Fleisig et al.,
63 2016) and that maximum-effort crow-hop throwing with the same implements can increase
64 shoulder internal rotation angular velocity and elbow varus torque (Fleisig et al., 2017); ~~Fleisig~~
65 ~~et al., 2016~~. ~~Additionally, another, while a study under current review published study~~
66 ~~(Reinold, M. 2017) indicates~~ weighted-baseball throwing can increase shoulder external
67 rotation in a six-week training period on high school athletes (Reinold, M. 2017).

68 There is also published research on heavier-weighted plyometric throws ~~as~~ used in training and
69 rehab programs, including but not limited to two handed chest passes and side throws of 8-
70 pound “plyoballs” or the more traditionally-named medicine balls (Wilk, Meister & Andrews,
71 2002). Further research has found eight weeks of plyometric training can improve shoulder
72 internal rotation power and throwing distance (Fortun, Davies & Kernozck, 1998). A different
73 study using plyoballs and “The Ballistic Six” found a significant improvement in throwing
74 velocity (Carter et al., 2007). While there is also research suggesting that throwing weighted
75 plyos from 2–8 lb. may improve proprioception (Swanik et al., 2002).

76 Major League Baseball teams have increasingly adopted the use of weighted-implement
77 training and have seen a large uptick in drafted players with prior experience using weighted
78 baseballs as amateurs. No conclusive evidence exists that truly explains the mechanism of the
79 velocity gains, but claims that weighted-ball training can increase “arm strength” may be false
80 (Cressey, E. 2013).

81 Increases in throwing shoulder external rotation and loss of throwing shoulder internal rotation
82 are potentially deleterious (Wilk et al., 2011), but, to our knowledge, no weighted-implement
83 training program in existing research combines a throwing program with other training
84 modalities to potentially reduce negative adaptive effects on the arm. There is evidence that
85 certain mobility programs can reduce the negative adaptive effects of throwing (Laudner et al.,
86 2008), and it is theorized that heavy resistance training and manual therapy can potentially may
87 aid in this regard as well.

88 in addition, collegiate and professional pitchers participating in Driveline Baseball’s summer
89 training programs have on average increased pitching velocity 2.7 MPH in 2016 and 3.3 MPH in
90 2017 (Driveline Baseball, 2016 and 2017). ~~Driveline Baseball’s summer training periods show~~

Commented [MH6]: Be careful with statements like this. “another published study”. Most readers will assume a peer reviewed scientific piece. You cite a website/blog. I get that this appears to be a solid study. Was it ever published? Can you cite that?

Commented [MH7]: What does this mean? Increase power?

Commented [MH8]: Increase....

Commented [MH9]: ?

Commented [MH10]: Is this anecdotal?

Commented [MH11]: Is this an opinion piece?

Commented [MH12]: Such as?

Commented [MH13]: Out of place. Avoid 1 sentence paragraphs.

91 documented increases of 2.7 MPH in 2016 and 3.26 MPH in 2017 pitching velocity (Driveline
92 Baseball, 2016 and 2017), so there is anecdotal case study support for the programs'
93 effectiveness.

94 The purpose of ~~the~~ this study ~~is~~ was to evaluate the training effects of a weighted-~~implement~~
95 ~~throwing program~~. The purpose of this study was to see what the positive training effects on
96 ~~throwing weighted implements would be and~~ plus individualized training routines
97 ~~geared focused on around~~ combating the negative effects of ~~if~~ throwing on pitching velocity,
98 external rotation and elbow varus torque that is demonstrated ~~shown~~ in other studies ~~on~~
99 ~~pitching velocity, external rotation and elbow varus torque~~. to see if the negative effects of
100 ~~throwing shown in other studies could be replicated or if the secondary exercises included in~~
101 ~~the training program could effectively combat these adaptations.~~ We hypothesize the
102 ~~previously described program will increase external rotation~~, it was hypothesized that the
103 ~~standard effects would be shown~~ — increased external rotation, increased ball velocity, and
104 increased elbow varus torque.

105

106 **Methods**

107 *Participants and Informed Consent*

108 Healthy and asymptomatic college and professional pitchers ~~were~~ recruited from the Driveline
109 Baseball 2017 training group via opt-in forms. ~~Before each pitcher was scheduled for their initial~~
110 ~~test~~ Prior to being included in the study, investigators asked the pitchers about their current
111 injury status. Pitchers were excluded if they had current symptoms of arm or shoulder pain or
112 fatigue, or any other pain or discomfort that would prohibit completion of the ~~testing~~
113 ~~period~~ study. Additionally, a prerequisite to train in the Driveline Baseball spring-summer group
114 required medical clearance and a certified athletic trainer's sign-off before throwing pitches off
115 a mound. Pitchers were not excluded based on previous history of injuries that did not
116 currently manifest themselves. ~~Pitchers were not excluded based on previous training history,~~
117 ~~although a few had trained at Driveline Baseball right before the study and most had~~
118 ~~experimented remotely with Driveline methods; the average time spent at Driveline right~~
119 ~~before the study's start was~~ around 16 ± 10 days, with a standard deviation of 10, a maximum
120 of 41 and a minimum of 3 days.

121 Pitchers were scheduled to come into the Driveline Baseball Research Facility (Kent, WA) for
122 one visit. Upon arrival, participants were provided a verbal explanation of the study and asked
123 to read and sign an Informed Consent document before beginning. The investigator verbally
124 confirmed the major bullet points of the Informed Consent document in addition to obtaining a
125 witnessed, legal signature from the pitcher, only proceeding if the pitcher submitted both a
126 valid signature *and* verbally confirmed acceptance of all the risks contained within the Informed
127 Consent document.

Commented [MH14]: You can be exact.

128 The study was approved by Hummingbird IRB, who granted ethical approval to carry out the
129 study at the author's facilities (Hummingbird IRB #: 2017-29, Protocol WB-DLR-115).

130 Twenty-one baseball pitchers (age range: 18-23) with high school and college pitching
131 experience met these criteria and agreed to participate. Four were excluded bringing the final
132 number to seventeen. The data on these pitchers is recorded in Table 1.

133 [TABLE 1]

134 ~~[TABLE 2]~~

135 *Testing Procedure and Measurements*

136 During the testing period, range of motion measurements were taken using a goniometer to
137 measure shoulder internal and external rotation in both the dominant and non-dominant arms.
138 The same investigator was used for each individual in the initial and final tests; previous
139 research has shown high intra-reliability ~~in-for~~ goniometer measurements (Boone et al., 1978).
140 Each pitcher was measured on the same day as the biomechanical screening.

Commented [MH15]: ?

141 Measurements were taken with each athlete lying in the lateral decubitus position (~~The~~
142 ~~specific testing position is displayed below in Figure 1~~). Testing was done in this position due to
143 the fact that when lying supine, the humeral head is more likely to glide forward in the socket,
144 causing irritation in the anterior shoulder and leading to more inaccurate measurements as the
145 athlete can compensate for a lack of range of motion by rolling ~~his-the~~ shoulder forward
146 anterior? or backwardposterior?. In the lateral decubitus plane, the humeral head is in a more
147 advantageous position to externally (Part A of Figure 1) and internally rotate (Part B of Figure 1)
148 without humeral head glide (Reinold et al., 2004).

149 [Figure 1]

150 The investigator performing this part of the study was a certified strength and conditioning
151 coach with seven years of experience and specifically trained in measuring range of motion of
152 the shoulder using standard tools. Once the athlete was in the appropriate position, the
153 investigator passively moved the arm until tension was reached and the measurement was
154 taken. The intraclass correlation coefficient (ICC) of a trained clinician performing total range of
155 motion tests of the shoulder have shown to be very reliable (Wilk et al., 2009).

156 The pitchers ~~were allowed to throw threw~~ as many warm up pitches as they liked ~~until they~~
157 ~~signalled/signaled to the main study investigators that they were ready to pitch prior to~~
158 beginning. ~~Next, P~~pitchers were fitted with reflective markers in preparation for three-
159 dimensional motion capture. ~~portion of the pre-test and post-test for later kinematic and~~
160 kinetic analyses. Forty-eight reflective markers were attached bilaterally on the third distal
161 phalanx, lateral and medial malleolus, calcaneus, tibia, lateral and medial femoral epicondyle,

162 femur, anterior and posterior iliac spine, iliac crest, inferior angle of scapula, acromial joint,
163 midpoint of the humerus, lateral and medial humeral epicondyle, midpoint of the ulna, radial
164 styloid, ulnar styloid, distal end of index metacarpal, parietal bone, and frontal bone, as well as
165 on the C7 and T10 vertebrae, the sternal end of the clavicle, and the xiphoid process.

166 Pitchers then threw between 3-8 maximum effort throws, with approximately 30-60 seconds of
167 rest between pitches. Fatigue was assumed to be negligible with such a low pitch count. Throws
168 were made using a 5-oz. (142g) regulation baseball off the mound to a strike zone target (Oates
169 Specialties, LLC, Huntsville, TX) located above home plate, which was 60' 6" (18.4 m) away.
170 Testing concluded when the investigators were satisfied they had recorded three successful
171 throws at least three clean takes for analysis. Sample photographs and high-speed videos
172 (Sanstreak Corp., San Jose, CA) of the setup and pitches in motion are shown in Supplemental
173 Photos and Videos 1-3.

174 For each trial, ball velocity was measured by a Doppler radar gun (Applied Concepts; Stalker
175 Radar, Richardson, Texas). Additionally, for all trials, the Three-dimensional kinematics
176 motions of the reflective markers were tracked with using an 8-camera automated motion-
177 capture system, sampling at 240 Hz (Prime 13 System, Natural Motion / Optitrack, Corvallis,
178 Oregon), shown in research to be comparable to more commonly-used high-end motion-
179 capture systems (Thewlis et al., 2013). A total of 8 cameras were placed symmetrically around
180 the capture volume, approximately 8 feet 2.4 meters from the center of the pitching mound, at
181 roughly 8 feet 2.4 meters high. One camera was lowered on the throwing arm side to avoid
182 collisions, and an additional camera was lowered to aid in marker tracking.

183 In total, 28 kinematic and kinetic values-measures (11 position, 6 velocity, and 11 kinetic) were
184 calculated using our personal code based using the ISB recommended model of joint coordinate
185 systems (Wu et al. 2005) with code based on Fleisig methods (Fleisig et al., 2017) in Visual3D (C-
186 Motion Inc., Germantown MD). Marker position data was filtered using a 20-Hz Butterworth
187 low-pass filter. The mean values for all variables were calculated for each participant based
188 upon their 3 best clearest throws, based upon marker and motion readability (Escamilla et al.,
189 1998).

190 Five joint angles, positional kinematic values were found calculated at the events of both foot
191 contact (FC) and ball release (BR), including: elbow flexion, shoulder horizontal abduction,
192 shoulder abduction, shoulder external rotation, and wrist extension. Additionally, as well as
193 maximum dynamic shoulder external rotation was measured as well. All kinematic measures
194 Measurements were all taken as their local joint angles, using local coordinate systems.

195 The Six velocity parameters included d mean pelvis angular velocity at FC and BR, and the
196 maximum values of of pelvis angular velocity, upper torso angular velocity, elbow extension
197 velocity, and shoulder internal rotation velocity. Pelvis and upper torso angular velocities were
198 measured by as their rotations in the global reference coordinate system frame. Elbow,

Commented [MH16]: Why?

Commented [MH17]: Specify order? Dual pass? 20 hz cutoff.

Commented [MH18]: ? does this mean there was marker drop out? Was there any gap filling or other procedures performed other than filtering the data?

Commented [MH19]:measured as.....

Expand. What is maximum dynamic shoulder external rotation?

Commented [MH20]: Why are some called "angular velocity" and others just "velocity"? Are they all the same?

199 shoulder, and wrist velocities were calculated as the rate of change in the joint angle and is
200 expressed as °/sec.

201 Maximum ~~values for~~ elbow and shoulder kinetics were calculated. ~~Values were reported~~ as
202 either a force or a torque applied ~~on~~ to the joint, ~~applied~~ by the proximal segment onto the
203 distal segment ~~at the joint~~. Six forces were calculated, including medial, anterior, and
204 compression forces on the elbow, and superior, anterior, and compression forces on the elbow.
205 Five joint torques were ~~also~~ computed: elbow flexion torque, elbow varus torque, shoulder
206 horizontal adduction torque, shoulder adduction torque, and shoulder internal rotation torque.

207 *Training Methods*

208 In between the two pre and post tests, pitchers were exposed to a six-week training program,
209 slightly individualized for each person-athlete based on their strengths and weaknesses, which
210 were determined vis-a-vis from a their biomechanical and performance assessment results.
211 Pitchers were slotted placed into one of three different categories for throwing programing.
212 These were velocity development, mound development, or a hybrid version of the two. All
213 athletes performed their training program six days a week with the seventh day being an off
214 day.

215 Each pitcher ~~began to~~ completed a warm-up ~~by~~ using foam rollers and lacrosse balls for self-
216 myofascial release (SMR) of various lower body and throwing arm muscles. Another option was
217 rolling out the forearm with Arm Aid Extreme devices (The Armaid Company, Inc., Blue Hill,
218 ME). Athletes were allowed ~~to roll out~~ SMR for a period of time that they determined necessary
219 and were able to use SMR on other body parts if necessary. The standard SMR exercises can be
220 found in the supplemental materials pages 1-7 of HTKC1.

221 Following SMR, athletes completed a set of exercises using Jaeger Band surgical tubing (Jaeger
222 Sports, Los Angeles, CA). Pitchers performed a forward fly to overhead reach, reverse fly to
223 overhead reach, bicep curl with supination, tricep extension with pronation, internal and
224 external rotations with elbow at shoulder height. Further details on the exercises can be found
225 on pages 8-12 of the supplemental materials of HTKC1

226 Although Jaeger bands use a wrist cuff, surgical-tubing exercises with a handle have been
227 shown to result in low to moderate EMG activation of the rotator cuff and surrounding
228 musculature (Myers et al., 2005). ~~A thesis has also stated~~ Surgical tubing exercises can improve
229 velocity and shoulder internal and external strength (Baheti, 2000).

230 Following band work, pitchers performed a series of exercises with an Oates Specialties
231 shoulder tube (info on product?). The tube is designed as-for oscillation work to warm up the
232 rotator cuff muscles. Pitchers performed shoulder flexion in front, shoulder abduction to the
233 side, external/internal rotations, pronation/supination twirls, and stride-length forward
234 shoulder rotations. More detail on these exercises can be found on pages 13-16 of the
235 supplemental materials of HTKC1.

Commented [MH21]: External force?

Commented [MH22]: ?

Commented [MH23]: Expand. How was this done? Inverse kinematics? Specify parameters.

What is a shoulder torque? Is it a resultant shoulder moment that was calculated?

Commented [MH24]: Reference?

236 The pitchers then performed a series of four exercises with 10-lb. wrist weights. The goals of
237 these exercises ~~are-were~~ to warm up the muscles of the forearm and ~~work~~ the back of the
238 shoulder eccentrically. The exercises were Pronated Swings (with two-arms), Two-Arm Throws,
239 modified Cuban Press, and Pivot-Pickoff Throws. Further details of the exercises can be found
240 on pages 17-24 in the supplemental materials of HTKC1.

Commented [MH25]: kg

Commented [MH26]: posterior?

241 Athletes then moved to a specific series of throws using plyometric PlyoBalls (~~custom made~~ soft
242 sand-filled weighted balls ranging from 100-2000 grams; ~~—sold out of Driveline Baseball, Kent,
243 WA~~). There ~~are-were~~ five exercises performed; each exercise ~~is-was~~ unique within the
244 constraints of the body's position to focus on different mechanical elements. Pitchers
245 performed Reverse Throws, Pivot Pickoffs, Roll-in Throws, Rockers, and Walking Windups.

246 The ball weights, sets, and reps were all standard across the participants, depending on the
247 training day. Pitchers completed the above warm-up six days a week with the volume and
248 intensity of PlyoCare throws varying on the day. The throwing schedules and explanations on
249 how to perform the exercises are listed on pages 25-36 in the supplemental materials of HTKC1.

250 On certain days pitchers were scheduled to long-toss. ~~This occurred either off-site at a local
251 park or inside while throwing into a net. Driveline utilizes tTwo different types of long toss days
252 were implemented. The first iswas a lower intensity day. Rate of perceived exertion (RPE) iswas
253 around 60-70% for the athlete accompanied by loose, relaxed throwing with a large arc as the
254 athlete backs up in distance. Maximum distance iswas determined by throwing ability and RPE
255 and as such will vary from athlete to athlete. This day doesdid not include any high intensity
256 compression throws.~~

Commented [MH27]: More specific.

257 ~~The second type of long toss day iswas similar to the first, except it is performed at an RPE of
258 80-90% and the athlete will carryies the extension throws out to roughly maximum throwing
259 distance. Upon reaching maximum throwing distance in as many or as few throws as required,
260 the athlete will come in andthen performs eight to twelve high intensity compression throws.
261 These compression throws remove the arc from the throw and are thrown roughly parallel to
262 the ground from the throwers release point. Number of throws will vary day to day for each
263 individual athlete as they are instructed to be extremely receptive to their body's response and
264 own personal comfort level.~~

265 Research on long-toss has largely focused on throws at max distance while throwing hard on-a-
266 line, with one study finding max distance throws resulted in more torque than in pitching
267 (Fleisig et al., 2011). Another study found that max distance, hard on-a-line throws resulted in
268 similar loads to pitching (Slenker et al., 2014).

269 Long-toss as described in the programming did not solely consist of max distance, hard on-a-
270 line throws. Most consisted of high-arc (extension) throws to a tolerable distance for the day,
271 otherwise described as catch-play to a distance that is tolerable. Certain training days did
272 consist of hard on-a-line (compression) throws, which are marked in the supplemental

273 materials. It is important to note these distinctions since a recent study showed that many
274 coaches, ATCs, and players define long-toss differently (Stone et al., 2017).

275 Each pitcher completed a post-throwing exercise circuit after each day of throwing workouts.
276 The circuit consisted of standing rebounders; the pitchers threw a 4- and 2-lb. PlyoCare ball at a
277 trampoline on the ground and were told to “stick” the catch of the ball or stop its upward
278 momentum right away.

279 ~~Following rebounders~~Next, ~~were~~ reverse scap pull-aparts, anterior band pull-aparts, and the no
280 money drill. After band exercises, pitchers performed waiter walks. The pitchers held a
281 kettlebell with their humerus at shoulder height and forearm facing vertically while walking.
282 The kettlebell ~~is-was~~ gripped by the handle with the weight facing the ceiling. More details of
283 the post throwing circuit can be found in the supplemental materials of HTKC2.

284 After the exercise circuit, each pitcher was able to use the Marc Pro EMS device (Marc Pro,
285 Huntington Beach, CA). The Marc Pro has been shown to improve muscle performance,
286 recovery, and reduce Delayed Onset Muscle Soreness (or-DOMS) caused by exercise (Westcott
287 et al., 2011, Westcott et al., 2013). It has been hypothesized that these results come from and
288 increase in blood flow (DiNubile et al., 2011).

289 In conjunction with the throwing program athletes were ~~also~~ involved in a strength and
290 conditioning program. This program included lifting weights, medicine ball throws, and mobility
291 work. This program was individualized to each athlete depending on a separate physical and
292 athletic screening.

293 Pitchers ~~also~~ saw a physical therapist during the training period. Trainers are also certified in
294 Functional and Kinetic Treatment with Rehabilitation (or-FAKTR), cupping, and other manual
295 therapy techniques. Athletes were able to receive treatment on an as-needed basis.

296 Each pitcher ~~in this study~~ had five- to six-throwing days scheduled a week. The throwing days
297 were classified as high-intent days, hybrid days (medium intent days), and recovery days (low
298 intent days), with the intensity and volume of throws changing per day. Athletes typically
299 performed two high intensity days, one moderate intensity day and three recovery days within
300 a given seven day cycle.

301 To be included in the final study post data collection, pitchers had to ~~have~~ participated in at
302 least 90% of the training days ~~laid out for them, or they were dropped from the study.~~ Four of
303 the twenty-one pitchers initially chosen for the study failed were chosen for failing to to meet
304 this criteriathis criterion.

305 Data from the training periods—including schedules, workloads, lifting programs, and
306 intermediate progress—can be found in the supplemental data as spreadsheets for all pitchers
307 ~~involved in the study.~~

Commented [MH28]: Scapular?
Pull aparts? Technical term? Scapular retraction?

Commented [MH29]: They held the kettlebell with their hand,
not humerus.
what is humerus at shoulder height? Shoulder flexed to 90deg?
Parallel with floor? What for how long?

Commented [MH30]: Able to. Did they? All?

Commented [MH31]: I would be careful with this. Mechanisms
of DOMS are widely debated.

308 All statistical analyses were performed using R (RStudio Team, Boston, MA). After cleaning and
309 preprocessing each individual athlete's data (the initial and post biomechanical parameters, the
310 range of motion measurements, the range of motion strength numbers, and velocity data),
311 means and standard deviations were calculated for each single metric measure, and then the
312 differences were computed, along with the subsequent t metric and p-value. A paired t-test
313 was used due to a relatively small sample size and unknown true population variances. To
314 calculate the t metric, the mean differences between observations were divided by the
315 standard error of these differences, which was calculated by the standard deviation of
316 differences divided by the square root of the sample size, n. An n-1 degree of freedom was
317 used, along with an alpha level of 0.05, leaving the pure probabilistic chance of any metric
318 being highlighted as a false positive as 5% or less. A post-hoc analysis with similar statistical
319 methods was also performed on both the subgroup of pitchers who saw a velocity increase
320 during the training period and those who saw a velocity decrease.

321 Results

322 Pre- and post-range of motion tests are shown in Table 23. Four arm range of motion arm laxity
323 measurements were significantly different after the training period: internal rotation range of
324 motion and total range of motion were higher (p values) for both dominant and non-dominant
325 arms. You say 4 were significant. What about the other 2 measures?

326 Perhaps notably, shoulder external-rotation range of motion did not was not significantly
327 increased after the training period. Of the entire sample size, passive external rotation saw a
328 gain of 1 degrees in the dominant arm, which was not significant.

329 [TABLE 23]

330 Splitting the groups post-hoc into the pitchers that gained velocity and those who did not gain
331 velocity did not did not may yield interesting results additional significant results differences
332 between the groups. For instance, when those who gained throwing velocity were split into
333 their own group, we might have expected to see the group that gained velocity (n=9) to have
334 statistically significant increases in post-training passive shoulder external-rotation range of
335 motion, but the gain in post-training passive shoulder external-rotation range of motion was
336 2.8 +/-? degrees, which was not statistically significant.

337 Range-of-motion changes of for the increase and decrease velocity groups can be found in
338 tables 3 and 4 below.

339 [TABLE 34]

340 [TABLE 45]

341 Mean kinematics values of for the pre and post-test are shown in Table 5. At front-foot contact,
342 there were no significant differences in any of the joint positions and velocities. During arm
343 cocking, maximum internal rotation velocity was higher. At ball release, shoulder abduction was

Commented [MH32]: This isn't outlined above at all. See my comment about the filter used.

Commented [MH33]: Rom strength?

Commented [MH34]: Ref?

Commented [MH35]: Why?

Commented [MH36]: 1 degree, given variability and no SD given, it probably is noise.

Commented [MH37]: How much? Give the data.

344 lower and external rotation decreased. ~~Meaning the arm was more fully internally rotated~~
345 ~~towards the target at the moment of ball separation in the delivery.~~

Commented [MH38]: How much? Give the data.

Commented [MH39]: Not results. Discussion.

346 [TABLE ~~56~~]

347 For the increased velocity group, no values were statistically significantly different at front foot
348 contact (~~as displayed below in Table 6~~). Maximum internal rotation velocity and maximum
349 elbow extension velocity were significantly higher in the arm cocking phase. External rotation
350 was significantly lower at ball release. No values were different for the velocity decrease group
351 at front foot contact, arm cocking, or ball release (as depicted in Table 7).

Commented [MH40]: P value. How much? Give #s.

353 [TABLE ~~67~~]^f

354 ~~No values were different for the velocity decrease group at front foot contact, arm cocking, or~~
355 ~~ball release, as depicted in Table 7.~~

357 [TABLE ~~78~~]

359 Maximum shoulder adduction torque was the only parameter to significantly increase during
360 the arm cocking phase for all athletes (~~as recorded in Table 8~~). ~~No values significantly changed~~
361 ~~in the deceleration phase. For the velocity increase group, no value kinetic measures were was~~
362 ~~significantly changed different in the arm cocking phase. Maximum shoulder superior force was~~
363 ~~the only variable significantly higher in the deceleration phase (with full detail in Table 9).~~
364 Maximum shoulder adduction torque was significantly higher in the velocity decrease group at
365 arm cocking. Elbow anterior force, elbow compressive force, elbow flexion torque, and
366 shoulder compressive force were all significantly lower in the arm deceleration phase (Table
367 10).

Commented [MH41]: Same. Give #'s. p-value.

Commented [MH42]: Give #'s.

370 [TABLE ~~89~~]

371 ~~For the velocity increase group, no value was significantly change in the arm cocking phase.~~
372 ~~Maximum shoulder superior force was the only variable significantly higher in the deceleration~~
373 ~~phase, with full detail in Table 9.~~

374 [TABLE ~~940~~]

375 Maximum shoulder adduction torque was the only value significantly higher in the velocity
376 decrease group at arm cocking. Elbow anterior force, elbow compressive force, elbow flexion
377 torque, and shoulder compressive force were all significantly lower in the arm deceleration
378 phase, as displayed in Table 10.

379 [TABLE 101]

380 Discussion

381 This study investigated the effects of.....The hypothesis that a baseball training program ~~that~~
382 featuring weighted implements would significantly increase shoulder external-rotation range
383 of motion was not supported by the current study. This was consistent for ~~—not in~~ the entire
384 subject pool ~~nor as well as in taking just those the sub grouping~~ who gained velocity, ~~—~~ despite
385 this phenomenon being posited as a way to enhance ball velocity (Matsuo et al., 2001).

386 It has generally been hypothesized ~~in research~~ that weighted balls work along the speed-
387 strength spectrum. One study found significant differences in maximal internal rotation (IR) and
388 elbow extension (EE) velocity when throwing different ball weights (Tillaar & Ettema, 2011).
389 With a second study finding 67% of ball velocity at release could be accounted for by internal
390 rotation and elbow extension (van den Tillaar & Ettema, 2004). In our work, for ~~Of~~ the entire
391 study sample, there was a significant change in IR velocity, ~~-~~ but not EE velocity.

392 When our sample was ~~B~~broken up into those who increased ~~d~~- and decreased ~~d~~-velocity groups,
393 we see ~~found~~ that the velocity-increase group saw statistically-significant increases in both
394 internal-rotation velocity and elbow-extension, whereas the velocity-decrease group saw no
395 significant change in either metric.

396 There was no significant change in elbow valgus torque, and the values reported in this study
397 are similar to previous studies (Feltner & Dapena, 1986; Fleisig et al., 2015)

398 No metrics were significantly different at front foot contact in any group.

399 A previous study found shoulder abduction angle at stride foot contact to be one of four
400 variables that could explain 97% of variance in valgus stress through a regression analysis
401 (Werner et al., 2002). In ~~this our~~ study, when comparing pre- and ~~postpost-training we -analysis~~
402 found no significant decrease in shoulder abduction angle at stride foot contact but a significant
403 change of abduction angle at ball release.

404 It has been suggested ~~previously~~ that the most optimal abduction angle at release is close to 90
405 degrees but may vary slightly depending on the individual- (Fortenbaugh, Fleisig & Andrews,
406 2009; Matsuo et al., 2002). The pitchers in ~~this our~~ study saw a significant change in shoulder
407 abduction angle at release ~~(from xx to xx)~~, moving closer to 90 degrees.

408 Notably, ~~ne of our sub-~~groups had significant changes in elbow valgus torque or shoulder
409 internal rotation torque ~~as a result of the training~~. The increase velocity group had a significant
410 increase in shoulder superior force, while the decrease velocity group had a significant increase
411 in shoulder adduction torque, and significant decreases in elbow anterior force, elbow
412 compressive force, elbow flexion torque, and shoulder compressive force.

Commented [MH43]: Increases? Decreases?

Commented [MH44]: How much?

Commented [MH45]: Abbreviated above. Consistency.

Commented [MH46]: This is not a paragraph.

Commented [MH47]: This is a statement. Tie it all together with a section on forces and or torques.

Commented [MH48]: Who's study? Werner or yours?

413 ~~The amount of e~~External rotation was not significantly different at front foot contact, but ~~was~~
414 significantly decreased at ball release, which may be a novel finding. This change was present
415 and significant in the combined and velocity increase group.

Commented [MH49]: Expand. Bring in other literature discuss why this may be novel and important.

416 Maximum shoulder adduction torque was significantly higher in the post-training group.
417 ~~Research has found that s~~Shoulder adduction torque ~~was-is~~ one of two variables related to
418 elbow valgus torque, along with maximum internal rotation torque (Sabick et al., 2004). ~~Sabick~~
419 ~~and colleagues~~ ~~The study~~ stated that maximum shoulder adduction torque and maximum
420 internal rotation torque were negatively correlated with elbow valgus torque, so as those two
421 values increased, elbow valgus torque tended to decrease.

422 Interestingly, ~~in our study~~, shoulder adduction torque only significantly increased in the group
423 that lost velocity. The group that increased velocity had an increase in shoulder adduction
424 torque, but it was not found to be significant.

Commented [MH50]: If it wasn't significant, can you really say they had an increase? Give p value. Was there a trend? If the increase was small and not significant be careful with your wording.

425 Maximum torso angular velocity and maximum pelvis angular velocity were not significantly
426 different in the pre- and post-group analysis. Split into increase and decrease velocity groups,
427 there were no significant changes in torso angular velocity or pelvis angular velocity.

Commented [MH51]: This doesn't contribute to the paper. Its just your results section repeated. At least combine it with below.

428 Previous research has shown mixed results on the relationship between pelvis- and torso-
429 angular velocity and throwing velocity, though none compared pre- and post-training periods
430 (Matsuo et al., 2001; Young, 2014; Dowling, 2016; Stodden et al., 2001). Theoretically,
431 increasing the rotational forces of the pelvis and torso allows energy to be transferred from the
432 trunk to the throwing arm and then to the ball, which should result in higher velocities.

Formatted: Space Before: 0 pt, After: 8 pt, Line spacing: Multiple 1.08 li

433 These studies would also suggest that peak torso and pelvis velocities play a role in increasing
434 velocity, but the timing of these forces ~~is~~ also vitally important. While the timing was not
435 examined in this study, further studies should examine the possible changes of constraint
436 training and weighted balls of the timing of hip and torso rotation. More research should also
437 be attempted at pre- and post-group analysis to look at hip and torso velocities.

Commented [MH52]: What forces? You are talking about torso and pelvis velocity.

Commented [MH53]: Of what?

438 ~~The degree of e~~Elbow flexion at ball release did not significantly change, even though a
439 previous study found significant differences in the angle of the elbow at ball release, depending
440 on ball weight (van den Tillaar & Ettema, 2004).

Commented [MH54]: More is needed in this section. You can't just state your findings, then list a study that found something similar (or different). Expand. Based on this, what do you guys think. Was the lack of differences all a result of your training program? Give some speculation.

441 It has also been postulated that training with weighted balls causes ~~gains-increases~~ in external
442 rotation, both passive and dynamic. Dynamic maximum shoulder ER has been associated with
443 ball velocity (Matsuo et al., 2001; Werner et al., 2008), but research looking within pitcher
444 variation found no significant association between maximum external rotation and ball velocity
445 (Stodden et al., 2005). The theory holds that weighted-ball use may result in velocity gains from
446 excess glenohumeral external rotation, which may be linked to increased elbow valgus load
447 (Aguinaldo & Chambers, 2009; Sabick et al., 2004).

Commented [MH55]: Again, expand on this. Why didn't you find a difference? What was different in the vanden Tillaar study that may have led to the discrepancy? Dig deeper.

Commented [MH56]: ER. Be consistent.

448 Although previous research on high-school pitchers did not find a significant correlation
449 between passive external rotation and pitch velocity (Keller, 2015), other research did see a
450 significant moderate correlation between passive external rotation and the degree of external
451 rotation seen in a throw (Miyashita et al., 2008).

452 It should be noted that the biomechanical measurement of external rotation cannot be
453 attributed only to changes of the glenohumeral joint. There can be changes in thoracic
454 extension or scapula position that can affect measurements. In addition, the possibility of
455 measurement error may also play a role, although the process was standardized in our work
456 during both the pre and post training/testing.

457 The Our subjects in this study did see a passive range-of-motion increase of 1.7 degrees in the
458 dominant arm, but the findings were statistically not insignificant. The non-dominant arm saw a
459 lesser, but still statistically insignificant, finding of 0.1 degrees.

461 Having broken up velocity into into increase and decrease groups, we can see the increase
462 group had an increase in external rotation of $2.8 \pm SD$ degrees while the decrease velocity
463 group saw an increase of $0.6 \pm SD$ degrees. Interestingly, there were wide swings in the non-
464 dominant arm external rotation. The velocity-increase group saw an increase in non-dominant
465 external rotation of $7.8 \pm SD$ degrees while the velocity decrease group saw a decrease of $8.6 \pm$
466 SD degrees. This may bring into question what part of the changes in the dominant arm can be
467 attributed to throwing and what parts can be attributed to non-throwing work, such as mobility
468 or strength work, as it seems the change in non-dominant ROM came from mobility or strength
469 work.

470 Although the finding of increased external rotation ER in the dominant arm was not statistically
471 significant, it should still be considered an interesting finding since it has been suggested that
472 humans have adapted to having more external rotation ER in order to better store elastic
473 energy and increase power (Roach et al., 2013).

474 It has been hypothesized that training with weighted baseballs would result in negative
475 anatomical effects, such as increased external rotation ER on top of other effects similar to
476 pitching. The Our findings of this study are interesting because the range-of-motion findings
477 results are dissimilar/reject our to that hypothesis and to most short- and long-term range-of-
478 motion studies.

479 Many of the pitchers in the study performed training days, which were either bullpens or
480 training with weighted balls, designed to replicate high-intent pitching. The acute effects of
481 range-of-motion on weighted balls have not been studied, but there has been research on
482 acute changes of pitching and bullpens. It has been hypothesized that range-of-motion changes

Commented [MH57]: SD?

Formatted: Space Before: 10 pt, After: 0 pt, Line spacing: Multiple 1.15 li

Commented [MH58]: Measurement error and likely no need to discuss.

Commented [MH59]: What does this mean? Large variability?

Commented [MH60]: Good. This type of thinking should be sprinkled throughout.

Commented [MH61]: Elbow? Shoulder?

Commented [MH62]: By who?

Commented [MH63]: ?

Commented [MH64]: By who?

483 that occur in the short-term may be exacerbated over the long-term. But the research
484 conclusions of both short- and long-term ROM changes vary.

485 Two studies ~~looking investigating at~~ the acute effects of pitching on range of motion found a
486 loss of internal rotation that was sustained for 24 or 72 hours (Reinold et al., 2008; Kibler,
487 Sciascia & Moore, 2012).

488 Counter to these ~~above~~ studies, Freehill et al. (2014) found ~~that~~ a single start resulted in no
489 significant change in IR but rather a significant increase in passive external rotation after
490 pitching in a game.

491 Another study on minor league pitching starts found both a significant decrease in internal
492 rotation, significant gain in external rotation, and significant gain in total arm range of motion
493 (Case et al., 2015). Twenty-four hours after pitching, IR returned to pre-game baseline while ER
494 was still significantly greater.

495 Long-term studies examining range of motion have also found conflicting results in internal
496 rotation and external rotation when compared to ~~this our work study~~. Freehill et al. (2011)
497 found an increase in external rotation and internal rotation that was not statistically significant.
498 This study has a similar sample size (21 pitchers, over 29 individual seasons) compared to the 17
499 pitchers in our study. Freehill et al. (2011) ~~study lasted was four months in duration, over the~~
500 ~~course of a baseball season, four months long,~~ compared to ~~the~~ six weeks ~~that in~~ our study
501 ~~lasted. Freehill and colleagues also had their pitchers. These pitchers also~~ performed a capsule-
502 stretching program during the season. Stretching programs have been seen to have positive
503 effects on pitchers, such as reducing the ~~chance likelihood~~ of ~~a loss of in~~ internal rotation
504 (Lintner et al., 2007).

505 ~~Additionally, in a follow up study, Freehill and colleagues A different study~~ found that
506 preseason and postseason measurements resulted in significantly more ER, significantly less IR,
507 and significantly less total range of motion (Freehill et al., 2014).

508 A study on baseball and softball athletes found no change in internal rotation over the course
509 of a season but did find increased external rotation and total range of motion (Dwelly et al.,
510 2009).

511 These long-term studies align with the acute studies, to the extent ~~in~~ that the most common
512 adaptations to throwing are a loss of internal rotation and a gain of external rotation, though
513 the ~~degree magnitude~~ of change varies.

514 It is unknown exactly why these long-term studies differ, but it could likely be attributed to
515 ~~something differences~~ in the training program outside of throwing. It should be noted that
516 none of these long-term studies found a significant increase in internal rotation in the throwing
517 arm.

Commented [MH65]: Make sure all abbreviations are consistent through paper.

Commented [MH66]: So, they found no increase.....

Considering stating is like this....

Freehill et al. (2011) found no significant increases in ER and IR.

518 This ~~w~~could suggest that range of motion is ~~a~~ fluid measurement. Further research should
519 attempt to examine if there is an acceptable range of internal and external measurements.

Commented [MH67]: What does this mean?

520 ~~It has been hypothesized that t~~A ~~he~~ loss of internal rotation may be caused by the eccentric
521 muscle contraction that occurs in the posterior shoulder during the follow-through of pitching
522 (Proske & Morgan, 2001). It is possible that no decreases were seen in ~~our work for~~ dominant
523 arm internal range of motion because of the daily soft-tissue work that each pitcher completed.
524 Although the exact causes of self-myofascial release are unknown, research has suggested SMR
525 has positive short-term effects on range of motion without negatively affecting muscle
526 performance (Cheatham et al., 2015).

Commented [MH68]: Great work.

527 As mentioned previously, the pitchers had access to instrument-assisted soft-tissue
528 mobilization (IASTM) on an as-needed basis. Previous research on baseball players found that
529 some acute ROM losses could be attributed to muscular/rotator-cuff stiffness, and IASTM plus
530 stretching displayed greater gains in internal rotation than in self-stretching alone (Bailey et al.,
531 2015). The gains in that study were attributed to decreased rotator-cuff stiffness and humeral
532 retrotorsion, but not joint translation.

533 More specifically, one study comparing IASTM and self-stretching saw a greater increase in
534 ~~shoulder~~ internal rotation and total range of motion when compared with self-stretching alone;
535 ~~gains which is~~ similar to those found in ~~this-our~~ study (Bailey et al., 2017). This would suggest
536 that soft-tissue work such as IASTM played a role in the increase in internal rotation and total
537 range of motion that was seen in ~~this-our~~ study.

538 Proske & Morgan (2001) also hypothesized that because injuries can occur from eccentric
539 exercise, a way to combat injury risk would be to perform an eccentric-exercise program to
540 strengthen and, therefore, protect the muscles. Eccentric training in this program ~~occurred~~
541 while using wrist weights, j-band external and internal rotations, rebounders, and upward
542 tosses. ~~But-However, to our knowledge,~~ wrist-weight exercises, and the other exercises, have
543 not been studied in the literature for their effects on strength or range-of-motion effects.

544 Pitchers also performed daily exercises in the warm-up and throwing program that are
545 designed to work the posterior shoulder concentrically: specifically, Jaeger band exercises and
546 reverse throws with PlyoCare balls. The effects that long-term concentric exercise has on
547 posterior shoulder strength and range of motion have also not been studied.

548 A previous study found that performing a series of short-duration stretching/calisthenics drills
549 (titled the Two-Out drill) resulted in short-term deficits in range of motion caused by pitching to
550 be restored to their pre-pitching levels (Rafael et al., 2017). The post-throwing exercise circuit
551 used in ~~this-our~~ study did not contain the same exercises; the exercises in ~~this-our~~ study
552 ~~were~~ strength-based, not stretching/calisthenic based. However, ~~it we do~~ show
553 evidence that possible deficits created by throwing may ~~be brought back to~~ return to baseline
554 by stretching or exercise. Further studies should examine the effect that the post-throwing

555 exercise circuit and the use of concentric and isometric exercise ~~on the shoulder used in this~~
556 ~~study have~~ might have on shoulder ~~on~~ range of motion.

557 It is unlikely that the use of the Marc Pro EMS device had an effect on range of motion. ~~Since it~~
558 has been ~~seen suggested~~ that pitchers see reduced blood flow in their throwing arms, ~~and the~~
559 Marc Pro is used to encourage blood flow, but that increase ~~in blood flow would not likely may~~
560 ~~not~~ result in changes in range of motion (Laudner et al., 2014). A study comparing different
561 recovery techniques found that EMS resulted in a lower rating of perceived exertion and blood-
562 lactate concentration, but no change in range-of-motion ~~testing~~ (Warren, Szymanski & Landers,
563 2015). It's unknown whether the different EMS devices used in the Warren et al. (2015) would
564 result in similar results.

565 A significant increase in internal rotation of the dominant arm may be seen as a positive since it
566 has been suggested that losses of internal rotation in the throwing arm may lead to a higher
567 risk of injury (Wilk et al., 2011; Myers et al., 2006; Dines et al., 2009). A study on pitchers in
568 Japan ~~did find~~ found a relationship between more IR range of motion in their dominant arms
569 and injury (Sueyoshi et al., 2017). Sueyoshi et al. included a wider range of athletes (Little
570 League to college age) than in this study, and younger athletes have been seen to have greater
571 IR ROM than older athletes, which may have affected the results (Astolfi et al., 2015). The
572 injured group in Sueyoshi et al. also pitched in more games and more innings than the no-injury
573 group.

574 The pitchers in both the pre and post measurements of ~~this our~~ study would not qualify for
575 either measurement of GIRD, even though the difference between non-dominant and
576 dominant arms increased (Burkhart, Morgan & Kibler, 2003). This increase in the difference
577 between internal rotation of the non-dominant and dominant arms was driven by larger
578 increases in internal rotation range of motion in the non-dominant arm than in the dominant
579 arm.

580 The concept of total range of motion (TROM) has also been introduced to examine whether
581 differences between arms may lead to injuries (Wilk, Meister & Andrews, 2002). In this study,
582 TROM saw significant increases in both the dominant and the non-dominant arm. Both arms
583 saw larger ~~gains increases~~ in internal rotation compared to external rotation.

584 Furthermore, neither the pre- or post-ROM measurements qualify for either external rotation
585 deficit (external rotation at least 5 degrees more in the dominant arm when compared to the
586 non-dominant arm) or TROM deficit (when TROM of the non-dominant arm is at least 5 degrees
587 more than that the dominant arm). Pitchers with insufficient external rotation (<5 greater
588 external rotation in throwing shoulder than non-dominant shoulder) have been seen to be
589 more likely to have a shoulder injury (Wilk et al., 2015). Pitchers with deficits equal to or
590 greater than 5 degrees in total rotation in their throwing shoulders compared to their non-
591 dominant arms have been viewed as at higher risk of injuries (Wilk et al., 2014).

Commented [MH69]: This is out of place. You discuss IR and ER above. Then bring in this discussion. Follow up paragraph goes back to IR.

Commented [MH70]: Glenohumeral Internal Rotation Deficit? This has not been defined yet.

592 One thing unknown about ~~these prospective studies~~ is if the problem of deficits, by comparing
593 the dominant to non-dominant arm, holds under longer term tracking and possible changes in
594 the non-dominant arm. For example, a pitcher may qualify for a deficit while having no change
595 of ROM in the dominant arm but see a significant change in the non-dominant arm. Even
596 though both dominant and non-dominant TROM gained in this study, the non-dominant arm
597 had a greater range of motion than the dominant arm post training.

Commented [MH71]: Who? Wilk?

598 When examining bilateral differences in range of motion over time, researchers should take
599 note of whether the changes are coming from the dominant or non-dominant arm. Many of the
600 changes in range of motion are focused on comparing from throwing and the dominant arm,
601 but significant changes in range of motion in the non-dominant arm, as seen in this study, show
602 that there can be large changes that don't come from throwing.

603 Humeral retroversion was not measured in this study, although ~~it has been said that~~ this could
604 partially explain the range-of-motion differences between the dominant and non-dominant arm
605 (Chant et al., 2007). There is also research suggesting that humeral torsion adaptations occur
606 pre-high school, suggesting that changes in this study came from soft tissue adaptations
607 (Oyama, Hibberd & Myers, 2013). Further research examining range-of-motion changes and
608 weighted-ball training should attempt to measure humeral retroversion, as well as range of
609 motion.

611 This study is one of ~~only a few that have included training programs, the first papers to be~~
612 ~~considered for publication regarding a training period,~~ and as such, there is little data to
613 compare ~~it to~~. The throwing velocity ~~of for our this group is was very competitive with studies~~
614 ~~comparing similar subject pools comparable to other work,~~ with ~~the an~~ average initial pitching
615 velocity ~~of the seventeen pitchers being of~~ 35.1 +/- 1.8 m/s ~~(78.6 mph)~~; Fleisig et al. (2017) had
616 a group of similar amateur pitchers (n=25) with an average pitching velocity of 34.2 +/- 2.0 m/s
617 ~~(76.5 mph)~~. Fleisig et al.'s study of underweight and overweight baseball throwing showed
618 variations in arm kinetics, variations in angular velocities, and relatively small changes in body
619 positions; ~~These changes could be reflective of could be considered~~ reasonable training
620 modalities for pitchers (Fleisig et al., 2017).

Commented [MH72]: You can't say things like this.

621 ~~This Our data study would~~ also suggests that pitching mechanics can be changed over a six-
622 week training period. A previous study by Fleisig et al. (2017b) found that pitchers can change
623 their mechanics based off a biomechanical observation over periods of time ranging from 2-48
624 months. In ~~this our~~ study, the initial screenings were not given to players with specific direction
625 to change mechanics; the screening was purposefully ~~observatory~~.

Commented [MH73]: More details here. You say your data suggests that pitching mechanics can be changed over a 6-week period. Why do you think this? How was it changed?

626 This paper included fourteen right-handed and three left-handed pitchers. Further research
627 should examine the differences of weighted-ball training between right- and left-handed
628 pitchers, as previous research has suggested differences in range of motion, humeral
629 retroversion, and biomechanics depending on the dominant throwing arm (Solomito, Ferreira &

630 Nissen, 2017; Werner et al., 2010; Takenaga et al., 2017). It is therefore possible that pitchers
631 should have different throwing, mobility, and strength programs depending on which arm is
632 dominant.

633 Limitations

634 The pitchers in this study were asked to throw as hard as comfortable on their testing days.
635 That, combined with the unfamiliarity of wearing biomechanical markers, resulted in lower
636 velocities than what would be seen in a game or training environment.

637 Range-of-motion measurements were taken during the training period, so there could be
638 unknown effects from measurements taken at different times. Range-of-motion measurements
639 were also taken in a way that differs from other papers studies. Since the same subject
640 measured every range-of-motion test, the results are should be reliable to each other, but may
641 not be directly comparable to other papers studies.

642 In addition, not every pitcher in our study had the same training background. ;sSome had been
643 training in-person at our facility Driveline for a few weeks while others were assessed within
644 their first week. However, the vast majority of them participants had previous experience
645 training with weighted balls so, while hard to quantify, previous training was less of a potential
646 confounding variable than it would might have been in other situations. for other research
647 questions.

648 Lastly, this study involved a small sample size of seventeen pitchers and references smaller
649 groups of velocity increase and velocity decrease groups.

650

651 *Conclusion*

652 This study contradicts the original hypothesis, which proposed -involved speculation that thea
653 6-week training program would increase pitching velocity, arm angular velocities, joint kinetics,
654 and arm range of motion. There were few changes comparing the pre- and post- groups, most
655 notably there was no significant increases in elbow valgus or shoulder internal rotation torque
656 and no significant increase in external rotation of the dominant arm. When sub-groups were
657 created based on velocity, Tthe velocity increase group had statistically significant increases in
658 internal rotation and elbow extension angular velocities.

659 This study contradicts the premise that weighted-implement training leads to rapid gains in
660 shoulder external range of motion (Reinold, M. 2017). Literature on the topic of restoring
661 shoulder internal rotation range of motion is supported (Laudner et al., 2008), but further
662 research is required into individual modalities that may be contributing to these physical
663 adaptations.

664 Limitations

Commented [MH74]: Was your study under powered?

665 ~~The pitchers in this study were asked to throw as hard as comfortable on their testing days.~~
666 ~~That, combined with the unfamiliarity of wearing biomechanical markers, resulted in lower~~
667 ~~velocities than what would be seen in a game or training environment.~~

668 ~~Range of motion measurements were taken during the training period, so there could be~~
669 ~~unknown effects from measurements taken at different times. Range of motion measurements~~
670 ~~were also taken in a way that differs from other papers. Since the same subject measured every~~
671 ~~range of motion test, the results are reliable to each other but may not be directly comparable~~
672 ~~to other papers.~~

673 ~~In addition, not every pitcher had the same training background; some had been training in~~
674 ~~person at Driveline for a few weeks while others were assessed within their first week.~~
675 ~~However, the vast majority of them had previous experience training with weighted balls so,~~
676 ~~while hard to quantify, previous training was less of a potential confounding variable than it~~
677 ~~would have in other situations.~~

678 ~~Lastly, this study involved a small sample size of seventeen pitchers and references smaller~~
679 ~~groups of velocity increase and velocity decrease groups.~~

680 **References**

- 681 Aguinaldo AL., Chambers H. 2009. Correlation of Throwing Mechanics With Elbow Valgus Load
682 in Adult Baseball Pitchers. *The American Journal of Sports Medicine* 37:2043–2048. DOI:
683 10.1177/0363546509336721.
- 684 Astolfi MM., Struminger AH., Royer TD., Kaminski TW., Swanik, CB. 2015. Adaptations of the
685 Shoulder to Overhead Throwing in Youth Athletes. *Journal of Athletic Training* 50:726–732. DOI:
686 10.4085/1062-6040-50.1.14.
- 687 Baheti N. 2000. The Effectiveness of a Surgical Tubing Strengthening Program in the
688 Maintenance of Performance Among High School Pitchers. Masters Thesis Thesis. Oregon State.
- 689 Bailey LB., Shanley E., Hawkins R., Beattie PF., Fritz S., Kwartowitz D., Thigpen CA. 2015.
690 Mechanisms of Shoulder Range of Motion Deficits in Asymptomatic Baseball Players. *The*
691 *American Journal of Sports Medicine* 43:2783–2793. DOI: 10.1177/0363546515602446.
- 692 Bailey LB., Thingpen CA., Hawkins RJ., Beattie PR., Shanley E. 2017 Effectiveness of Manual
693 Therapy and Stretching for Baseball Players with Shoulder Range of Motion Deficits *Sports*
694 *Health* 9:320-237 DOI: 10.1177/1941738117702835
- 695 Boone DC., Azen SP., Lin CM., Spence C., Baron C., Lee L. 1978. Reliability of goniometric
696 measurements. *Physical Therapy* 58:1355–1360.
- 697 Burkhart SS., Morgan CD., Kibler WB. 2003. The disabled throwing shoulder: Spectrum of
698 pathology part I: Pathoanatomy and biomechanics. *Arthroscopy: The Journal of Arthroscopic &*
699 *Related Surgery* 19:404–420. DOI: 10.1053/jars.2003.50128.

700 Carter AB., Kaminski TW., Douex Jr AT., Knight CA., Richards JG. 2007. Effects of high volume
701 upper extremity plyometric training on throwing velocity and functional strength ratios of the
702 shoulder rotators in collegiate baseball players. *Journal of Strength and Conditioning Research*
703 21:208.

704 Case JM., Mannava S., Fallin JH., Stone AV., Freehill MT. 2015. Acute changes in glenohumeral
705 range-of-motion following in-season minor league pitching starts. *The Physician and*
706 *Sportsmedicine* 43:360–365. DOI: 10.1080/00913847.2015.1059249.

707 Chant CB., Litchfield R., Griffin S., Thain LMF. 2007. Humeral Head Retroversion in Competitive
708 Baseball Players and Its Relationship to Glenohumeral Rotation Range of Motion. *Journal of*
709 *Orthopaedic & Sports Physical Therapy* 37:514–520. DOI: 10.2519/jospt.2007.2449.

710 Cheatham SW., Kolber MJ., Cain M., Lee M. 2015. THE EFFECTS OF SELF-MYOFASCIAL RELEASE
711 USING A FOAM ROLL OR ROLLER MASSAGER ON JOINT RANGE OF MOTION, MUSCLE
712 RECOVERY, AND PERFORMANCE: A SYSTEMATIC REVIEW. *International Journal of Sports*
713 *Physical Therapy* 10:827–838.

714 DeRenne C, Buxton BP, Hetzler RK, and Ho KW. 1994. Effects of under- and overweighted
715 implement training on pitching velocity. *Journal of Strength and Conditioning Research* 8: 247-
716 250.

717 DeRenne C. 1985. Increasing throwing velocity. *Athletic Journal* 65: 36-39.

718 DeRenne C, Ho KW, and Blitzblau A. 1990. Effects of weighted implement training on throwing
719 velocity. *Journal of Applied Sports Science Research* 4: 16-19.

720 Dines JS., Frank JB., Akerman M., Yocum LA. 2009. Glenohumeral Internal Rotation Deficits in
721 Baseball Players with Ulnar Collateral Ligament Insufficiency. *The American Journal of Sports*
722 *Medicine* 37:566–570. DOI: 10.1177/0363546508326712.

723 DiNubile N., Westcott WL., Reinl G., Bajaj A., Braverman ER., Madigan MA., Giordano J., Blum K.
724 2011. The Marc Pro Device is a Novel Paradigm Shift in Muscle Conditioning, Recovery and
725 Performance: Induction of Nitric Oxide (NO) Dependent Enhanced Microcirculation Coupled
726 with Angiogenesis Mechanisms. *Journal of Exercise Physiology Online* 14.

727 Dowling, Brittany & Pearl, Caitlin & Laughlin, Walter & Fleisig, Glenn. (2016). Relationship of
728 pelvis and trunk kinematics to ball velocity in professional baseball pitchers.

729 Dwelly PM., Tripp BL., Tripp PA., Eberman LE., Gorin S. 2009. Glenohumeral Rotational Range of
730 Motion in Collegiate Overhead-Throwing Athletes During an Athletic Season. *Journal of Athletic*
731 *Training* 44:611–616. DOI: 10.4085/1062-6050-44.6.611.

732 Egstrom GH, Logan GA, and Wallis EL. 1960. Acquisition of throwing skill involving projectiles of
733 varying weight. *Res Q* 31: 420-425.

734 Escamilla RF, Fleisig GS, Barrentine SW, Zheng N, Andrews JR. 1998. Kinematic Comparisons of
735 Throwing Different Types of Baseball Pitches. *Journal of Applied Biomechanics* 14:1-23 DOI:
736 10.1123/jab.14.1.1

737 Escamilla RF, Speer KP, Fleisig GS, Barrentine SW, Andrews JR. 2000. Effects of throwing
738 overweight and underweight baseballs on throwing velocity and accuracy. *Sports Med.* 2000;
739 29:259-272.

740 Feltner M., Dapena J. 1986. Dynamics of the shoulder and elbow joints of the throwing arm
741 during a baseball pitch. *International Journal of Sport Biomechanics* 2:235-259.

742 Fleisig GS., Bolt B., Fortenbaugh D., Wilk KE., Andrews JR. 2011. Biomechanical Comparison of
743 Baseball Pitching and Long-Toss: Implications for Training and Rehabilitation. *Journal of*
744 *Orthopaedic & Sports Physical Therapy* 41:296-303. DOI: 10.2519/jospt.2011.3568.

745 Fleisig GS., Diffendaffer AZ., Aune KT., Ivey B., Laughlin WA. 2017. Biomechanical Analysis of
746 Weighted-Ball Exercises for Baseball Pitchers. *Sports Health: A Multidisciplinary Approach*
747 9:210-215. DOI: 10.1177/1941738116679816.

748 Fleisig GS., Diffendaffer AZ., Ivey B., Aune KT. 2017. Do baseball pitchers improve mechanics
749 after biomechanical evaluations? *Sports Biomechanics*:1-8. DOI:
750 10.1080/14763141.2017.1340508.

751 Fleisig GS., Leddon CE., Laughlin WA., Ciccotti MG., Mandelbaum BR., Aune KT., Escamilla RF.,
752 MacLeod TD., Andrews JR. 2015. Biomechanical Performance of Baseball Pitchers With a
753 History of Ulnar Collateral Ligament Reconstruction. *The American Journal of Sports Medicine*
754 43:1045-1050. DOI: 10.1177/0363546515570464.

755 Fortenbaugh D., Fleisig GS., Andrews JR. 2009. Baseball Pitching Biomechanics in Relation to
756 Injury Risk and Performance. *Sports Health: A Multidisciplinary Approach* 1:314-320. DOI:
757 10.1177/1941738109338546.

758 Freehill MT., Ebel BG., Archer KR., Bancells RL., Wilckens JH., McFarland EG., Cosgarea AJ. 2011.
759 Glenohumeral Range of Motion in Major League Pitchers. *Sports Health* 3:97-104. DOI:
760 10.1177/1941738110374627.

761 Freehill MT., Archer KR., Diffenderfer BW., Ebel BG., Cosgarea AJ., McFarland EG. 2014. Changes
762 in Collegiate Starting Pitchers' Range of Motion After Single Game and Season. *The Physician*
763 *and Sportsmedicine* 42:69-74. DOI: 10.3810/psm.2014.02.2049.

764 Fortun CM., Davies GJ., Kernozck TW. 1998. The effects of plyometric training on the shoulder
765 internal rotators. *Phys Ther* 78:S87.

766 Keller R., Marshall N., Mehran N., Moutzouros V., 2015 Pitching Speed and Glenohumeral
767 Adaptation in High School Pitchers. *Orthopedics* 38 e668-e672 DOI: 10.3928/01477447-
768 20150804-52.

769 Kibler WB., Sciascia A., Moore S. 2012. An Acute Throwing Episode Decreases Shoulder Internal
770 Rotation. *Clinical Orthopaedics and Related Research* 470:1545–1551. DOI: 10.1007/s11999-
771 011-2217-z.

772 Laudner KG., Selkow NM., Burke NC., Lynall RC., Meister K. 2014. Decreased blood flow in the
773 throwing arm of professional baseball pitchers. *Journal of Shoulder and Elbow Surgery* 23:1753–
774 1756. DOI: 10.1016/j.jse.2014.03.007.

775 Laudner KG, Sipes RC, Wilson JT. 2008. The acute effects of sleeper stretches on shoulder range
776 of motion. *Journal of Athletic Training*. 43:359-363. doi:10.4085/1062-6050-43.4.359

777 Lintner D., Mayol M., Uzodinma O., Jones R., Labossiere D. 2007. Glenohumeral Internal
778 Rotation Deficits in Professional Pitchers Enrolled in an Internal Rotation Stretching Program.
779 *The American Journal of Sports Medicine* 35:617–621. DOI: 10.1177/0363546506296736.

780 Matsuo T., Escamilla RF., Fleisig GS., Barrentine SW., Andrews JR. 2001. Comparison of
781 Kinematic and Temporal Parameters between Different Pitch Velocity Groups. *Journal of*
782 *Applied Biomechanics* 17:1–13. DOI: 10.1123/jab.17.1.1.

783 Matsuo T., Matsumoto T., Mochizuki Y., Takada Y., Saito K. 2002. Optimal Shoulder Abduction
784 Angles during Baseball Pitching from Maximal Wrist Velocity and Minimal Kinetics Viewpoints.
785 *Journal of Applied Biomechanics* 18:306–320. DOI: 10.1123/jab.18.4.306.

786 Miyashita K., Urabe Y., Kobayashi H., Yokoe K., Koshida S., Kawamura M., Ida K. 2008.
787 Relationship Between Maximum Shoulder External Rotation Angle During Throwing and
788 Physical Variables. *Journal of Sports Science & Medicine* 7:47–53.

789 Myers JB., Pasquale MR., Laudner KG., Sell TC., Bradley JP., Lephart SM. 2005. On-the-Field
790 Resistance-Tubing Exercises for Throwers: An Electromyographic Analysis. *Journal of Athletic*
791 *Training* 40:15–22.

792 Myers JB., Laudner KG., Pasquale MR., Bradley JP., Lephart SM. 2006. Glenohumeral Range of
793 Motion Deficits and Posterior Shoulder Tightness in Throwers with Pathologic Internal
794 Impingement. *The American Journal of Sports Medicine* 34:385–391. DOI:
795 10.1177/0363546505281804.

796 Oyama S., Hibberd EE., Myers JB. 2013. Changes in humeral torsion and shoulder rotation range
797 of motion in high school baseball players over a 1-year period. *Clinical Biomechanics* 28:268–
798 272. DOI: 10.1016/j.clinbiomech.2013.01.014.

799 Proske U., Morgan DL. 2001. Muscle damage from eccentric exercise: mechanism, mechanical
800 signs, adaptation and clinical applications. *The Journal of Physiology* 537:333–345. DOI:
801 10.1111/j.1469-7793.2001.00333.x.

802 Rafael E., Kyle Y., Tony M., Jeff C., Keith L., James A. 2017. Effects of a Short-Duration Stretching
803 Drill After Pitching on Elbow and Shoulder Range of Motion in Professional Baseball Pitchers.
804 *American Journal of Sports Medicine* 45:692–700.

805 Reinold, Mike & E Wilk, Kevin & Fleisig, Glenn & Zheng, Naiquan & W Barrentine, Steven &
806 Chmielewski, Terri & C Cody, Rayden & G Jameson, Gene & Andrews, James. (2004).
807 Electromyographic Analysis of the Rotator Cuff and Deltoid Musculature During Common
808 Shoulder External Rotation Exercises. *The Journal of orthopaedic and sports physical therapy*.
809 34. 385-94. 10.2519/jospt.2004.34.7.385.

810 Reinold MM., Wilk KE., Macrina LC., Sheheane C., Dun S., Fleisig GS., Crenshaw K., Andrews JR.
811 2008. Changes in Shoulder and Elbow Passive Range of Motion after Pitching in Professional
812 Baseball Players. *The American Journal of Sports Medicine* 36:523–527. DOI:
813 10.1177/0363546507308935.

814 Roach NT., Venkadesan M., Rainbow MJ., Lieberman DE. 2013. Elastic energy storage in the
815 shoulder and the evolution of high-speed throwing in Homo. *Nature* 498:483–486. DOI:
816 10.1038/nature12267.

817 Robertson D., E Caldwell G., Hamill J., Kamen G., N Whittlesey S. *Research Methods in*
818 *Biomechanics*. 2nd ed., Human Kinetics, 2014.

819 Sabick MB., Torry MR., Lawton RL., Hawkins RJ. 2004. Valgus torque in youth baseball
820 pitchers: a biomechanical study. *Journal of Shoulder and Elbow Surgery* 13:349–355. DOI:
821 10.1016/j.jse.2004.01.013.

822 Slenker NR., Limpisvasti O., Mohr K., Aguinaldo A., ElAttrache NS. 2014. Biomechanical
823 Comparison of the Interval Throwing Program and Baseball Pitching: Upper Extremity Loads in
824 Training and Rehabilitation. *The American Journal of Sports Medicine* 42:1226–1232. DOI:
825 10.1177/0363546514526152.

826 Solomito MJ., Ferreira JV., Nissen CW. 2017. Biomechanical differences between left- and right-
827 handed baseball pitchers. *Sports Biomechanics* 16:143–151. DOI:
828 10.1080/14763141.2016.1186725.

829 Stodden DF., Fleisig GS., McLean SP., Andrews JR. 2005. Relationship of biomechanical factors
830 to baseball pitching velocity: within pitcher variation. *Journal of applied biomechanics* 21:44–
831 56.

832 Stodden DF., Fleisig GS., McLean SP., Lyman SL., Andrews JR. 2001. Relationship of pelvis and
833 upper torso kinematics to pitched baseball velocity. *Journal of applied biomechanics* 17:164–
834 172.

835 Stone AV., Mannava S., Patel A., Marquez-Lara A., Freehill MT. 2017. Defining the Long-Toss: A
836 Professional Baseball Epidemiological Study. *Orthopaedic Journal of Sports Medicine* 5. DOI:
837 10.1177/2325967116686773.

838 Sueyoshi T., Nakatani T., Tsuruta T., Emoto G. 2017. Upper Extremity Range of Motion and
839 Pitching Profile of Baseball Pitchers in Japan. *Orthopaedic Journal of Sports Medicine* 5. DOI:
840 10.1177/2325967117715436.

841 Swanik KA., Lephart SM., Swanik CB., Lephart SP., Stone DA., Fu FH. 2002. The effects of
842 shoulder plyometric training on proprioception and selected muscle performance
843 characteristics. *Journal of Shoulder and Elbow Surgery* 11:579–586. DOI:
844 10.1067/mse.2002.127303.

845 Takenaga T., Goto H., Sugimoto K., Tsuchiya A., Fukuyoshi M., Nakagawa H., Nozaki M.,
846 Takeuchi S., Otsuka T. 2017. Left-handed skeletally mature baseball players have smaller
847 humeral retroversion in the throwing arm than right-handed players. *Journal of Shoulder and*
848 *Elbow Surgery* 26:2187–2192. DOI: 10.1016/j.jse.2017.07.014.

849 Tillaar R. van den, Ettema G. 2004. A Force-Velocity Relationship and Coordination Patterns in
850 Overarm Throwing. *Journal of Sports Science & Medicine* 3:211–219.

851 Tillaar R van den., Ettema G. 2011. A comparison of kinematics between overarm throwing with
852 20% underweight, regular, and 20% overweight balls. *Journal of applied biomechanics* 27:252–
853 257.

854 Thewlis D., Bishop C., Daniell N., Paul G. 2013. Next-generation low-cost motion capture
855 systems can provide comparable spatial accuracy to high-end systems. *Journal of applied*
856 *biomechanics* 29:112–117.

857 Warren CD., Szymanski DJ., Landers MR. 2015. Effects of Three Recovery Protocols on Range of
858 Motion, Heart Rate, Rating of Perceived Exertion, and Blood Lactate in Baseball Pitchers During
859 a Simulated Game. *The Journal of Strength & Conditioning Research* 29:3016–3025.

860 Werner SL., Guido JA., Delude NA., Stewart GW., Greenfield JH., Meister K. 2010. Throwing Arm
861 Dominance in Collegiate Baseball Pitching: A Biomechanical Study. *The American Journal of*
862 *Sports Medicine* 38:1606–1610. DOI: 10.1177/0363546510365511.

863 Werner SL., Murray TA., Hawkins RJ., Gill TJ. 2002. Relationship between throwing mechanics
864 and elbow valgus in professional baseball pitchers. *Journal of Shoulder and Elbow Surgery*
865 11:151–155. DOI: 10.1067/mse.2002.121481.

866 Werner SL., Suri M., Guido JA., Meister K., Jones DG. 2008. Relationships between ball velocity
867 and throwing mechanics in collegiate baseball pitchers. *Journal of Shoulder and Elbow Surgery*
868 17:905–908. DOI: 10.1016/j.jse.2008.04.002.

869 Westcott WL., Chen T., Neric FB., DiNubile N., Bowirrat A., Madigan M., Downs BW., Giordano
870 JMS., Chen A., Bajaj A. 2011. The Marc Pro™ device improves muscle performance and
871 recovery from concentric and eccentric exercise induced muscle fatigue in humans: A pilot
872 study. *JEPonline* 14:55–67.

873 Westcott W., Han D., DiNubile N., Neric F., Loud RLR., Whitehead S., Blum K. 2013. Effects of
874 Electrical Stimulation Using the Marc Pro™ Device during the Recovery Period on Calf Muscle
875 Strength and Fatigue in Adult Fitness Participants. *Journal of Exercise Physiology Online* 16:40–
876 50.

877 Wilk KE., Macrina LC., Fleisig GS., Porterfield R., Charles D. Simpson I., Harker P., Paparesta N.,
878 Andrews JR. 2011. Correlation of glenohumeral internal rotation deficit and total rotational
879 motion to shoulder injuries in professional baseball pitchers. *American Journal of Sports*
880 *Medicine*. 39:329-335. doi:10.1177/036354651038422325.

881 Wilk KE., Macrina LC., Fleisig GS., Aune KT., Porterfield RA., Harker P., Evans TJ., Andrews JR.
882 2014. Deficits in Glenohumeral Passive Range of Motion Increase Risk of Elbow Injury in
883 Professional Baseball Pitchers: A Prospective Study. *The American Journal of Sports Medicine*
884 42:2075–2081. DOI: 10.1177/0363546514538391.

885 Wilk KE., Macrina LC., Fleisig GS., Aune KT., Porterfield RA., Harker P., Evans TJ., Andrews JR.
886 2015. Deficits in Glenohumeral Passive Range of Motion Increase Risk of Shoulder Injury in
887 Professional Baseball Pitchers: A Prospective Study. *The American Journal of Sports Medicine*
888 43:2379–2385. DOI: 10.1177/0363546515594380.

889 Wilk KE., Meister K., Andrews JR. 2002. Current concepts in the rehabilitation of the overhead
890 throwing athlete. *The American journal of sports medicine* 30:136–151.

891 Wilk KE., Reinold MM., Macrina LC., Porterfield R., Devine KM., Suarez K., Andrews JR. 2009.
892 Glenohumeral Internal Rotation Measurements Differ Depending on Stabilization Techniques.
893 *Sports Health* 1:131–136. DOI: 10.1177/1941738108331201.

894 Young J. 2014. Trunk Contributions to Baseball Pitching Velocity. PhD Thesis. The Ohio
895 State University.

896

897 **Unpublished Data**

898 Cressey, E. 2013. It Needs to Be Said: Throwing Doesn't Build Arm "Strength." Accessed
899 December 30th, 2017. [https://ericcressey.com/it-needs-to-be-said-throwing-doesnt-build-arm-](https://ericcressey.com/it-needs-to-be-said-throwing-doesnt-build-arm-strength)
900 [strength](https://ericcressey.com/it-needs-to-be-said-throwing-doesnt-build-arm-strength)

901 Reinold, M. 2017. Are Weighted Baseball Velocity Programs Safe and Effective? Accessed
902 December 10th, 2017. [https://elitebaseballperformance.com/are-weighted-baseball-velocity-](https://elitebaseballperformance.com/are-weighted-baseball-velocity-programs-safe-and-effective/)
903 [programs-safe-and-effective/](https://elitebaseballperformance.com/are-weighted-baseball-velocity-programs-safe-and-effective/)

904 Driveline Baseball 2017. Driveline Baseball Review - College Summer Training Results 2017
905 Accessed April 1st, 2018.
906 [https://www.drivelinebaseball.com/2017/09/driveline-baseball-review-college-summer-](https://www.drivelinebaseball.com/2017/09/driveline-baseball-review-college-summer-training-results-2017/)
907 [training-results-2017/](https://www.drivelinebaseball.com/2017/09/driveline-baseball-review-college-summer-training-results-2017/)

908 Driveline Baseball 2016. Driveline Baseball Review - College Summer Training Results 2016
909 Accessed April 1st, 2018.
910 <https://www.drivelinebaseball.com/2016/08/college-summer-wrap/>

911