

Decline in coral cover and flattening of the reefs around Mauritius (1998 - 2010)

Jennifer A Elliott ^{Corresp., 1}, Mark R Patterson ¹, Caroline G Staub ², Meera Koonjul ³, Stephen M Elliott ⁴

¹ Marine Science Center, Northeastern University, Nahant, Massachusetts, United States

² Agricultural and Biological Engineering, University of Florida, Gainesville, Florida, United States

³ Albion Fisheries Research Center, Ministry of Fisheries, Petite Riviere, Mauritius

⁴ Woods Hole Oceanographic Institute, Woods Hole, Massachusetts, USA

Corresponding Author: Jennifer A Elliott
Email address: jahking.elliott@gmail.com

Coral reefs are degrading through the impacts of multiple anthropogenic stressors. How are coral reef communities going to change and how to protect them for future generations are important conservation questions. Using coral reef data from Mauritius, we examined changes in cover in 23 benthic groups for a 13-yr period and at 15 sites. Moreover, we determined which land-based stressor out of four (human population, agriculture, tourism, rainfall) correlated the most with the observed changes in coral reef cover. Among the stony corals, *Acropora* corals appeared to be the most impacted, decreasing in cover at many sites. However, the non-*Acropora* encrusting group increased in cover at several sites. The increase in abundance of dead corals and rubble at some sites also supported the observations of stony coral decline during the study period. Additionally, the decline in stony corals appeared to be more important in second half of the study period for all sites suggesting that a global factor rather than a local factor was responsible for this decline. There was little change in cover for the other benthic groups, some of which were quite rare. Human population was significantly correlated with changes in coral reef cover for 11 sites, followed by tourism and agriculture. Rainfall, a proxy for runoff, did not appear to affect coral reef cover. Overall, our results showed that there has been a decline of stony coral especially the ones with complex morphologies, which in turn suggest that coral reefs around Mauritius have experienced a decline in habitat complexity during the study period. Our study also suggests that humans are an important factor contributing to the demise of coral reefs around the island.

Title

Decline in coral cover and flattening of the reefs around Mauritius (1998 - 2010)

Jennifer A. Elliott (corresponding author)

Marine Science Center
Northeastern University
430 Nahant Road
Nahant, MA 01908 USA
j.ahking.elliott@gmail.com

Mark R. Patterson

Marine Science Center
Northeastern University
430 Nahant Road
Nahant, MA 01908 USA

Caroline G. Staub

Climate Science and Climate Science Applications
Agricultural and Biological Engineering Department
University of Florida
220 Frazier-Rogers Hall
Gainesville, FL 32611 USA

Meera Koonjul

Albion Fisheries Research Centre, Ministry of Fisheries
Petite Riviere, Mauritius

Stephen M. Elliott

Woods Hole Oceanographic Institute
266 Woods Hole Road,
Woods Hole, MA 02543 USA

Abstract

Coral reefs are degrading through the impacts of multiple anthropogenic stressors. How are coral reef communities going to change and how to protect them for future generations are important conservation questions. Using coral reef data from Mauritius, we examined changes in cover in 23 benthic groups for a 13-yr period and at 15 sites. Moreover, we determined which land-based stressor out of four (human population, agriculture, tourism, rainfall) correlated the most with the observed changes in coral reef cover. Among the stony corals, *Acropora* corals appeared to be the most impacted, decreasing in cover at many sites. However, the non-*Acropora* encrusting group increased in cover at several sites. The increase in abundance of dead corals and rubble at some sites also supported the observations of stony coral decline during the study period. Additionally, the decline in stony corals appeared to be more important in second half of the study period for all sites suggesting that a global factor rather than a local factor was responsible for this decline. There was little change in cover for the other benthic groups, some of which were quite rare. Human population was significantly correlated with changes in coral reef cover for 11 sites, followed by tourism and agriculture. Rainfall, a proxy for runoff, did not appear to affect coral reef cover. Overall, our results showed that there has been a decline of stony coral especially the ones with complex morphologies, which in turn suggest that coral reefs around Mauritius have experienced a decline in habitat complexity during the study period. Our study also suggests that humans are an important factor contributing to the demise of coral reefs around the island.

59 **Introduction**

60 Coral reefs worldwide are declining through the impacts of many different types of natural and
 61 anthropogenic stressors (Pandolfi et al., 2003). The planet's climate is changing at a pace faster
 62 than ever observed before primarily due to the extensive burning of fossil fuels. As a result, the
 63 oceans are warming (IPCC, 2014) and stony corals, the foundation species of coral reefs, have
 64 been observed to bleach and die on a large scale (Roberts et al., 2002; Gardner et al., 2003;
 65 Pandolfi et al., 2003; Bellwood et al., 2004; Graham et al., 2006; Bruno & Selig, 2007; Heron et
 66 al., 2016; Hughes et al., 2017). Severe bleaching events are happening five times more often
 67 today than in the 1980s (Hughes et al., 2018).

68 Over 60% of the world's coral reefs have been estimated to be threatened by local stressors
 69 arising from anthropogenic activities (Burke et al., 2012). Land clearing for agriculture and
 70 coastal urbanization have led to sediments, nutrients, and pollutants being discharged through
 71 runoff into these coastal ecosystems (Smith et al., 1981; Fabricius, 2005; Markey et al., 2007;
 72 Brodie et al., 2012). Overfishing has also caused significant changes in ecological communities
 73 and contributed to the collapses of some coastal ecosystems (Jackson et al., 2001).

74 How are coral reef communities going to change in the face of multiple anthropogenic stressors
 75 and what makes some species more resilient than others are critical conservation questions.

76 Inherent biological traits (e.g. morphology, growth rate and reproductive mode) and
 77 physiological mechanisms of coral species are thought to play an important role in their
 78 capabilities to respond to repeated assaults from multiple stressors (Edmunds & Gates, 2008;
 79 Darling et al., 2012). For example, a study on coral community structure reported that coral
 80 colonies with branching morphologies bleached and died before those with massive and
 81 encrusting morphologies after a bleaching event (Loya et al., 2001). Theoretical work has also

shown that at the same flow rate, an organism with a flat morphology will have a higher mass transfer or diffusion rate (of gases and nutrients to and from the organism) than an organism with a branched morphology (Patterson, 1992). This could in turn also lead to faster removal of harmful superoxides and toxic radicals produced during stressful thermal events (Lesser, 1997), providing a competitive advantage to species with a flat morphology.

Physical external factors can also play an important role in helping coral reef organisms face multiple stressors. Localized upwelling (Glynn & D'Croz, 1990), water-flow rates and flushing of toxins have been reported to help stony corals weather thermal stresses (Nakamura & Van Woesik, 2001; 2005). Decreased light stress, e.g., in the form of shading or cloud cover or turbidity can also reduce some of the impacts from stressors (Hoegh-Guldberg, 1999; Moothien Pillay, Terashima & Kawasaki, 2002; Jokiel & Brown, 2004; Coelho et al., 2017).

Both global and local efforts are needed to conserve coral reefs worldwide as they continue to degrade because of increasing human population and their activities (Mora, 2008; Sandin et al., 2008; Smith et al., 2016; Crane et al., 2017). Long-term coral reef monitoring and studies disentangling the impacts of diverse anthropogenic activities are especially important to inform conservation actions. We investigated the effects of four land-based stressors on the coral reefs around Mauritius. This Indian Ocean island has one of the densest human population on earth, with 618 individuals km⁻² in 2017 (data.worldbank.org). Moreover, a large area of the island has been converted to agricultural land and the island is a prime tourist destination in the Indian Ocean. The two primary goals of this study were to examine changes in coral reef cover over a 13-yr period, and to investigate the relationship between the land-based stressors and changes in benthic communities around Mauritius.

Materials and Methods

Study sites

This study was carried out at the southwestern Indian Ocean island of Mauritius (20° 10' S, 57° 31' E; Fig. 1a). The island has a surface area of 1,869 km² and possesses 240 km² of reef habitats (Turner & Klaus, 2005). It also lies within the cyclonic belt and is visited by cyclones seasonally, typically from January to March. The land area is also divided into 42 watersheds (or catchment areas; Fig. 1b) (Proag, 1995). A total of 15 sites, 11 back- and four fore-reef sites, were investigated. Back-reef sites (circles) were found all around the island and fore-reefs sites (stars) only on the west coast, which is sheltered from the prevailing southeast trade winds (Fig. 1b). The reefs around Mauritius are broken up by natural breaks that result in a series of distinct lagoons. Back-reef sites were found in the lagoons that are < 3 m deep and 1 – 8 km wide. Additionally, the island has one 10 km long barrier reef on the southeast coast where one of the back reef site was found. This barrier reef is also next to a deep channel (15 – 30 m) that flushes seawater offshore (Fig. 1b) (Turner & Klaus, 2005). The fore-reef sites were found outside the lagoons in 8 – 20 m deep waters.

Some 160 stony corals species have been reported in Mauritius (Moothien Pillay et al., 2002). The back-reefs of Mauritius consist mainly of branching and tabular *Acropora*, massive *Porites*, foliaceous *Montipora* and *Pavona*, and sand consolidated with seagrass. The narrow reef flats are made up mostly of dead coral platform, stony corals and macroalgae, and the outer slopes have a spur and groove system that is colonized by a diverse range of hard and soft coral genera (e.g. *Pocillopora*, *Favia*, *Porites*, *Sarcophyton* and *Lobophyton*). For a more comprehensive description of the coral reefs of Mauritius, see Turner and Klaus (2005).

Datasets

(a) Benthic communities

Benthic coral reef monitoring data spanning over 13 years (1998 to 2010) were obtained from the Albion Fisheries Research Centre (AFRC), Government of Mauritius. The data were collected from three 20 m long transects at each site; each transect was marked by two metal stakes. The line intercept method (Kaiser, 1983) was used to survey these transects. A measuring tape was laid down along each transect; all benthic groups touching the tape were recorded and the number of centimeters that they covered along the tape was also recorded. The number in centimeters for each benthic group was then converted to percentages such that each transect would be equal to 100%. E.g. if sand was found from 0 cm to 23 cm, then sand would be equal to $(23-0)/(20*100)*100=1.15\%$ of the transect. The three transects were considered as three replicates for each site.

The data were originally collected twice a year, once in summer (December to May) and once in winter (June to November). However, later in the study period, data were sometimes only collected once a year and not all sites were surveyed each year during the study period (Table S1 – study years by site). During this 13-yr period, the data were collected by a small group of staff from the AFRC, which minimized observer error in the data.

The back-reef sites surveyed were: Albion (ALB), Anse La Raie (ALR), Baie du Tombeau (BDT), Bambous Virieux (BVX), Bel Ombre (BOM), Belle Mare (BME), Île aux Bénitiers (BEN), Poudre d’Or (PDO), Pte aux Sables (PTE), Trou aux Biches (TBI), Trou d’Eau Douce (TDO). The four fore-reefs sites were: Albion (ALB), Île aux Bénitiers (BEN), Pte aux Sables

(PTE) and Trou aux Biches (TBI). The back-reefs sites were in waters < 3m deep; the fore-reefs sites ALB, BEN, PTE and TBI were set up respectively at 10m, 10m, 8m and 12 m deep.

The benthic cover data were recorded as one of the following 23 groups: *Acropora* branching, *Acropora* digitate, *Acropora* encrusting, *Acropora* submassive, *Acropora* tabulate, non-*Acropora* branching, non-*Acropora* encrusting, non-*Acropora* foliaceous, non-*Acropora* submassive, non-*Acropora* massive, fungiid coral, turf algae, macroalgae, coralline algae, *Millepora* spp., sponges, soft corals, zoanthids, other live, dead corals, rubble, sand and rock. Dead corals consisted of calcium carbonate coral skeletons that were more or less intact that would eventually become rubble when they finally break down into smaller pieces.

(b) Land-based stressors

Four land-based stressors on the coral reefs, human population, tourism, agriculture, and rainfall were investigated. Spatial data for the selected stressors were gathered for the period of 1998 to 2010 so that they would match the time period for which the benthic data were available. Data for the first three stressors were obtained from the governmental organization ‘Statistics Mauritius’ (statsmauritius.govmu.org). Rainfall data were obtained from the monthly meteorological summaries (Mauritius Meteorological Services) that were made available by the Mauritius Sugar Industry Research Institute (MSIRI) archives department.

Data for human population, agriculture and tourism were spatially reorganized using a geographic information system (GIS; QGIS Lyon 2.12) such that each stressor would have one value per watershed per study year (Fig. 1b). The human population data were organized as the total number of people per watershed per year. The agricultural data were organized as total surface area under cultivation per watershed per year. The agricultural data consisted of both land under sugarcane and other food crop cultivation during the study period. For the tourism

data, a tourism index was created because the available data were rather coarse; only the number of hotels per watershed per year and total number of tourist nights for the whole island per year were available. This tourism index was created by dividing the number of hotels per watershed by the total number of hotels for the whole island and multiplied by total number of tourist nights. The resulting index gave an approximate number of tourist nights per watershed per year.

A more extensive spatial analysis was done on the rainfall data using ArcView GIS. Simulated monthly rainfall totals generated by Staub et al. 2014 were averaged across months and then by watershed for each study year. In their study, monthly rainfall totals (2000-2011) from 85 stations, a digital elevation model (DEM) and location-specific landscape characteristics were used to calculate linear relationships between mean rainfall and landscape characteristics on Mauritius. The time and space-varying nature of the modeled relationships were then explored by estimating separate models for each month (Staub, Stevens & Waylen, 2014). Finally, monthly rainfall totals for 1998 to 2011 were generated by removing the variability characteristic to each month from the observed data, interpolating these values across space using ordinary kriging, and adding these back to the spatial mean model for the corresponding month (Staub, Stevens & Waylen, 2014). Since rainfall was a proxy for runoff in this study the rainfall data (mm) per watershed per year was multiplied by their respective watershed areas (m^2) and divided by 1000 to obtain a volume of rainfall per watershed (m^3) per year.

(c) Linking the stressor effects arising within watersheds to study sites

Landscape characteristics including mountains and riverbeds were used to determine which watershed(s) drained into which lagoons. This was done as multiple back-reef sites were found in different lagoons and were thus impacted by different levels of land-based stressors. Study sites and their respective watersheds were color-coded and are displayed in Fig. 1b. When sites

from different lagoons shared a common watershed, one of the sites was colored white and the watersheds to which it was linked were shaded (Fig. 1b). The fore-reefs sites were linked to the same watersheds as their complementary back-reef sites. Table S2 in Supplemental Materials provides the long-term mean of each of the four local stressors by site for the period spanning 1998 to 2010.

Statistical analysis

All statistical tests in this study were done using R (R Core Team, 2015).

(a) Trends in benthic cover for 1998 - 2010

Temporal changes in benthic cover on the back- and fore-reefs were examined. A stacked bar graph was made for 1999 to show how the benthic cover varies by site. The benthic groups were aggregated into eight groups as it was difficult to visualize 23 groups (23 colors). The eight groups were: *Acropora* corals, non-*Acropora* corals, macroalgae, turf algae, coralline algae, dead corals, other living organisms and non-living organisms (included rubbles).

A mean percent cover per year for each benthic group was calculated for each season and for each study site. This was done by averaging data from the three transects. These mean values were then used in Spearman's rank correlation tests to determine whether there were any correlations between the benthic group cover and year. Only sites that had at least five or more years of observation were considered for this study. Year 2001 was also not considered as only four back- and two fore-reefs sites were surveyed, and therefore the data for this year were not representative of the whole island.

Temporal changes in total stony coral percent cover were also studied. A long-term mean for each site was calculated by averaging the annual percent cover of total live stony coral for the 13-year period for each site. This long-term mean for each site was subtracted from their respective annual means and the resulting values were used to create anomaly plots. These plots showed the mean annual deviations in total live stony coral cover for the 1998 to 2010 period. Two sets of anomaly plots were created, one from summer data and one from winter data.

(b) Land-based stressors – relationship with benthic cover trends

Temporal trends in land-based stressors were first examined by plotting total human population, total land area under agriculture, total number of tourists and total rainfall per year for the whole island against time. A Z-score was used on the y-axis to facilitate comparisons between stressors, which had widely different ranges.

To investigate the effects of the land-based stressors on the benthic communities, we used a combination of non-Metric Multidimensional Scaling analyses (nMDS) on Bray-Curtis distances (Borcard, Gillet & Legendre, 2011) and the ‘envfit’ function of the package vegan in R (Oksanen et al., 2015). The mean percent cover per benthic group per year were first calculated using data from both summer and winter; the seasonal data were combined since we only had one set of annual data for the land-based stressors (no seasonal data) to which the benthic data were tested with. The mean percent cover values were then used in nMDS to generate ordination plots for each site. The ordination plots indicate the scale of changes in community structure at each site. Then, using the ‘envfit’ function, data from the four stressors were fitted onto these ordinations to determine which ones had a significant correlation with the placement of benthic community observations in ordination space. The ‘envfit’ function considered the effects of each stressor individually.

238 Results

239 1. Changes in benthic cover

240 (i) Individual sites

241 Stony corals were the most abundant group at all sites. The back-reef sites had in general more
 242 *Acropora* corals while the fore-reef sites had more of the non-*Acropora* corals (Fig. 2). All sites
 243 had rubble and/or rocks (non-living category). The fore-reefs had in general more rocks. Turf
 244 algae and macroalgae were present at most sites albeit in varying amounts. Coralline algae were
 245 rarely detected. All sites had some dead corals; back-reef sites tended to have more dead corals
 246 than the fore-reef sites. Very little of other living organisms were recorded at the study sites.

247 Changes in benthic cover varied a lot among sites (Table 1). The 'X' was used to indicate
 248 benthic groups not observed during the survey. The *Acropora* coral groups experienced a
 249 decrease in cover over time. Although some non-*Acropora* coral groups also showed a
 250 significant decrease, a few groups increased in abundance. In particular, non-*Acropora*
 251 encrusting increased in cover at four back-reefs sites. Non-*Acropora* foliaceous, massive and
 252 submassive also increased at one site each.

253 Other benthic groups studied showed variable responses (Table 1). The cover of dead corals
 254 increased at four sites and decreased at one site. The cover of rubble increased at one site and
 255 decreased at three sites. Turf algae cover increased at one and three sites. The cover of
 256 macroalgae increased at three sites and decreased at two sites. Coralline algae increased at two
 257 sites and soft corals at one site. Sand increased at two sites and decreased at three sites.

258 *Millepora* and sponges respectively decreased at one site and increased at one site. The group
 259 'other live' decreased at two sites. No clear seasonal signal was detected except for turf algae
 260 and macroalgae, which appeared to be more abundant in summer than in winter.

(ii) Anomaly plots - stony corals

The summer anomaly plots showed that a large number of sites had a mean annual cover above their respective long-term mean (more circles with warm colors) till 2005, after which this trend was reversed; most sites afterwards had a mean annual cover below their respective long-term mean (more circles with cold colors; Fig. 3). Not a single site in 2009 and 2010 had a mean annual cover above their respective long-term mean. A similar trend was observed from the winter anomaly plots, although not as clear as those from the summer data (Fig. 4).

2. Island-wide trends for land-based stressors

When local stressor trends for the whole island were examined, there were clear increases in human population and the tourism index over the study time (Fig. 5). There was a general decline in land area under agriculture from 2001 and 2008, after which more land were cultivated. The big dip in cultivated land area in 1999 coincided with the big dip in annual rainfall, the year with the lowest rainfall during the study period. There was more rain after 1999, but it fluctuated quite a bit over time.

3. Land-based stressors - relationship with benthic cover

Significant effects of human population were detected at 11 of the 15 study sites (Table 2). Tourism and agriculture had significant effects respectively at five and four study sites. No significant effects of rainfall were detected at any of the study sites. There were insufficient data to carry out this test at Belle Mare (BME). The 'X' in the table indicates that there were either only a few years or no tourism data in the watersheds linked to these study sites. Therefore, it was not possible to test for the effects of tourism at these sites.

Discussion

Some significant changes in benthic cover occurred around the island of Mauritius during the period of 1998 to 2010. Among the stony corals, *Acropora* coral groups appeared to have been the most impacted, decreasing in abundance at many sites. While the non-*Acropora* coral groups also decreased in abundance at many sites, some groups however experienced an increase over time. The increase in abundance of dead corals and rubble at some sites also supported the observations of stony coral decline during the study period. Additionally, the decline in stony corals appeared to be more important in second half of the study period for all sites. There was little change in cover for the other benthic groups, some of which were quite rare. Our study also suggests that humans are an important factor contributing to the demise of coral reefs around the island.

The decline of *Acropora* corals around the island was expected as *Acropora* corals have been reported to be most sensitive to natural and anthropogenic stressors (Marshall & Baird, 2000; Loya et al., 2001; Rachello-Dolmen & Cleary, 2007; Cleary et al., 2008; Côté & Darling, 2010). In contrast, the non-*Acropora* coral groups, particularly ones with encrusting morphologies, appeared to be more successful competitors on the reefs. Their morphology could be a key aspect allowing them to have a higher mass transfer rate for gases and nutrients (Patterson, 1992), which in turn could lead to faster removal of harmful superoxides and toxic radicals produced during stressful events (Lesser, 1997). A study done in Japan also reported the greater survival of corals with massive and encrusting morphologies after a bleaching event (Loya et al., 2001). This increase of encrusting corals coupled with the loss of branching, digitate and tabulate *Acropora* corals also suggest that the coral reefs around Mauritius have flattened and

304 experienced a decline in habitat complexity. Similar flattening of reefs has been observed in the
305 Caribbean (Alvarez-Filip et al., 2009).

306 Interestingly, all study sites appeared to have experienced a more significant decline in stony
307 coral cover during the second half of the study period, i.e. around 2005 and onwards. These
308 observations suggest that a global rather than a local factor was involved. The 50-km Degree
309 Heating Weeks (DHW) data from Coral Reef Watch NOAA website for 2005 showed that the
310 reefs around Mauritius experienced severe heat stress (4°C-weeks and above) for three
311 consecutive summer months (ending 29 March, 30 April and 31 May; Table S3). Therefore,
312 temperature was a most likely stressor contributing to impact severely the corals during 2005,
313 which led to subsequent declines in cover. This temperature stress from 2005 was so severe that
314 the corals had not recovered by 2010.

315 The cover of other benthic groups such as macroalgae and turf algae did not change much at
316 most of the study sites, but they did appear to grow more in summer. These observations suggest
317 that there was in general a good top-down control by herbivores at these study sites. Herbivore
318 biomass is known to play an important role in coral reef dynamics and in mediating reef regime-
319 shifts (Jouffray et al., 2014). As observed on the Great Barrier Reef (De'ath, Lough & Fabricius,
320 2009), herbivorous fish on the reefs of Mauritius also face less significant fishing pressure as
321 artisanal fisheries target mainly piscivorous fishes.

322 Coralline algae is important for coral larval settlement (Harrington et al., 2004) and their cover
323 did not change much at most sites; they only increased at two sites. *Millepora* spp, sponges, soft
324 corals, zooanthids and other live group were either quite rare at most sites and/or did not appear
325 to experience much change during the study. The lumping of species into larger groups could,

however, have obscured temporal trends for individual species. For these rare groups, more targeted approaches are needed to study them.

Human population was the land-based stressor that appeared to impact the most sites around Mauritius. High human densities have previously been reported to impact negatively coral reef communities (Mora, 2008; Sandin et al., 2008; Smith et al., 2016; Crane et al., 2017). However, as also argued, it is not the number of people, but rather their activities that affect the coral reefs (Sanderson et al., 2002; Mora, 2008). Tourism was the second most common stressor to impact the reefs. Therefore, it appeared that the activities of the growing local population and the tourism industry have contributed the most to affect the coral reefs. The fore-reef sites located on the west coast also appeared to be primarily impacted by human population. These observations suggest that fore-reefs may not be far enough from land to be isolated from the impacts of the local human population. However, we had data for only four fore-reefs sites all located on the west coast of the island, and this small sample size may not be adequate to draw broad conclusions.

Agriculture appeared to affect only four study sites, however total land area under agriculture was used as a proxy for its effects and may not be representative of its negative impacts, e.g., the excessive use fertilizers and their detrimental effects are well documented (Markey et al., 2007; Brodie et al., 2012). Similarly, rainfall did not appear to affect the coral reefs; the negative impacts of runoff bringing pollutants to lagoons are also well documented (Fabricius, 2005; Brodie et al., 2012). Therefore, investigating the impacts of specific human activities would provide greater insights on the root causes of reef degradation around Mauritius.

The coral reefs as we know them are changing rapidly and will continue to degrade under the pressure of multiple natural and anthropogenic stressors. How coral reefs are going to change and how to protect them so they maintain important ecosystem functions for future generations remain a challenge. Our results showed that *Acropora* corals have been declining and suggest that corals reefs are flattening. Moreover, our results also suggest that human are a contributing factor to coral reef degradation and loss. Therefore, reducing the impacts of human activities remains very important. Other studies focusing on biological traits that convey resilience in the face of multiple stressors are needed better understand how benthic communities are going to change in the future.

Acknowledgements

We would like to thank the Albion Fisheries Research Centre, Ministry of Fisheries, Government of Mauritius for providing us with the benthic community dataset without which this work would not have been possible. Jennifer Elliott expresses her deepest gratitude to the members of her advisory committee, Peter Edmunds, Tarik Gouhier, Brian Helmuth and Steve Vollmer for their thoughtful inputs during the execution of this work. This is contribution number XXX from the Marine Science Center at Northeastern University.

References

- Alvarez-Filip L, Dulvy NK, Gill JA, Cote IM, Watkinson AR 2009. Flattening of Caribbean coral reefs: region-wide declines in architectural complexity. *Proceedings of the Royal Society B: Biological Sciences* 276:3019–3025. DOI: 10.1098/rspb.2009.0339.
- Bellwood D, Hughes T, Folke C, Nyström M 2004. Confronting the coral reef crisis. *Nature* 429:827–833.
- Borcard D, Gillet F, Legendre P 2011. *Numerical Ecology with R*. New York, NY: Springer Science & Business Media. DOI: 10.1007/978-1-4419-7976-6.
- Brodie JE, Kroon FJ, Schaffelke B, Wolanski EC, Lewis SE, Devlin MJ, Bohnet IC, Bainbridge ZT, Waterhouse J, Davis AM 2012. Terrestrial pollutant runoff to the Great Barrier Reef: An update of issues, priorities and management responses. *Marine Pollution Bulletin* 65:81–100. DOI: 10.1016/j.marpolbul.2011.12.012.

- 375 Bruno J, Selig E 2007. Regional decline of coral cover in the Indo-Pacific: timing, extent, and
376 subregional comparisons. *PLoS ONE* 2:e711.
- 377 Burke LM, Reynter K, Spalding M, Perry A 2012. *Reefs at risk revisited in the Coral Triangle*.
378 World Resource Institute, Washington DC.
- 379 Cleary DFR, De Vantier L, Giyanto, Vail L, Manto P, de Voogd NJ, Rachello-Dolmen PG, Tuti
380 Y, Budiyanto A, Wolstenholme J, Hoeksema BW, Suharsono 2008. Relating variation in
381 species composition to environmental variables: a multi-taxon study in an Indonesian coral
382 reef complex. *Aquatic Sciences* 70:419–431. DOI: 10.1007/s00027-008-8077-2.
- 383 Coelho VR, Fenner D, Caruso C, Bayles BR, Huang Y, Birkeland C 2017. Shading as a
384 mitigation tool for coral bleaching in three common Indo-Pacific species. *Journal of*
385 *Experimental Marine Biology and Ecology* 497:152–163. DOI:
386 10.1016/j.jembe.2017.09.016.
- 387 Côté I, Darling E 2010. Rethinking Ecosystem Resilience in the Face of Climate Change. *PLoS*
388 *Biology* 8:765–781.
- 389 Crane NL, Nelson P, Abelson A, Precoda K, Rulmal J, Bernardi G, Paddock M 2017. Atoll-scale
390 patterns in coral reef community structure: Human signatures on Ulithi Atoll, Micronesia.
391 *PLoS ONE* 12:e0177083–19. DOI: 10.1371/journal.pone.0177083.
- 392 Darling ES, Alvarez-Filip L, Oliver TA, McClanahan TR, Côté IM 2012. Evaluating life-history
393 strategies of reef corals from species traits. *Ecology Letters* 15:1378–1386. DOI:
394 10.1111/j.1461-0248.2012.01861.x.
- 395 De'ath G, Lough J, Fabricius K 2009. Declining coral calcification on the Great Barrier Reef.
396 *Science* 323:116.
- 397 Edmunds PJ, Gates RD 2008. Acclimatization in tropical reef corals. *Marine Ecology Progress*
398 *Series* 361:307–310. DOI: 10.3354/meps07556.
- 399 Fabricius KE 2005. Effects of terrestrial runoff on the ecology of corals and coral reefs: review
400 and synthesis. 50:125–146. DOI: 10.1016/j.marpolbul.2004.11.028.
- 401 Gardner TA, Côté IM, Gill JA, Grant A, Watkinson AR 2003. Long-Term Region-Wide
402 Declines in Caribbean Corals. *Science* 301:958–960. DOI: 10.1126/science.1086050.
- 403 Glynn PW, D'Croz L 1990. Experimental evidence for high temperature stress as the cause of El
404 Nino-coincident coral mortality. *Coral Reefs* 8:181–191. DOI: 10.1086/511756?ref=search-
405 gateway:e1b1fe7c141c6f73655a09e8dff067c.
- 406 Graham N, Wilson S, Jennings S, Polunin N, Bijoux J, Robinson J 2006. Dynamic fragility of
407 oceanic coral reef ecosystems. *Proceedings of the National Academy of Sciences* 103:8425.
- 408 Harrington L, Fabricius K, De'ath G, Negri A 2004. Recognition and selection of settlement
409 substrata determine post-settlement survival in corals. *Ecology* 85:3428–3437. DOI:
410 10.1890/04-0298.
- 411 Heron SF, Maynard JA, van Hooideonk R, Eakin CM 2016. Warming Trends and Bleaching
412 Stress of the World's Coral Reefs 1985–2012. *Scientific Reports* 6:1–14. DOI:
413 10.1038/srep38402.
- 414 Hoegh-Guldberg O 1999. Climate change, coral bleaching and the future of the world's coral
415 reefs. *Marine and freshwater research*:839–866.
- 416 Hughes TP, Anderson KD, Connolly SR, Heron SF, Kerry JT, Lough JM, Baird AH, Baum JK,
417 Berumen ML, Bridge TC, Claar DC, Eakin CM, Gilmour JP, Graham NAJ, Harrison H,
418 Hobbs J-PA, Hoey AS, Hoogenboom M, Lowe RJ, McCulloch MT, Pandolfi JM, Pratchett
419 M, Schoepf V, Torda G, Wilson SK 2018. Spatial and temporal patterns of mass bleaching
420 of corals in the Anthropocene. *Science* 359:80–83. DOI: 10.1126/science.aan8048.

Hughes TP, Kerry JT, Álvarez-Noriega M, Álvarez-Romero JG, Anderson KD, Baird AH, Babcock RC, Beger M, Bellwood DR, Berkelmans R, Bridge TC, Butler IR, Byrne M, Cantin NE, Comeau S, Connolly SR, Cumming GS, Dalton SJ, Diaz-Pulido G, Eakin CM, Figueira WF, Gilmour JP, Harrison HB, Heron SF, Hoey AS, Hobbs J-PA, Hoogenboom MO, Kennedy EV, Kuo C-Y, Lough JM, Lowe RJ, Liu G, McCulloch MT, Malcolm HA, McWilliam MJ, Pandolfi JM, Pears RJ, Pratchett MS, Schoepf V, Simpson T, Skirving WJ, Sommer B, Torda G, Wachenfeld DR, Willis BL, Wilson SK 2017. Global warming and recurrent mass bleaching of corals. *Nature* 543:373–377. DOI: 10.1038/nature21707.

IPCC 2014. *Climate change 2014: synthesis report. contribution of working groups I, II and III to the fifth assessment report of the Intergovernmental Panel on Climate Change*. IPCC, Geneva, Switzerland.

Jackson J, Kirby M, Berger W, Bjorndal K, Botsford L, Bourque B, Bradbury R, Cooke R, Erlandson J, Estes J 2001. Historical overfishing and the recent collapse of coastal ecosystems. *Science* 293:629–637.

Jokiel PL, Brown EK 2004. Global warming, regional trends and inshore environmental conditions influence coral bleaching in Hawaii. *Global Change Biology* 10:1627–1641. DOI: 10.1111/j.1365-2486.2004.00836.x.

Jouffray JB, Nyström M, Norstrom AV, Williams ID, Wedding LM, Kittinger JN, Williams GJ 2014. Identifying multiple coral reef regimes and their drivers across the Hawaiian archipelago. *Philosophical Transactions of the Royal Society of London. Series B: Biological Sciences* 370:20130268–20130268. DOI: 10.1098/rstb.2013.0268.

Kaiser L 1983. Unbiased estimation in line-intercept sampling. *Biometrics* 39:965–976. DOI: 10.2307/2531331.

Lesser MP 1997. Oxidative stress causes coral bleaching during exposure to elevated temperatures. *Coral Reefs* 16:187–192. DOI: 10.1007/s003380050073.

Loya Y, Sakai K, Yamazato K, Nakano Y, Sambali H, Van Woesik R 2001. Coral bleaching: the winners and the losers. *Ecology Letters* 4:122–131. DOI: 10.1046/j.1461-0248.2001.00203.x.

Markey KL, Baird AH, Humphrey C, Negri AP 2007. Insecticides and a fungicide affect multiple coral life stages. *Marine Ecology* 330:127–137. DOI: 10.3354/meps330127.

Marshall PA, Baird AH 2000. Bleaching of corals on the Great Barrier Reef: differential susceptibilities among taxa. *Coral Reefs* 19:155–163. DOI: 10.1007/s003380000086.

Moothien Pillay R, Terashima H, Kawasaki H 2002. The extent and intensity of the 1998 mass bleaching event on the reefs of Mauritius, Indian Ocean. 4:43–52.

Moothien Pillay R, Terashima H, Venkatasami A, Uchida H 2002. *Field guide to corals of Mauritius*. Albion Fisheries Research Center, Ministry of Fisheries, Mauritius.

Mora C 2008. A clear human footprint in the coral reefs of the Caribbean. *Proceedings of the Royal Society B: Biological Sciences* 275:767.

Nakamura T, Van Woesik R 2001. Water-flow rates and passive diffusion partially explain differential survival of corals during the 1998 bleaching event. *Marine Ecology Progress Series* 212:301–304.

Nakamura T, Van Woesik R 2005. Photoinhibition of photosynthesis is reduced by water flow in the reef-building coral *Acropora digitifera*. *Marine Ecology Progress Series* 301:109–118.

Oksanen J, Blanchet FG, Kindt R, Legendre P, Minchin PR, O'Hara RB, Simpson GL, Solymos P, Stevens MH, Szoecs E, Wagner H 2015. *Vegan: Community Ecology Package. R package version 2.3-2*. <http://CRAN.R-project.org/package=vegan>.

- Pandolfi JM, Bradbury RH, Sala E, Hughes TP, Bjorndal KA, Cooke RG, McArdle D, Mcclenachan L, Newman MJH, Paredes G, Warner RR, Jackson JBC 2003. Global trajectories of the long-term decline of coral reef ecosystems. *Science* 301:955–958. DOI: 10.1126/science.1085706.
- Patterson MR 1992. A mass transfer explanation of metabolic scaling relations in some aquatic invertebrates and algae. *Science* 255:1421–1423.
- Proag V 1995. *The geology and water resources of Mauritius*. Mahatma Gandhi Institute Press, Mauritius.
- R Core Team 2015. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. <https://www.r-project.org/>.
- Rachello-Dolmen PG, Cleary DFR 2007. Relating coral species traits to environmental conditions in the Jakarta Bay/Pulau Seribu reef system, Indonesia. *Estuarine Coastal and Shelf Science* 73:816–826. DOI: 10.1016/j.ecss.2007.03.017.
- Roberts CM, McClean CJ, Veron JEN, Hawkins JP, Allen GR, McAllister DE, Mittermeier CG, Schueler FW, Spalding M, Wells F, Vynne C, Werner TB 2002. Marine biodiversity hotspots and conservation priorities for tropical reefs. *Science* 295:1280–1284. DOI: 10.1126/science.1067728.
- Sanderson EW, Jaiteh M, Levy MA, Redford KH, Wannebo AV, Woolmer G 2002. The human footprint and the last of the wild. *Science* 25:891–904.
- Sandin SA, Smith JE, DeMartini EE, Dinsdale EA, Donner SD, Friedlander AM, Konotchick T, Malay M, Maragos JE, Obura D, Pantos O, Paulay G, Richie M, Rohwer F, Schroeder RE, Walsh S, Jackson JBC, Knowlton N, Sala E 2008. Baselines and degradation of coral reefs in the Northern Line Islands. *PLoS ONE* 3:e1548. DOI: 10.1371/journal.pone.0001548.
- Smith JE, Brainard R, Carter A, Grillo S, Edwards C, Harris J, Lewis L, Obura D, Rohwer F, Sala E, Vroom PS, Sandin S 2016. Re-evaluating the health of coral reef communities: baselines and evidence for human impacts across the central Pacific. *Proceedings of the Royal Society B: Biological Sciences* 283:20151985–9. DOI: 10.1098/rspb.2015.1985.
- Smith SV, Kimmerer WJ, Laws EA, Brock RE, Walsh TW 1981. Kaneohe Bay sewage diversion experiment: perspectives on ecosystem responses to nutritional perturbation. *Pacific Science* 35:279–395.
- Staub CG, Stevens FR, Waylen PR 2014. The geography of rainfall in Mauritius: Modelling the relationship between annual and monthly rainfall and landscape characteristics on a small volcanic island. *Applied Geography* 54:222–234. DOI: 10.1016/j.apgeog.2014.08.008.
- Turner J, Klaus R 2005. Coral reefs of the Mascarenes, Western Indian Ocean. *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences* 363:229–250.

Captions for figures and tables

Figure 1. Mauritius. (a) The island of Mauritius in the southwestern Indian Ocean. (b) Map of the island showing back- (circles) and fore- (stars) reef sites. Land area is divided into several watersheds (catchment areas). Study sites were connected to adjacent watersheds(s) and are

508 color-coded identically. The back-reef site colored white is connected to adjacent shaded
509 watersheds.

510 **Figure 2.** Benthic cover by site for 1999. The benthic cover data (23 categories) were aggregated
511 into eight groups to facilitate visualization of the data.

512 **Figure 3.** Anomaly plots for stony coral cover– Summer data. a) Long-term means for each site
513 for the period of 1998 – 2010; year 2001 was not included as very few sites were surveyed
514 during that year. (b-m) Data for each year excluding 2001. Warm colors indicate cover was
515 above long-term mean and cold colors indicate that cover was below long-term mean.

516 **Figure 4.** Anomaly plots for stony coral cover – Winter data. a) Long-term means for each site
517 for the period of 1998 – 2010; year 2001 was not included as very few sites were surveyed
518 during that year. (b-m) Data for each year excluding 2001. Warm colors indicate cover was
519 above long-term mean and cold colors indicate that cover was below long-term mean.

520 **Figure 5.** Trends in land-based stressors for 1998 – 2010. The total human population (total
521 number of local people), total number of tourists, total surface area under agriculture and the
522 total rainfall for each year are shown.

523 **Table 1.** Temporal trends in benthic cover at individual sites. Spearman rank correlation results.
524 Only significant results are shown ($p < 0.05$). The ‘X’ indicates that benthic group was not
525 observed. Only sites that had at least five years of observation were used in this test. See M&M
526 for full site names.

527 **Table 2.** Land-based stressors and their relationship with benthic cover.

528

529 **Supplementary materials**

530 **Table S1.** Number of study years available by site. See M&M for site names.

531

532 **Table S2.** Long-term mean of local stressor by site for 1998-2010. Numbers in brackets indicate

533 rank of each station. Rank goes from least to most abundant. See M&M for full site names.

534 **Table S3.** Degree Heating Weeks data from NOAA. The seawater around Mauritius experienced

535 severe temperature stress during the months of March to April 2005.

Figure 1

Mauritius

(a) The island of Mauritius in the southwestern Indian Ocean. (b) Map of the island showing back- (circles) and fore- (stars) reef sites. Land area is divided into several watersheds (catchment areas). Study sites were connected to adjacent watersheds(s) and are color-coded identically. The back-reef site colored white is connected to adjacent shaded watersheds.




Figure 2(on next page)

Benthic cover by site for 1999

The benthic cover data (23 categories) were aggregated into eight groups to facilitate visualization of the data.




Figure 3

Anomaly plots for stony coral cover– Summer data

(a) Long-term means for each site for the period of 1998 – 2010; year 2001 was not included as very few sites were surveyed during that year. (b-m) Data for each year excluding 2001. Warm colors indicate cover was above long-term mean and cold colors indicate that cover was below long-term mean.




Figure 4

Anomaly plots for stony coral cover – Winter data

a) Long-term means for each site for the period of 1998 – 2010; year 2001 was not included as very few sites were surveyed during that year. (b-m) Data for each year excluding 2001. Warm colors indicate cover was above long-term mean and cold colors indicate that cover was below long-term mean.




Figure 5(on next page)

Trends in land-based stressors for 1998 – 2010

The total human population (total number of local people), total number of tourists, total surface area under agriculture and the total rainfall for each year are shown.




Table 1(on next page)

Temporal trends in benthic cover at individual sites

Spearman rank correlation results. Only significant results are shown ($p < 0.05$). The 'X' indicates that benthic group was not observed. Only sites that had at least five years of observation were used in this test. See M&M for full site names.

Table 1. Temporal trends in benthic cover at individual sites. Spearman rank correlation results. Only significant results are shown (p < 0.05). The 'X' indicates that benthic group was not observed. Only sites that had at least five years of observation were used in this test. See M&M for full site names.

Reef (Back or Fore)	Stations	Season (S or W)	n	<i>Acropora</i> branching (ab)	<i>Acropora</i> digitate (ad)	<i>Acropora</i> encrusting (ae)	<i>Acropora</i> submassive (as)	<i>Acropora</i> tabulate (at)	non- <i>Acropora</i> branching (nb)	non- <i>Acropora</i> encrusting (ne)	non- <i>Acropora</i> foliaceous (nf)	non- <i>Acropora</i> massive (nm)	non- <i>Acropora</i> mushroom (nu)	non- <i>Acropora</i> submassive (ns)
B	ALB	W	12	-0.87	X	X	X				-0.58			-0.80
B	ALR	S	7		X		X	X			-0.96			
B	ALR	W	6		X		X	X						
B	BDT	W	6		X	X	X		X	+0.87				
B	BEN	S	8		-0.76	X	X	-0.91			X			
B	BEN	W	7			X	X		-0.80		X			
B	BME	W	6		X	X	X				X	X	X	X
B	BOM	S	8	-0.71	X	X	X					X		
B	BOM	W	7		X	X	X					X		
B	BVX	S	8			X				+0.87	+0.70			
B	BVX	W	8	-0.88		X		-0.73	-0.85		+0.95			
B	PDO	S	6	X	X	X	X		X		-0.93	-0.84		-0.82
B	PSA	S	8	-0.78	X	X		X		X	X			
B	PSA	W	5		X	X		X		X	X			
B	TBI	S	9			X	X	-0.78	X	+0.78				
B	TBI	W	5			X	X		X					-0.86
B	TDO	S	9			X	X	-0.73	-0.73	+0.83		+0.67		+0.82
B	TDO	W	6			X	X							
F	ALB	W	10					X		-0.76			X	
F	BEN	S	5					X					X	
F	BEN	W	7		-0.97			X	-0.80			-0.82	X	
F	PSA	S	8				-0.76	X						-0.80
F	TBI	S	6	-0.84		X			-0.84				X	-0.86
F	TBI	W	6			X							X	

6

7 Table 1 continued

Reef (Back or Fore)	Stations	Season (S or W)	n	Turf algae (ta)	Macroalgae (ma)	Coraline algae (ca)	Millepora spp. (mi)	Sponges (sp)	Soft corals (sc)	Zoanthids (zo)	Dead corals (de)	Rubble (ru)	Sand (sa)	Rock (ro)	Other (ot)
B	ALB	W	12	-0.81			X			X		+0.83			
B	ALR	S	7	+0.79						X	+0.85			X	
B	ALR	W	6							X				X	
B	BDT	W	6				X		X			-0.94		X	X
B	BEN	S	8				X			X	+0.83				X
B	BEN	W	7	-0.93		+0.75	X		X	X	+0.82		+0.86		-0.80
B	BME	W	6		+0.94	X	X	X	X	X		-0.83			X
B	BOM	S	8	X			X	X	X				-0.79	X	
B	BOM	W	7				X	X	X					X	
B	BVX	S	8					X	X					X	
B	BVX	W	8					X	X		+0.71			X	
B	PDO	S	6		+0.82		X		X	X				X	X
B	PSA	S	8		+0.91		X			X	-0.90	+0.95			
B	PSA	W	5				X			X		+0.90	-0.90		
B	TBI	S	9			X		X	X	X	+0.78		-0.68		X
B	TBI	W	5	-0.94	-0.93	X		X	X	X					X
B	TDO	S	9				X					-0.82			
B	TDO	W	6				X					-0.93			
F	ALB	W	10		-0.78		-0.70		+0.84						
F	BEN	S	5				X	X	X					X	
F	BEN	W	7				X	X	X					X	
F	PSA	S	8					X		X					
F	TBI	S	6				X			X					
F	TBI	W	6			+0.97	X	-0.83		X			+0.81		-0.82

8

Table 2(on next page)

Land-based stressors and their relationship with benthic cover.

nMDS results.

1 **Table 2.** Land-based stressors and their relationship with benthic cover.

Station	N	Reef	2-D Stress	Population	Tourism	Agriculture	Rainfall
ALB	12	Back	0.03	*	X	*	
ALR	11	Back	0.09	*	*		
BDT	8	Back	0.06				
BEN	11	Back	0.06	*	*	*	
BME	7	Back	Insufficient data				
BOM	12	Back	0.20		X		
BVX	11	Back	0.09	*	X		
PDO	8	Back	0.04	*	*	*	
PSA	11	Back	0.07	*		*	
TBI	11	Back	0.11	*	*		
TDO	11	Back	0.10	*	X		
ALB	11	Fore	0.09	*	X		
BEN	10	Fore	0.11				
PSA	10	Fore	0.10	*			
TBI	10	Fore	0.09	*	*		
TOTAL				11	5	4	0

2