

Evidence for the Cretaceous shark *Cretoxyrhina mantelli* feeding on the pterosaur *Pteranodon* from the Niobrara Formation

David W E Hone ^{Corresp., 1}, Mark P Witton ², Michael B Habib ³

¹ School of Biological and Chemical Sciences, Queen Mary University of London, London, United Kingdom

² School of Earth and Environmental Sciences, University of Portsmouth, Portsmouth, United Kingdom

³ Keck School of Medicine, University of Southern California, Los Angeles, United States

Corresponding Author: David W E Hone

Email address: d.hone@qmul.ac.uk

A cervical vertebra of the large, pelagic pterodactyloid pterosaur *Pteranodon* sp. from the Late Cretaceous Niobrara Formation of Kansas, U.S.A is significant for its association with a tooth from the large lamniform shark, *Cretoxyrhina mantelli*. Though the tooth does not pierce the vertebral periosteum, the intimate association of the fossils – in which the tooth is wedged below the left prezygapophysis – suggests their preservation together was not mere chance, and the specimen is evidence of *Cretoxyrhina* biting *Pteranodon*. It is not possible to infer whether the bite reflects predatory or scavenging behaviour from the preserved material. There are several records of *Pteranodon* having been consumed by other fish, including other sharks (specifically, the anacoracid *Squalicorax kaupi*), and multiple records of *Cretoxyrhina* biting other vertebrates of the Western Interior Seaway, but until now interactions between *Cretoxyrhina* and *Pteranodon* have remained elusive. The specimen increases the known interactions between large, pelagic, vertebrate carnivores of the Western Interior Seaway of North America during the Late Cretaceous, in addition to bolstering the relatively small fossil record representing pterosaurian interactions with other species.

Evidence for the Cretaceous shark *Cretoxyrhina mantelli* feeding on the pterosaur *Pteranodon* from the Niobrara Formation

David W. E. Hone^{1*}

Mark P. Witton ²

Michael B. Habib³

1. School of Biological and Chemical Sciences, Queen Mary University of London, London, E1 4NS, UK.

2. School of Earth and Environmental Sciences, University of Portsmouth, Burnaby Road, Portsmouth PO1 3QL, UK.

3. Keck School of Medicine, University of Southern California, Los Angeles, CA, United States of America

Corresponding author: David Hone, d.hone@qmul.ac.uk

Abstract:

A cervical vertebrae of the large, pelagic pterodactyloid pterosaur *Pteranodon* sp. from the Late Cretaceous Niobrara Formation of Kansas, U.S.A is significant for its association with a tooth from the large lamniform shark, *Cretoxyrhina mantelli*. Though the tooth does not pierce the vertebral periosteum, the intimate association of the fossils – in which the tooth is wedged below the left prezygapophysis – suggests their preservation together was not mere chance, and the specimen is evidence of *Cretoxyrhina* biting *Pteranodon*. It is not possible to infer whether the bite reflects predatory or scavenging behaviour from the preserved material. There are several records of *Pteranodon* having been consumed by other fish, including other sharks (specifically, the anacoracid *Squalicorax kaupi*), and multiple records of *Cretoxyrhina* biting other vertebrates of the Western Interior Seaway, but until now interactions between *Cretoxyrhina* and *Pteranodon* have remained elusive. The specimen increases the known interactions between large, pelagic, vertebrate carnivores of the Western Interior Seaway of North America during the Late Cretaceous, in addition to bolstering the relatively small fossil record representing pterosaurian interactions with other species.

Introduction:

Pteranodon is a large pterodactyloid pterosaur from the Late Cretaceous (Coniacian-Campanian) of North America with an estimated maximum wingspan of 7.25 m (Bennett, 2001). The genus was among the first pterosaurs reported from North America (Marsh, 1876 – see Bennett, 2001 and Witton, 2010 for context of its discovery) and has become one of the

best known flying reptiles thanks to a representation of over 1100 specimens – the highest sample size for any pterosaur genus. Although most specimens are incomplete and crushed, every component of its osteology is known and has been described in detail (Eaton, 1910; Bennett, 1991, 1994, 2001, 2017, 2018; Bennett & Penkalski, 2018). As a result of the number of available specimens, its long research history and comprehensive documentation, the genus has become a cornerstone of pterosaur research. *Pteranodon* has been an important animal for understanding pterosaur flight (Hankin & Watson, 1914; Bramwell & Whitfield, 1974; Stein, 1975), the evolution of giantism in flying animals (Witton & Habib, 2010), pterosaur ontogeny (Bennett, 1993), and palaeoecology (Bennett, 2001; Witton, 2018).

The majority of *Pteranodon* specimens are known from the Late Cretaceous Niobrara Formation from Kansas, U.S.A., a marine deposit created by the Western Interior Seaway, though other specimens also occur in additional formations in Wyoming and South Dakota (Bennett, 1994, 2001). Niobrara specimens of *Pteranodon* occur in localities that were hundreds of kilometres from the palaeocoastline and this, along with a number of aspects of functional anatomy, has seen the genus long interpreted as a seagoing, pelagic animal (e.g., Bennett, 2001; Witton, 2013, p. 179).

Pteranodon was likely an important component of the Western Interior Seaway ecosystem. It seems to have been relatively abundant, being known from both a large number of fossils and making up some 97% of Niobrara Formation pterosaur finds. It was also a large animal - Bennett (1993) identified a bimodal size distribution among the large *Pteranodon* sample where two thirds of individuals were c. 3.5 m in wingspan, and the remaining third were much larger, some exceeding 6 m across the wings (Bennett, 2001). Larger specimens likely

exceed the masses of any flying bird, extant or extinct, with estimated body masses of 35-50 kg for animals of 6 m wingspan (Paul, 2002; Witton, 2008; Henderson 2010), compared to 21.9–40.1 kg in the largest fossil flying birds, the pelagornithids (Mayr & Rubilar-Rogers, 2010; Ksepka, 2014). *Pteranodon* populations may therefore have been major consumers in the Western Interior Seaway ecosystem, as well as potentially sources of food for other animals.

However, our understanding of interactions between *Pteranodon* and other Seaway taxa are limited. As with other pterosaur species, few *Pteranodon* fossils preserve remains of ingested content and they only rarely record evidence of consumption by other animals (Witton, 2018). Regurgitated fish are preserved in the gular region of one *Pteranodon* specimen (Brown, 1943; Bennett, 2001, 2018) and some palaeoecological significance has been ascribed to small fish vertebrae found in association with *Pteranodon* fossils (Bennett, 2001; Hargrave, 2007; Ehret & Harrell, 2018). Biting traces on *Pteranodon* elements show that some individuals were eaten by the anacoracid shark *Squalicorax kaupi* as well as a saurodontid fish, most likely *Saurodon* or *Saurocephalus* (Witton, 2018; Ehret & Harrell, 2018). The record of pterosaur ecological interactions is sufficiently sparse that any fossilised interactions with other species should be put on record, so we hereby report on a series of *Pteranodon* cervical vertebrae, LACM 50926, associated with a tooth of the lamniform shark *Cretoxyrhina mantelli*. This is first documented occurrence of this large shark interacting with any pterosaur.

Systematic nomenclature:

The taxonomy of *Pteranodon* is a matter of recent dispute. For the last two decades most workers have followed the treatment of the genus outlined by Bennett (1994), who made

80 a case for reducing the 11 binomials associated with *Pteranodon* (excluding those names
81 related to *Nyctosaurus*) to two sexually dimorphic chronospecies: the older *Pteranodon*
82 *sternbergi* and the younger *P. longiceps*. In this scheme, the skulls of these species are
83 distinguished by details of their cranial crests, and (more tentatively) occiput orientation and
84 mandibular ramus depth. Postcranial bones of these specimens are nearly identical and of little
85 taxonomic utility (Bennett, 1994). More recently, Kellner (2010) argued for *Pteranodon sensu*
86 Bennett (1994) being comprised of four species in three genera. While agreeing with Bennett
87 (1994) that all ‘historic’ *Pteranodon* species were problematic excepting *longiceps* and
88 *sternbergi*, Kellner (2010) created a multi-taxic pteranodontid assemblage for the Niobrara
89 specimens comprising *Pteranodon longiceps*, *Geosternbergia* (rather than *Pteranodon*)
90 *sternbergi*, and two novel species, *Geosternbergia maiseyi* and *Dawndraco kanzai*. These taxa
91 are primarily distinguished by headcrest morphology and details of the posterior skull, as well
92 as finer stratigraphic divisions of the Niobrara Formation (Kellner, 2010) than the broader
93 ‘upper’ and ‘lower’ divisions of the Smoky Hill Chalk *Pteranodon* fauna recognised by other
94 workers (e.g. Bennett, 1994; Everhart, 2005; Carpenter, 2008). Subsequent criticism of this
95 proposal has questioned the validity of the proposed differences between at least *Dawndraco*
96 and *Pteranodon sensu* Bennett, noted incongruence between the stratigraphic divisions
97 signified by Kellner (2010) against other Niobrara Formation taxa, as well as the lack of
98 statistical support for splitting *Pteranodon* into multiple genera, compared to the strong
99 statistical support for Bennett’s interpretation (Martin-Silverstone et al., 2017; Acorn et al.,
100 2017). We thus follow several other works (Witton, 2013, 2018; Bennett 2016, 2017, 2018) in

retaining Bennett's (1994) treatment of *Pteranodon* here. Note however that discussion of *Pteranodon* taxonomy is ongoing (Brandão & Rodrigues, 2018).

Materials and Methods:

LACM 50926 (Los Angeles County Museum of Natural History, USA) is a specimen of *Pteranodon* mounted in a large glass case for public display at the Los Angeles County Museum of Natural History and is difficult to access directly (Fig 1). The specimen has a large *Cretoxyrhina mantelli* tooth intimately associated with the fourth cervical vertebra (Fig 2). Parts of the mount are genuine, well preserved fossils showing only limited crushing compared to many specimens of the genus. However, several elements are reconstructed to replace missing parts and the mount is a composite of material from several individuals (see Bennett, 1991, 2001), as is obvious from bone size discrepancies between several neighbouring elements (Fig. 1). Bennett (pers. comm. 6/ 2016) also notes material accessioned under this number (much of it in collections space and not in the exhibit mount) includes three mandibular rami, confirming the multi-individual nature of this specimen. An alternate specimen number (65218) occurs on the mandible and the cervical bearing the shark tooth, but this cannot be seen on other elements. This may indicate that the mandible and cervical were associated when discovered. Bennett (2001) was able to identify many of the LACM 50926 forelimb elements as belonging to a single individual, although there are no records to indicate which parts of the mounted specimen might relate directly to the cervical series. The preservation quality and size of the vertebrae correspond well to the other elements (including the forelimb bones) and this implies that LACM 50926 may represent much of a skeleton. However, the absence of both

anteriormost and posterior cervical vertebrae means no anatomical continuity links the vertebrae with the rest of the material, and their association to the rest of the skeleton cannot be confidently assumed.

Notes held at the LACM show that the specimen was collected in 1965 by M.C. Bonner from Niobrara Chalk 23, Niobrara Formation, Logan County, Kansas. Bennett (1991) refers to two specimens under this number (LACM 50926 and 50926 "A") and concurs with this locality, adding that they were collected between Marker Units 14 and 19. This makes a Santonian age likely for LACM 50926 (Hattin, 1982; Bennett, 1994).

Description:

The anatomy of *Pteranodon* has been described in detail elsewhere (Bennett, 2001) and we will therefore focus exclusively on the association between the shark tooth and pterosaur material. The cervical vertebra bearing the shark tooth is preserved in contact with two other cervicals as a series of three elements. Thus, within the composite context of the LACM specimen, these vertebrae at least can be safely considered part of a single individual. The cervicals are preserved with contact between the successive post- and prezygapophyses. These are identified by Bennett (2001) as cervical vertebrae 4-6, and he also identified a preceding, though not articulated, cervical in the LACM 50926 mount as a cervical 3. The vertebrae retain some three-dimensionality, although they are partially crushed at an oblique angle, shearing them along their midline such that the left sides are depressed and right sides elevated (Figs 1-2). The neural spine is missing (now restored) from cervical 4 and parts of the neural spines of

cervicals 5 and 6 are damaged. Damage to the bone cortex reveals the internal structure of the bones in all three vertebrae.

The centrum lengths of the three cervical vertebrae in the series have been measured as 69.0, 77.8, and 71.5 mm respectively (Bennett, 2001). Based on comparisons to other specimens (see Bennett, 2001) this would correspond to a *Pteranodon* with a c. 5 m wingspan, and was presumably therefore osteological adult or near adult in size. The embedded shark tooth is approximately 24 mm in total height (root plus crown) (this was measured from photographs as it was impossible to measure the tooth given its location and the mount of the specimen), subtriangular in shape and highly compressed labiolingually. A wide, lunate root is formed from two obtusely angled, swollen root lobes. The termination of the left lobe (viewed from lingual aspect) forms a broad, somewhat rounded surface, but the termination of the right lobe is missing (Fig. 2). The crown is swollen on the labial surface, c. 12 mm long (measured from the base of the root to apex of the crown), almost symmetrical but not significantly recurved with respect to the root. No serrations are apparent on the tooth crown but the lateral and medial crown edges are somewhat worn with chipped margins. The tooth enameloid is bright white with grey to brown patches, and the root is pale grey-brown and close in colour to that of the pterosaur elements.

The tooth lies between the left prezygapophysis of cervical 4 and the centrum. In some aspects it appears that the tooth is wedged or has cut into the base of the prezygapophysis and the centrum however, it lies medial to the prezygapophysis and does not directly contact this projection. The tooth is preserved at a shallow angle to the long axis of the vertebra (though this may reflect the crushing of the specimen rather than its original orientation) and the apex

of the crown faces posteriorly and ventrally with respect to the vertebral corpus. Although the tip of the tooth does not appear to penetrate the centrum, the tip of the tooth contacts it.

Results:

Taxonomic identities:

The composite nature of LACM 50926 complicates discussions of its affinities, but there is no doubt that the specimen can be referred to *Pteranodon* given its provenance and matching anatomy to this pterosaur (Eaton, 1910; Bennett, 2001). Identification to species level is more problematic as *Pteranodon* taxonomy is exclusively informed by the posterior skull region (e.g. Eaton, 1910; Bennett, 1994; Kellner, 2010), and the vertebra is not associated with any skull material. Following Bennett's (1994) tentative suggestion that *P. sternbergi* may have a shallower mandible than *P. longiceps* we compared the LACM 50926 mandibular ramus with specimens referred to these species. However, we were unable to determine a significant match with either taxon. Hargrave (2007) suggested that the tomial margins of posterior *P. longiceps* mandibles are curved, and this morphology is present in the LACM 50926 mandible. However, while we agree this can be seen in some *P. longiceps* (e.g. YPM 2594 - YPM, Yale Peabody Museum, USA) it does not seem to be a universal trait (e.g. YPM 1177).

The recovery of LACM 50926 from marker units 14-19 of Hattin's (1982) Smoky Hill Chalk stratigraphy gives it provenance among younger Niobrara beds yielding *Pteranodon longiceps* rather than *P. sternbergi* (Bennett, 1994; Carpenter, 2008, although Kellner, 2010 argues that species more closely related to *P. sternbergi* than *P. longiceps* may persist into

younger deposits). This indicates that LACM 50926 probably represents *P. longiceps* but, in lieu of diagnostic fossil material, we regard the specimen as *Pteranodon* sp.

A number of medium- to large-sized, sharp-toothed sharks are known from the Niobrara Formation, and they have left an extensive record of tooth traces and shed teeth among other vertebrates of the Smoky Hill Chalk Member (Everhart, 2005). The Niobrara species best known for leaving bite traces is *Squalicorax kaupi*, but this identification can be excluded for the LACM tooth because it lacks the asymmetrical crown, notched cutting edge and serrations characterising the dentition of this genus (e.g. see Everhart, 2005; Becker & Chamberlain, 2012). The tooth is a good match for the large lamniform shark *Cretoxyrhina mantelli* (Fig. 3), which has subtriangular, relatively broad, symmetrical and short crowns without serrated margins (e.g. Shimada, 1997; Siverson & Lindgren, 2005, their fig 2; Bourdon & Everhart, 2011). In particular, the morphology of the tooth in LACM 50926 matches teeth recovered from anterior positions of *Cretoxyrhina* jaws (Shimada, 1997; Bourdon & Everhart, 2011, their figs 2, 5). The identification of the shark tooth as belonging to *Cretoxyrhina* was also independently made by Konuki (2008) and Witton (2018). Comparison of the LACM tooth size with a superb *C. mantelli* skeleton, FHSM VP-2187 (Shimada, 1997), suggests that the individual was c. 2.5 m long. This is little more than one third of length of the largest known individuals of this species (Everhart, 2005).

Discussion:

Significance of association of Pteranodon and Cretoxyrhina

Ecological interactions between pterosaurs and other species are rarely represented in fossil specimens, despite vast increases in pterosaur specimen numbers in recent years (Witton, 2018). Data on diet from stomach contents is sparse, limited to a handful of taxa known to have eaten fish (e.g. *Eudimorphodon* – Wild, 1978, *Pteranodon*, *Rhamphorhynchus* – Wellnhofer, 1991). Coprolites are also scarce, with only one record for pterosaurs known to date (Hone et al., 2014). A number of animals are recorded as pterosaur consumers, including fish (e.g. Frey & Tischlinger, 2012), dinosaurs (e.g. Hone et al., 2012), Crocodyliformes (Vremir et al., 2013) and possibly plesiosaurs (Cicimurri & Everhart, 2001, although also see Witton, 2018), but they remain very rare despite the good fossil records of these ‘consumer’ taxa. Thus, this additional potential record of a pterosaur-carnivore association is significant.

The taphonomic history and association of LACM 50926 is unknown so it is difficult to draw firm conclusions about the action that left the shark tooth in situ. However, we rule out abiotic association of the pterosaur and shark tooth for several reasons: 1) embedded *Cretoxyrhina* teeth and feeding traces are known from numerous Smoky Hill vertebrate fossils, and are widely interpreted as related to feeding behaviour (Shimada, 1997; Everhart, 2004, 2005); 2) although isolated *Cretoxyrhina* teeth are common fossils in the Smoky Hill Chalk Member (Everhart, 2005), its teeth have not been reported in association with any *Pteranodon* fossils in the past, despite the large sample size of this pterosaur and the fact that other fish remains (e.g. vertebrae) are not uncommonly associated with their remains (Bennett, 2001; Hargrave, 2007); 3) the spatial relationship between the tooth and the vertebra is complex and intimate, and unlike that expected to have occurred by chance association in a low energy

deposit such as the Niobrara Chalk. We thus prefer an interpretation of the tooth becoming associated with the vertebra though the biting action of a small *Cretoxyrhina*.

We were unable to find additional indications of bite traces on LACM 50926. There is a small and almost perfectly circular puncture on the neural arch of cervical four, behind the left prezygapophysis but this is most likely a preparation trace or damage derived from a previous museum mount. The damaged and missing neural spines of the cervical series may be linked to the shark bite, but other pterosaur fossils show that these elements are prone to damage and/or poor preservation, so other causes cannot be excluded.

Cretoxyrhina was a large (up to 7 m in length) and powerful carnivore, perhaps one of the top predators of the Smoky Hill Chalk fauna (Everhart, 2005). Shimada (1997) compared its likely ecological feeding guild to larger modern species of lamnid and carcharhinid sharks, and there is fossil evidence that it consumed a variety of large vertebrates including mosasaurs, plesiosaurs and large teleost fish (Shimada, 1997; Everhart, 2004, 2005). LACM 50926 is the first palaeoecological link between this shark genus and a pterosaur however, this rarity perhaps reflecting the relatively delicate nature of pterosaur skeletons against the evident bite force of *Cretoxyrhina*. Extremely hollow bones such as those characterising most of the *Pteranodon* skeleton are especially prone to failure against buckling forces (Currey, 2004) and likely broke easily under strong bites from large predators.

Both Bennett (2001) and Hargrave (2007) have noted that *Pteranodon* may have been consumed destructively by large aquatic carnivores, their relatively muscular torsos being targeted, perhaps explaining why wing skeletons (which had considerably less soft-tissue, see Bennett, 2008) are the most common form of associated pterosaur fossil in the Smoky Hill

Chalk Member. It should be noted however, that articulated wings are also common in the Late Jurassic Solnhofen fauna where this may reflect decay and the loss of wings from intact and floating corpses of pterosaurs (Beardmore, Lawlor & Hone, 2017), although this is not mutually exclusive with the effects of predation and scavenging. Witton (2016) noted that, to date, only the larger, more robust elements – limb bones and neck vertebrae – of larger pterosaur species are known to preserve embedded teeth, and speculated that small pterosaurs and/or more gracile pterosaur bones were probably too easily destroyed to record evidence of carnivore bites. It may be that pterosaurs were not rare dietary components of *Cretoxyrhina* or other animals, but that their anatomy precludes common fossilisation of evidence for these acts.

There is limited potential for knowing whether the LACM 50926 association reflects predatory or scavenging behaviour from *Cretoxyrhina*. *Pteranodon* is widely considered to have been a pelagic pterosaur species which foraged for small aquatic prey by means of dip-feeding, fishing from an alighted position on the water surface or diving after food (Wellnhofer, 1991; Bennett, 2001; Witton, 2013, 2016). Adaptations to aquatic launch (identified by Habib & Cunningham, 2010) are apparent in *Pteranodon* and suggest that it may have routinely entered (and thus needed to launch from) bodies of water. Thus, there are good reasons to think living *Pteranodon* could have been within reach of predatory sharks, and the likely pterodactyloid floating posture places their head and neck close to the waters' surface (Hone & Henderson, 2014). Various seabirds are known to be predated by pelagic predators, including sharks, in modern times (Wetherbee, Cortés & Bizzarro, 2004; Johnson et al., 2006) and we cannot exclude this possibility for the LACM *Pteranodon*. Witton (2018) noted that even moderately-sized sharks, akin to the 2.5 m long *Cretoxyrhina* indicated by the LACM tooth, would vastly

outweigh the largest *Pteranodon* (35-50 kg – see Paul, 2002; Witton, 2008; Henderson, 2010 for *Pteranodon* mass estimates), and we have little doubt that such predators could subdue these pterosaurs if they caught them (Fig. 4). Conversely, *Pteranodon* likely had a relatively low body density and their carcasses may have floated for sustained periods (Hone & Henderson, 2014). This would make them obvious targets for scavenging marine animals. Ultimately, LACM 50926 preserves no evidence to falsify any of these hypotheses.

Evidence of the anacoracid shark *Squalicorax* consuming *Pteranodon* is known in the Niobrara (e.g. KU 972 - KU, Kansas University, USA; YPM 2597, YPM 42810 – Bennett, pers. comm. 06/16), and recent finds of Mooreville Chalk Formation *Pteranodon* also have bite marks attributed to *Squalicorax kaupi* (RMM 3274 and ALMNH 8630) (Ehret & Harrell, 2018). This body of evidence, augmented with the *Cretoxyrhina*-*Pteranodon* association described here, and the recovery of fish remains within the gular region of *Pteranodon* specimens (Brown, 1943; Bennett, 2001, 2018) makes the trophic interactions of *Pteranodon* well understood compared to most other pterosaurs (Witton, 2018). However, such finds are still relatively rare occurrences - these seven associations are less than 1% of the >1100 specimens of *Pteranodon* on record. In contrast, at least ten palaeoecologically significant fossil associations are known for the Late Jurassic Solnhofen pterosaur *Rhamphorhynchus muensteri* (including five associations with the carnivorous fish *Aspidorhynchus acutirostris* (e.g. Frey & Tishchlinger, 2012) and four examples of consumed items – see Witton, 2018 for a recent review). There are perhaps 150 specimens of *Rhamphorhynchus* in public collections, suggesting that recording of palaeoecological events is several times higher than in *Pteranodon* (>6%) despite a considerably

smaller sample size. The taphonomic factors contributing to this difference may be worthy of further study.

Acknowledgements:

We thank Luis Chiappe and Maureen Walsh for access to the specimen and information on its history and Stephanie Abramowicz for access to archival photographs of the specimen. We would like to thank Steffi Klug for assistance on identification of the shark tooth, Dana Ehret and Chris Bennett for discussions on the composite mount and bite marks on *Pteranodon* bones and providing key literature. We thank both Dana Ehret and Chris Bennett for their reviews and Mark Young for his handling of the manuscript as editor that helped improve this work.

References:

- Acorn JH, Martin-Silverstone E, Glasier JR, Mohr S, Currie PJ. 2017. Response to Kellner (2017) 'Rebuttal of Martin-Silverstone, E., JRN Glasier, JH Acorn, S. Mohr, and PJ Currie, 2017'. *Vertebrate Anatomy Morphology Palaeontology* 3.
- Beardmore SR, Lawlor E, Hone DWE. 2017. Using taphonomy to infer differences in soft tissues between taxa: an example using basal and derived forms of Solnhofen pterosaurs. *The Science of Nature* 104:65.
- Becker MA, Chamberlain JA. 2012. *Squalicorax* chips a tooth: a consequence of feeding-related behavior from the lowermost Navesink Formation (Late Cretaceous: Campanian-Maastrichtian) of Monmouth County, New Jersey, USA. *Geosciences* 2:109–129.

- 315 Bennett SC. 1991. Morphology of the Late Cretaceous pterosaur *Pteranodon* and systematics of
- 316 the Pterodactyloidea. Doctoral dissertation, University of Kansas, Systematics and Ecology.
- 317 Bennett SC. 1992. Sexual dimorphism of *Pteranodon* and other pterosaurs, with comments on
- 318 cranial crests. *Journal of Vertebrate Paleontology* 12:422–434
- 319 Bennett SC. 1993. The ontogeny of *Pteranodon* and other pterosaurs. *Paleobiology* 19:92–106
- 320 Bennett SC. 1994. Taxonomy and systematics of the Late Cretaceous pterosaur *Pteranodon*
- 321 (Pterosauria, Pterodactyloidea). *Occasional Papers of the Museum of Natural History,*
- 322 *University of Kansas, Lawrence* 169:1–70.
- 323 Bennett SC. 2001. The osteology and functional morphology of the Late Cretaceous pterosaur
- 324 *Pteranodon*. (parts 1 and 2). *Paleontographica Abteilung A*, 260:1–153.
- 325 Bennett SC. 2003. A survey of pathologies of large pterodactyloid pterosaurs. *Palaeontology*
- 326 46:185–198.
- 327 Bennett SC. 2018. New smallest specimen of the pterosaur *Pteranodon* and ontogenetic niches
- 328 in pterosaurs. *Journal of Paleontology* 92:254–271.
- 329 Bennett SC, Penkalski P. 2017. Waves of bone deposition on the rostrum of the pterosaur
- 330 *Pteranodon*. Geological Society of London, Special Publications, 455:69.
- 331 Bourdon J, Everhart MJ. 2011. Analysis of an associated *Cretoxyrhina mantelli* dentition from
- 332 the Late Cretaceous (Smoky Hill Chalk, Late Coniacian) of western Kansas. *Transactions of*
- 333 *the Kansas Academy of Science* 114:15–32.
- 334 Bramwell CD, Whitfield GR. 1974. Biomechanics of *Pteranodon*. *Philosophical Transactions of*
- 335 *the Royal Society of London: Series B* 267:503–581.

336 Brandão RS, Rodrigues T. 2018. Reappraisal of *Dawndraco kanzai* as a valid taxon. *Flugsaurier*
337 *2018 conference abstracts*, 17.

338 Brown B. 1943. Flying reptiles. *Natural History* 52:104–111.

339 Carpenter K. 2008. Vertebrate biostratigraphy of the Smoky Hill Chalk (Niobrara Formation) and
340 the Sharon Springs Member (Pierre Shale). In: Harries PJ. ed. *High-Resolution Approaches in*
341 *Stratigraphic Paleontology: Topics in Geobiology Series*, 21:421–437.

342 Cicimurri DJ, Everhart MJ. 2001. An elasmosaur with stomach contents and gastroliths from the
343 Pierre Shale (Late Cretaceous) of Kansas. *Transactions of the Kansas Academy of Science*
344 104:129–143.

345 Eaton GF. 1910. Osteology of *Pteranodon*. *Memoirs of the Connecticut Academy of Arts and*
346 *Sciences* 2:1–38.

347 Ehret DJ, Harrell Jr, TR. 2018. Feeding traces on a Pteranodon (Reptilia: Pterosauria) bone from
348 the Late Cretaceous (Campanian) Mooreville Chalk in Alabama, USA. *Palaios*, 33:414–418.

349 Everhart MJ. 2004. Late Cretaceous interaction between predators and prey. Evidence of
350 feeding by two species of shark on a mosasaur. *PalArch, vertebrate palaeontology*
351 *series* 1:1–7.

352 Everhart MJ. 2005. *Oceans of Kansas*. Indiana University Press.

353 Frey E, Tischlinger H. 2012. The Late Jurassic Pterosaur *Rhamphorhynchus*, a frequent victim of
354 the ganoid fish *Aspidorhynchus*? *PLoS ONE* 7:e31945.

355 Habib M, Cunningham J. 2010. Capacity for water launch in *Anhanguera* and *Quetzalcoatlus*.
356 *Acta Geoscientica Sinica* 31:24–25.

- 357 Hankin EH, Watson DMS. 1914. On the flight of pterodactyls. *The Aeronautical Journal* 18:324-
358 335.
- 359 Hargrave JE. 2007. *Pteranodon* (Reptilia: Pterosauria): Stratigraphic distribution and taphonomy
360 in the lower Pierre Shale Group (Campanian), western South Dakota and eastern Wyoming.
361 In: Martin JE, Parris DC. eds. *The Geology and Paleontology of the Late Cretaceous Marine*
362 *Deposits of the Dakotas: Geological Society of America Special Paper* 427:215–225.
- 363 Hattin DE. 1982. Stratigraphy and depositional environment of Smoky Hill Chalk Member,
364 Niobrara Chalk (Upper Cretaceous) of the type area, western Kansas. *Kansas Geological*
365 *Survey Bulletin* 225:108.
- 366 Henderson DM. 2010. Pterosaur body mass estimates from three-dimensional mathematical
367 slicing. *Journal of Vertebrate Paleontology* 30:768–785.
- 368 Hone DWE, Henderson DM. 2014. The posture of floating pterosaurs: ecological implications for
369 inhabiting marine and freshwater habitats. *Palaeogeography, Palaeoclimatology,*
370 *Palaeoecology* 398:89–98.
- 371 Hone DWE, Tsuchiji T, Watabe M, Tsogbataar K. 2012. Pterosaurs as a food source for small
372 dromaeosaurs. *Palaeogeography, Palaeoclimatology, Palaeoecology* 331:27–30.
- 373 Hone DWE, Henderson DM, Therrien F, Habib MB. 2015. A specimen of *Rhamphorhynchus* with
374 soft tissue preservation, stomach contents and a putative coprolite. *PeerJ* 3:e1191.
- 375 Johnson RL, Venter A, Bester MN, Oosthuizen WH. 2006. Seabird predation by white shark,
376 *Carcharodon carcharias*, and Cape fur seal, *Arctocephalus pusillus pusillus*, at Dyer
377 Island. *South African Journal of Wildlife Research* 36:23–32.

Kellner AWA. 2010. Comments on the Pteranodontidae (Pterosauria, Pterodactyloidea) with the description of two new species. *Anais de Academia Brasileira de Ciencias* 82:1063–1084.

Konuki Reira. 2008. Biostratigraphy of Sea Turtles and Possible Bite Marks on a *Toxochelys* (Testudine, Chelonioidae) from the Niobrara Formation (Late Santonian), Logan County, Kansas, and Paleoecological Implications for Predator-prey Relationships Among Large Marine Vertebrates. Masters dissertation, Fort Hays State University,

Marsh OC. 1876. Notice of a new sub-order of Pterosauria. *American Journal of Sciences* 11:507–509.

Martin-Silverstone E, Glasier JRN, Acorn JH, Mohr S, Currie PJ. 2017. Reassessment of *Dawndraco kanzai* Kellner, 2010 and reassignment of the type specimen to *Pteranodon sternbergi* Harksen, 1966. *Vertebrate Anatomy Morphology Palaeontology* 3:47–59.

Paul GS. 2002. *Dinosaurs of the air: the evolution and loss of flight in dinosaurs and birds*. JHU Press.

Shimada K. 1997. Paleoecological relationships of the Late Cretaceous lamniform shark, *Cretoxyrhina mantelli* (Agassiz). *Journal of Paleontology* 71:926–933.

Siverson M, Lindgren J. 2005. Late Cretaceous sharks *Cretoxyrhina* and *Cardabiodon* from Montana, USA. *Acta Palaeontologica Polonica* 50:301–314.

Stein RS. 1975. Dynamic analysis of *Pteranodon ingens*: a reptilian adaptation to flight. *Journal of Paleontology* 49:534–548.

Vremir M, Kellner AW, Naish D, Dyke GJ. 2013. A new azhdarchid pterosaur from the Late Cretaceous of the Transylvanian Basin, Romania: implications for azhdarchid diversity and distribution. *PLoS One* 8:e54268.

Wellnhofer P. 1991. The Illustrated Encyclopedia of Pterosaurs. Salamander Books, London.

Wetherbee BM, Cortés E, Bizzarro JJ. 2004. Food consumption and feeding habits. *Biology of Sharks and their Relatives*, 225–246.

Wild R. 1978. Die Flugsaurier (Reptilia, Pterosauria) aus der Oberen Trias von Cene bei Bergamo, Italien. *Bollettino della Società Paleontologica Italiana* 17:176–256

Witton MP. 2008. A new approach to determining pterosaur body mass and its implications for pterosaur flight. *Zitteliana B* 28:143–158.

Witton MP. 2010. *Pteranodon* and beyond: the history of giant pterosaurs from 1870 onwards. *Geological Society, London, Special Publications* 343:313–323.

Witton MP. 2013. *Pterosaurs: natural history, evolution, anatomy*. Princeton University Press.

Witton MP. 2018. Pterosaurs in Mesozoic food webs: a review of fossil evidence. *Geological Society, London, Special Publications* 455:7–23.

Witton MP, Habib M. 2010. On the size and flight diversity of giant pterosaurs, the use of birds as pterosaur analogues and comments on pterosaur flightlessness. *PLoS ONE* 5:e13982.

Figure captions.

Fig 1. A, mounted *Pteranodon* sp. skeleton LACM 50926 on display in the Los Angeles county museum with highlighted section of the vertebrae shown below; B, Close up of the vertebral series and shark tooth (indicated by an arrow). Cervical vertebrae III-VII as mounted on the display are indicated. Scale bar is 50 mm – this is an approximate value based on published measurements of the vertebrae. Image credit: A, Stephanie Abramowicz, courtesy Dinosaur Institute, Natural History Museum of Los Angeles County, B, David Hone.

Fig 2. Two close up views of the *Cretoxyrhina mantelli* tooth with tracings. A, left dorsolateral view; B, left dorsoventral view showing its intimate association with cervical vertebra IV. The tooth is highlighted in medium grey, the 4th cervical vertebra in pale grey and the 5th cervical in dark grey. Abbreviations: ns neural spine, prz prezygopophysis, psz postzygopophysis, st shark tooth. Image credit: David Hone.

Fig 3. Tracing of *Cretoxyrhina mantelli* anterior teeth from Bourdon and Everhart (2011, their fig 5, mirrored from their original). A, position 3 in the jaw; B, position 4; C, LACM 50926 tooth. The roots of the teeth are shaded in pale grey and the enamel is dark grey. Image credit: David Hone.

Fig. 4. Life reconstruction of a c. 2.5 m long breaching *Cretoxyrhina mantelli* biting the neck of a 5 m wingspan *Pteranodon longiceps*, a scene inspired by LACM 50926. The predatory behaviour of this scene is speculative with respect to the data offered by the specimen, but reflects the fact that *Cretoxyrhina* is generally considered a predatory species, the vast weight advantage of the shark against the pterosaur (see text), and the juvenile impulse of the artist to draw an explosive predatory scene. Image credit: Mark Witton.

Figure 1

Mounted Pteranodon and close up of the neck

Fig 1. A, mounted *Pteranodon* sp. skeleton LACM 50926 on display in the Los Angeles county museum with highlighted section of the vertebrae shown below; B, Close up of the vertebral series and shark tooth (indicated by an arrow). Cervical vertebrae III-VII are indicated. Scale bar is 50 mm – this is an approximate value based on published measurements of the vertebrae. Image credit: A, Stephanie Abramowicz, courtesy Dinosaur Institute, Natural History Museum of Los Angeles County, B, David Hone.




Figure 2

Two close up views of the *Cretoxyrhina mantelli* tooth with tracings.

Fig 2. Two close up views of the *Cretoxyrhina mantelli* tooth with tracings. A, left dorsolateral view; B, left dorsoventral view showing its intimate association with cervical vertebra IV. The tooth is highlighted in medium grey, the 4th cervical vertebra in pale grey and the 5th cervical in dark grey. Abbreviations: ns neural spine, prz prezygopophysis, psz postzygopophysis, st shark tooth. Image credit: David Hone.




Figure 3

Cretoxyrhina mantelli anterior teeth

Fig 3. Tracing of *Cretoxyrhina mantelli* anterior teeth from Bourdon and Everhart (2011, their fig 5, mirrored from their original). A, position 3 in the jaw; B, position 4; C, LACM 50926 tooth. The bases of the teeth are shaded in pale grey and the enamel is dark grey. Image credit: David Hone.




Figure 4

Life reconstruction of a *Cretoxyrhina mantelli* attacking a *Pteranodon longiceps*

Fig. 4. Life reconstruction of a c. 2.5 m long breaching *Cretoxyrhina mantelli* biting the neck of a 5 m wingspan *Pteranodon longiceps*, a scene inspired by LACM 50926. The predatory behaviour of this scene is speculative with respect to the data offered by the specimen, but reflects the fact that *Cretoxyrhina* is generally considered a predatory species, the vast weight advantage of the shark against the pterosaur (see text), and the juvenile impulse of the artist to draw an explosive predatory scene. Image credit: Mark Witton.

