

High tempo music prolongs high intensity exercise (#29630)

1

First submission

Editor guidance

Please submit by **23 Aug 2018** for the benefit of the authors (and your \$200 publishing discount).



Structure and Criteria

Please read the 'Structure and Criteria' page for general guidance.



Custom checks

Make sure you include the custom checks shown below, in your review.



Raw data check

Review the raw data. Download from the [materials page](#).



Image check

Check that figures and images have not been inappropriately manipulated.

Privacy reminder: If uploading an annotated PDF, remove identifiable information to remain anonymous.

Files

Download and review all files from the [materials page](#).

2 Figure file(s)

1 Table file(s)

! Custom checks

Human participant/human tissue checks


- ! Have you checked the authors [ethical approval statement](#)?
- ! Does the study meet our [article requirements](#)?
- ! Has identifiable info been removed from all files?
- ! Were the experiments necessary and ethical?



Structure your review

The review form is divided into 5 sections. Please consider these when composing your review:

1. BASIC REPORTING
2. EXPERIMENTAL DESIGN
3. VALIDITY OF THE FINDINGS
4. General comments
5. Confidential notes to the editor






 You can also annotate this PDF and upload it as part of your review

When ready [submit online](#).





Editorial Criteria

Use these criteria points to structure your review. The full detailed editorial criteria is on your [guidance page](#).





BASIC REPORTING

-  Clear, unambiguous, professional English language used throughout.
-  Intro & background to show context. Literature well referenced & relevant.
-  Structure conforms to [PeerJ standards](#), discipline norm, or improved for clarity.
-  Figures are relevant, high quality, well labelled & described.
-  Raw data supplied (see [PeerJ policy](#)).

EXPERIMENTAL DESIGN

-  Original primary research within [Scope of the journal](#).
-  Research question well defined, relevant & meaningful. It is stated how the research fills an identified knowledge gap.
-  Rigorous investigation performed to a high technical & ethical standard.
-  Methods described with sufficient detail & information to replicate.

VALIDITY OF THE FINDINGS

-  Impact and novelty not assessed. Negative/inconclusive results accepted. *Meaningful* replication encouraged where rationale & benefit to literature is clearly stated.
-  Speculation is welcome, but should be identified as such.
-  Conclusions are well stated, linked to original research question & limited to supporting results.
-  Data is robust, statistically sound, & controlled.

Standout reviewing tips

3



The best reviewers use these techniques

Tip

Support criticisms with evidence from the text or from other sources

Example

Smith et al (J of Methodology, 2005, V3, pp 123) have shown that the analysis you use in Lines 241-250 is not the most appropriate for this situation. Please explain why you used this method.

Give specific suggestions on how to improve the manuscript

Your introduction needs more detail. I suggest that you improve the description at lines 57- 86 to provide more justification for your study (specifically, you should expand upon the knowledge gap being filled).

Comment on language and grammar issues

The English language should be improved to ensure that an international audience can clearly understand your text. Some examples where the language could be improved include lines 23, 77, 121, 128 – the current phrasing makes comprehension difficult.

Organize by importance of the issues, and number your points

1. Your most important issue
2. The next most important item
3. ...
4. The least important points

Please provide constructive criticism, and avoid personal opinions

I thank you for providing the raw data, however your supplemental files need more descriptive metadata identifiers to be useful to future readers. Although your results are compelling, the data analysis should be improved in the following ways: AA, BB, CC

Comment on strengths (as well as weaknesses) of the manuscript

I commend the authors for their extensive data set, compiled over many years of detailed fieldwork. In addition, the manuscript is clearly written in professional, unambiguous language. If there is a weakness, it is in the statistical analysis (as I have noted above) which should be improved upon before Acceptance.

High tempo music prolongs high intensity exercise

Meaghan Maddigan¹, Kathleen M Sullivan¹, Fabien A Basset¹, Israel Halperin¹, David G Behm^{Corresp. 1}

¹ School of Human Kinetics and Recreation, Memorial University of Newfoundland, St. John's, Newfoundland/Labrador, Canada

Corresponding Author: David G Behm
Email address: dbehm@mun.ca

Music has been shown to reduce rating of perceived exertion, increase exercise enjoyment and enhance exercise performance, mainly in low-moderate intensity exercises. However, the effects of music are less conclusive with high-intensity activities. The purpose of this study was to compare the effects of high tempo music (130 bpm) to a no-music condition during repeated high intensity cycling bouts (80% of peak power output (PPO)) on the following measures: time to task failure (TTF), rating of perceived exertion (RPE), heart rate (HR), breathing frequency, ventilatory kinetics and blood lactate (BL). Under the music condition, participants exercised 10.7% longer ($p = 0.035$; Effect size (ES)= 0.28) (increase of one minute) and had higher HR (4%; $p = 0.043$; ES= 0.25), breathing frequency (11.6%; $p = 0.0006$; ES= 0.57), and RER (7% at TTF; $p = 0.021$; ES=1 .1) during exercise. Trivial differences were observed between conditions in RPE and other ventilatory kinetics during exercise. Interestingly, HR recovery was 13.0% faster following the music condition ($p < 0.05$). These results strengthen the notion that music can alter the association between central motor drive, central cardiovascular command and perceived exertion, and contribute to prolonged exercise duration at higher intensities along with a quicken HR recovery.

HIGH TEMPO MUSIC PROLONGS HIGH INTENSITY EXERCISE

Authors: Meaghan E. Maddigan, Kathleen M. Sullivan, Fabien A. Basset, Israel Halperin
David G. Behm

Institution: School of Human Kinetics and Recreation
Memorial University of Newfoundland
St. John's, Newfoundland, Canada

Corresponding Author: Dr. David Behm
School of Human Kinetics and Recreation
Memorial University of Newfoundland
St. John's, Newfoundland, Canada
dbehm@mun.ca
tel: 709-864-3408
fax: 709-864-3979

Running Title: Effect of Music on Exercise

Funding: There was no direct funding received for this research.

Conflict of Interest: There were no conflicts of interests with any of the authors.

ABSTRACT

Music has been shown to reduce rating of perceived exertion, increase exercise enjoyment and enhance exercise performance, mainly in low-moderate intensity exercises. However, the effects of music are less conclusive with high-intensity activities. The purpose of this study was to compare the effects of high tempo music (130 bpm) to a no-music condition during repeated high intensity cycling bouts (80% of peak power output (PPO)) on the following measures: time to task failure (TTF), rating of perceived exertion (RPE), heart rate (HR), breathing frequency, ventilatory kinetics and blood lactate (BL). Under the music condition, participants exercised

10.7% longer ($p = 0.035$; Effect size (ES)= 0.28) (increase of one minute) and had higher HR (4%; $p= 0.043$; ES= 0.25), breathing frequency (11.6%; $p= 0.0006$; ES= 0.57), and RER (7% at TTF; $p= 0.021$; ES=1 .1) during exercise. Trivial differences were observed between conditions in RPE and other ventilatory kinetics during exercise. Interestingly, HR recovery was 13.0% faster following the music condition ($p< 0.05$). These results strengthen the notion that music can alter the association between central motor drive, central cardiovascular command and perceived exertion, and contribute to prolonged exercise durations at higher intensities along with a quicken HR recovery.

Key Words: aerobic exercise, cycling, heart rate, lactate

INTRODUCTION

Music has long been thought to affect the senses (Szmedra and Bacharach 1998) and can act as an external distracting stimulus during exercise and thus enhance exercise enjoyment and performance (Karageorghis and Priest 2012A/B). Music is able to promote ergogenic and psychological benefits during exercise due to three proposed explanations (Karageorghis and Priest 2012A/B). First, music may allow individuals to separate thoughts from feelings. This divergence can change one's perception of unpleasant feelings, narrowing the performer's attention and reducing the sensations of fatigue during exercise (Atkinson et al. 2004; Edworthy and Haring 2006; Yamashita et al. 2006; Murrock and Higgins 2009). Second, the divergent stimulus (i.e. music) will somehow alter psychomotor arousal (movement or muscular activity associated with mental processes) and therefore can act as either a stimulant or a sedative prior to and during physical activity (Szmedra and Bahanach 1998; Yamamoto et al. 2003; Schucker et al. 2009). The third explanation postulates that during continual submaximal activity, an

individual is predisposed to respond to rhythmical elements (music being one of many rhythmical patterns); the result being synchronization between the tempo and the performer's movement making physical activity or exercise a more harmonious or less stressful experience (Rendi et al. 2008; Waterhouse et al. 2009).

The available evidence on this topic is congruent and demonstrates that music can and does have a consistent and measurable effect on attention, the ability to trigger a range of emotions, affect mood, increase work output, and encourage rhythmic movement (Atkinson et al. 2004; Karageorghis 2008; Scherer 2004; Terry and Karageorghis 2011; Yamashita et al. 2006). The 'psychophysical' effects primarily examine the perception of effort (Pandolf 1978), which in almost all cases involves the Borg's Ratings of Perceived Exertion scale (RPE).

The effect of music on exercise in the low-to-moderate range of exercise intensities is a common trend in the literature (Karageorghis and Priest 2012). Since very high exercise intensities are affected to a high degree by muscle metabolite-induced failures (peripheral fatigue), there is a lesser influence of the central nervous system (central fatigue) compared to lower exercise intensities (Rejeski 1985; Tenenbaum 2004). However, it has been shown that peripheral fatigue alone is not able to explain the fatigue induced with higher intensity exercise (Noakes and Gibson 2004, Halperin et al. 2014). The few studies that have examined the effect of music on higher intensity exercise bouts show methodological weaknesses and a misuse of music terminology resulting in some incongruent results (Bharani, Sahu and Matthew 2004; Crust and Clough 2006; Nakamura et al. 2010; Schie, Stewart, Becker and Rogers 2008). In contrast, some recent studies using 'supramaximal' exercise, such as a Wingate test, have shown significant improvements in peak power output and decreases in fatigue index with the use of music (Brohmer and Becker 2006; Haluk, Turchian, and Adnan 2009). This seems to contradict

the previous conclusions drawn by researchers that the ‘distraction effect’ of music is attenuated at higher exercise intensities ($>70\%$ maximal oxygen uptake – $\text{VO}_{2\text{max}}$) due to the internal feedback dominating the capacity of the respective afferent nervous system (Karageorghis, Terry, Lane, Bishop and Priest 2011). More importantly, it highlights some significant gaps in the literature with regards to the so-called intensity limitations of music’s benefits and the actual mechanisms that result in music’s ergogenic effects on exercise performance. Further research is still necessary in order to draw decisive conclusions.

Therefore, the primary goal of this study was to examine if listening to high tempo music (130 bpm) while performing high intensity cycling bouts would improve a participant’s time to task failure (TTF) and have a positive effect on the commonly seen physiological signs of fatigue.

METHODS


Subjects

Sixteen healthy and recreationally active individuals (Eight males: age 24.5 ± 3.4 yrs., mass 75.2 ± 7.4 kg, height 178.3 ± 6.2 cm, $\text{VO}_{2\text{max}}$ 4.1 ± 0.4 $\text{L}\cdot\text{min}^{-1}$ and eight females: age 23.1 ± 3.0 yrs., mass 65.7 ± 4.7 kg, height 163.9 ± 5.3 cm, $\text{VO}_{2\text{max}}$ 3.4 ± 0.3 $\text{L}\cdot\text{min}^{-1}$) volunteered from the university community to participate in a counterbalanced randomized cross-over design study consisting of a preliminary testing session and two experimental sessions separated by a minimum of two days. All the participants filled out a Physical Activity Readiness Questionnaire from the Canadian Society for Exercise Physiology to determine physical activity level and to screen for a history of cardiovascular, pulmonary, metabolic and orthopedic conditions. The participants read and signed a consent form prior to the study. Subjects were blind to the

101 hypotheses of the study. The Memorial University of Newfoundland Human Investigations
102 Committee approved the study [IRB approval number: 11.26].

103 **Experimental Design**

104 *Preliminary testing session*

105 Upon the completion of questionnaire filling and anthropometric measurements, participants
106 performed a ramp protocol starting at 50 watts at a self-selected cadence (> 60 revolutions per
107 minute – RPM) with increment of 1 watt every 3 second to determine VO_{2max} and peak power
108 output (PPO). The test was terminated when the participants reached one of the following
109 criteria: 1) volitional exhaustion, 2) RPE value ≥ 19 , or 3) RPM ≤ 60 . 

110 *Experimental sessions.*

111 Participants sat on a stationary bike with feet secured while wearing an oro-nasal facemask to
112 record cardiorespiratory parameters for the duration of the experiment. Every minute during the
113 4-min of the high intensity cycling bout, participants were asked to rate their perceived exertion
114 (RPE) on the Borg scale, (6 – 20). After each 4-min high intensity cycling bout, HR and BL were
115 recorded. HR was continuously recorded throughout the session using a Polar® HR monitor.
116 Upon exercise termination, four of the five pre-test measurements (minus blood pressure) were
117 repeated (See description below). Participants were then asked to remain in position and relax for
118 5-min to ensure proper recording of HR and BL recovery (Lambert, 2012). Prior to being
119 released from the experiment an adequate recovery period (resting heart rate ≤ 100 bpm) was
120 given to all participants.

121 **Instrumentation**

Cardiorespiratory Measurements. Oxygen uptake (VO_2), carbon dioxide output (VCO_2), breathing frequency and tidal volume were continuously collected with an automated breath-by-breath system (Metamax, Sensor Medics® version Vmax ST 1.0) using a nafion filter tube and a turbine flow meter (opto-electric). Respiratory exchange ratio (RER) and minute ventilation (VE) were calculated as the quotient of VCO_2 on VO_2 and as the product of breathing frequency by tidal volume, respectively. HR values were transmitted with a Polar HR monitor (PolarElectro, Kempele, Finland). Prior to testing, gas analyzers and volume were calibrated with medically certified calibration gases (16.0% O_2 and 3.98% CO_2) and with a three-liter calibration syringe. In addition, a propane gas calibration was performed to assess the sensitivity of the oxygen and carbon dioxide analysers.

Lactate Measurements. All lactate measurements were taken using the Lactate Pro (LP, Arkay KDK, Japan) hand-held portable analyzer. A blood sample of $\geq 5 \mu\text{L}$ was taken from the participant's fingertip using a spring loaded lancet and then blood lactate values were recorded. The company supplied a check strip to confirm that the analyzer operated correctly, and a calibration strip that provided a non-quantitative indication of analyzer accuracy, which was used at the beginning of each testing session to ensure validity of the measures.

Cycle Ergometer. All exercise protocols were performed on the Velotron Dynafit Pro cycle ergometer (RacerMate, Inc., Seattle, WA). Factory calibration of the cycle ergometer was performed using Velotron CS software (RacerMate, Inc.) and the Accuwatt rundown verification procedure. Individual positional adjustments (saddle and handlebar height) were made before the first exercise test and were replicated for all subsequent exercise tests. Visual feedback of pedalling rate (RPM) was available to the participants during each exercise session.

Music. All participants listened to the exact same playlist of popular music which was set to a 130 bpm tempo; meaning each song was not originally 130 bpm but was altered to keep a consistent tempo throughout the playlist. Music was played through an Ipod® nano using ‘earbud’ type head phones and volume was held constant for each participant at 50% of the maximum volume approximately 65 decibels, based on manufacturer specifications of maximum volume being 130 decibels.

Exercise Bouts. Participants were given an explanation of the 6-20 point Borg RPE scale and told that if at any time they wished to stop exercising they could do so. After participants were fitted with an armband that held an Ipod© nano and ear bud type headphones (headphones and armband were worn regardless of condition), the facemask and mesh headpiece were secured next and hooked up to the indirect calorimetric system. Each participant then had a 5-min warm up period where they were instructed to keep a cadence of 60-70 rpm for the duration of the exercise protocol, a parameter displayed on a large computer screen. In both music and no music conditions, the experimental sessions started at 40% of PPO (active recovery), followed by a 4-min high intensity cycling bout at 80% of PPO. The participants were asked to report the RPE score every minute of the 4-min cycling bout. In addition, at the end of the 4-min high intensity cycling bout BL was assessed. Participants repeated the cycle [4-min work load and 2-min active recovery] until reaching one of the following criteria: 1) volitional exhaustion, 2) RPE value ≥ 19 , or 3) RPM ≤ 60 . At that point of exhaustion, time elapsed from the start of the cycling bouts and final RPE values were recorded. A final BL sample was also taken. Afterwards, they removed the Ipod© and quickly and carefully moved back to the bench to perform subsequent MVC's.

Data Reduction

All data sets were analysed using Sigmaplot (version 10.0; Systat Software Inc). First, cardiorespiratory parameters of the incremental test and of the high intensity cycling bouts were smoothed using second-order polynomial function to determine VO_2 , and its corresponding values of VCO_2 , breathing frequency, and tidal volume. Second, HR was time-aligned with the cardiorespiratory parameters and smoothed using the same data reduction technique. Third, RPE scores were interpolated to produce a continuous linear even data point distribution using a two-dimensional interpolation function and were then time-aligned as above-mentioned. Fourth, the high intensity cycling bout epochs were summed to represent time-to-fatigue. Finally, all above time-aligned parameters were expressed in a relative form to correspond to 25, 50, 75 and 100% of TTF.

Statistical Analysis

All statistical analyses were conducted using Jamovi (version 0.8). Differences between music and no-music conditions in time to task failure (measured in seconds and excluding recovery period) were examined using paired t-test. Cardio-respiratory parameters (Bf , V_T , V_E , VO_2 , VCO_2) and rating of perceived exertion were analyzed with a 2-way analysis of variance [ANOVA] (2 conditions x 4 time [25%, 50%, 75% and 100% of time of failure]). Differences between conditions in blood lactate were measured with 2-way ANOVA (2 conditions x 3 time [pre-test, immediate post-test, 5-min post-test]). HR was analyzed in two ways. First, with a 2-way ANOVA (2 conditions x 4 time [25%, 50%, 75% and 100% of time of failure]). Second, the effect of music upon recovery at the same absolute time period for all subjects (pre-test and 5 min post-test) was analyzed with paired t-tests. Gender was treated as a between-subjects variable in all tests. Differences were considered significant at $p < 0.05$. If significant main effects or interactions were present, a Bonferroni (Dunn) procedure was conducted. Cohen d effect sizes



were also calculated using the following equation ($d = \frac{\text{Mean differences}}{\text{SD average}}$) in which SD average is $\frac{\sqrt{\text{SD condition 1} + \text{SD condition 2}}}{2}$ to provide qualitative descriptors of standardized effects using these criteria: trivial < 0.2, small 0.2-0.5, moderate 0.5-0.8, and large > 0.8 (Cohen 1988). Note that since each participant exercised to volitional fatigue, each testing session was a different length of time; therefore, some variables were collapsed over time or reported as a percentage of TTF.

RESULTS

Time to Task Failure (TTF)

The average TTF in the music condition ($10:30 \pm 3:38$ mins:secs) was 10.7 % longer than the average TTF in the no-music condition ($9:33 \pm 3:42$ mins:secs) ($p = 0.035$; $ES = 0.28$). These averages reported times did not include the time spent in active recovery period (Figure 1A). Therefore on average participants in the music condition completed approximately 2.6 intervals and exercised for one minute longer while participants in the no-music condition only completed approximately 2.3 intervals.

Rate of Perceived Exertion (RPE)

Small magnitude and non-statistically significant ($p \geq 0.26$, $ES = \sim 0.15$) differences were observed between conditions in absolute RPE across the four time points (Figure 1B). A main effect for time was identified in which RPE increased over the course of the exercise across both conditions ($p < 0.001$).

Blood Lactate (BL)

While no statistical differences were observed between conditions at any time point ($p = 0.250$), the average BL levels post-exercise in the music condition at TTF exhibited a moderate magnitude 12.5% higher blood concentration than those at post-exercise in the no-music condition ($13.5 \pm 2.7 \text{ mmol} \cdot \text{L}^{-1}$ vs. $12.0 \pm 3.0 \text{ mmol} \cdot \text{L}^{-1}$; $ES = 0.51$) (Figure 1C).

Heart Rate (HR)

First, a small magnitude and statistically insignificant ($p = 0.223$; $ES = 0.27$) difference was identified between conditions at resting baseline. Second, a main effect for condition was observed in which HR was 4% higher in the music condition across the four time points ($p = 0.043$; $ES = 0.25$) (Figure 1D). HR increased in both conditions over time ($p < 0.001$; $ES = 1.84$). At post 5-minutes HR was 13% lower in the music condition ($99.6 \pm 7.6 \text{ bpm}$) than in the no-music condition ($112.6 \pm 10.6 \text{ bpm}$) ($p < 0.001$; $ES = 1.40$) (Figure 1D).

VO_2 and VCO_2

Small magnitude and statistically insignificant differences were identified in both VO_2 and VCO_2 between conditions and across time ($p \geq 0.192$; $ES \leq 0.31$) (Figure 2A and B).

Ventilation and Tidal Volume

Small magnitude and statistically insignificant differences were identified in ventilation and tidal volume between conditions ($p \geq 0.012$; $ES \leq 0.27$) (Figure 2C and E). A main effect for time was observed in which ventilation increased from beginning to the end of the cycling bout ($p < 0.001$; $ES = 0.91$).

Breathing Frequency (Bf)

A medium magnitude, statistically significant ($ES = 0.57$; $p = 0.006$) main effect for condition was observed in which average Bf was 11.6% higher in the music condition (43.2 ± 8.7 breath \cdot min $^{-1}$) compared to the no-music condition (38.7 ± 8.2 breath \cdot min $^{-1}$) (Figure 2F). A large magnitude and statistically significant main effect ($p < 0.001$; $ES = 1.17$) for time was identified in which Bf increased by 23% over time in both conditions from 34.7 ± 6.2 breath \cdot min $^{-1}$ to 45.5 ± 11.5 breath \cdot min $^{-1}$.

Respiratory Exchange Ratio (RER)

A significant interaction was observed between conditions and time ($p = 0.040$). While post-hoc testing only revealed statistical differences favoring the music condition at TTF (7%; $p = 0.021$; $ES = 1.1$) (Figure 2D). RER was also 5% greater (non-significant with a large effect size magnitude) at 75% of TTF ($p = 0.12$; $ES = 0.85$). RER increased over time across both conditions by 5% ($p = 0.006$; $ES = 0.78$).

DISCUSSION

The primary purpose of this study was to determine the effects of listening to high tempo music (130 bpm) on physical performance and acute physiological responses during high intensity interval bouts of cycling. Listening to music led to a small increased TTF, accompanied by an increase in BF, HR, as well as a steeper heart rate recovery post-exercise in the high tempo music condition. Yet, RPE scores were similar between conditions although participants exercised longer and recovered faster in high tempo music condition.

A possible explanation for the participant's increased TTF while experiencing a build-up of peripheral metabolites (increased BL), albeit still reporting comparable RPE scores, can be

attributed to the CNS regulation and its impacts on exercise performance. Models that emphasise the role of the brain in exercise regulation have pointed to the lack of clear peripherally based fatigue reasons to explain why participants reach the point of exercise failure (i.e., inability to continue exercising) since not all muscle fibers are recruited at the point of exhaustion (Kayser 2003; Noakes 2000). As such, even during high intensity exercise efforts, the brain has a role in regulating effort despite the accumulating metabolites in the muscles. Henceforth, the CNS must have some control over exercise performance (Kayser 2003; Noakes 2000). The increased breathing frequency response of our participants implies that central command cardiovascular drive regulates through medullar integration the dual-talk between the sympathetic and parasympathetic nervous systems even with high intensity exercise. Breathing frequency is controlled via the autonomic nervous system, by the respiratory centers located in the medulla oblongata in the lower brainstem, functioning largely below the level of consciousness (Bechbach and Duffin 1977; Williamson 2010). Hence, at some point during the cycling intervals, performed with high tempo music, the central neural command seemed to modulate the cardiovascular centres. As a result, the participants breathing frequency was higher during high intensity cycling bouts, which could be an indication of the involvement of the cerebral cortex in cardiovascular control mechanisms (Williamson, 2010). Although the exact mechanism for this physiological response is unclear, these findings do support the idea that further research into the underlying cortical modulation mechanisms involved with the beneficial effects of music on exercise duration are warranted.

Similar to most of the current literature on this topic, HR during exercise was only slightly affected by the music. However, there was an anomaly in the present study, which has to do with post-exercise recuperative effects of music. Participants who completed the cycling intervals

with music had a steeper HR recovery. There is very little known about the impact of music on post-exercise performance. Music seems to relieve stress and improve affective states in non-exercise settings (Särkämö et al. 2008). Note that participants in this study had the music stimulus removed immediately at the end of high intensity cycling bouts, however, five minutes post-exercise HR were significantly lower compared to no-music condition. This translates again to a possible effect of distracting stimuli on the autonomic nervous system; whereby attention to the external environment seems to reduce the awareness of physiological sensations and negative emotions, which may have triggered a quicker parasympathetic response leading to steeper post-exercise HR recovery. These results further support the theory that the CNS has the ability to control some types of exercise performance and aid in the ability to exercise longer at higher intensities; unfortunately the verification of these theories were not within the scope of this investigation.

Although the research is somewhat conflicting when it comes to measuring the extent to which music can enhance exercise performance at maximal or near-maximal levels, this study demonstrated that listening to high tempo music (via headphones) during high intensity cycling intervals can lead to greater efforts without increasing individual's perceived exertion. The ability to alter perception is consistent with a vast majority of the current research, which is focused on the psychological effects of music on exercise, mood, and emotion and affect. It is the 'psychophysical' effects or more prudently the physiological effects that have been noted in this study that should motivate future research endeavours. If music can truly distract or disguise the peripheral signs of fatigue it may increase exercise duration, and perhaps enjoyment and adherence. In view of the results presented here, the central motor drive can be uncoupled from the central cardiovascular command to evoke different circulatory responses (Williamson 2010);

indeed, the individuals worked harder while breathing at a higher frequency, and experiencing greater muscle fatigue, all whilst diminishing the feeling of discomfort. The association between central motor drive, central cardiovascular command and perceived exertion was clearly altered by distracting stimuli (high music tempo).

CONCLUSIONS

The music condition in the present study elicited an increase in TTF, breathing frequency, and HR while not influencing RPE. Additionally, a steeper heart rate recovery post-exercise compared to the control condition was observed. These results support the notion that music can modify the interplay between central motor drive, central cardiovascular command and perceived exertion. Changes in factors such as breathing frequency suggest that this modification may occur at a subconscious level. Therefore, further studies of the effect of music on exercise performance should focus on the physiological mechanisms responsible for the beneficial acute physiological responses, including potentially more complex procedures (i.e. fMRI) as opposed to the psychological mechanisms and responses.

REFERENCES

- Atkinson, G., Wilson, D., Eubank, M. 2004. Effect of music on work rate distribution during a cycle time trial. *Int. J. Sports. Med.* **62**: 413-419.
- Barwood, M.J., Westson, N.J.V., Thelwell, R., Page, J. 2009. A motivational music and video improves high-intensity exercise performance. *J. Sports. Sci. Med.* **8**: 435-442.
- Bechbache, R.R., Duffin, J. 1977. The entrainment of breathing frequency by exercise rhythm. *The Journal of Physiology.* **272**(3): 553-561.
- Bharani, A., Sahu, A., Mathew, V. 2004. Effect of passive distraction on treadmill exercise test

322 performance in healthy males using music. *Int. J. Cardiol.* **97**: 305-306.

323 Blanc, Y., Dimanico, U. 2010. Electrode Placement in Surface Electromyography (sEMG)

324 “Minimal Crosstalk Area” (MCA). *The Open Rehabilitation Journal.* **3**: 110 – 126.

325 Boutcher, S.H., Trenske, M. 1990. The effects of sensory deprivation and music on perceived

326 exertion and affect during exercise. *J. Sport. Exerc. Psychol.* **12**(2): 167-176.

327 Cohen J. 1988. *Statistical Power Analysis for the Behavioral Sciences.* 2nd ed. Hillsdale (NJ):

328 Lawrence Earlbaum Associates.

329 Copeland, B.L., Franks, B.D. 1991. Effects of types and intensities of background music on

330 treadmill endurance. *J. Sports. Med. Phys. Fitness.* **31**(1): 100-103.

331 Crust, L., Clough, P.J. 2006. The influence of rhythm and personality in the endurance response

332 to motivational asynchronous music. *J. Sports. Sci.* **24**: 187-195.

333 Dudley, G.A., Djamil, R. 1985. Incompatibility of endurance- and strength-training modes of

334 exercise. *J. Appl. Physiol.* **59**: 1446–1451.

335 Edworthy, J., Waring, H. 2006. The effects of music tempo and loudness level on treadmill

336 exercise. *Ergonomics.* **49**: 1597-1610.

337 Elliott, D., Carr, S., Savage, D. 2004. Effects of motivational music on work output and affective

338 responses during sub-maximal cycling of a standardized perceived intensity. *J. Sport. Behav.*

339 **27**(2): 134-147.

340 Gandevia, S.C. 2001. Spinal and supraspinal factors in human muscle fatigue. *Physiol. Rev.* **81**:

341 1725–1789.

342 Ghaderi, M., Rahimi, R., Azarbayjani, A.M. 2009. The effect of motivational and relaxation

343 music on aerobic performance, rating perceived exertion and salivary cortisol in athlete meals.

344 *South African Journal for Research in Sport, Physical Education and Recreation.* **31**(2): 29-38.

345 Hakkinen, K., Alen, M., Kraemer, W.J., Gorostiaga, E., Izquierdo, Z., Rusko, H., Mikkola, J.,
 346 Hakkinen, A., Valkeinen, H., Kaarakainen, E.K., Romu, S., Erola, V., Ahtianinen, J.,
 347 Paavolainen, L. 2003. Neuromuscular adaptations during strength training and concurrent
 348 strength and endurance training in previously untrained men. *Eur. J. Appl. Physiol.* **89**: 42–52.
 349 Hardy, C.J., Rejeski, W.J. 1989. Not what, but how one feels: the measurement of affect during
 350 exercise. *J. Sport. Exerc. Psychol.* **11**: 304-317.
 351 Hickson, R.C. 1980. Interference of strength development by simultaneously training strength
 352 and endurance. *Eur. J. Appl. Physiol.* **45**: 255–263.
 353 Hill, A.V. 1924. Muscular activity and carbohydrate metabolism. *Science*. **60**: 505-514.
 354 Hunter, G., Demment, R., Miller, D. 1987. Development of strength and maximum oxygen
 355 uptake during simultaneous training for strength and endurance. *J. Sports. Med.* **27**: 269–275.
 356 Iwanaga, I. 1995. Relationship between heart rate and performance for tempo music. *Percept.*
 357 *Mot. Skills.* **81**: 67-71.
 358 Johnson, J., Siegel, D. 1987. Active vs. passive attentional manipulation and multidimensional
 359 perceptions of exercise intensity. *Can. J. Sport. Sci.* **12**: 41-44.
 360 Karageorghis, C.I., Jones, L., Low, D.C. 2006. Relationship between exercise heart rate and
 361 music tempo preference. *Res. Q. Exerc. Sport.* **26**: 240-250.
 362 Karageorghis, C.I., Jones, L., Priest, D.L., Akers, R.I., Clarke, A., Perry, J., Lim, H.B.T. 2011.
 363 Revisiting the exercise heart rate-music tempo preference relationship. *Res. Q. Exerc. Sport.* **82**:
 364 274-284.
 365 Karageorghis, C.I., Jones, L., Stuart, D.P. 2008. Psychological effects of music tempi during
 366 exercise. *Int. J. Sports. Med.* **29**: 613-619.
 367 Karageorghis, C.I., Mouzourides, D., Priest, D.L., Sasso, T., Morrish, D., Whalley, C. 2009.

368 Psychophysical and ergogenic effects of synchronous music during treadmill walking.
369 J. Sport. Exerc. Psychol. **31**: 18-36.

370 Karageorghis, C.I., Priest, D.L. 2012A. Music in the exercise domain: a review and synthesis
371 (Part I). Int. Rev. Sport. Exerc. Psychol. **5**(1): 44-66.

372 Karageorghis, C.I., Priest, D.L. 2012B. Music in the exercise domain: a review and synthesis
373 (Part II). Int. Rev. Sport. Exerc. Psychol. **5**(1): 67-84.

374 Karageorghis, C.I., Terry, P.C. 1997. The psychophysical effects of music in sport and exercise:
375 A review. J. Sport. Behav. **20**: 54-68.

376 Karageorghis, C.I., Terry, P.C., Lane, A.M. 1999. Development and initial validation of an
377 instrument to assess the motivational qualities of music in exercise and sport: The Brunel Music
378 Rating Inventory. J. Sports. Sci. **17**: 713-724.

379 Karageorghis, C.I., Terry, P.C., Lane, A.M., Bishop, D.T., Priest, D.L. 2011. British Association
380 of Sport and Exercise Sciences expert statement on the use of music in exercise. The Sport and
381 Exercise Scientist. **28**: 18-19.

382 Kayser, B. 2003. Exercise begins and ends in the brain. Eur. J. Appl. Physiol. **90**: 411-419.

383 Leveritt, M., Abernethy, P., Barry, B., Logan, P. 1999. Concurrent strength and endurance
384 training: A review. Sports Med. **28**: 413-427.

385 Lindstedt, S.L., Conley, K.E. 2001. Human aerobic performance Too much ado about limits
386 VO₂. J. Exp. Biol. **204**: 3195-3199.

387 Lucaccini, L.F., Kreit, L.H. 1972. Music. In W.P. Morgan (Ed.), Ergogenic aids and muscular
388 performance. New York: Academic Press. pp. 240-245.

389 Macone, D., Baldari, C., Zelli, A., Guidetti, L. 2006. Music and physical activity in
390 psychological well-being. Percept. Mot. Skills. **103**: 285-295.

391 Marcora, S.M. 2008. Do we really need a central governor to explain brain regulation of exercise
392 performance? *Eur. J. Appl. Physiol.* **104**: 929–931.

393 Michel, W., Wanner, H. U. 1973. Effect of music on sports performance. *Schweizerische*
394 *Zeitschrift für Sportmedizin.* **2**: 141-159.

395 Murrock, C.J., Higgins P.A. 2009. The theory of music, mood and movement to improve health
396 outcomes. *J. Adv. Nurs.* **65**(10): 2249-57.

397 Nakamura, P.M., Pereira, G., Papini, C.B., Nakamura, F.Y., Kokubum, E. 2010. Effects of
398 preferred and nonpreferred music on continuous cycling exercise performance. *Percept.*
399 *Mot. Skills.* **110**: 257-264.

400 Nethery, V.M. 2002. Competition between internal and external sources of information during
401 exercise: Influence on RPE and the impact of exercise load. *J. Sports. Med. Phys. Fitness.* **42**:
402 172-178.

403 Noakes, T.D. 2000. Physiological models to understand exercise fatigue and the adaptations that
404 predict or enhance athletic performance. *Scand. J. Med. Sci. Sports.* **10**: 123-125.

405 Noakes, T.D., Gibson, A.S.C. 2004. Logical limitations to the “catastrophe” models of fatigue
406 during exercise in humans. *Br. J. Sports. Med.* **38**: 1-30.

407 Pujol, T.J., Langenfeld, M.E. 1999. Influence of music on Wingate anaerobic test performance.
408 *Percept. Mot. Skills.* **88**: 292-296.

409 Rejeski, W. J. 1985. Perceived exertion: an active or passive process? *J. Sport. Psychol.* **7**: 371-
410 378.

411 Rendi, M.R., Szabo, A., Szaba, T.S. 2008. Performance Enhancement with Music in Rowing
412 Sprint. *Sport Psychologist.* **22**(2): 175-182.

413 Särkämö, T., Tervaniemi, M., Laitinen, S., Forsblom, A., Soinila, S., Mikkonen, M., Autti, T.,

414 Silvennoinen, H.M., Erkkilä, J., Laine, M., Peretz, I., Hietanen M. 2008. Music listening
415 enhances cognitive recovery and mood after middle cerebral artery stroke. *Brain*. **131**: 866-876.

416 Scherer, K.R. 2004. Which emotions can be induced by music? What are the underlying
417 mechanisms? And how can we measure them? *Journal of New Music Research*. **33**: 239-251.

418 Schie, N.A., Stewart, A., Becker, P., Rogers, G.G. 2008. Effect of music on submaximal cycling.
419 *Scand. J. Med. Sci. Sports*. **20**: 28-31.

420 Schücker, L., Hagemann, N., Strauss, B., Völker K. 2009. The effect of attentional focus on
421 running economy. *J. Sports. Sci.* **27**(12): 1241-1248.

422 Schwartz, S. E., Femhall, B., Plowman, S. A. 1990. Effects of Music on Exercise Performance. *J.*
423 *Cardiopulm. Rehabil.* **10**: 312-316.

424 Simpson, S.D., Karageorghis, C.I. 2006. The effects of synchronous music on 400-m sprint
425 performance. *J. Sports. Sci.* **24**: 1095-1102.

426 Szmedra, L., Bacharach, B.W. 1998. Effect of music on perceived exertion, plasma lactate,
427 norepinephrine and cardiovascular hemodynamics during treadmill running. *Int. J. Sports. Med.*
428 **19**(1): 32-37.

429 Tenenbaum, G., Lidor, R., Lavyan, N., Morrow, K., Tonnel, S., Gershgoren, A., Johnson, M.
430 2004. The effect of music type on running perseverance and coping with effort sensations.
431 *Psychol. Sport. Exerc.* **5**: 89-109.

432 Terry, P.C., Karageorghis, C.I. 2011. Music in sport and exercise. In T. Morris and P.C. Terry
433 (Eds.). *The new sport and exercise psychology companion*. Morgantown, WV: Fitness
434 Information Technology. pp. 359-380.

435 Uppal, A. K., Datta, U. 1990. Cardiorespiratory response of junior high school girls to exercise
436 performed with and without music. *Journal of Physical Education and Sport Science*. **2**: 52-56.

Waterhouse, J., Hudson, P., Edwards, B. 2009. Effects of music tempo upon submaximal cycling performance. *Scand. J. Med. Sci. Sports.* **20**(4), 662-669.

Yamamoto, T., Ohkuwa, T., Itoh, H., Kitoh, M., Terasawa, J., Tsuda, T., Sato, Y. 2003. Effects of pre-exercise listening to slow and fast rhythm music on supramacimal cycle performance and selected metabolic variables. *Arch. Physiol. Biochem.* **111**: 211-214.

Yamashita, S., Iwai, K., Akimoto, T., Sugawara, J., Kono, I. 2006. Effects of music during exercise on RPE, heart rate and the autonomic nervous system. *J. Sports. Med. Phys. Fitness.* **46**: 425-43.

Figure captions

Figure 1. All figures include means+SD. **A:** TTF= Time to failure. **B:** RPE= Rate of perceived exertion. **C:** BL= Blood lactate. **D:** HR= Heart rate.

Figure 2. All figures include means+SD. **A:** VO_2 = Maximal oxygen uptake. **B:** VCO_2 = Carbon dioxide output. **C:** V_T = Tidal volume. **D:** RER= Respiratory exchange ratio. **E:** VE = Minute ventilation. **F:** **Bf** = Breathing frequency.

Figure 1

Performance measures

All figures include means+SD. **A:** TTF= Time to failure. **B:** RPE= Rate of perceived exertion. **C:** BL= Blood lactate. **D:** HR= Heart rate.




Figure 2

Metabolic data

All figures include means+SD. **A:** $\dot{V}O_2$ = Maximal oxygen uptake. **B:** $\dot{V}CO_2$ = Carbon dioxide output. **C:** V_T = Tidal volume. **D:** RER= Respiratory exchange ratio. **E:** $\dot{V}E$ = Minute ventilation. **F:** Bf = Breathing frequency.

