

Taxifolin protects rat against myocardial ischemia/reperfusion injury by modulating the mitochondrial apoptosis pathway

Zhenqiu Tang¹, Chunjuan Yang^{2,3}, Baoyan Zuo⁴, Yanan Zhang¹, Gaosong Wu¹, Yudi Wang¹, Zhibin Wang

Corresp.¹

¹ Key Laboratory of Chinese Materia Medica, Heilongjiang University of Traditional Chinese Medicine, Harbin, Heilongjiang, China

² College of Pharmacy, Harbin Medical University, Harbin, Heilongjiang, China

³ Beijing Shunyue Technology Co., Ltd., Beijing, China

⁴ School of Pharmacy, Shenyang Pharmaceutical University, Shenyang, Liaoning, China

Corresponding Author: Zhibin Wang

Email address: wz@mailbox@hljucm.net

Background: Taxifolin (TAX), is an active flavonoid, that plays an underlying protective role on the cardiovascular system. This study aimed to evaluate its effect and potential mechanisms on myocardial ischemia/reperfusion (I/R) injury. **Methods:** Healthy rat heart was subjected to I/R using the Langendorff apparatus. Hemodynamic parameters, including heart rate (HR), left ventricular developed pressure (LVDP), maximum/minimum rate of the left ventricular pressure rise ($+dp/dt_{max}$ and $-dp/dt_{min}$) and rate pressure product (RPP) were recorded during the perfusion. Histopathological examination of left ventricular was measured by hematoxylin-eosin (HE) staining. Creatine kinase-MB (CK-MB) and lactate dehydrogenase (LDH) activities in the effluent perfusion, and the levels of malondialdehyde (MDA), superoxide dismutase (SOD) and glutathione peroxidase (GSH-PX) in the tissue were assayed. Apoptosis related proteins, such as B-cell lymphoma-2 (Bcl-2), Bcl2-associated X (Bax) and cytochrome c (Cyt-c) were also assayed by ELISA. Western blot was employed to determine apoptosis-executive proteins, including caspase 3 and 9. TUNEL assay was performed to evaluate the effect TAX on myocardial apoptosis. **Results:** TAX significantly improved the ventricular functional recovery, as evident by the increase in LVDP, $+dp/dt_{max}$, $-dp/dt_{min}$ and RPP, The levels of SOD, GSH-PX were also increased, but those of LDH, CK-MB, and MDA were decreased. Furthermore, TAX up-regulated the Bcl-2 protein level but down-regulated the levels of Bax, Cyt-c, caspase 3 and 9 protein, thereby inhibits the myocardial apoptosis. **Discussion:** TAX treatment remarkably improved the cardiac function, regulated oxidative stress and attenuated apoptosis. Hence, TAX has a cardioprotective effect against I/R injury by modulating mitochondrial apoptosis pathway.

Taxifolin protects rat against myocardial ischemia/reperfusion injury by modulating the mitochondrial apoptosis pathway

Zhenqiu Tang^{1,†}, Chunjuan Yang^{2,3,†}, Baoyan Zuo⁴, Yanan Zhang¹, Gaosong Wu¹, Yudi Wang¹, Zhibin Wang¹, *

¹ Key Laboratory of Chinese Materia Medica, Heilongjiang University of Chinese Medicine, Harbin, Heilongjiang, China

² College of Pharmacy, Harbin Medical University, Harbin, Heilongjiang, China

³ Beijing Shunyu Technology Co., Ltd., Beijing, China

⁴ School of Pharmacy, Shenyang Pharmaceutical University, Shenyang, Liaoning, China

† These authors contribute equally to this paper.

Corresponding Author:

Zhibin Wang

No. 24 Heping road, Harbin, Heilongjiang, 150040, China

Email address: wzmailbox@hljucm.net

ABSTRACT

Background: Taxifolin (TAX), is an active flavonoid, that plays an underlying protective role on the cardiovascular system. This study aimed to evaluate its effect and potential mechanisms on myocardial ischemia/reperfusion (I/R) injury.

Methods: Healthy rat heart was subjected to I/R using the Langendorff apparatus. Hemodynamic parameters, including heart rate (HR), left ventricular developed pressure (LVDP), maximum/minimum rate of the left ventricular pressure rise (+dp/dt_{max} and -dp/dt_{min}) and rate pressure product (RPP) were recorded during the perfusion. Histopathological examination of left ventricular was measured by hematoxylin-eosin (HE) staining. Creatine

kinase-MB (CK-MB) and lactate dehydrogenase (LDH) activities in the effluent perfusion, and the levels of malondialdehyde (MDA), superoxide dismutase (SOD) and glutathione peroxidase (GSH-PX) in the tissue were assayed. Apoptosis related proteins, such as B-cell lymphoma-2 (Bcl-2), Bcl2-associated X (Bax) and cytochrome c (Cyt-c) were also assayed by ELISA. Western blot was employed to determine apoptosis-executive proteins, including caspase 3 and 9. TUNEL assay was performed to evaluate the effect TAX on myocardial apoptosis.

Results: TAX significantly improved the ventricular functional recovery, as evident by the increase in LVDP, $+dp/dt_{max}$, $-dp/dt_{min}$ and RPP. The levels of SOD, GSH-PX were also increased, but those of LDH, CK-MB, and MDA were decreased. Furthermore, TAX up-regulated the Bcl-2 protein level but down-regulated the levels of Bax, Cyt-c, caspase 3 and 9 protein, thereby inhibits the myocardial apoptosis.

Discussion: TAX treatment remarkably improved the cardiac function, regulated oxidative stress and attenuated apoptosis. Hence, TAX has a cardioprotective effect against I/R injury by modulating mitochondrial apoptosis pathway.

INTRODUCTION

Ischemic heart disease is a threat to human health. Extracorporeal circulation and coronary bypass surgery are usually employed to improve myocardial ischaemia after myocardial infarction occurs. The timely restoration of blood flow to the ischemic myocardium (reperfusion) became the standard treatment for these patients. However, reperfusion may cause additional heart damage. This condition is referred to as cardiac ischaemia/reperfusion (I/R) injury ([Braunwald et al., 2012](#)). The reperfusion of ischaemic tissues is often associated with microvascular dysfunction. The mechanisms may involve the release of oxygen radicals and inflammatory mediators ([Carden et al., 2000](#)). Nowadays, tissue injury induced by I/R is a major factor, which often cause death. During myocardial I/R injury, cardiomyocytes undergo death at an increased frequency, mainly including necrosis and apoptosis ([Gottlieb et al., 1994](#)). Apoptosis involves programmed cell death, which is the vital pathological process in acute reperfusion injury ([Konstantinidis et al., 2012](#)). When the amount of cardiomyocyte decreases, the heart may undergo ventricular remodelling, compensatory cardiac hypertrophy and eventually lead to heart failure ([Pangonyte et al., 2008](#); [Du et al., 2010](#)). Therefore, exploring the detailed mechanisms that trigger cardiomyocyte death and the means to prevent it during I/R injury is still a public issue.

In I/R injury, morphological changes in myocardial tissue can be observed, including microvascular damage and myocardial cell edema. The symptoms of I/R include myocardial enzyme release, arrhythmias and weak systolic function ([Naito et al., 2000](#)). Generally, this reperfusion damage is caused by increased free radical activity. When circulating blood decreases, the level of oxygen supply cannot sustain the oxygen demand by cardiomyocytes, and

the aerobic metabolism turns into anaerobic metabolism (*Giordano et al., 2005*). Anaerobic metabolism leads to the production of lactic acid which results in disturbances in ionic homeostasis. A timely reperfusion is crucial for the recovery of an ischemic myocardium, but by the sudden re-appearance of circulating blood to the dying myocardium, oxygen species (ROS) will be produced as a response to hyperoxia which can worsen the functional situation of the organization (*Akhlaghi et al., 2009*).

Flavonoid is the most prevalent class of naturally occurring compound and is ubiquitous in woody and herb plants. It exerts multiple biochemical properties and wide pharmacological effects (*Moon et al., 2006*). Epidemiological studies have shown that flavonoid is associated with a reduced risk of cardiovascular diseases (*Raj Narayana et al., 2001; Bjorklund et al., 2017*). Fisetin, a plant-derived bioflavonoid, significantly attenuated I/R-induced tissue injury, blunted the oxidative stress, and restored mitochondrial structure and function (*Shanmugam et al., 2018*). Quercetin has been demonstrated to improve post ischemic recovery of the isolated heart of rats after global ischemia (*Barteková et al., 2010*). TAX exerts anti-inflammatory effects and prevents oxidative stress-induced injury in human endothelial cells (*Guo et al., 2015*) and rat peritoneal macrophages (*Arutyunyan et al., 2016*). It also possesses free radical scavenging, antioxidant and anti-inflammatory effects (*Sun et al., 2014; Xie et al., 2017*). TAX is structurally similar to quercetin. Hence, we suspect that it also has a beneficial effect on the cardiovascular system. Recent studies demonstrated that TAX could inhibit cardiac hypertrophy and attenuate ventricular fibrosis after pressure overload. These beneficial effects were at least mediated by suppressing oxidative stress and the excess production of ROS. (*Guo et al., 2015; Sun et al., 2014*). However, the potential of TAX for I/R protection remains unclear. TAX is a potential candidate for the prevention or treatment for I/R injury. However, the influence of TAX on the injury of I/R in isolated rat hearts has not been reported. In this study, we aimed to evaluate the cardioprotective effects of TAX and investigated the mechanisms underlying these effects in isolated hearts of rats.

Myocardial ischemic events are unpredictable. The clinical application of pre-conditioning drugs is limited. Therefore, researchers turned to a new endogenous protective strategy, which is post-conditioning. In 2006, the protective effect of post-conditioning in ischemic reperfusion on cerebral ischemia was first reported (*Zhao et al., 2006*). Subsequent studies have validated the protective effect of post-conditioning in various global ischemia models *ex vivo* and ischemia and hypoxia models *in vitro*. Flavonoids reduce the injury of myocardial ischemia on perfusion in a post-conditioning way (*Wang et al., 2017; Xuan et al., 2016*). Therefore, we adopted the post-conditioning method in this study.

MATERIALS & METHODS

Experimental animals and treatment

Male SD rats (280-300 g each) were obtained from the Laboratory Animal Center of

Heilongjiang Medicine University Medical (License Number: SCXK (hei) 2013-004). The rats were housed under standard conditions with natural light (12 hr) and dark (12 hr) at 22 ± 2 °C. Rats were fed with common laboratory chow and allowed to drink tap water ad libitum during the experimental period. The investigation conformed to Guide for the Care and Use of Laboratory Animals (revised, 1996, <http://dels.nas.edu/resources/static-assets/ilar/miscellaneous/GUIDE1996.pdf>). All animal experiments were approved by the College of Pharmacy of Heilongjiang University of Chinese Medicine, Animal Ethics Committee (Approval number: SYXK (hei)-2013-012).

Reagents and antibodies

TAX (purity $\geq 98\%$) was purchased from Sigma-Aldrich (St. Louis, MO, USA). Creatine kinase-MB (CK-MB), lactate dehydrogenase (LDH), malondialdehyde (MDA) glutathione peroxidase (GSH-PX) and superoxide dismutase (SOD) assay kit were obtained from Nanjing Jiancheng Bioengineering Institute (Jiangsu, China). The enzyme-linked immunosorbent assay kit about mitochondrial apoptosis [B-cell lymphoma-2 (Bcl-2), Bcl2-associated X (Bax) and cytochrome c (Cyt-c)] was obtained from Cloud-Clone Corp (Houston, TX, USA). Monoclonal primary antibodies anti- β -actin, anti-active caspase 3 and 9 were purchased from Abcam (MA, USA). Fluorescent-labeled goat anti-rabbit IgG secondary antibody was obtained from LI-COR Biosciences (Lincoln, Nebraska, USA).

Experimental protocol

TAX was dissolved in ethanol (15.21, 45.63 mg/ml) and then transferred into Krebs–Henseleit (K–H) solution. Ethanol solution (0.1 mL) was added into 1 L K–H solution. The final concentration was 5 or 15 μ M. Rats were randomly divided into four groups (n=8): Normal control group (Control); Myocardial I/R control group (I/R); I/R + TAX 5 μ M treatment group (TAX 5 μ M); I/R + TAX 15 μ M treatment group (TAX 15 μ M). The experimental protocol is shown in Fig. 1. Control group: The hearts were subjected a continuous perfusion of K–H solution for 120 min. I/R group: The hearts were perfused for 30 min to stabilization. Subsequently, global ischemia was performed at 37 °C for 30 min, followed by reperfusion with K–H solution for 60 min. TAX 5 μ M group: The hearts were perfused for 30 min to stabilization. Subsequently, global ischemia was performed for 30 min at 37 °C, followed by reperfusion with 5 μ M of TAX-saturated K–H solution for 60 min. TAX 15 μ M group: The hearts were perfused for 30 min to stabilization. Subsequently, global ischemia was performed for 30 min, followed by reperfusion perfused with 15 μ M of TAX-saturated K–H solution for 60 min.

Langendorff preparation

After anesthetization via intraperitoneal injection (chloral hydrate solution, 300 mg/kg), rat hearts were quickly removed and subsequently perfused in the Langendorff apparatus. The perfusion was performed for 30 min in a modified K–H buffer gassed with 95 % O₂ and 5 % CO₂ at a constant flow velocity and constant temperature (37 °C). The composition of K–H buffer as

the following (mM): NaCl 118, MgSO₄ 1.2, KCl 3.2, NaHCO₃ 25, KH₂PO₄ 1.18, CaCl₂ 2.5 and glucose 5.5. After equilibration, 30 min global ischemia was induced followed by 60 min of reperfusion. Control group utilized the same protocol, but no ischemic induction was used. Water-filled balloon that is inserted into the left ventricular cavity was used to monitor hemodynamic parameters. The left ventricular end-diastolic pressure (LVEDP) was maintained at 5-10 mmHg by adjusting the size and position of the balloon. The whole procedure was completed within 2 min. The inclusion criteria of experimental samples were a heart rate (HR) of > 250 beats/min and a left ventricular developed pressure (LVDP) of > 75 mmHg in equilibrium period. In the experiment, we prepared 12 rats in each group with an average of three or four failures. Finally, only eight rats from each group were used in the subsequent experiment. The hemodynamic parameters were recorded during perfusion, including heart rate (HR), LVDP, the maximum/minimum rate of left ventricular pressure rise (+dp/dt_{max} and -dp/dt_{min}), which are important indices to evaluate the left ventricular systolic and diastolic function. Rate pressure product (RPP) = HR × LVDP. After 60 minutes of reperfusion, heart tissue was taken from Langendorff apparatus. The left ventricles of three hearts were cut from each group and then fixed in 10 % neutral formalin. The other tissues were immediately placed in the freezer at -80 °C.

Histopathological evaluation of left ventricle sections

For histopathological examination, the cut left ventricle of heart tissues were fixed in 10 % neutral formalin at room temperature. After 2 hours, the tissue piece was embedded in paraffin. Next, the piece was cut into 3 μm thick tissue sections, after which it was subjected to hematoxylin-eosin (H&E) staining. At least three samples from each group were evaluated. The tissue sections were visualized under light microscope (Olympus BX60, Tokyo, Japan).

Estimation of cardiac damage

In present study, heart tissue injury was assessed by determining the concentration of LDH and CK-MB in the perfusate. The LDH and CK-MB content in the perfusate were measured using the assay kit following the manufacturer's instructions. Samples of the perfusate were collected from the isolated heart at 25, 63, 90 and 120 min of perfusion.

Measurements of anti-oxidant indices

The hearts tissue was cut into small pieces of tissue and then was ground with lysate buffer by using a glass homogenizer. Supernatant of tissue homogenate was frozen for each tissue analysis. MDA, SOD and GSH-PX activity were assessed using commercial ELISA kits following the manufacturer's instructions. All enzyme activities were normalized to the total protein concentrations, which were determined using a bicinchoninic acid (BCA) protein assay kit (Beyotime, Shanghai, China).

Estimation of Cyt-c, Bcl-2 and Bax levels

Following the instructions of nuclear and cytoplasmic protein extraction kit (Beyotime, Shanghai, China), heart tissue samples were weighed, minced into small pieces and homogenized using a glass homogenizer on ice (w: v = 1:30, 1 mL lysis buffer was added in 30 mg tissue sample). The homogenates were centrifuged at 1500× g for 5 min at 4 °C and the cytoplasmic protein was obtained from the supernatant. The Cyt-c, Bcl-2 and Bax protein levels were measured according to the manufacturer's instructions of commercial kits through an ELISA-type method (Cloud Clone Corp, Houston, USA).

Western blotting analysis

Myocardial tissue samples were lysed with RIPA buffer containing protease inhibitors for 15 min on ice. The total lysates were clarified by centrifugation, and supernatants were collected. Protein samples (20-25 mg per lane) were loaded on the gels and then separated by 10 % sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) under reducing conditions and transferred onto the nitrocellulose membrane (Roche, Mannheim, Germany). The membrane was washed with PBS with 0.1% Tween-20 (PBST) and blocked with 5% skim milk in shaking table for 2 h at room temperature. Then the membrane was washed with PBST and incubated with antigen-specific rabbit IgG antibodies (anti-caspase 3 and 9, anti-β-actin; Abcam) diluted to 1:1000 in PBST. Next, the membrane was washed with PBST and incubated with fluorescent-labeled goat anti-rabbit secondary antibodies (LI-COR, Lincoln, Nebraska, USA) diluted to 1:2500 in PBST for 2 h at 4 °C. The target protein bands were scanned using the blot imaging system GelLogic 212 PRO (Carestream, Rochester, NY, USA) after washing with PBST. The obtained images were quantified as the final results by image J 1.4.3 (www.imagej.nih.gov/ij). The results were expressed as the fold induction, which were than compared with the normal control.

TUNEL assay

Heart sections of 3 μm thickness were obtained using a microtome. The sections were deparaffinized in xylene and rehydrated in concentration gradient of ethanol (100%, 95%, 75%). Following this, sections were then incubated with proteinase K and with 30% H₂O₂ to enhance tissue permeability and diminish any endogenous peroxidase activity respectively. Apoptosis was determined using a terminal deoxynucleotidyl transferase-mediated dUTP-X nick end labeling (TUNEL) assay kit (Roche, Basel, Switzerland) according to the manufacturer's protocol. The sections were incubated with complete labeling reaction buffer for 60 min and antibody solution for 30 min. Chromogenic reaction was visualized using 3,3-diaminobenzidine (DAB). Sections were visualized under light microscope (Olympus BX60, Tokyo, Japan).

Statistical Analysis

SPSS16.0 for Windows (SPSS Inc., Chicago, USA) was used for statistical analysis. All data were expressed as mean ± standard deviation (SD). For comparisons between groups, the one-

way ANOVA or student T-test was used where appropriate. Statistical differences were considered significant at $P < 0.05$. # $P < 0.05$ and ## $P < 0.01$ vs. control group. * $P < 0.01$ and ** $P < 0.001$ vs. I/R group.

RESULTS

Effects of TAX on Cardiac Parameters of Isolated Hearts

We examined whether TAX could protect the hearts of rat against *ex vivo* I/R injury. Results showed no obvious alteration in the average HR of isolated heart during reperfusion with or without TAX. In addition, no significant HR change was observed between I/R and normal groups during the 30 min of ischemia and 60 min of reperfusion (Fig. 2A). After reperfusion, LVDP, $+dp/dt_{max}$ and $-dp/dt_{min}$ from different treatment groups decreased in varying degrees. For instance, LVDP was significantly increased in TAX 15 μ M group compared with I/R group at the end periods of reperfusion (LVDP = 68 mmHg vs. 52 mmHg, $P < 0.05$, Fig. 2B). In comparison with I/R group, treatment with 15 μ M TAX significantly improved the RPP in rat at 60 min of reperfusion (RPP = 15294 mmHg \times beats/min in TAX vs. 10643 mmHg \times beats/min in I/R, $P < 0.01$, Fig. 2E). Results showed that TAX treatment improved the cardiac function recovery of rats during myocardial I/R injury.

TAX Down-regulated the Release of LDH and CK-MB

At different time points of perfusion, heart effluents were collected. The LDH level in the whole perfusion process was not conspicuously altered in control group. The perfusate LDH activity of I/R group was improved compared with that of control group after reperfusion and was significantly increased at 60 min of reperfusion ($P < 0.01$). TAX highly reduced the LDH levels compared with I/R group at 60 min of reperfusion (Fig. 3A). CK-MB release was similar to the LDH release. The expression in the whole perfusate process was not conspicuously changed in the control group. In the perfusion stabilization, no significant difference was observed in the CK-MB level among the four groups. However, the levels in I/R group were markedly higher after 30 min of reperfusion compared with perfusion stabilization ($P < 0.05$). Interestingly, the CK-MB release in both TAX 5 and 15 μ M group was significantly decreased at end of reperfusion compared with I/R group ($P < 0.05$ or $P < 0.01$) (Fig. 3B). These results indicated that TAX could protect the cardiac function against I/R injury.

Effect of TAX on Myocardial Morphology

Histopathological examination of myocardial tissue was assessed by H&E staining. Typical micrographs of the myocardial structure are shown in Fig. 4. In control group (Fig. 4A), The morphology of the myocardial tissue was normal. Cardiomyocytes are arranged closely, the intercellular space is small, and edema does not exist between cells. By contrast, I/R group (Fig. 4B) showed degenerated muscle fibers and obvious contraction band, severe obvious cells edema, many infiltrated inflammatory cells. Fig. 4C shows that the TAX 5 μ M group maintained the myocardium with only slightly irregularly arranged fibers and a few contraction bands. Fig. 4D shows that the TAX 15 μ M group showed orderly cardiomyocytes but a few cell dissolution and degeneration. Results showed that treatment with 15 μ M TAX significantly reduced I/R injury compared with I/R group.

Effect of TAX on I/R-induced Oxidative Stress in the Myocardium

To explore the cardio-protective mechanism of TAX, the effects of TAX on SOD, GSH-PX and MDA activity were investigated in myocardial tissue in response to I/R injury. Fig. 5 shows that TAX 15 μ M group, the SOD and GSH-PX activity were increased significantly compared with those of in I/R group ($P < 0.01$), whereas no significant difference was observed in the TAX 5 μ M group. Conversely, these TAX treatment groups showed that MDA production was significantly reduced ($P < 0.01$) compared with I/R group.

TAX Protects Myocardial Cell from I/R-induced Mitochondrial Damage

To evaluate if the effect of TAX is mediated through attenuation of the mitochondrial damage, we determined cytochrome c (Cyt-c) in cytosol. Fig. 6A shows that I/R increased the cytosolic Cyt-c levels ($P < 0.01$). By comparison, TAX at different doses could reduce I/R-induced increase of Cyt-c levels ($P < 0.01$). The result suggested that TAX attenuated I/R induced Cyt-c release to the cytosol. The Bcl-2 family of proteins modulates the release of Cyt-c by regulating mitochondrial transmembrane potential. We also determined Bcl-2 and Bax protein levels. Compared with the control group, I/R down-regulated Bcl-2 but up-regulated Bax protein expression (Figs. 6B-6C). TAX increased Bcl-2 levels but decreased Bax levels compared with I/R. These results indicate that TAX may protect mitochondria against apoptosis by regulating the expression of the Bcl-2 family proteins.

TAX Attenuates Myocardial I/R-induced Apoptosis

To explore the potential mechanism of TAX in rats subjected to I/R-induced myocardial injury, we investigated the protein expressions of active-caspase 3 and 9. Compared with control

group, the expression of active-caspase 3 was up-regulated in I/R group. While compared with I/R group, TAX treatment group significantly reduced the level of active-caspase 3 (Fig. 7C). In TAX treatment group, the expression of active-caspase 9 was down-regulated in 15 μ M ($P < 0.05$), but did not change significantly in 5 μ M (Fig. 7D). In addition, TUNEL assay was performed to evaluate the effect TAX on myocardial apoptosis. An increased number of TUNEL positive cells were observed in I/R group in comparison to control group. Furthermore, a less number of TUNEL positive cell was present in the TAX treatment group (Fig. 8). Results indicated that TAX inhibited apoptotic level in heart injury induced by I/R.

DISCUSSION

Growing evidence indicates a therapeutic action of TAX in cardiovascular disease. However, the implications of TAX in I/R injury remain unclear. To the best of our knowledge, this study was the first to evaluate the cardioprotective effects of TAX in isolated rat heart subjected to I/R injury. We demonstrated that an important role of TAX in improving cardiac function, and inhibiting oxidative stress and apoptosis in a model of I/R injury *in vitro*.

In the late 19th century, Oscar Langendorff pioneered the isolated perfused heart model. Since then, the procedure has been used to probe the pathophysiology of I/R with the dawn of molecular biology (Bell *et al.*, 2011). Today, the Langendorff heart assay is a predominant technique *in vitro*, which is used in physiological and pharmacological research. It allows the examination of cardiac contractile strength and heart rate without complications of an intact animal (Herr *et al.*, 2015). Therefore, we determined the effect of TAX to the cardiac function parameters of isolated heart in myocardial I/R injury by using the Langendorff equipment. Cardiac functions mainly depend on the contraction and relaxation properties of the ventricular muscle. Changes in cardiac function are strongly linked to the severity of I/R injury (Mehdizadeh *et al.*, 2013). The data from our analyses showed that I/R can cause marked myocardial dysfunction, including the reduction of LVDP, $+dp/dt_{max}$ and $-dp/dt_{min}$. TAX treatment significantly improved cardiac diastolic dysfunction but did not alter the average HR in isolated heart.

LDH is a marker of cellular damage and common disease due to its mass release to plasma during tissue injuries, such as heart failure. CK-MB, which is expressed extensively in cardiac myocyte, was often tested in the serum as an indicator of rhabdomyolysis damage, myocardial damage and acute kidney injury in clinic (Moghadam-Kia *et al.*, 2016). The increase in LDH and CK-MB levels in the organ perfusate after ischemia is a direct evidence of cardiac damage (Houshmand *et al.*, 2009). In comparison with control group, LDH and CK-MB activity were significantly increased in I/R group during myocardial I/R injury. Perfusate LDH and CK-MB activity in the TAX treatment groups, particularly in TAX 15 μ M group, were remarkably reduced compared with those in I/R group, which is consistent with the observation of changing cardiac function parameters. In addition, histopathological examination was implemented to

assess the effect of TAX on myocardial morphology. The results of pathomorphological research in the heart samples in I/R group show acute myocardial damage, and TAX causes favorable morphological changes in the heart during I/R injury. These results supported the potential application of TAX as a cardioprotective agent in myocardial I/R injury.

Under normal conditions, tissues could maintain the balance between generation and clearance of ROS. However, the balance is disrupted during I/R and causes significant increase in ROS (Becker *et al.*, 2004). Excess ROS can oxidize lipids, proteins and DNA, which cause dysfunction of these molecules, resulting in the degeneration of tissue function (Kleikers *et al.*, 2012). Minimizing the ROS production is an important strategy to prevent cardiomyocyte I/R injury (He *et al.*, 2016). Therefore, the activation of the anti-oxidant enzyme system is necessary to reduce oxidative stress-induced tissue damage (Matsushima *et al.*, 2013). The SOD and GSH-PX rate are used to evaluate tissue per-oxidative injury (Maciejczyk *et al.*, 2017). In addition, MDA is an index to evaluate the severity of lipid peroxidation, which is produced by lipid peroxidation, resulting in the destruction of structural proteins and cellular structures (Pizzimenti *et al.*, 2013). Our results showed that SOD and GSH-PX activities were conspicuously increased, whereas MDA level was dramatically decreased by TAX, especially in TAX 15 μ M group. Therefore, TAX exhibited the cardioprotective effects by enhancing the antioxidant activity and inhibiting free radical peroxidation.

Mitochondrial damage plays an important role in I/R-induced injury. It is the final arbitrator for I/R-induced cell apoptosis (Powers *et al.*, 2007). During ischemia and mainly during the early period of reperfusion, excessive ROS causes myocardial Ca^{2+} overload and the opening of the mitochondrial permeability transition pore (mPTP), which can reduce mitochondrial function and finally result in an increase in myocardial cell apoptosis (Garciaarena *et al.*, 2011; Halestrap *et al.*, 2015). One of the ways of cell apoptosis is activated by the release of Cyt-c from the mitochondria to the cytosol. In our study, results showed that TAX can weaken the observed increase in the expression of Cyt-c in cytosol. It is very likely that the increased cytosolic content of Cyt-c, which mediates apoptosis, while its expression in mitochondria was not changed (Lundberg *et al.*, 2004). Further investigation of the pathological changes in myocardial tissues by TUNEL assays showed different degrees of apoptosis. Results indicated a positive effect of TAX in the inhibition of apoptosis. Therefore, it can make an assumption that down-regulation of Cyt-c result from TAX attenuated apoptotic processes.

As an important mitochondrial regulator during myocardial apoptosis, Bcl-2 exerts anti-apoptotic effects by blocking the release of Cyt-c and reducing caspase activity. Apoptosis-related proteins, caspase 3 and 9, also play crucial roles in apoptosis. The caspase apoptotic pathway responds to death signals by releasing apoptosis-inducing factor from the mitochondria, which were then translocated to the nucleus (D'Amelio *et al.*, 2012). In this study, Bcl-2, an anti-apoptotic protein and Bax, a pro-apoptotic protein were used to assess the effects of TAX on cardiomyocytes apoptosis. The result demonstrated that TAX treatment increased the protein expression of Bcl-2, and significantly reduced the Bax expression compared with I/R group.

Caspase 3 and 9 were tested to measure the apoptotic level in the isolated heart after I/R injury. We found that the increased expression of the active form of caspase 3 and 9 under ischemic conditions and their expression were decreased in TAX group. Consistent with these results, treatment with TAX significantly decreased myocardial apoptosis by regulating the expression of apoptosis-related proteins, including Bax, Bcl-2 and caspase 3 and 9. These findings suggest that the inhibition of apoptosis is closely related to the underlying beneficial effect of TAX in I/R injury. Cardiomyocytes death occurs during I/R injury by apoptosis, by necrosis and in association with autophagy (*Whelan et al., 2010; Gatica D et al., 2015*). In this study, we studied the effect of apoptosis on myocardial injury, but the effect of other pathways on myocardial injury was not excluded, which requires further exploration.

CONCLUSIONS

In conclusion, TAX exerted cardioprotective effects against I/R injury by inhibiting oxidative stress and cardiac myocyte apoptosis. The underlying mechanism for these phenomena may involve modulation of mitochondrial apoptosis pathway. Our finding provides a novel thought for therapeutic development as an adjuvant therapy to I/R injury.

REFERENCES

- Akhlaghi M, Bandy B. 2009.** Mechanisms of flavonoid protection against myocardial ischemia–reperfusion injury. *Journal Molecular Cellular Cardiology* **46(3)**:309–317.
- Arutyunyan TV, Korystova AF, Kublik LN, Levitman MKh, Shaposhnikova VV, Korystov YN. 2016.** Taxifolin and fucoidin abolish the irradiation-induced increase in the production of reactive oxygen species in rat aorta. *Bulletin of Experimental Biology and Medicine* **160(5)**:635–638.
- Barteková M, Carnická S, Pancza D, Ondrejčáková M, Breier A, Ravingerová T. 2010.** Acute treatment with polyphenol quercetin improves postischemic recovery of isolated perfused rat heart after global ischemia. *Canadian Journal of Physiology and Pharmacology* **88(4)**:465–471.
- Becker LB. 2004.** New concepts in reactive oxygen species and cardiovascular reperfusion physiology. *Cardiovascular Research* **61(3)**:461–470.
- Bell RM, Mocanu MM, Yellon DM. 2011.** Retrograde heart perfusion: the Langendorff technique of isolated heart perfusion. *Journal of Molecular and Cellular Cardiology* **50(6)**:940–950.
- Bjorklund G, Dadar M, Chirumbolo S, Lysiuk R. 2017.** Flavonoids as detoxifying and pro-survival agents: what’s new? *Food and Chemical Toxicology* **110**:240–250.
- Braunwald E. 2012.** The treatment of acute myocardial infarction: the past, the present, and the

403 future. *European Heart Journal of Acute Cardiovascular Care* **1**(1):9–12.

404 **Carden DL, Granger DN. 2000.** Pathophysiology of ischaemia-reperfusion injury. *Journal of*
 405 *Pathology* **190**: 255–266.

406 **Chong SJ, Low IC, Pervaiz S. 2014.** Mitochondrial ROS and involvement of Bcl-2 as a
 407 mitochondrial ROS regulator. *Mitochondrion* **19**:39–48.

408 **D'Amelio M, Sheng M, Cecconi F. 2012.** Caspase-3 in the central nervous system: beyond
 409 apoptosis. *Trends in Neurosciences* **35**(11):700–709.

410 **Dianzani C, Lepore A, Gentile F, Barrera G. 2013.** Interaction of aldehydes derived from lipid
 411 peroxidation and membrane proteins. *Frontiers in Physiology* **4**:242.

412 **Du Y, Plante E, Janicki JS, Brower GL. 2010.** Temporal evaluation of cardiac myocyte
 413 hypertrophy and hyperplasia in male rats secondary to chronic volume overload. *American*
 414 *Journal of Pathology* **177**(3):1155–1163.

415 **Garciarena CD, Fantinelli JC, Caldiz CI, Chiappe de Cingolani G, Ennis IL, Perez NG,**
 416 **Cingolani HE, Mosca SM. 2011.** Myocardial reperfusion injury: reactive oxygen species vs.
 417 NHE-1 reactivation. *Cellular Physiology and Biochemistry* **27**(1):13–22.

418 **Gatica D, Chiong M, Lavandero S, Klionsky DJ. 2015.** Molecular mechanisms of autophagy
 419 in the cardiovascular system. *Circulation Research* **116**:456–467.

420 **Giordano FJ. 2005.** Oxygen, oxidative stress, hypoxia, and heart failure. *Journal Clinical*
 421 *Investigation* **115**(3):500–508.

422 **Gottlieb RA, Burleson KO, Kloner RA, Babior BM, Engler RL. 1994.** Reperfusion injury
 423 induces apoptosis in rabbit cardiomyocytes. *Journal of Clinical Investigation* **94**(4):1621–1628.

424 **Guo H, Zhang X, Cui Y, Zhou H, Xu D, Shan T, Zhang F, Guo Y, Chen Y, Wu D. 2015.**
 425 Taxifolin protects against cardiac hypertrophy and fibrosis during biomechanical stress of
 426 pressure overload. *Toxicology and Applied Pharmacology* **287**(2):168–177.

427 **Halestrap AP, Richardson AP. 2015.** The mitochondrial permeability transition: a current
 428 perspective on its identity and role in ischaemia/reperfusion injury. *Journal of Molecular and*
 429 *Cellular Cardiology* **78**:129–141.

430 **He F, Li J, Liu Z, Chuang CC, Yang W, Zuo L. 2016.** Redox mechanism of reactive oxygen
 431 species in exercise. *Frontiers in Physiology* **7**:486.

432 **Herr DJ, Aune SE, Menick DR. 2015.** Induction and Assessment of Ischemia-reperfusion
 433 Injury in Langendorff-perfused Rat Hearts. *Journal of Visualized Experiments* **101**:e52908.

434 **Houshmand F, Faghihi M, Zahediasl, S. 2009.** Biphasic protective effect of oxytocin on
 435 cardiac ischemia/reperfusion injury in anaesthetized rats. *Peptides* **30**(12):2301–2308.

436 **Kleikers PW, Wingler K, Hermans JJ, Diebold I, Altenhöfer S, Radermacher KA, Janssen**
 437 **B, Görlach A, Schmidt HH. 2012.** NADPH oxidases as a source of oxidative stress and
 438 molecular target in ischemia/reperfusion injury. *Journal of Molecular Medicine (Berlin)*
 439 **90**(12):1391–1406.

440 **Konstantinidis K, Whelan RS, Kitsis RN. 2012.** Mechanisms of cell death in heart disease.
 441 *Arteriosclerosis Thrombosis Vascular Biology* **32**(7):1552–1562.

442 **Lundberg KC, Szweda LI. 2004.** Initiation of mitochondrialmediated apoptosis during cardiac
443 reperfusion. *Archives of Biochemistry and Biophysics* **432(1)**:50–57.

444 **Maciejczyk M, Mikoluc B, Pietrucha B, Heropolitanska-Pliszka E, Pac M, Motkowski R,**
445 **Car H. 2017.** Oxidative stress, mitochondrial abnormalities and antioxidant defense in Ataxia-
446 telangiectasia, Bloom syndrome and Nijmegen breakage syndrome. *Redox Biology* **11**:375–383.

447 **Matsushima S, Kuroda J, Ago T, Zhai P, Ikeda Y, Oka S, Fong GH, Tian R, Sadoshima J.**
448 **2013.** Broad suppression of NADPH oxidase activity exacerbates ischemia/reperfusion injury
449 through inadvertent downregulation of HIF-1 α and upregulation of PPAR- α . *Circulation*
450 *Research* **112(8)**:1135–1149.

451 **Mehdizadeh R, Parizadeh MR, Khooei, AR, Mehri S, Hosseinzadeh H. 2013.**
452 Cardioprotective effect of saffron extract and safranal in isoproterenol-induced myocardial
453 infarction in wistar rats. *Iranian Journal of Basic Medical Sciences* **16(1)**:56–63.

454 **Moghadam-Kia S, Oddis CV, Aggarwal R. 2016.** Approach to asymptomatic creatine kinase
455 elevation. *Cleveland Clinic Journal of Medicine* **83(1)**:37–42.

456 **Moon YJ, Wang X, Morris ME. 2006.** Dietary flavonoids: Effects on xenobiotic and
457 carcinogen metabolism. *Toxicology in Vitro* **20(2)**:187–210.

458 **Naito H, Furukawa Y, Chino D, Yamada C, Hashimoto K. 2000.** Effects of zatebradine and
459 propranolol on canine ischemia and reperfusion-induced arrhythmias. *European Journal of*
460 *Pharmacology* **388(2)**:171–176.

461 **Pangonyte D, Stalioraityte E, Ziuraitiene R, Kazlauskaite D, Palubinskiene J, Balnyte I.**
462 **2008.** Cardiomyocyte remodeling in ischemic heart disease. *Medicina (Kaunas)* **44(11)**:848–854.

463 **Pizzimenti S, Ciamporcerio E, Daga M, Pettazzoni P, Arcaro A, Cetrangolo G, Minelli R,**
464 **Powers SK, Murlasits Z, Wu M, Kavazis AN. 2007.** Ischemia-reperfusion-induced
465 cardiainjury: a brief review. *Medicine and Science in Sports and Exercise* **39(9)**:1529–1536.

466 **Raj Narayana K, Sripal Reddy M, Chaluvadi MR, Krishna DR. 2001.** Bioflavonoids
467 classification, pharmacological, biochemical effects and therapeutic potential. *Indian Journal of*
468 *Pharmacology* **33**:2–16

469 **Shanmugam K, Ravindran S, Kurian GA, Rajesh M. 2018.** Fisetin confers cardioprotection
470 against myocardial ischemia reperfusion injury by suppressing mitochondrial oxidative stress
471 and mitochondrial dysfunction and inhibiting glycogen synthase kinase 3 β activity. *Oxidative*
472 *Medicine and Cellular Longevity*

473 **Sun X, Chen RC, Yang ZH, Sun GB, Wang M, Ma XJ, Yang LJ, Sun XB. 2014.** Taxifolin
474 prevents diabetic cardiomyopathy in vivo and in vitro by inhibition of oxidative stress and cell
475 apoptosis. *Food and Chemical Toxicology* **63**:221–232.

476 **Wang YY, Yuan Y, Wang XC, Wang YF, Cheng J, Tian L, Guo XH, Qin DM, Cao WJ.**
477 **2017.** Tiliarin post-conditioning attenuates myocardial ischemia/reperfusion injury via
478 mitochondrial protection and inhibition of apoptosis. *Medical Science Monitor* **18(23)**:4490–4499.

479 **Whelan RS, Kaplinskiy V, Kitsis RN. 2010.** Cell death in the pathogenesis of heart disease:
480 mechanisms and significance. *Annual Review of Physiology* **72**:19–44.

481 **Xie X, Feng J, Kang Z, Zhang S, Zhang L, Zhang Y, Li X, Tang Y. 2017.** Taxifolin protects
482 RPE cells against oxidative stress-induced apoptosis. *Molecular Vision* **23**:520–528.


483 **Xuan FF, Jian J. 2016.** Epigallocatechin gallate exerts protective effects against myocardial
484 ischemia/reperfusion injury through the PI3K/Akt pathway-mediated inhibition of apoptosis and
485 the restoration of the autophagic flux. *International Journal of Molecular Medicine* **38(1)**:328-
486 336.

487 **Zhao H, Sapolsky RM, Steinberg G. 2006.** Interrupting reperfusion as a stroke therapy:
488 ischemic postconditioning reduces infarct size after focal ischemia in rats. *Journal of Cerebral*
489 *Blood Flow and Metabolism* **26(9)**:1114-1121.

Figure 1

Figure 1 Experimental protocol

Figure 1. Schematic diagram of experimental protocol. Normal control group (Control); Myocardial I/R control group (I/R); I/R + TAX treatment group (TAX 5 μ M); I/R + TAX treatment group (TAX 15 μ M).



★ Taxflion and K-H solution for 60 min after 30 min of whole heart ischemia

Figure 2

Figure 2. TAX treatment improved the cardiac function recovery of rats during myocardial I/R injury in vitro model.

(A) Heart rate (HR, beat/per min); (B) left ventricular developed pressure (LVDP, mmHg); (C) maximum rate of left ventricular pressure ($+dp/dt_{max}$, mmHg/s); (D) minimum rate of increase of left ventricular pressure ($-dp/dt_{min}$, mmHg/s); (E) Rate pressure product (RPP, mmHg \times bpm); (F) representative left ventricular pressure records in experimental protocol form different experiment groups.




Figure 3

Figure 3. Effect of TAX on injury of cardiomyocytes by measurement of LDH and CK-MB.

The levels of perfusate LDH and CK at different time points in the control, I/R and TAX-treat group (5 μ M and 15 μ M) are shown. # $P < 0.05$ and ## $P < 0.01$ compared with the control group; * $P < 0.05$ and ** $P < 0.01$ compared with the I/R group; U/L: international enzyme activity unit per liter.




Figure 4

Figure 4. Representative micrographs of HE staining results in various experimental groups.

(A) control group; (B) I/R group; (C) TAX 5 μ M group; (D) TAX 15 μ M group, n=3 per group, (magnification, $\times 400$), (\rightarrow) tissue damage and edema.




Figure 5

Figure. 5. Effect of TAX on cardiac the activity of SOD and GSH-PX, contents of MDA.

Values are presented as mean \pm SD. # $P < 0.05$ and ## $P < 0.01$ compared with the control group; * $P < 0.05$ and ** $P < 0.01$ compared with the I/R group. U/mgprot: international enzyme activity unit per milligram tissue protein.




Figure 6

Figure 6. Effect of TAX on the expression of Cyt-c, Bax and Bcl-2 protein.

$P < 0.05$, ## $P < 0.01$ vs. Control group; * $P < 0.05$, ** $P < 0.01$ vs. IR group. ng/mgprot indicate the nanogram level of the target protein per milligram total protein.




Figure 7

Fig. 7. The changes in the levels of caspase 3, and caspase 9 at the end of reperfusion.




Figure 8

Figure 8. Effect of taxifolin on cardiomyocytes apoptosis in different experimental groups.

(A) control group; (B) I/R group; (C) I/R + TAX 5 μ M group; (D) I/R + TAX 15 μ M group. n=3 per group, 20X, Scale bar 50 μ m, (\rightarrow) positive stain

