

Quantitative heterodonty in Crocodyliformes: assessing size and shape across modern and extinct taxa (#29741)

1

First revision

Editor guidance

Please submit by **30 Nov 2018** for the benefit of the authors (and your \$200 publishing discount).



Structure and Criteria

Please read the 'Structure and Criteria' page for general guidance.



Author notes

Have you read the author notes on the [guidance page](#)?



Raw data check

Review the raw data. Download from the location [described by the author](#).



Image check

Check that figures and images have not been inappropriately manipulated.

Privacy reminder: If uploading an annotated PDF, remove identifiable information to remain anonymous.

Files

Download and review all files from the [materials page](#).

- 1 Tracked changes manuscript(s)
- 1 Rebuttal letter(s)
- 7 Figure file(s)
- 1 Table file(s)
- 1 Other file(s)



Structure your review

The review form is divided into 5 sections. Please consider these when composing your review:

1. BASIC REPORTING
2. EXPERIMENTAL DESIGN
3. VALIDITY OF THE FINDINGS
4. General comments
5. Confidential notes to the editor

 You can also annotate this PDF and upload it as part of your review

When ready [submit online](#).

Editorial Criteria

Use these criteria points to structure your review. The full detailed editorial criteria is on your [guidance page](#).

BASIC REPORTING

-  Clear, unambiguous, professional English language used throughout.
-  Intro & background to show context. Literature well referenced & relevant.
-  Structure conforms to [PeerJ standards](#), discipline norm, or improved for clarity.
-  Figures are relevant, high quality, well labelled & described.
-  Raw data supplied (see [PeerJ policy](#)).

EXPERIMENTAL DESIGN

-  Original primary research within [Scope of the journal](#).
-  Research question well defined, relevant & meaningful. It is stated how the research fills an identified knowledge gap.
-  Rigorous investigation performed to a high technical & ethical standard.
-  Methods described with sufficient detail & information to replicate.

VALIDITY OF THE FINDINGS

-  Impact and novelty not assessed. Negative/inconclusive results accepted. *Meaningful* replication encouraged where rationale & benefit to literature is clearly stated.
-  Data is robust, statistically sound, & controlled.
-  Speculation is welcome, but should be identified as such.
-  Conclusions are well stated, linked to original research question & limited to supporting results.

Standout reviewing tips

3



The best reviewers use these techniques

Tip

Support criticisms with evidence from the text or from other sources

Example

Smith et al (J of Methodology, 2005, V3, pp 123) have shown that the analysis you use in Lines 241-250 is not the most appropriate for this situation. Please explain why you used this method.

Give specific suggestions on how to improve the manuscript

Your introduction needs more detail. I suggest that you improve the description at lines 57- 86 to provide more justification for your study (specifically, you should expand upon the knowledge gap being filled).

Comment on language and grammar issues

The English language should be improved to ensure that an international audience can clearly understand your text. Some examples where the language could be improved include lines 23, 77, 121, 128 – the current phrasing makes comprehension difficult.

Organize by importance of the issues, and number your points

1. Your most important issue
2. The next most important item
3. ...
4. The least important points

Please provide constructive criticism, and avoid personal opinions

I thank you for providing the raw data, however your supplemental files need more descriptive metadata identifiers to be useful to future readers. Although your results are compelling, the data analysis should be improved in the following ways: AA, BB, CC

Comment on strengths (as well as weaknesses) of the manuscript

I commend the authors for their extensive data set, compiled over many years of detailed fieldwork. In addition, the manuscript is clearly written in professional, unambiguous language. If there is a weakness, it is in the statistical analysis (as I have noted above) which should be improved upon before Acceptance.

Quantitative heterodonty in **Crocodyliformes**: assessing size and shape across modern and extinct taxa

Domenic C D'Amore ^{Corresp., 1}, Megan Harmon ¹, Stephanie K Drumheller ², Jason Testin ³

¹ Department of Natural Sciences, Daemen College, Amherst, NY, United States

² Department of Earth and Planetary Sciences, University of Tennessee - Knoxville, Knoxville, Tennessee, United States

³ Iowa Western Community College, Council Bluffs, IA, United States

Corresponding Author: Domenic C D'Amore

Email address: ddamore@daemen.edu

Heterodonty in Crocodylia and associated taxa has not been defined quantitatively, as the teeth rarely have been measured. This has resulted in a range of qualitative descriptors, with little consensus on the condition of dental morphology in the clade. The purpose of this study is to present a method for the quantification of both size- and shape-heterodonty in members of Crocodyliformes. Data were collected from dry skeletal and fossil specimens of **34 crown crocodylians and one crocodyliform**. Digital photographs were taken of each tooth and the skull, and the margins of both were converted into both landmarks and semilandmarks. We expressed heterodonty through Foote's morphological disparity, and a Principal Components Analysis visualized shape variance. This analysis reveals that all **Crocodyliformes** sampled are heterodont to varying degrees. The majority of the shape variance was represented by a 'caniniform' to 'molariform' transition. Heterodonty varied significantly between positions; size undulated whereas shape was significantly linear from mesial to distal. Size and shape, although significantly correlated, appear to be primarily decoupled. Skull shape correlated significantly with tooth shape. High size-heterodonty often correlated with very large caniniform teeth, reflecting a prioritization of securing prey. Large, highly molariform, distal teeth may be a consequence of high-frequency durophagy combined with prey size. The slender-snouted skull shape correlated with a caniniform arcade with low heterodonty. This was reminiscent of other underwater-feeding tetrapods, as they focus on small prey that requires minimal processing. Several extinct taxa were very molariform, which was associated with low heterodonty. **Hamadasuchus rebouli** may have dealt with vertebrate prey similar to large modern taxa, but prey processing may have been different due to its terrestrial habitat. Disparity measures can be inflated or deflated if numerous teeth are absent from the tooth row, and regression analysis may not best apply to strongly slender-snouted taxa. Nevertheless, when these methods are used in tandem they can give a

complete picture of crocodylian heterodonty. Future researchers may apply our proposed method to most crocodyliform specimens with an intact enough tooth row regardless of age, species, or rearing conditions, as this will add rigor to many life history studies of the clade.

TITLE: Quantitative heterodonty in Crocodyliformes: assessing size and shape across modern and extinct taxa.

AUTHORS: Domenic C. D’Amore*¹, Megan Harmon¹, Stephanie K. Drumheller², Jason Testin³

1. Department of Natural Sciences, Daemen College, 4380 Main Street, Amherst, New York 14226 United States of America

2. Department of Earth and Planetary Sciences, University of Tennessee, 1621 Cumberland Avenue, 602 Strong Hall, Knoxville, Tennessee 37996 United States of America

3. Iowa Western Community College, Council Bluffs, Iowa, 51503 United States of America

***Corresponding Author:** Domenic C. D’Amore, ddamore@daemen.edu

13 ABSTRACT

14 Heterodonty in Crocodylia and associated taxa has not been defined quantitatively, as the teeth
 15 rarely have been measured. This has resulted in a range of qualitative descriptors, with little
 16 consensus on the condition of dental morphology in the clade. The purpose of this study is to
 17 present a method for the quantification of both size- and shape-heterodonty in members of
 18 Crocodyliformes. Data were collected from dry skeletal and fossil specimens of 34 crown
 19 crocodylians and one crocodyliform. Digital photographs were taken of each tooth and the
 20 skull, and the margins of both were converted into both landmarks and semilandmarks. We
 21 expressed heterodonty through Foote's morphological disparity, and a Principal Components
 22 Analysis visualized shape variance. This analysis reveals that all Crocodyliformes sampled are
 23 heterodont to varying degrees. The majority of the shape variance was represented by a
 24 'caniniform' to 'molariform' transition. Heterodonty varied significantly between positions; size
 25 undulated whereas shape was significantly linear from mesial to distal. Size and shape
 26 appeared to be primarily decoupled. Skull shape correlated significantly with tooth shape. High
 27 size-heterodonty often correlated with very large caniniform teeth, reflecting a prioritization of
 28 securing prey. Large, highly molariform, distal teeth may be a consequence of high-frequency
 29 durophagy combined with prey size. The slender-snouted skull shape correlated with a
 30 caniniform arcade with low heterodonty. This was reminiscent of other underwater-feeding
 31 tetrapods, as they focus on small prey that requires minimal processing. Several extinct taxa
 32 were very molariform, which was associated with low heterodonty. *Hamadasuchus rebouli* may
 33 have dealt with vertebrate prey similar to large modern taxa, but prey processing may have
 34 been different due to its terrestrial habitat. Disparity measures can be inflated or deflated if

numerous teeth are absent from the tooth row, and regression analysis may not best apply to strongly slender-snouted taxa. Nevertheless, when these methods are used in tandem they can give a complete picture of crocodylian heterodonty. Future researchers may apply our proposed method to most crocodyliform specimens with an intact enough tooth row regardless of age, species, or rearing conditions, as this will add rigor to many life history studies of the clade.

INTRODUCTION

What constitutes heterodonty often seems like a moving target, with different qualitative definitions in place depending on the clade being studied (Shimada, 2001). Kieser et al. (1993, p.195) referred to the definition of heterodonty as “a bone of contention,” and since then the issue never has been fully resolved. Arguably, this lack of clarity is most pronounced within members of Crocodylia. Researchers have often referred to crocodylians as homodont (Langston, 1973; Osborn, 1998; Larsson & Sidor, 1999; Zahradnicek et al., 2014). Peyer (1968 p.17) defined the term as lacking the discrete dental categories seen in mammals (incisors, canines, premolars, molars), even though he admitted, “a sharp distinction between homodont and heterodont is not possible.” Ferguson (1981) referred to *Alligator mississippiensis* as “pseudoheterodont,” because it showed a gradual, as opposed to punctuated, change in tooth shape along the tooth row (see also Grigg and Gans, 1993; Hendrickx, Mateus, & Araújo, 2015a). Size variability along the tooth row has motivated the term “heterometric homodonty”

for *Crocodylus niloticus* (Fruchard, 2012). Others have applied anisodonty to the clade, which is apparent size, but not shape, heterodonty (Vullo et al., 2016). Certain fossil crocodylians, often interpreted as herbivores or omnivores, exhibit multi-cusped and/or grinding teeth, and are specifically called “heterodont crocodylians” by researchers (e.g. Martin, 2007; Ósi, Clark, & Weishampel, 2007; Novas et al., 2009). Lastly, some researchers have argued certain modern crocodylians are heterodont, and argue dental categories do in fact exist (Aoki, 1989; Kieser et al., 1993).

Semantics aside, one reason for the lack of resolution concerning crocodylian heterodonty is that their teeth rarely have been measured. Few studies have performed quantitative shape analyses of crocodylian teeth. Of these, linear-distance measures have been used for fossil identification (Frey & Monninger, 2010), replacement rates (Bennett, 2012), and biomechanical analyses (Monfroy, 2017). Aside from a study evaluating two fossil notosuchians (Lecuona & Pol, 2008), and a preliminary geometric morphometric investigation of *Crocodylus niloticus* (Farrugia, Polly, & Njau, 2016), no studies have quantitatively investigated heterodonty either within or between species. Typically, crocodylian dentition is described qualitatively, with the goal of characterization for phylogenetic analysis, or paleoecological inference (e.g. Schwarz-Wings, Rees, & Lindgren, 2009; Young et al., 2012; Salas-Gismondi et al., 2015; Adams, Noto, & Drumheller, 2017). Qualitative descriptors of crocodylian tooth morphology are numerous, and include terms such as “blunt, bulbous, broadened, button-shaped, conical, globular, fang, kidney-shaped, lanceolate, needle-like, procumbent, pseudocanine, robust, short, slender, spike-like,” and “thick” (e.g. Brazaitis, 1973; Groombridge, 1982; Aoki, 1989; Brochu, 1999; Erickson, Lappin, & Vliet, 2003; Ósi, Clark, & Weishampel, 2007; Schwarz-Wings,

78 Rees, & Lindgren, 2009; Fruchard, 2012; Gignac & Erickson, 2014; Salas-Gismondi et al., 2015;
 79 Berkovitz and Shellis, 2017). There is clearly a gap in our knowledge concerning the nature of
 80 dental morphology in this clade, and closing this gap may be crucial for a more complete
 81 understanding of performance, behavior, and trophic ecology within Crocodylia, as well as
 82 more distantly related, crocodylian-line archosaurs.

83 The lack of quantitative studies on heterodonty in crocodylians and associated taxa is
 84 not due to a lack of applicable methodology, as there has been a burst of morphometric
 85 research in non-mammalian teeth in the past decade. Dinosaur teeth have probably received
 86 the most attention, with multiple studies using linear-distance measures for the identification
 87 of loose fossil crowns or to infer functional paleoecology (D'Amore, 2009; Larson & Currie,
 88 2013; Buckley and Currie, 2014; Hendrickx and Mateus, 2014; Torices, Reichel, & Currie, 2014;
 89 Hendrickx, Mateus, & Araújo, 2015b, Gerke and Wings, 2016; Larson, Brown, & Evans, 2016).
 90 Extant reptiles have been investigated quantitatively as well, including colubrid snakes (Britt,
 91 Clark, & Bennett, 2009) and varanid lizards (D'Amore, 2015). Prior to this, lamniform sharks
 92 were studied heavily (Shimada, 2002b, 2004; Shimada and Seigel, 2005). These morphometric
 93 analyses have shed light on the nature of heterodonty, dental allometry, and ecomorphology in
 94 these vertebrates, and similar methods may be applied to Crocodyliformes in the hopes to
 95 elaborate upon the state of heterodonty in this taxon.

96 The purpose of this study is to present a method for the quantification of both size- and
 97 shape-heterodonty in members of Crocodyliformes. Data were collected from a multispecific
 98 sample of both extant and extinct specimens housed in museum collections, and their tooth

morphology was assessed through two-dimensional geometric morphometrics. In addition to this major goal, we also 1) outline and describe dental morphology within the specimens sampled; 2) report any morphological consistencies found within the members of our sample; and 3) present the advantages, limitations, and potential future uses of the method. Our intention is to put forward a method for assessing heterodonty that may be applicable to most crocodyliform specimens.

MATERIALS AND METHODS

Institutional abbreviations

American Museum of Natural History, New York, NY (AMNH); Royal Ontario Museum, Toronto, ON (ROM); University of California Museum of Paleontology, Berkeley, CA (UCMP)

Nomenclature

Crocodyliform teeth have very few discrete homologous anatomical loci, but, because they exhibit thecodont dentition (sensu Edmund, 1962, 1969), we defined them as having a crown with an apex, a neck, and a root within an alveolus. Nomenclature for tooth morphology used here was proposed by Smith & Dodson (2003; Figure 1A-B): mesial, towards the central premaxilla and mandibular symphysis; distal, away from the central premaxilla and mandibular symphysis; lingual, towards the tongue; labial, towards the lips; basal, towards the base of the tooth or alveolus; apical, away from the alveolus or towards the apex. An ‘enlarged’ tooth is the largest tooth in a given section of the jaw independent of shape. This type of tooth is typically referred to as a procumbent tooth (Gignac & Erickson, 2014) or a pseudocanine (Brochu, 1999).

Tooth position was indicated by either the presence of a tooth or an empty alveolus in the host bone (Figure 1A). Teeth were lettered based on the host bone (premaxilla = P, maxilla = M, dentary = D), and numbered in ascending order from mesial to distal positions (the mesial-most dentary tooth was D1, followed by D2, D3, etc.). For consistency, all specimens were assumed to have 5 premaxillary positions (P1–P5) (Berkovitz & Shellis, 2017). Members of *Paleosuchus* and *Osteolaemus* have only 4 premaxillary teeth during early stages of ontogeny (Brochu and Storrs, 2012; Narvaez et al., 2015), and an alveolus may atrophy in certain species (usually P2) as they grow (Webb & Messel, 1978; Brown et al., 2015; DC D’Amore, personal observation). If only four premaxillary positions were present, position P2 was assumed absent, and skipped over during numbering. In one case, a specimen had 6 premaxillary teeth (*Alligator mississippiensis*, ROM 4408). This tooth position was omitted for consistency. Our *Alligator prenasalis* specimen (ROM 1375) had its distal-most cranial positions obscured by poor preservation and matrix. We therefore based its maxillary tooth count on previous osteological accounts (a total of 15 maxillary teeth based on Harvard Museum of Comparative Zoology specimen #1015, Mook, 1932).

Specimens

Data were collected from 27 extant, and 8 extinct, crocodyliform specimens (Supplemental Information Table S1). This resulted in a total of 21 species. Although *Caiman crocodilus* is an extant species, a fossil specimen was also included. From these we measured 1,263 teeth in total. Although we did not use any distinct criteria to distinguish juveniles from adults, larger specimens were selected when possible. Specimens with the most complete tooth rows in the

collections were selected, in that they had the **most of their tooth positions** represented by measurable teeth on at least one side of the mouth. Certain fossil specimens only had cranial (*Alligator prenasalis* ROM 1375, *Borealosuchus sternbergii* UCMP 126099, “*Crocodylus*” *affinis* UCMP 131090, *Hamadasuchus rebouli* ROM 52620, *Leidyosuchus canadensis* ROM 1903) or dentary (*Borealosuchus sternbergii* UCMP 131769) material to sample.

Tooth data collection

Methods were similar to those proposed by D’Amore (2015). We photographed each tooth using either an Olympus Stylus or a Canon Rebel T3 EOS camera with a non-zoom lens. Skulls were positioned on a flat surface with a dark background such as a tabletop or camera stand, and held stationary by an available prop such as a box or sandbag if necessary. A scale was positioned at the same distance from the camera as the tooth. The camera was either mounted on a camera stand, or was held stationary by a researcher (for larger specimens). Digital photographs were taken from the labial perspective (Figure 1B). For each tooth, we positioned the specimen so the camera lens was parallel to the host bone adjacent to the tooth. This resulted in both carinae being visible in the shot (if present). We simultaneously positioned the lens parallel to the apical-basal long axis, determined qualitatively as when the tooth looked its tallest to the photographer regardless of any labio-lingual curvature. Each tooth was photographed separately, and teeth from both sides were photographed if available. Only fully erupted teeth with the neck visible were included (Figure 1B). Tooth quality was variable in extant specimens. Teeth with slightly worn apices were included. As the outline of the tooth margin was our basis of measurement, we omitted any teeth with large wear facets or chips

that largely interrupted this margin. Cracks down the long axis of the teeth were common, and were omitted if the crack distorted the shape of the tooth or resulted in a space where light could be seen from the other side.

We used a sliding semilandmark analysis (Bookstein, 1997; Sheets, Kim, & Mitchell 2004; Zelditch et al., 2004; Mitteroecker et al., 2013) to derive shape measurements from each tooth's outline. Photographs were entered in TpsDig 2.16, and the margin of the tooth was traced using the curve drawing tool (Rohlf, 2010) (Figure 1B). Because the enamel margin was not always clear, each tooth was traced from apex to the point where the tooth ceased to taper on the neck for both the mesial and distal side. TpsDig then transformed each of the two traced margins into 30 equidistant coordinates, and we combined the apical-most coordinates. This resulted in 3 discrete landmarks (two at the base and one at the apex) and 56 semilandmarks (Figure 1C). This number of coordinates has been used in previous studies of both dinosaur (Smith, Vann, & Dodson, 2005) and monitor lizard (D'Amore, 2015) dentition, as well as claw morphology (Tinius and Russel, 2016), in which it has been shown to accurately represent the totality of two-dimensional shape (Tinius and Russel, 2016). We performed a generalized least squares Procrustes (GLSP) superimposition on the data, calculated centroid size (CS), and slid the semilandmarks to minimize the total bending energy (Perez, Bernal, & Gonzalez, 2006; Gunz & Mitteroecker, 2013) using the program TpsRelw 1.53 (Rohlf, 2013).

Skull data collection

The shape of the skull, and particularly the rostrum, has long been considered both an important phylogenetic and ecomorphological feature in crocodylians (Busbey, 1995; Daniel &

McHenry, 2001; Brochu, 2001; Sadleir & Makovicky, 2008; Salas-Gismondi et al., 2016; Drumheller, Wilberg, & Sadlier, 2016; Wilberg, 2017). We attempted to determine if there was a correlation between tooth morphology and head shape, as these traits may be linked. All specimens' skulls were photographed from the dorsal perspective using the same cameras as above (Figure 1E). Each skull was positioned so the palate was parallel with the tabletop, and the camera was positioned with a camera stand and leveled. A scale was included. We derived skull shape data using a modified version of our technique for tooth outlines. Using TpsDig again, we traced the skull margin from the rostral-most point of contact between the premaxillae to the caudal-most quadratojugal along the margin on each side. We chose this margin because it outlined head-shape as close to as it would have appeared in life as possible, but avoided internal structures such as the jaw articulations or occipital condyles. Each margin was broken into 50 equidistant coordinates, and the rostral-most coordinates were combined. This resulted in 3 landmarks (two at the quadratojugals and one at the premaxillary junction) and 97 semilandmarks (Figure 1E). These also underwent a GLSP superimposition and the semilandmarks were slid to minimize the total bending energy using TpsRelw 1.53. In specimens with damaged or missing bones on one side, bilateral symmetry was assumed and the coordinates on the intact side were mirrored.

A body-size metric was needed for several of the following analyses, but unfortunately few were available for all specimens. Commonly used metrics such as snout-vent length and mass were not recorded for most dry skeletons prior to cataloging, and many specimens (especially fossils) lacked femora (see Farlow et al., 2005). Therefore, the length of the skull was used as a body size metric (see Fukuda et al. 2013 for potential influences on this

measurement). We derived skull length from the same landmarks outlining the skull above; it was the linear distance from the rostral-most landmark to the posterior-most landmarks along the mid-sagittal plane (Figure 1D). (Note: *Borealosuchus sternbergii* UCMP 131769 and *Crocodylus niloticus* AMNH 142494 did not have intact skulls, and were therefore omitted from all analyses involving skull data.)

Statistical approaches

All analyses were conducted in MorphoJ v. 106d (Klingenberg 2011), SPSS Version 19.0 [IBM Corp, Armonk, NY], and PAST (Hammer, Harper, & Ryan, 2001). If both left and right teeth were available at a given position, we averaged them. For size, CS values were simply averaged together. For shape, each x-y coordinate of the GLSP superimposed landmarks and semilandmarks was averaged with its counterpart for both teeth. To ensure that the left and right sides were not significantly different, we ran a 10,000 permutations test on the Procrustes distances between left and right teeth at positions that had both. The null hypothesis of bilateral symmetry was confirmed ($p = 0.6785$). If only one tooth was available for a given position, that tooth alone represented said position. We also tested the null hypothesis that there was no statistical difference between tooth rows in Crocodyliiformes, by running a 10,000 permutations test on the Procrustes distance between cranial (premaxilla and maxilla) and dentary teeth. The specimens that only had one tooth row available were excluded from this. No significant difference was found ($p = 0.2455$).

A singular measure of heterodonty was derived for each specimen in the form of Foote's morphological disparity [$MD = (\sum_{i=1}^m D_i^2)/(m - 1)$] (Foote, 1993; Zelditch, Sheets, &

Fink, 2003; Sheets & Zelditch, 2013). Disparity (MD) was the sum of the differences of the values of a given tooth (i) from the mean for all teeth from that specimen (Di , also known as the grand mean) squared, with the number of tooth positions (m) factored in. We calculated disparity for all tooth positions for each specimen. For size-heterodonty, Di was simply the difference in CS of a tooth from the mean of the specimen (Zelditch et al., 2004). For shape-heterodonty Di was the Procrustes distance between the tooth and the mean, and was calculated using DisparityBox7 (Sheets, 2012). Heterodonty then was regressed with a reduced major axis against head length to determine if there was a significant allometric change in the clade.

We attempted to correlate skull shape to tooth shape between individuals by using a two-block partial least squares (PLS) analysis in MorphoJ. Skull shape represented one block, and average tooth shape represented the other. Average tooth shape was constructed by averaging the corresponding GLSP superimposed landmarks and semilandmarks of every tooth from an individual. The scores for the first PLS of each shape block were plotted against one another and regressed with a reduced major axis. Visualization of variation along each PLS axis was depicted through vector diagrams (Figure 1F).

To determine if size and shape were coupled in Crocodyliformes, shape coefficients generated by MorphoJ were regressed against CS using a reduced major axes. Significance and a high goodness of fit would be indicative of strong coupling between size and shape. A Principal Components Analysis (PCA) was then conducted to visualize the degree of shape

variance within all cranial and dentary teeth. We only considered any PC representing over 5% of the shape variance, which we then visualized using vector diagrams (Figure 1D).

We plotted shape and size against tooth position to visualize heterodonty along the tooth row. For size CS was normalized by dividing it by head length, and then was plotted against tooth position in a series of box plots. For shape, PC scores of PCs over 5% were plotted against tooth position in a similar fashion. Each box represented a position. Note that we did not do any adjustments to these positions, so specimens with more tooth positions will be the only occupants of the distal-most categories. To determine if significant differences in these size and shape metrics existed between these positions, we used separate analyses of variance (ANOVA) for each superfamily. Both metrics had unequal variances according to Levene's test, so we specifically ran Welch's ANOVAs in SPSS. Any positions represented by less than 7 teeth were excluded from the ANOVAs.

Preliminary quantitative work has suggested a linear transition in tooth shape along the arcade (Farrugia, Polly, & Njau, 2016). We test this by regressing shape data against tooth position using ordinary least squares regressions for each individual. To standardize these regressions, we normalized tooth position into a percentage. We numbered the positions along the tooth row starting with 1 at the mesial-most position, divided each by the total number of positions along the arcade, and then subtracted 0.5 (this subtraction placed the y-intercept halfway along the arcade). PCs for each tooth were then regressed against this, and regression statistics were collected. Several factors may be implied by a significantly linear crocodyliform tooth row. Slope may be linked to heterodonty, as a steeper slope would imply more shape

change along the PC1 scores at y-axis and, consequently, greater shape-heterodonty. The y-intercept would represent shape value for the median position, as the intercept is located half-way along the tooth row. To visualize these coefficients, we plotted both slope and y-intercept for each regression in scatterplots for both the cranium and the dentary.

Results:

Shape variability in the sample

There was statistically significant coupling between size and shape in Crocodyliformes, but to a very weak degree overall. When shape coefficients were plotted against CS, they formed a regression with a goodness of fit accounting for less than 10% of the variance ($y=0.313x-0.981$; $r^2=0.09$; $p<0.0001$; 95% =0.296,0.328).

Most of the shape variance in Crocodyliformes was along a single axis. The only PC accounting for over 5% of the shape variance was the first (PC1). It accounted for over 92.11% of the variance, and is the only PC considered further [see supplemental information for Eigenvalues and variances (Table S2), as well as a discussion on PC2 and PC3 (Figure S1)]. ~~PC1 scores ranged from -0.3173 to 0.3715 within our sample.~~ The negative-most condition involved apical-basal elongation, narrowing at the base, and a gentle concavity on the distal margin (Figure 2). For simplicity, we will refer to this extreme as ‘caniniform’ (Erickson et al., 2012; 2014; Gignac & Erickson, 2014). The positive-most values depicted an apical-basal shortening and mesial-distal broadening, and we will refer to the extreme as ‘molariform’ (Figure 2) (Erickson et al., 2012; 2014; Gignac & Erickson, 2014).

Skull vs. tooth morphology

Skull and average tooth shape were significantly correlated. For the two-block test, PLS1 encompassed 99.96% of shape covariance. Shape variability within the skull shape block showed the snout transitioning from narrow to broad (Figure 3). Taxa that occur below a PLS1 score of -0.15 are the slender-snouted taxa as defined by Brochu (2001), including *Gavialis*, *Mecistops*, and *Tomistoma*. The remainder of the species, defined as either generalized or blunt-snouted (also by Brochu, 2001), occurred around the mean and positive half mixed together. Shape variability within the tooth shape block was similar to the above PCA of tooth shape, displaying a transition from caniniform to molariform with increasing values. These blocks regressed significantly against one another, with slender-snouted taxa separating out with the most caniniform teeth. Scatter increased around the means, indicating the correlation was not as strong among the generalized-to-blunt snouted taxa. *Alligator prenasalis*, *Brachychampsia* sp., and “*Crocodylus*” *affinis* were all relatively blunt-snouted, but rose noticeable above the regression. This indicated they possessed much more molariform teeth on average than their counterparts of similar skull shape.

Foote’s disparity and heterodonty

Size heterodonty was significantly correlated with head length, with an r^2 of ~0.76. The largest individuals according to head length (members of *Crocodylus niloticus* and *Crocodylus porosus*) possessing the greatest unadjusted size heterodonty (Figure 4A). The slope indicated that heterodonty increased at over double the rate of size within the clade. Alligatoroids occur on both sides of the regression. Members of *Alligator* had negative residual size heterodonty, with

Alligator prenasalis as the lowest. On the other side of the regression, caimanine (*Caiman*, *Paleosuchus*) residuals were all positive with the exception of one individual. Several members of *Crocodylus* had values around zero, but one *Crocodylus porosus* specimen had a high residual and a *Crocodylus palustris* had a low residual. One *Osteolaemus tetraspis* individual had the highest size-heterodonty residual, with the other around zero. *Hamadasuchus rebouli* had positive residuals, similar to the greater *Crocodylus porosus* and the caimanines. The slender-snouted taxa (*Gavialis*, *Mecistops*, *Tomistoma*) had some of the most negative residuals.

Shape heterodonty was highly variable, and correlated significantly, albeit rather poorly ($r^2 \sim 0.16$), with head length (Figure 4B). Because of this poor correlation, we do not consider residuals as very biologically meaningful. Similar to size heterodonty, the slender-snouted taxa had some of the lowest shape heterodonty, although *Tomistoma* was relatively greater than the others. In addition, members of *Brachychampsia* sp. and *Alligator prenasalis* also had some of the lowest shape heterodonty in our sample. *Crocodylus siamensis* specimens were more shape heterodont than their congeners, with one individual being the most shape heterodont in our sample. Several caimanine individuals, and both *Osteolaemus tetraspis* specimens, also had relatively high shape heterodonty.

Heterodonty along the tooth row

Tooth position count varied between species (see Supplemental Information Table S3). Most alligatoroids had between 19-20 positions on the cranial tooth row. Many had a similar number on the dentary, except members of *Paleosuchus* had 22 positions. *Leidyosuchus canadensis* had the most (23) cranial positions of the alligatoroids. *Hamadasuchus* (20 cranial) and

328 *Borealosuchus* (23 cranial and 20 dentary) specimens fit within ranges of alligatoroids.
 329 Members of *Crocodylus* and *Mecistops* had between 18-19 cranial and 15 dentary positions.
 330 *Osteolaemus* specimens had the least tooth positions for any crocodyloid (17 cranial and 14
 331 dentary), and *Tomistoma* had the most (21 cranial and 19 dentary). The *Gavialis* specimen had
 332 more positions than any other species sampled (28 cranial and 26 dentary). These tooth counts
 333 are similar to previous published accounts (Brown et al., 2015; Berkovitz & Shellis, 2017). The
 334 vast majority of the modern taxa tooth rows had over 60% their tooth positions represented by
 335 measureable teeth (Supplemental Information Table S3). Fossil taxa ranged from having ~30%
 336 to ~78% of their tooth row represented.

337 In both extant members of Alligatoroidea and Crocodyloidea, size varied significantly
 338 between positions (Figure 5A-B). Size undulated three times along the dental arcade resulting in
 339 significant differences between positions for both the cranium and mandible. Each undulation
 340 peaked with an enlarged tooth. These were typically represented by P4 for both clades, and M4
 341 for alligatoroid and M5 for crocodyloid specimens (sensu Brochu and Storrs, 2012). In addition,
 342 members of *Paleosuchus* had very large P3 and M3. "*Crocodylus*" *affinis* also had a large P3.
 343 Unlike other alligatoroid specimens, the *Leidyosuchus* specimen had both M4 and M5 enlarged,
 344 and the *Brachychampsa* sp. had an enlarged M5 like crocodyloids (sensu Norell, Clark, &
 345 Hutchison, 1994). A final undulation resulted in an enlarged tooth at M9-11 (Figure 5A-B).
 346 Interspersed between these were smaller teeth, with the distal-most tooth often the smallest.
 347 The dentary was similar to the cranium, with three undulations in size. Enlarged teeth were
 348 found at positions D1 and D4, with a third size-peak between D11 and D14. Note that the
 349 position of the enlarged teeth along the cranial tooth row tended to align with smaller teeth

along the dentary tooth row, and vice versa. This resulted in an ‘interlocking’ pattern between the size peaks of one arcade and the valleys of the other. The gavialoid specimen differed markedly by having the two mesial-most teeth enlarged, and the remainder showed a gradual decrease in size distally (Figure 5C). *Hamadasuchus rebouli* had some of the largest teeth for its head length with a dramatic variation in size.

Alligatoroids and crocodyloids both showed a similar trend concerning tooth shape. Mesial teeth are typically the most caniniform in the mouth and distal teeth the most molariform, resulting in positions being significantly different (Figure 5D-E). In cranial teeth P1-M4 tend to exist primarily between PC scores and -0.25 and 0.00, followed by a gradual increase in score values as positions become more distal. Dentary teeth represented a more uniformly gradual caniniform-to-molariform transition. Both superfamilies were highly variable. Alligatoroid had teeth generally more molariform, with upper outliers almost entirely represented by *Alligator prenasalis*, *Alligator sinensis*, and *Brachychampsia* sp. (Figure 5D). Crocodyloids were generally more caniniform, with mesial upper outliers represented by “*Crocodylus*” *affinis* and lower outliers represented primarily by *Tomistoma* (Figure 5E). Both *Borealosuchus* tooth rows had PC1 scores between -0.22 and 0.20, and *Hamadasuchus* ranged between -0.19 and 0.23 (Figure 5F). Both taxa showed a similar progression from caniniformy to molariformy as the alligatoroids and crocodyloids. *Gavialis* deviated from the others the most, where most teeth had scores of < -0.20 with a steep increase towards the average in the distal-most fifth of the arcade (Figure 5F).

Shape heterodonty was found to be strongly linear. When each modern individual's PC1 values were regressed against position, all regressions were significant (Supplemental Information Table S3). The vast majority of tooth row regressions had r^2 values above 80%. *Gavialis* had the lowest r^2 values (cranium=0.495, mandible=0.616), followed by the crania of the fossil *Caiman crocodilus* ($r^2=0.728$) and *Tomistoma* ($r^2=0.747$). Both cranial and dentary tooth rows typically had slopes between 0.25-0.55 (Figure 6). More shape heterodont taxa typically had greater slopes, with *Crocodylus siamensis* specimens having some of the steepest slopes (0.39-0.68). All the slender-snouted specimens had the y-intercepts between -0.25 and -0.14, indicating strong caniniformy at the median position. The *Gavialis* specimen also had very shallow slopes (<0.25). Living members of *Alligator* had y-intercepts between 0.029 and 0.085, indicating molariformy at the median (Figure 6). *Alligator prenasalis*, *Brachychampsa* sp., and "*Crocodylus*" *affinis* had the shallowest slopes (0.22-0.32) and the greatest y-intercepts (0.15-0.24) in our sample, indicating molariform teeth consistent along the tooth row. *Hamadasuchus rebouli*'s regression characteristics are similar to members of *Caiman* and *Osteolaemus*. The slope of the fossil *Caiman crocodilus* differed from modern members of *Caiman* by being much shallower (Figure 6).

DISCUSSION

Defining heterodonty within Crocodyliformes

The methods proposed here offer a multi-faceted approach to quantifying heterodonty in Crocodyliformes. Foote's morphological disparity allows for a discrete measure of heterodonty that may be used for comparison, or normalized with other variables (as was done here with a

body size metric). This is not limited to Crocodyliformes, and may be used to compare distantly related taxa and/or a wide variety of dental morphotypes. This measure is ideal if one is interested in *how much* heterodonty is apparent. Alternatively, if one is interested in what characteristics make up tooth heterodonty, more traditional geometric morphometrics suffice in describing shape variability. It is opportune that only PC1 accounted for over 90% of the variance, allowing for us to use it as the sole measure of shape here. If more PCs accounted for over 5% of the shape variance, it would be appropriate to depict those other PCs in the same manner as PC1 for a comprehensive understanding of shape.

Regression analysis of shape is appropriate based on significance and high r^2 values, and the coefficients associated with it are useful characters for comparison. Certain caveats should be considered though. Regression may be more appropriate for the dentary than the cranium, as the premaxillary teeth do not appear to differ from one another as much as the maxillary. In most crocodylians this effect is mild, and r^2 values are still high. The effect is very pronounced in *Gavialis* though, as both the premaxillary and the mesial maxillary teeth are similar in shape. It is also interesting that this happens in the dentary as well. Future studies should consider this when applying this method to Gavialoidea or dentally analogous taxa.

Although the task of assigning a singular dental morphotype to any one species of crocodyliform is beyond the scope of the study, our data suggests that it would be potentially difficult. Heterodonty seems to vary within species, making the assignment of a singular heterodonty measure to an entire species dubious. As far as biological explanations for this, tooth form is almost certainly influenced by allometry. Ontogenetic shifts in feeding niche have

been documented in modern crocodylian species (e.g. Groombridge, 1982; Webb, Manolis, & Buckworth, 1982; Pooley & Gans, 1976; Pooley, 1989; Delany, 1990; Santos et al., 1996; Da Silveira and Magnusson, 1999; Subalussy, A. L., Fitzgerald, L. A., & Smith, 2009 Wallace and Leslie, 2008; Borteiro et al., 2009; Hanson et al., 2014), and allometric changes in the feeding apparatus with size are often explained as a structural consequence of this (e.g. Dodson, 1975; Webb and Messel, 1978; Hutton, 1987; Erickson, Lappin, & Vliet, 2003; Verdade, 2000; Wu et al., 2006; Watanabe & Slice, 2014; Gignac and Erickson, 2016; Gignac & O'Brien, 2016).

Concerning teeth, a qualitative increase in overall molariformy was observed in *Alligator mississippiensis*, and functioned to meet the mechanical demands of increased durophagy (Erickson, Lappin, & Vliet, 2003; Gignac & Erickson, 2014). Although our sample size is too low to confidently assess dental ontogeny within each species, we did see a similar general trend in conspecifics of different sizes. In particular, the larger of our two *Crocodylus porosus* had a greater y-intercepts indicating greater molariformy. In addition to allometry, phenotypic changes due to environmental factors may also influence teeth. Skull shape and tooth orientation are irregularly influenced by captive rearing (Erickson et al., 2004; Drumheller, Wilberg, & Sadleir, 2016), and how this may also influence tooth shape has yet to be determined. Many of our specimens had 'no data' concerning their rearing, so we do not know if captivity influenced either tooth or skull morphology.

Morphological trends within Crocodyliformes

All crocodyliform specimens measured here were heterodont to varying degrees, and these data showed significant variability of morphotypes along the dental arcade for all specimens

(Figure 7). Although dentition varied between species, certain consistencies were seen throughout the clade:

1. Similar teeth occurred on both the cranial and dentary dental arcades.
2. As body size increased, size-heterodonty increased reliably with it. Shape-heterodonty shows a much less reliable negative correlation with body size.
3. The vast majority of shape variance from the labial perspective occurred along a single shape axis, representing the transition from caniniform to molariform. Only minor distal curvature was apparent in very caniniform crowns.
4. There was serial homology in tooth shape from-mesial-to-distal along the tooth row, and molariformy increased in this direction. The transition was significantly linear for both dental arcades for all specimens.
5. Size variability consisted of an undulating pattern with three peaks that interlock with the opposing row, with enlarged crowns interspersed within smaller crowns. This corresponded with the festooning pattern seen in the tooth bearing bones, and was less apparent in slender-snouted taxa.

Shape- and size-heterodonty were very loosely coupled in Crocodyliformes as they changed in dramatically different, and primarily independent, fashions along the arcade. Some correlation did occur; the regression's significance was probably the result of the fact that the distal-most crowns were typically both the smallest crowns as well as the most molariform. Nevertheless, the undulating pattern of tooth size did not align with linear shape heterodonty for the vast majority of the tooth row, as indicated by the very low r^2 .

The low degree of coupling begs the question; do developmental agents influence size and shape separately? Although quite a bit of research has looked at how crocodylian teeth grow and replace themselves (Edmund, 1962; Westergaard and Ferguson, 1986, 1987, 1990; LeBlanc et al., 2017), surprisingly little has been done on what developmental influences affect tooth size and shape. Modern crocodylians replace their teeth in waves, or Zahnreihe (Edmund, 1962; Westergaard and Ferguson, 1990; Osborn, 1998), but it is unclear how the nature of these waves relate to the morphological variables investigated here. Keiser et al., (1993) compartmentalized the dentition along the tooth row for *Crocodylus niloticus*, grouping teeth into ‘incisor,’ ‘premolar,’ and ‘molar’ regions. These designations attempted to account for both size and shape heterodonty; each was defined by an enlarged tooth, and each become progressively more molariform. They did not offer a developmental mechanism that differentiates these categories though. Fruchard (2012, p.7) suggested that the only difference between enlarged teeth and their smaller counterparts was that the former was “programmed to be bigger,” suggesting some sort of additional developmental signaling to enlarge teeth. More research is needed on how tooth shape and size are established developmentally in order to truly understand what generates heterodonty.

Adaptive explanations for morphological variability in modern taxa

There is a wide range of tooth morphologies present in modern Crocodylia, and, as teeth are anatomical units used for feeding and aggression, functional inferences may be drawn based on our present understanding of behavior and performance. Bite force in crocodylians is primarily influenced by size (Erickson et al., 2012, 2014), and our data set shows that similarly sized

crocodylians may have very different tooth dimensions. This rules out maximum bite force as the sole limiting factor dictating tooth form. Although we are reluctant to associate specific prey items with specific tooth forms, size and shape will influence how a tooth interacts with food items possessing certain physical properties. We therefore suggest that a biomechanical link should exist between the structural limits imposed by tooth form and the material properties of the substrates with which it interacts.

As with all jawed vertebrates, crocodylian teeth will succumb to different speeds and stresses based on their respective position along the arcade. Caniniform mesial teeth are ideal for the initial acquisition of prey. Pointed apices reduce surface area to puncture compliant foods that deform under pressure, such as muscle, fat, and fibrous connective tissue (Frazetta, 1988). Being farther from the hinge, these teeth move faster during a strike and are more likely to contact prey trying to escape (Busbey, 1989). They will also will endure less force based on their position, and can afford to be relatively elongate and gracile. On the other end, distal teeth need to withstand greater forces due to their close proximity to the hinge (Cleuren, Aerts, & Vree, 1995; Erickson, Lappin, & Vliet, 2003; McHenry et al., 2006; Erickson et al., 2012). This explains why these teeth are typically on the molariform half of the shape spectrum; the larger base-to-height ratio gives them greater relative bending strengths (Van Valkenburgh and Ruff, 1987; Gignac & Erickson, 2014; Monfroy, 2017). Because force is highest in this region, it is ideal for processing food items after they are acquired (Busbey, 1989; Davenport et al., 1990; Cleurens and de Vree, 2000). The reduced height of these teeth also ensures they do not impede jaw closure. This necessity is very apparent in our representative member of *Gavialis*, and provides a functional explanation for the poor linear shape relationship along the tooth

row in this individual. Having all the teeth be highly caniniform except for the distal-most region may be interpreted as an attempt to reduce heterodonty as much as possible (Grigg and Gans, 1993), while ensuring the distal crowns do not impede jaw closure or break when processing food.

Particular attention should be paid to the relative size of the distal-most crowns, as they vary considerably within our sample. Most modern alligatoroids and crocodyloids have a single enlarged tooth followed distally by several smaller teeth. These typically were represented by positive PC1 scores, especially concerning *Alligator mississippiensis* and *Crocodylus siamensis*, and were also some of the smallest teeth in its arcade (Figure 7A). Both members of *Alligator sinensis* differed from this though, in that they had a row of 4-5 relatively large, high-molariform crowns (followed by only one crown reduced in size). Probably the most extreme condition, *Osteolaemus tetraspis* specimens had distal crowns that were exceptionally large; the largest relative crowns at positions M10-12 and D11-13 for modern taxa all belonged to members of this species. These two species also have the lowest number of teeth for modern alligatoroids and crocodyloids respectively, a reduction potentially based on the need to fit these enlarged teeth. Aoki (1989) qualitatively noted these unique conditions, and suggested they facilitated durophagy. All alligatoroids and crocodyloids sampled here have been recorded to consume at least some hard prey items though (e.g. Brazaitis, 1973; McIlhenny, 1976; Taylor, 1979; Groombridge, 1982; Ross and Magnusson, 1989; Santos et al., 1996; Selvaraj, 2012; Nifong & Silliman, 2013), so it is unclear what selection pressure resulted in these particular morphologies. It may be a result of body size. Bite force tests of *Alligator mississippiensis* showed the pressure produced at its enlarged M11 to be adequate to crush its harder prey

items (Erickson, Lappin, & Vliet, 2003; Gignac & Erickson, 2014). If this is the case in most of the large crocodylians, enlarging the distal-most crowns would be unnecessary. *Alligator sinensis* and *Osteolaemus tetraspis*, on the other hand, may need more extreme dentition closer to the hinge; their smaller size would make it more difficult to process foods with similar mechanical properties. Another explanation for this may be the frequency of consuming hard prey. Although both these species have broad diets, studies have shown certain (but not all) populations to consume disproportionately large numbers of shelled mollusks and crustaceans (Cheng-Kuan, 1957; Groombridge, 1982; Ross and Magnusson, 1989; Luiselli, Akani, & Capizzi, 1999; Pauwels et al., 2007).

All taxa measured here also have two sets of enlarged mesial teeth on both arcades. These teeth are well built for puncturing, likely make first contact with prey during jaw closure, and are resilient against struggling prey (Iordansky, 1964). An apparent trade-off to enlarging these teeth is the need to reduce the size of teeth on the opposing tooth row. This character played a large role in size-heterodonty, with different crocodylians undulating their tooth sizes to different degrees. High relative size-heterodonty in caimanine specimens was typically a consequence of the dramatic size difference between the enlarged teeth and the small remaining crowns, (Figure 7B). Their dentary crowns in particular became so large they often grew entirely through the cranial rostrum in adults (as mentioned in Brazaitis, 1973), which suggests securing prey takes priority. The remaining crowns were rather small by comparison, including the distal crowns: the teeth with the greatest mechanical advantage when processing hard prey. This arrangement may be a specialization for hunting more mobile and/or compliant prey (Sampaio et al., 2013), as insects and fish can make up a large portion of the caimanine

diet (Santos et al., 1996). The *Crocodylus porosus* specimens had the largest M5 crowns in our sample, which may also show a prioritization for puncturing and securing soft-bodied prey in a larger context (Figure 7B). This species is notorious for actively hunting large vertebrates such as sharks, cattle, horses, and humans (e.g. Taylor, 1979; Kar & Bustard, 1981; Groombridge, 1982; Doody, 2009; Hanson et al., 2015), and these teeth are ideal for puncturing and securing such prey. Similar to caimanines, this species atrophies position P2 to make room for its enlarged D1 crowns (Brown et al., 2015)

The slender-snouted species possessed generally more caniniform teeth, which may be a consequence of feeding habitat and prey preference. These taxa have a reputation for eating small, aquatic prey with a focus on fish (Peyer, 1968; Webb, Manolis, & Buckworth, 1982), and multiple lines of evidence suggest the feeding apparatus is well suited for this function. The slender shape reduces resistance during both lateral motion and jaw adduction when feeding underwater, and the increased snout length allows for a faster strike (Taylor, 1987; Pooley, 1989; Thorbjarnarson, 1990; McHenry et al., 2006; Pierce et al., 2008). Highly caniniform teeth can quickly puncture fast-moving, compliant prey, and their elongate shape may also lower their mechanical resistance (Figure 7C). The longirostrine condition, defined as a snout that is both slender and elongate (Brochu, 2001), resulted in increased tooth positions; *Tomistoma* had more teeth than any other crocodyloid, and *Gavialis* has the most teeth out of all crocodylians sampled. This cranio-dental morphotype may be prey-size prohibitive though, as larger prey could damage the slender rostrum while struggling. Their elongate mandibular symphysis results in a mechanical disadvantage against the forces produced by shaking and twisting prey (Walmsley et al., 2013). The gracile nature of the dentition means a lower bending

strength, making them more susceptible to breakage from larger and/or harder prey as well. On rare occasions, large individuals have been known to take land-based, vertebrate prey (Thorbjarnarson, 1990; Selvaraj, 2012). This is most likely because the overwhelming size of these crocodylians allows their feeding apparatus to withstand the forces exerted by said prey.

The slender-snouted taxa had some of the lowest size- and shape-heterodonty of modern crocodylians, which is reminiscent of several other clades of aquatic predators. They share certain traits with the anisodont plesiosauiromorphs (Sassoon et al., 2015; Kear et al., 2017). Although these crocodylians are not anisodont in the strict sense (they all have some shape heterodonty), both taxa have elongate mesial crowns transitioning to smaller distal ones. These taxa also reflect similarities with the ‘homodont’ condition apparent in odontocete whales (Rommel, 1990), where all the teeth in the arcade possess a similar, peg-like shape. This condition is believed to be ideal for catching and holding, but not processing, small aquatic prey (MacLeod et al., 2007), as most prey items consumed are under 10% of their body length (MacLeod et al., 2006). A convergent reduction in size- and shape-heterodonty within these independently aquatic groups may indicate a transition from a multi-functional dental arcade to one almost exclusively for prey capture. This is clearly the condition in members of *Gavialis*, as it is almost entirely caniniform along its tooth row and eats primarily fish (Groombridge, 1982, Figure 7C). Members of *Mecistops* and *Tomistoma*, although also primarily caniniform, still displayed the linear shape change typical of other crocodylioids. These species may consume prey that require relatively more processing with their distal crowns, and there are numerous reports of them eating prey such as crustaceans, turtles, and immature primates (Brazaitis, 1973; Groombridge, 1982; Galdikas and Yeager, 1984; Selvaraj, 2012).

Tooth shape may indicate differences in feeding behavior and processing ability, even though overlap exists in prey selection. *Alligator mississippiensis* and *Crocodylus niloticus* both consume a wide variety of prey, including both large and small mammals, crustaceans, fish, water fowl, snakes, turtles, and conspecifics (McIlenny, 1976; Pooley & Gans, 1976; Groombridge, 1982; Delany and Abercrombie, 1986; Hutton, 1987; Shoop & Ruckdeschel, 1990; Rootes & Chabreck, 1993; Elsey, Trosclair III, & Linscombe, 2004; Wallace and Leslie, 2008; Gabrey, 2010). A comparison of controlled feedings of each of these species showed *Alligator mississippiensis* to fracture and consume noticeably more bovine skeletal elements than *Crocodylus niloticus* (Njau & Blumenschine, 2006; Drumheller & Brochu, 2014; 2016). Our *Alligator mississippiensis* specimen was generally more molariform than *Crocodylus niloticus*. These teeth would have greater bending strengths to resist breakage when processing hard material such as bone.

Fossil taxa and the appropriateness of analogues

Certain fossil taxa were reminiscent of modern counterparts. We expected the fossil *Caiman crocodilus* to be similar to its congeners, due to the fact that these specimens are closely related and all consume insects, crustaceans, and fish (Brazaitis, 1973; Groombridge, 1982; Da Silveira and Magnusson, 1999). Any differences in size and shape ranges appear to simply be a consequence of the former's incomplete arcades; no distal maxillary or any dentary crowns were available (see **Limitations** below). *Alligator mississippiensis* specimens have similar shape regression statistics to our *Leidyosuchus* specimen, but, unlike members of *Alligator*, this specimen lacked enlarged distal teeth (Figure 7D). This caused size heterodonty to differ

noticeably, and may be indicative of a difference in the degree these taxa process hard materials (although no taphonomic evidence for this currently exists associated with *Leidyosuchus*). The two specimens of *Borealosuchus* differed from one another in median shape as indicated by y -intercepts, which may due to an allometric increase in molariformity. The best analogue for this species may be a member of *Crocodylus* with similar slopes, but more data is necessary to confirm this (Figure 7D).

The inclusion of *Hamadasuchus*, and the fact that it showed similar trends in heterodonty to crown crocodylians, was particularly revealing. Peirosaurids are believed to be primarily terrestrial crocodyliforms (Tavares et al., 2017), and they most likely did not occupy the semi-aquatic, sit-and-wait predator niche dominated by modern crocodylians (Larsson & Sues, 2007). Nevertheless, the nature of size- and shape-heterodonty is similar between the two groups, indicating the methods proposed here are transferable outside of Crocodylia. The fact that the *Hamadasuchus* specimen had similar relative size heterodonty and relative maximum tooth size to the larger *Crocodylus porosus* specimen indicates that it may have dealt with similar prey from a mechanical standpoint (Figure 7E). Its greatly enlarged mesial teeth would puncture vertebrate tissue with similar effectiveness to those of a large *Crocodylus porosus* (Figure 7B). The *Hamadasuchus* specimen differed in that it had very large distal crowns, which, unlike members of *Osteolaemus*, are laterally flattened and considered ziphodont (Larsson & Sidor, 1999). This suggests potential differences in prey processing. It may have used these for either sheering soft tissue or breaking bone similar to modern mammalian carnassials, as rolling on land is not an effective means of dismemberment (Fish et al., 2007).

Several authors have stated that modern taxa do not have, or have secondarily lost, an extreme degree of molariformy commonly found in extinct representatives. ‘Globidonty’ describes the enlarged, highly molariform crowns in fossil taxa potentially used for durophagy (Norell, Clark, & Hutchison, 1994; Brochu, 1999; 2001; Ősi & Barrett, 2011). Species of *Brachychampsa* are textbook examples of a globidont taxon (Case, 1925; Carpenter and Lindsey, 1980, Figure 7F), and our specimen is the only one in the sample with distal teeth so molariform their PC1 scores exceed 0.349. Although we agree with Brochu (2001, 2004) that *Osteolaemus tetraspis* is not as extreme, the PC1 scores of its enlarged distal teeth are closer to our *Brachychampsa* than another other taxon sampled (0.322-0.341). *Alligator prenasalis* and “*Crocodylus*” *affinis* distal crowns are similar to *Alligator sinensis* in shape, and also create a ridge of robust teeth (Mook, 1932). The mechanical capabilities of these particular crowns in modern taxa should be similar to the extinct, which suggests similar processing abilities in the distal regions of the skull. The similarities break down when the rest of the jaw is considered though. In addition to these highly molariform teeth, modern taxa also possess caniniform mesial teeth suggesting a division of labor along the tooth row. Contrarily, almost all teeth of members of *Alligator prenasalis*, *Brachychampsa* sp., and “*Crocodylus*” *affinis* are on the molariform half of the shape-spectrum (Figure 7F), making both their size- and shape-heterodonty rather low. These extinct taxa probably did not need to do as much puncturing of compliant substrate, which supports the argument that they may have foraged for mollusks and slow moving turtles (Carpenter and Lindsey, 1980; similar to Salas-Gismondi et al., 2015) rather than being ambush predators.

Limitations and future work

Footnote's morphological disparity is a reliable method for assessing heterodonty if the tooth row is near complete, but some of the variability in heterodonty seen here is the result of incompleteness. This measure of disparity relies on, among other things, the grand mean and the sample size. Size heterodonty may be underrepresented if, for example, an enlarged tooth is missing. This tooth would deviate greatly from the grand mean if present, so its exclusion would deflate size heterodonty. As shape is linear in Crocodyliformes, missing the mesial- or distal-most teeth would deflate size heterodonty. This was apparent in the fossil *Caiman crocodilus*; even though it shared almost identical tooth morphology with modern congenics when positions were compared, its shape heterodonty was much lower because the distal 30% of its teeth were missing. This also affected the shape regression, as the fossil specimen's cranium slope was much shallower than the modern members of *Caiman*.

We did not consider all three dimensions here. Living crocodylian teeth are often discussed as conical (Edmund, 1969) or conodont (Hendrickx, Mateus, & Araújo, 2015b). Studies of bending strengths show variation between mesial-distal and labial-lingual axes (Monfrey, 2017), indicating that functional information may be drawn from the dimension not measured here. This is especially important concerning fossil taxa, as pronounced lateral compression is commonplace. *Hamadasuchus* distal teeth have been referred to as zipodont (Larsson & Sidor, 1999), but, as this is defined by lateral flattening, our method did not register this character. Future studies should consider this third dimension at least qualitatively, in order to avoid conflating disparate tooth morphotypes such as these.

Principal component scores as shape measurements are very much dependent on the nature of the sample. Although they are very revealing concerning shape variability, they are not transferable between different data sets. A potential method for creating transferable shape metrics is using our PC1 axis as a guide to derive linear distance measures that would account for the serial shape homology seen in Crocodyliformes. Since PC1 essentially represents molariformy vs. caniniformy from the labial perspective, it could possibly be simplified into a comparison of linear distance measures such as maximum mesial-distal widths and apical-basal heights. These metrics would be not only easy to collect, but also transferable between data sets.

These limitations aside, future researchers may apply our proposed method to any crocodyliform specimen with an intact enough tooth row. We limit our evaluation of interspecific differences, and make no attempt to analyze other factors such as ontogenetic changes or the effect of captive rearing (Erickson, Lappin, & Vliet, 2003; Erickson et al., 2004; Gignac & Erickson, 2014; Drumheller, Wilberg, & Sadleir, 2016). All of these variables may be investigated in the future using our method, as there is nothing to suggest that crocodyliform individuals of most species, ages, and/or rearing conditions would not be able to be quantified in a similar manner. This method could be very useful in dealing with incomplete fossils. It is common for fossil crocodylian specimens to be lacking many, or even most, of their teeth. The linear nature of tooth shape can predict the shape of these missing teeth. A record of the ranges of slopes may be accumulated for fossil specimens with intact teeth. These slopes may then be used as a reference, and be applied to a fossil with the missing teeth. The preserved teeth can be plugged into the linear equation, and the shapes of missing teeth may be

predicted with a high degree of certainty. This would result in a more complete representation of the extinct animal's anatomy, useful from the standpoint of both anatomical science and paleontological reconstruction.

Quantifying the teeth of Crocodyliformes will add rigor to future life history studies of the clade. First and foremost, values may be applied to the plethora of qualitative terms used by researchers (see **Introduction**). This would allow for stricter definitions of the terms when used in the future. As a quantifiable trait, both tooth shape in a single position and heterodonty as a whole may be incorporated into character matrices for phylogenetic analyses. Quantitative descriptors of dentition can describe a numerical range of morphology as opposed to cherry-picking an average tooth or single position. The teeth of fossil taxa can be compared statistically to modern taxa to determine the best analogue, and rigorous hypotheses about paleobehavior and paleoecology may be drawn. Crocodyliformes, both living and extinct, may be grouped into dental categories, allowing for species and specimens to be compared to one another (similar to snouts in Brochu, 2001). Frequency, size, and hardness of food items may be compared to these categories to determine if a link exists between dental morphotypes and dietary patterns (similar to Aoki, 1989). Crocodylians are used in both performance and actualistic taphonomy studies frequently (Njau & Blumenschine, 2006; Erickson et al., 2012, 2014; Drumheller & Brochu, 2014; 2016), and the output of these studies could be correlated with tooth dimensions. Tooth shape may also be compared to bite-force, death-rolling, bone-modification.

CONCLUSION

Multiple measures of morphology have allowed us to describe heterodonty in a thorough manner across a number of both extinct and extant crocodyliform specimens. Through a combination of Foote's morphological disparity and regression analysis along the tooth row, our data indicated that crocodylians are indeed heterodont non-mammals with a number of dental morphotypes available spanning from extreme cases of caniniform to molariform. This variability may be functional in nature, and relate to the size, frequency, and compliance of certain prey in their typically generalist diets. The methods used here should be applied in the future to most crocodylian specimens to investigate dental morphology in the context of a number of natural history related questions.

Acknowledgements

The American Museum of Natural History Department of Herpetology, Royal Ontario Museum Vertebrate Paleontology, and University of California Museum of Paleontology curatorial staff allowed on-site access to dry skull specimens. We would like to particularly thank D. Evans, P. Holroyd, D. Kizirian, C. Raxworthy, K. Seymour for their assistance.

References:

- Adams TL, Noto CR, Drumheller S. 2017. A large neosuchian crocodyliform from the Upper Cretaceous (Cenomanian) woodbine formation of North Texas. *Journal of Vertebrate Paleontology* 37(4):e1349776.
- Aoki R. 1989. The jaw mechanics in the heterodont crocodilians. *Current Herpetology in East Asia* 1:17-21.
- Bennett GE. 2012. Community structure and paleoecology of crocodyliforms from the upper Hell Creek Formation (Maastrichtian), eastern Montana, based on shed teeth. *Jeffersoniana* 28:1-15.
- Berkovitz BK, Shellis RP. 2017. *The Teeth of Non-mammalian Vertebrates*. Cambridge, MA: Academic Press.
- Bookstein FL. 1997. *Morphometric tools for landmark data: geometry and biology*. Cambridge: Cambridge University Press.
- Borteiro C, Gutiérrez F, Tedros M, Kolenc F. 2009. Food habits of the Broad-snouted Caiman (*Caiman latirostris*: Crocodylia, Alligatoridae) in northwestern Uruguay. *Studies on Neotropical Fauna and Environment* 44(1):31-36.
- Brazaitis P. 1973. *The Identification of Living Crocodilians*. New York: New York Zoological Society.

- 745 Britt EJ, Clark AJ, Bennett AF. 2009. Dental morphologies in gartersnakes (*Thamnophis*) and
746 their connection to dietary preferences. *Journal of Herpetology* 43(2):252-259.
- 747 Brochu CA. 1999. Phylogenetics, taxonomy, and historical biogeography of Alligatoroidea.
748 *Journal of Vertebrate Paleontology* 19(S2):9-100.
- 749 Brochu CA. 2001. Crocodylian snouts in space and time: phylogenetic approaches toward
750 adaptive radiation. *American Zoologist* 41(3):564-585.
- 751 Brochu CA. 2004. Alligatorine phylogeny and the status of *Allognathosuchus* Mook, 1921.
752 *Journal of Vertebrate Paleontology* 24(4):857-873.
- 753 Brochu CA, Storrs W. GW. 2012. A giant crocodile from the Plio-Pleistocene of Kenya, the
754 phylogenetic relationships of Neogene African crocodylines, and the antiquity of
755 *Crocodylus* in Africa. *Journal of Vertebrate Paleontology* 32(3): 587-602.
- 756 Brown CM, VanBuren CS, Larson DW, Brink KS, Campione NE, Vavrek MJ, Evans DC. 2015. Tooth
757 counts through growth in diapsid reptiles: implications for interpreting individual and
758 size-related variation in the fossil record. *Journal of Anatomy*, 226(4):322-333.
- 759 Buckley LG, Currie PJ. 2014. Analysis of intraspecific and ontogenetic variation in dentition of
760 *Coelophysis bauri* (Late Triassic), and implications for the systematics of isolated
761 theropod teeth. *New Mexico Museum of Natural History and Science Bulletin* 63.
- 762 Busbey AB. 1995. The structural consequences of skull flattening in crocodilians. *Functional*
763 *Morphology in Vertebrate Paleontology* 173-192.

- 764 Busbey AB. 1989. Form and function of the feeding apparatus of *Alligator mississippiensis*.
765 *Journal of Morphology* 202(1):99-127.
- 766 Carpenter K, Lindsey D. 1980. The dentary of *Brachychampsa montana* Gilmore (Alligatorinae;
767 Crocodylidae), a Late Cretaceous turtle-eating alligator. *Journal of Paleontology*
768 54(6):1213-1217.
- 769 Case EC. 1925. Note on a new species of the Eocene crocodilian *Allognathosuchus*, *A. wartheni*.
770 Ann Arbor, MI: University of Michigan.
- 771 Cheng-kuan C. 1957. Observations on the life history of Chinese alligator (*Alligator sinensis*
772 Fauvel). *Acta Zoologica Sinica* 2(004).
- 773 Cleuren J, Aerts P, Vree FD. 1995. Bite and joint force analysis in *Caiman crocodilus*. Belgian
774 *Journal of Zoology* 125(1):79-94.
- 775 Cleuren J, de Vree F. 2000. Feeding in crocodilians. In Schwenk K. ed. *Feeding: Form, Function,*
776 *and Evolution in Tetrapod Vertebrates*, Cambridge, MA: Elsiever, 359-394.
- 777 Da Silveira R, Magnusson WE. 1999. Diets of spectacled and black caiman in the Anavilhanas
778 Archipelago, Central Amazonia, Brazil. *Journal of Herpetology* 33(2):181-192.
- 779 D'Amore DC. 2009. A functional explanation for denticulation in theropod dinosaur teeth. *The*
780 *Anatomical Record* 292(9):1297-1314.

781 D'Amore DC. 2015. Illustrating ontogenetic change in the dentition of the Nile monitor lizard,
782 *Varanus niloticus*: a case study in the application of geometric morphometric methods
783 for the quantification of shape–size heterodonty. *Journal of Anatomy* 226(5):403-419.

784 Daniel WJT, McHenry C. 2001. Bite force to skull stress correlation-modelling the skull of
785 *Alligator mississippiensis*. In Grigg GC, Seebacher F, Franklin C. (Eds.), *Crocodilian Biology*
786 *And Evolution*, New South Wales: Surrey Beatty and Sons, Chipping Norton, 135-143.

787 Davenport J, Grove DJ, Cannon J, Ellis TR, Stables R. 1990. Food capture, appetite, digestion rate
788 and efficiency in hatchling and juvenile *Crocodylus porosus*. *Journal of Zoology*
789 220(4):569-592.

790 Delany MF. 1990. Late summer diet of juvenile American alligators. *Journal of Herpetology*
791 24(4):418-421.

792 Delany MF, Abercrombie CL. 1986. American alligator food habits in northcentral Florida. *The*
793 *Journal of Wildlife Management* 50(2):348-353.

794 Dodson P. 1975. Functional and ecological significance of relative growth in *Alligator*. *Journal of*
795 *Zoology* 175(3):315-355.

796 Doody S. JS. 2009. Eyes bigger than stomach: prey caching and retrieval in the saltwater
797 crocodile, *Crocodylus porosus*. *Herpetological Review* 40(1):26.

798 Drumheller SK, Brochu CA. 2014. A diagnosis of *Alligator mississippiensis* bite marks with
799 comparisons to existing crocodylian datasets. *Ichnos* 21(2):131-146.

- 800 Drumheller SK, Brochu CA. 2016. Phylogenetic taphonomy: a statistical and phylogenetic
801 approach for exploring taphonomic patterns in the fossil record using crocodylians.
802 *Palaaios* 31(10):463-478.
- 803 Drumheller SK, Wilberg EW, Sadleir RW. 2016. The utility of captive animals in actualistic
804 research: a geometric morphometric exploration of the tooth row of *Alligator*
805 *mississippiensis* suggesting ecophenotypic influences and functional constraints. *Journal*
806 *of Morphology* 277(7):866-78.
- 807 Edmund AG. 1962. Sequence and rate of tooth replacement in the Crocodilia. *Royal Ontario*
808 *Museum, Life Science Division Contributions* 56: 1-42.
- 809 Edmund AG. 1969. Dentition. In: Gans C. d'A. Bellairs A, Parsons TS, eds. *Biology of the Reptilia*,
810 *Volume 1: Morphology A*. Cambridge, MA: Academic Press, 117-200.
- 811 Elsey RM, Trosclair III PL, Linscombe JT. 2004. The American alligator as a predator of mottled
812 ducks. *Southeastern Naturalist* 3(3), 381-390.
- 813 Erickson GM, Lappin AK, Vliet KA. 2003. The ontogeny of bite-force scaling among Crocodylia.
814 *Journal of Zoology* 260(3):317-327.
- 815 Erickson GM, Lappin AK, Parker T, Vliet KA. 2004. Comparison of bite-force performance
816 between long-term captive and wild American alligators (*Alligator mississippiensis*).
817 *Journal of Zoology* 262(1):21-28.

818 Erickson GM, Gignac PM, Steppan SJ, Lappin AK, Vliet KA, Brueggen JD, Inouye BD, Kledzik D,
819 Webb GJ. 2012. Insights into the ecology and evolutionary success of crocodilians
820 revealed through bite-force and tooth-pressure experimentation. *PLoS One* 7(3):e31781.

821 Erickson GM, Gignac PM, Lappin AK, Vliet KA, Brueggen JD, Webb GJW. 2014. A comparative
822 analysis of bite-force scaling among Crocodylia. *Journal of Zoology* 292(1):48-55.

823 Farlow JO, Hurlburt GR, Elsey RM, Britton AR, Langston Jr2005. Femoral dimensions and body
824 size of *Alligator mississippiensis*: estimating the size of extinct mesoeucrocodylians.
825 *Journal of Vertebrate Paleontology* 25(2):354-69.

826 Farrugia P, Polly PD, Njau JK. 2016. An analysis of shape differences in crocodylian dentition
827 using geometric morphometrics. *Journal of Vertebrate Paleontology*, Program and
828 Abstracts 133.

829 Ferguson MWJ. 1981. The structure and development of the palate in *Alligator mississippiensis*.
830 *Archives of Oral Biology* 26(5):427-443.

831 Fish FE, Bostic SA, Nicastro AJ, Beneski JT. 2007. Death roll of the alligator: mechanics of twist
832 feeding in water. *Journal of Experimental Biology* 210(16):2811-2818.

833 Foote M. 1993 Contributions of individual taxa to overall morphological disparity. *Paleobiology*
834 19(4):403-19.

835 Frey E, Monninger S. 2010. Lost in action—the isolated crocodilian teeth from Enspel and their
836 interpretive value. *Palaeobiodiversity and Palaeoenvironments* 90(1):65-81.

- Fruchard C. 2012. The Nile crocodile, a new model for investigating heterodonty and dental continuous renewal in vertebrates. *BioSciences Master Reviews*
- Multivariate and cladistic analyses of isolated teeth reveal sympatry of theropod dinosaurs in the Late Jurassic of northern Germany. Frazzetta TH. 1988. The mechanics of cutting and the form of shark teeth (Chondrichthyes, Elasmobranchii). *Zoomorphology* 108(2):93-107.
- Fukuda Y, Saalfeld K, Lindner G, Nichols T. 2013. Estimation of total length from head length of saltwater crocodiles (*Crocodylus porosus*) in the Northern Territory, Australia. *Journal of Herpetology* 47(1):34-40.
- Gabrey SW. 2010. Demographic and geographic variation in food habits of American alligators (*Alligator mississippiensis*) in Louisiana. *Herpetological Conservation and Biology* 5(2):241-250.
- Galdikas BM, Yeager CP. 1984. Brief report: Crocodile predation on a crab-eating macaque in Borneo. *American Journal of Primatology* 6(1):49-51.
- Gerke O, Wings O. 2016. *PloS One* 11(7):e0158334.
- Gignac PM, Erickson GM. 2014. Ontogenetic changes in dental form and tooth pressures facilitate developmental niche shifts in American alligators. *Journal of Zoology* 295(2):132-142.

- 855 Gignac PM, Erickson GM. 2016. Ontogenetic bite-force modeling of *Alligator mississippiensis*:
856 implications for dietary transitions in a large-bodied vertebrate and the evolution of
857 crocodylian feeding. *Journal of Zoology* 299(4):229-38.
- 858 Gignac P, O'Brien H. 2016. Suchian feeding success at the interface of ontogeny and
859 macroevolution. *Integrative and Comparative Biology* 56(3):449-458.
- 860 Grigg G, Gans C. 1993. Morphology and physiology of the Crocodylia. *Fauna of Australia Volume*
861 *2A, Amphibia and Reptilia*. Canberra: Australian Government Publishing Service, 326-
862 336.
- 863 Groombridge B. 1982. *The IUCN Amphibia-Reptilia Red Data Book, Part 1: Testudines,*
864 *Crocodylia, Rhynchocephalia*. Gland, Switzerland: IUCN.
- 865 Gunz P, Mitteroecker P. 2013. Semilandmarks: a method for quantifying curves and surfaces.
866 *Hystrix, the Italian Journal of Mammalogy* 24(1):103-109.
- 867 Hammer Ø, Harper DAT, Ryan PD. 2001. PAST: Paleontological statistics software package for
868 education and data analysis. *Palaeontologia Electronica* 4(1):9pp.
- 869 Hanson JO, Salisbury SW, Campbell HA, Dwyer RG, Jardine TD, Franklin CE. 2014. Feeding across
870 the food web: The interaction between diet, movement and body size in estuarine
871 crocodiles (*Crocodylus porosus*). *Austral Ecology* 40(3):275-286.
- 872 Hendrickx C, Mateus O. 2014. Abelisauridae (Dinosauria: Theropoda) from the Late Jurassic of
873 Portugal and dentition-based phylogeny as a contribution for the identification of
874 isolated theropod teeth. *Zootaxa* 3759(1):1-74.

- 875 Hendrickx C, Mateus O, Araújo R. 2015a The dentition of megalosaurid theropods. *Acta*
876 *Palaeontologica Polonica*. A proposed terminology of theropod teeth (Dinosauria,
877 Saurischia). *Journal of Vertebrate Paleontology* 35(5):e982797.
- 878 Hendrickx C, Mateus O, Araújo R. 2015b. 60(3):627-642.
- 879 Hutton JM. 1987. Growth and feeding ecology of the Nile crocodile *Crocodylus niloticus* at
880 Ngezi, Zimbabwe. *The Journal of Animal Ecology* 56(1):25-38.
- 881 Iordansky NN. 1964. The jaw muscles of the crocodiles and some relating structures of the
882 crocodilian skull. *Anatomischer Anzeiger* 115:256-280.
- 883 Kar SK, Bustard HR. 1983. Attacks on domestic livestock by juvenile saltwater crocodile,
884 *Crocodylus porosus*, in Bhitarkanika Wildlife Sanctuary, Orissa India. *Amphibia-Reptilia*
885 4(1):81-83.
- 886 Kear BP, Larsson D, Lindgren J, Kundrát M. 2017. Exceptionally prolonged tooth formation in
887 elasmosaurid plesiosaurs. *PloS One* 12(2):e0172759.
- 888 Kieser JA, Klapálský C, Law L, Marion M. 1993. Heterodonty and patterns of tooth replacement
889 in *Crocodylus niloticus*. *Journal of Morphology* 218(2): 195-201.
- 890 Klingenberg CP. 2011. MorphoJ: an integrated software package for geometric morphometrics.
891 *Molecular Ecology Resources* 11(2):353-357.
- 892 Langston W. 1973. The crocodilian skull in historical perspective. *Biology of the Reptilia* 4:263-
893 284.

- 894 Larson DW, Brown CM, Evans DC. 2016. Dental disparity and ecological stability in bird-like
895 dinosaurs prior to the end-Cretaceous mass extinction. *Current Biology* 26(10):1325-
896 1333.
- 897 Larson DW, Currie PJ. 2013. Multivariate analyses of small theropod dinosaur teeth and
898 implications for paleoecological turnover through time. *PLoS One* 8(1):e54329.
- 899 Larsson HCE, Sidor CA. 1999. Unusual crocodyliform teeth from the Late Cretaceous
900 (Cenomanian) of southeastern Morocco. *Journal of Vertebrate Paleontology* 19(2):398-
901 401.
- 902 Larsson HC, Sues HD. 2007. Cranial osteology and phylogenetic relationships of *Hamadasuchus*
903 *rebouli* (Crocodyliformes: Mesoeucrocodylia) from the Cretaceous of Morocco.
904 *Zoological Journal of the Linnean Society* 149(4):533-67.
- 905 LeBlanc AR, Brink KS, Cullen TM, Reisz RR. 2017. Evolutionary implications of tooth attachment
906 versus tooth implantation: a case study using dinosaur, crocodilian, and mammal teeth.
907 *Journal of Vertebrate Paleontology* 37(5):e1354006.
- 908 Lecuona A, Pol D. 2008. Tooth morphology of *Notosuchus terrestris* (Notosuchia:
909 Mesoeucrocodylia): new evidence and implications. *Comptes Rendus Palevol* 7(7):407-
910 417.
- 911 Luiselli L, Akani GC, Capizzi D. 1999. Is there any interspecific competition between dwarf
912 crocodiles (*Osteolaemus tetraspis*) and Nile monitors (*Varanus niloticus ornatus*) in the

- 913 swamps of central Africa? A study from south-eastern Nigeria. *Journal of Zoology*
- 914 247(1):127-131.
- 915 MacLeod CD, Reidenberg JS, Weller M, Santos MB, Herman J, Goold J, Pierce GJ. 2007. Breaking
- 916 symmetry: The marine environment, prey size, and the evolution of asymmetry in
- 917 cetacean skulls. *The Anatomical Record* 290(6):539-545.
- 918 MacLeod CD, Santos MB, Lopez A, Pierce GJ. 2006. Relative prey size consumption in toothed
- 919 whales: implications for prey selection and level of specialisation. *Marine Ecology*
- 920 *Progress Series* 326:295-307.
- 921 Martin JE. 2007. New material of the Late Cretaceous globidontan *Acynodon iberoccitanus*
- 922 (Crocodylia) from southern France. *Journal of Vertebrate Paleontology* 27(2):362-372.
- 923 McHenry CR, Clausen PD, Daniel WJ, Meers MB, Pendharkar A. 2006. Biomechanics of the
- 924 rostrum in crocodylians: a comparative analysis using finite-element modeling. *The*
- 925 *Anatomical Record* 288(8):827-849.
- 926 McIlhenny EA. 1976. The Alligator's Life History. Society for the Study of Amphibians and
- 927 Reptiles.
- 928 Mitteroecker P, Gunz P, Windhager S, Schaefer K. 2013. A brief review of shape, form, and
- 929 allometry in geometric morphometrics, with applications to human facial morphology.
- 930 *Hystrix, the Italian Journal of Mammalogy* 24(1):59-66.
- 931 Monfroy QT. 2017. Correlation between the size, shape and position of the teeth on the jaws
- 932 and the bite force in Theropoda. *Historical Biology* 29(8):1089-1105.

- 933 Mook CC. 1932. A study of the osteology of *Alligator prenasalis* (Loomis). *Bulletin of the*
934 *Museum of Comparative Zoology* 74(2).
- 935 Narváez I, Brochu CA, Escaso F, Pérez-García A, Ortega F. 2015. New crocodyliforms from
936 southwestern Europe and definition of a diverse clade of European Late Cretaceous
937 basal eusuchians. *PloS One* 10(11):e0140679.
- 938 Nifong JC, Silliman BR. 2013. Impacts of a large-bodied, apex predator (*Alligator mississippiensis*
939 Daudin 1801) on salt marsh food webs. *Journal of Experimental Marine Biology and*
940 *Ecology* 440:185-191.
- 941 Njau JK, Blumenschine RJ. 2006. A diagnosis of crocodile feeding traces on larger mammal
942 bone, with fossil examples from the Plio-Pleistocene Olduvai Basin, Tanzania. *Journal of*
943 *Human Evolution* 50(2):142-162.
- 944 Norell M, Clark JM, Hutchison JH. 1994. The Late Cretaceous alligatoroid *Brachychampsa*
945 *montana* (Crocodylia): new material and putative relationships. *American Museum*
946 *Novitates* 3116.
- 947 Novas FE, Pais DF, Pol D, Carvalho IDS, Scanferla A, Mones A, Riglos, MS. 2009. Bizarre
948 notosuchian crocodyliform with associated eggs from the Upper Cretaceous of Bolivia.
949 *Journal of Vertebrate Paleontology* 29(4):1316-1320.
- 950 Osborn JW. 1998. Relationship between growth and the pattern of tooth initiation in alligator
951 embryos. *Journal of Dental Research* 77(9):1730-1738.

- 952 Ósi A, Barrett PM. 2011. Dental wear and oral food processing in *Caiman latirostris*: analogue
953 for fossil crocodylians with crushing teeth. *Neues Jahrbuch für Geologie und*
954 *Paläontologie-Abhandlungen* 261(2):201-207.

- 955 Ósi A, Clark JM, Weishampel DB. 2007. First report on a new basal eusuchian crocodyliform
956 with multicusped teeth from the Upper Cretaceous (Santonian) of Hungary. *Neues*
957 *Jahrbuch für Geologie und Paläontologie-Abhandlungen* 243(2):169-177.

- 958 Pauwels OS, Barr B, Sanchez ML, Burger M. 2007. Diet records for the dwarf crocodile
959 (*Osteolaemus tetraspis tetraspis*) in Rabi Oil Fields and Loango National Park,
960 Southwestern Gabon. *Hamadryad* 31(2):258-264.

- 961 Perez SI, Bernal V, Gonzalez PN. 2006. Differences between sliding semi-landmark methods in
962 geometric morphometrics, with an application to human craniofacial and dental
963 variation. *Journal of Anatomy* 208(6):769-784.

- 964 Peyer B. 1968. *Comparative odontology*. Chicago, IL: University of Chicago Press.

- 965 Pierce SE, Angielczyk KD, Rayfield EJ. 2008. Patterns of morphospace occupation and
966 mechanical performance in extant crocodilian skulls: a combined geometric
967 morphometric and finite element modeling approach. *Journal of Morphology*
968 269(7):840-864.

- 969 Pooley AC. 1989. Food and feeding habits. In Ross CA, and Garnet S. eds. *Crocodiles and*
970 *Alligators* New York, NY: Facts on File, 76-91.

- 971 Pooley AC, Gans C. 1976. The Nile crocodile. *Scientific American* 234(4):114-125.

- 972 Rohlf FJ. 2010. Tps Series. Department of Ecology and Evolution, State University of New York,
973 Stony Brook.
- 974 Rohlf FJ. 2013. TpsRelw 1.53. Dept. of Ecology and Evolution, State Univ. of New York at Stony
975 Brook, Stony Brook.
- 976 Rootes WL, Chabreck RH. 1993. Cannibalism in the American alligator. *Herpetologica* 49(1):99-
977 107.
- 978 Rommel S. 1990. Osteology of the bottlenose dolphin. In Leatherwood S, Reeves RR. Eds, *The*
979 *Bottlenose Dolphin*, Cambridge, MA: Elsevier, 29-49.
- 980 Ross CA, Magnusson WE. 1989. Living crocodilians. In Ross CA, and Garneteds. *Crocodiles and*
981 *Alligators* New York, NY: Facts on File, 58-73.
- 982 Sadleir RW, Makovicky PJ. 2008. Cranial shape and correlated characters in crocodilian
983 evolution. *Journal of Evolutionary Biology* 21(6):1578-1596.
- 984 Salas-Gismondi R, Flynn JJ, Baby P, Tejada-Lara JV, Wesselingh FP, Antoine, PO. 2015. A
985 Miocene hyperdiverse crocodylian community reveals peculiar trophic dynamics in
986 proto-Amazonian mega-wetlands. *Proceedings of the Royal Society of London B:*
987 *Biological Sciences* 282(1804):20142490.
- 988 Salas-Gismondi R, Flynn JJ. Baby P, Tejada-Lara JV, Claude J, Antoine PO. 2016. A new 13 million
989 year old gavialoid crocodylian from proto-Amazonian mega-wetlands reveals parallel
990 evolutionary trends in skull shape linked to longirostry. *PloS One* 11(4):e0152453.

- 991 Sampaio PRM, da Silva MN, de Matos SA, de Matos LRA, Acosta M. 2013. First report of
992 predation by a caiman (*Paleosuchus trigonatus*, Crocodylia: Alligatoridae) on a caecilian
993 (*Caecilia marcusii*, Gymnophiona: Caeciliidae). *Salamandra* 49(4):227-228.
- 994 Santos SA, Nogueira MS, Pinheiro MS, Campos Z, Magnusson WE, Mourao GM. 1996. Diets of
995 *Caiman crocodilus yacare* from different habitats in the Brazilian pantanal.
996 *Herpetological Journal* 6:111-117.
- 997 Sassoon J, Foffa D, Marek R. 2015. Dental ontogeny and replacement in Pliosauridae. *Royal*
998 *Society Open Science* 2(11):150384.
- 999 Selvaraj G. 2012. Herpetological notes: *Tomistoma schlegelii* (False Gharial). Diet.
1000 *Herpetological Review*, 43:608-609.
- 1001 Schwarz-Wings D, Rees J, Lindgren J. 2009. Lower cretaceous mesoeucrocodylians from
1002 Scandinavia (Denmark and Sweden). *Cretaceous Research* 30(5):1345-1355.
- 1003 Sheets HD. 2012. IMP software series. Computer software and manual. Buffalo, NY: Canisius
1004 College.
- 1005 Sheets HD, Kim K, Mitchell CE. 2004. A combined landmark and outline-based approach to
1006 ontogenetic shape change in the Ordovician trilobite *Triarthrus becki*. In *Morphometrics*
1007 Berlin: Springer, 67-82.
- 1008 Sheets HD, Zelditch ML. 2013. Studying ontogenetic trajectories using resampling methods and
1009 landmark data. *Hystrix, the Italian Journal of Mammalogy* 24(1):67-73.

- 1010 Shimada, K. 2001. On the concept of heterodonty. *Journal of Fossil Research* 34(2):52-54.
- 1011 Shimada K. 2002. Dental homologies in lamniform sharks (Chondrichthyes: Elasmobranchii).
- 1012 *Journal of Morphology* 251(1):38-72.
- 1013 Shimada K. 2004. The relationship between the tooth size and total body length in the sandtiger
- 1014 shark, *Carcharias taurus* (Lamniformes: Odontaspidae). *Journal of Fossil Research*
- 1015 37:76–81.
- 1016 Shimada K, Seigel JA. 2005. The relationship between the tooth size and total body length in the
- 1017 goblin shark, *Mitsukurina owstoni* (Lamniformes: Mitsukurinidae). *Journal of Fossil*
- 1018 *Research* 38:49–56.
- 1019 Shoop CR, Ruckdeschel CA. 1990. Alligators as predators on terrestrial mammals. *American*
- 1020 *Midland Naturalist* 124(2):407-412.
- 1021 Smith JB, Dodson P. 2003. A proposal for a standard terminology of anatomical notation and
- 1022 orientation in fossil vertebrate dentitions. *Journal of Vertebrate Paleontology* 23(1):1-
- 1023 12.
- 1024 Smith JB, Vann DR, Dodson P. 2005. Dental morphology and variation in theropod dinosaurs:
- 1025 implications for the taxonomic identification of isolated teeth. *The Anatomical Record*
- 1026 285(2):699-736.
- 1027 Subalusky AL, Fitzgerald LA, Smith LL. 2009. Ontogenetic niche shifts in the American Alligator
- 1028 establish functional connectivity between aquatic systems. *Biological Conservation*
- 1029 142(7):1507-1514.

- 1030 Tavares SAS, Branco FR, de Souza Carvalho I, Maldanis L. 2017. The morphofunctional design of
1031 *Montealtosuchus arrudacamposi* (Crocodyliformes, Upper Cretaceous) of the Bauru
1032 Basin, Brazil. *Cretaceous Research* 79:64-76.
- 1033 Taylor JA. 1979. The foods and feeding habits of subadult *Crocodylus porosus* Schneider in
1034 northern Australia. *Wildlife Research* 6(3):347-359.
- 1035 Thorbjarnarson JB. 1990. Notes on the feeding behavior of the gharial (*Gavialis gangeticus*)
1036 under semi-natural conditions. *Journal of Herpetology* 24(1):99-100.
- 1037 Tinius A, Russell AP. 2017. Points on the curve: An analysis of methods for assessing the shape
1038 of vertebrate claws. *Journal of Morphology* 278(2):150-69.
- 1039 Torices A, Reichel M, Currie PJ. 2014. Multivariate analysis of isolated tyrannosaurid teeth from
1040 the Danek Bonebed, Horseshoe Canyon Formation, Alberta, Canada. *Canadian Journal*
1041 *of Earth Sciences* 51(11):1045-1051.
- 1042 Valkenburgh BV, Ruff CB. 1987. Canine tooth strength and killing behaviour in large carnivores.
1043 *Journal of Zoology* 212(3):379-397.
- 1044 Verdade LM. 2000. Regression equations between body and head measurements in the broad-
1045 snouted caiman (*Caiman latirostris*). *Revista Brasileira de Biologia* 60(3):469-482.
- 1046 Vullo R, Allain R, Cavin L. 2016. Convergent evolution of jaws between spinosaurid dinosaurs
1047 and pike conger eels. *Acta Palaeontologica Polonica* 61(4):825-828.

- 1048 Wallace KM, Leslie AJ. 2008. Diet of the Nile crocodile (*Crocodylus niloticus*) in the Okavango
1049 Delta, Botswana. *Journal of Herpetology* 42(2):361-368.
- 1050 Walmsley CW, Smits PD, Quayle MR, McCurry MR, Richards HS, Oldfield CC, ... & McHenry CR.
1051 2013. Why the long face? The mechanics of mandibular symphysis proportions in
1052 crocodiles. *PLoS One* 8(1):e53873.
- 1053 Watanabe A, Slice DE. 2014. The utility of cranial ontogeny for phylogenetic inference: a case
1054 study in crocodylians using geometric morphometrics. *Journal of Evolutionary Biology*
1055 27(6):1078-1092.
- 1056 Webb GJW, Manolis SC, Buckworth R. 1982. *Crocodylus johnstoni* in the McKinlay river area, NTI
1057 Variation in the diet, and a new method of assessing the relative importance of prey.
1058 *Australian Journal of Zoology* 30(6):877-899.
- 1059 Webb GJW, Messel H. 1978. Movement and dispersal patterns of *Crocodylus porosus* in some
1060 rivers of Arnhem Land, Northern Australia. *Wildlife Research* 5(2):263-283.
- 1061 Westergaard B, Ferguson MWJ. 1986. Development of the dentition in *Alligator*
1062 *mississippiensis*. Early embryonic development in the lower jaw. *Journal of Zoology*
1063 210(4):575-597.
- 1064 Westergaard B, Ferguson MWJ. 1987. Development of the dentition in *Alligator*
1065 *mississippiensis*. Later development in the lower jaws of embryos, hatchlings and young
1066 juveniles. *Journal of Zoology* 212(2):191-222.

- 1067 Westergaard B, Ferguson MWJ. 1990. Development of the dentition in *Alligator*
1068 *mississippiensis*: upper jaw dental and craniofacial development in embryos, hatchlings,
1069 and young juveniles, with a comparison to lower jaw development. *American Journal of*
1070 *Anatomy* 187(4):393-421.
- 1071 Wilberg EW. 2017. Investigating patterns of crocodyliform cranial disparity through the
1072 Mesozoic and Cenozoic. *Zoological Journal of the Linnean Society* 181(1):189-208.
- 1073 Wu XB, Xue H, Wu LS, Zhu JL, Wang RP. 2006. Regression analysis between body and head
1074 measurements of Chinese alligators (*Alligator sinensis*) in the captive population. *Animal*
1075 *Biodiversity and Conservation* 29(1):65-71.
- 1076 Young MT, Brusatte SL, Beatty BL, De Andrade MB, Desojo JB. 2012. Tooth-on-tooth
1077 interlocking occlusion suggests macrophagy in the Mesozoic marine crocodylomorph
1078 *Dakosaurus*. *The Anatomical Record* 295(7):1147-1158.
- 1079 Zahradnick O, Buchtova M, Dosedelova H, Tucker AS. 2014. The development of complex
1080 tooth shape in reptiles. *Frontiers in Physiology* 5:74.
- 1081 Zelditch ML, Sheets HD, Fink WL. 2003. The ontogenetic dynamics of shape disparity.
1082 *Paleobiology* 29(1):139-156.
- 1083 Zelditch M, Swiderski D, Sheets DH, Fink W. 2004. *Geometric Morphometrics for Biologists: A*
1084 *Primer*. Waltham, MA: Elsevier Academic Press.

1085 **FIGURE LEGENDS**

1086 **Figure 1: Data collection methods.** A) We numbered teeth based on position and host bone
 1087 (only the left teeth are labeled). Teeth P4, M5, and M10 are defined as the enlarged teeth. B)
 1088 We photographed each tooth individually, and traced the margins. C) The software converted
 1089 each outline into 30 equidistant coordinates. Three coordinates were transformed into
 1090 landmarks (magenta) and the rest into semilandmarks (green). D) We represented tooth shape
 1091 variance through vector diagrams, with points representing the mean and vectors representing
 1092 shape deviation. E) For skulls, we produced an outline from the dorsal perspective, with 50
 1093 coordinates on each margin that were transformed into landmarks (magenta) and
 1094 semilandmarks (green). F) We represented shape variance with vector diagrams similar to
 1095 above. (Specimen depicted: *Crocodylus palustris* AMNH 96134.) [Half page width]

1096 **Figure 2: Variability within the first Principal Component for teeth.** Vector diagrams indicate
 1097 the maximum range of variance (vectors) from the mean (points) for both cranial and dentary
 1098 teeth. Landmarks are in magenta and semilandmarks are in green. [Half page width]

1099 **Figure 3: Partial Least Squared (PLS) two-block analysis of shape.** A) Vector diagrams indicate
 1100 shape variance of PLS1 for skull shape (Block 1) and average tooth shape (Block 2). B) PLS1
 1101 scores for both Blocks were regressed, with colors representing major taxonomic groups. Solid
 1102 markers depict extant specimens, and hollow markers depict extinct. [Full page width]

1103 **Figure 4: Heterodonty represented by Foote's morphological disparity.** Ln scaling of
 1104 Morphological Disparity (MD) for size (A) and shape (B) are plotted against the Ln of skull length.

1105 Colors represent major taxonomic groups, solid markers depict extant specimens, and hollow
1106 markers depict extinct. [Half page width]

1107 **Figure 5: Heterodonty by tooth position.** Centroid Size (CS) and Principal Component one (PC1)
1108 for extant Alligatoroidea, Crocodyloidea, and remaining taxa, plotted against position along the
1109 arcade. Welch's ANOVA output comparing positions is listed for each graph with multiple
1110 specimens. Colors represent major taxonomic groups. See Figure 2 for a visual representation
1111 of shape change depicted by PC1 scores. [Full page width]

1112 **Figure 6: Regression information for shape heterodonty.** Slope (m) and y-intercept (b) data for
1113 regressions of the first Principal Component plotted against tooth position for individuals. Error
1114 bars indicate 95% confidence intervals. Colors represent major taxonomic groups, with solid
1115 markers depicting extant specimens and hollow markers depicting extinct. Regression statistics
1116 are available in Table S3. [Full page width]

1117 **Figure 7: Direct comparisons between selected extant and extinct taxa.** The size axis
1118 represents normalized centroid size (ranging from 0.00-0.20), and the shape axis represents
1119 scores from the first principal component (ranging from -0.04-0.04). Taxa are grouped by A)
1120 specimens with high molariform distal teeth, B) modern taxa that display high size heterodonty,
1121 C) slender-snouted taxa, D) *Hamadasuchus rebouli*, E) *Borealosuchus sternbergii* and
1122 *Leidyosuchus canadensis*, and F) fossil globidont taxa. (Note: *B. sternbergii* teeth were not size
1123 normalized by its own skull length, as indicated by hollow bars.) Scale = 5 cm. [Full page width]

Figure 1

Data collection methods

A) We numbered teeth based on position and host bone (only the left teeth are labeled). Teeth P4, M5, and M10 are defined as the enlarged teeth. B) We photographed each tooth individually, and traced the margins. C) The software converted each outline into 30 equidistant coordinates. Three coordinates were transformed into landmarks (magenta) and the rest into semilandmarks (green). D) We represented tooth shape variance through vector diagrams, with points representing the mean and vectors representing shape deviation. E) For skulls, we produced an outline from the dorsal perspective, with 50 coordinates on each margin that were transformed into landmarks (magenta) and semilandmarks (green). F) We represented shape variance with vector diagrams similar to above. (Specimen depicted: *Crocodylus palustris* AMNH 96134.)




Figure 2

Variability within the first Principal Component for teeth

Vector diagrams indicate the maximum range of variance (vectors) from the mean (points) for both cranial and dentary teeth. Landmarks are in magenta and semilandmarks are in green.




Figure 3

Partial Least Squared (PLS) two-block analysis of shape

A) Vector diagrams indicate shape variance of PLS1 for skull shape (Block 1) and average tooth shape (Block 2). B) PLS1 scores for both Blocks were regressed, with colors representing major taxonomic groups. Solid markers depict extant specimens, and hollow markers depict extinct.




Figure 4

Heterodonty represented by Foote's morphological disparity

Ln scaling of Morphological Disparity (MD) for size (A) and shape (B) are plotted against the Ln of skull length. Colors represent major taxonomic groups, solid markers depict extant specimens, and hollow markers depict extinct.





Figure 5

Heterodonty by tooth position

Centroid Size (CS) and Principal Component one (PC1) for extant Alligatoroidea, Crocodyloidea, and remaining taxa, plotted against position along the arcade. Welch's ANOVA output comparing positions is listed for each graph with multiple specimens. Colors represent major taxonomic groups. See Figure 2 for a visual representation of shape change depicted by PC1 scores.




Figure 6

Regression information for shape heterodonty

Slope (m) and y-intercept (b) data for regressions of the first Principal Component plotted against tooth position for individuals. Error bars indicate 95% confidence intervals. Colors represent major taxonomic groups, with solid markers depicting extant specimens and hollow markers depicting extinct. Regression statistics are available in Table S3.




Figure 7

Direct comparisons between selected extant and extinct taxa

The size axis represents normalized centroid size (ranging from 0.00-0.20), and the shape axis represents scores from the first principal component (ranging from -0.04-0.04). Taxa are grouped by A) specimens with high molariform distal teeth, B) modern taxa that display high size heterodonty, C) slender-snouted taxa, D) *Hamadasuchus rebouli*, E) *Borealosuchus sternbergii* and *Leidyosuchus canadensis*, and F) fossil globidont taxa. (Note: *B. sternbergii* teeth were not size normalized by its own skull length, as indicated by hollow bars.) Scale = 5 cm.

